U.S. Department of Veterans Affairs

Public Access Author manuscript

Mol Cell Endocrinol. Author manuscript; available in PMC 2017 November 30.

Published in final edited form as:

Mol Cell Endocrinol. 2014 May 25; 389(1-2): 31–39. doi:10.1016/j.mce.2014.01.002.

Estrogen and the Female Heart

A. A. Knowlton^{1,2} and D. H. Korzick³

¹The Department of Veteran's Affairs, Northern California VA, Sacramento, CA, USA

²Molecular & Cellular Cardiology, Departments of Medicine and Pharmacology, University of California, Davis, USA

³Intercollege Program in Physiology and Department of Kinesiology, The Pennsylvania State University, University Park, PA, USA

Abstract

Estrogen has a plethora of effects in the cardiovascular system. Studies of estrogen and the heart span human clinical trials and basic cell and molecular investigations. Greater understanding of cell and molecular responses to estrogens can provide further insights into the findings of clinical studies. Differences in expression and cellular/intracellular distribution of the two main receptors, estrogen receptor (ER) α and β , are thought to account for the specificity and differences in responses to estrogen. Much remains to be learned in this area, but cellular distribution within the cardiovascular system is becoming clearer. Identification of GPER as a third ER has introduced further complexity to the system. 17β -estradiol (E₂), the most potent human estrogen, clearly has protective properties activating a signaling cascade leading to cellular protection and also influencing expression of the protective heat shock proteins (HSP). E₂ protects the heart from ischemic injury in basic studies, but the picture is more involved in the whole organism and clinical studies. Here the complexity of E₂'s widespread effects comes into play and makes interpretation of findings more challenging. Estrogen loss occurs primarily with aging, but few studies have used aged models despite clear evidence of differences between the response to E₂ deficiency in adult and aged animals. Thus more work is needed focusing on the effects of aging vs. estrogen loss on the cardiovascular system.

Keywords

estradiol; 17β -estradiol; ER α ; ER β ; GPER; cardioprotection; HSF1; heat shock proteins; HSP72; HRT; TNF α ; inflammation; heart; cardiovascular; mitochondria; PKC

Introduction

Estrogens are potent steroid hormones present in high levels in women from adolescence to menopause. Estrogens have many properties, both protective and deleterious, and there have been numerous reviews on estrogen and cardiovascular disease.(1–6) The pace and depth of

Address for Correspondence: A.A. Knowlton, M.D., Molecular & Cellular Cardiology, University of California, Davis, One Shields Avenue, Davis, CA 95616, Tel. - 530-752-5461, Fax - 530-754-7167, aaknowlton@ucdavis.edu.

Disclaimer: The contents reported do not represent the views of the Department of Veterans Affairs or the United States Government.

estrogen research has steadily increased since the identification of the estrogen receptors and the development of knockout models, facilitating investigation of the many functions of estrogens.

Estrogen Receptors, Genomic/Nongenomic Effects—Estrogen binds two intracellular receptors, estrogen receptor (ER) α and ER β . These two receptors are primarily found in the cytosol and in the nucleus (figure 1). They are also loosely associated with the plasma membrane, rather than transmembrane like many other receptors. Mitochondrial localization has also been reported. Genomic actions mediated by nuclear ER α are well-described (7;8) and involve ligand binding at E₂ response elements. Non-genomic (rapid) effects of E₂ are thought to be mediated by ER α and/or ER β localized to the plasma membrane (9–11) and associated functions include Ca²⁺ homeostasis, anti-apoptotic effects and mitochondrial metabolism.(10) Rapid ER signals are also known to regulate ER gene transcription in the myocardium.(12) In this regard, ERs are subject to post-translational modification (PTM) through phosphorylation, acetylation and sumoylation, which not only has the potential to influence ER activity, but may also influence ER stability and localization, particularly with aging (for review see 13–20).

A third receptor, G-protein-coupled estrogen receptor (GPER) or GPR30, has been more recently described. GPER is a classic membrane receptor with 7 transmembrane spanning domains.(21) This receptor is discussed below.

It is thought that at least part of the specificity of E₂ signaling is the result of differential expression and cellular localization of the ERs in different tissues and cell types. As ER α and ER β have been studied the longest, much more is known about their localization than that of GPER. Deconvolution of rat endothelium (cerebral and coronary arteries) images demonstrated that about a third of ER α co-localized with caveolin-1 at the plasma membrane.(22) In this study, ER β was predominantly nuclear. In studies of the ovine uterine artery endothelial cells, ER α was present in the nucleus and associated with caveolin-1 at the plasma membrane.(23) In contrast, ER β mRNA levels were much less than those of ER α , and ER β did not localize with caveolin-1. Thus in adult endothelial cells, ER α is nuclear and membrane associated, co-localizing with caveolin.

Considerably less is known about the ERs in cardiac myocytes and vascular smooth muscle cells (VSMCs), and few studies have investigated their precise subcellular localization in these specific tissues.(24;25) For cardiac myocytes, ER α has been reported to localize to nuclear, cytosolic, mitochondrial, and plasma membrane fractions, respectively.(25) ER β expression has been reported in human cardiac mitochondria,(10) however controversy exists regarding antibody quality and species dependency.(26) Indeed, we were unable to identify ER β mRNA in hearts of adult or aged Fisher 344 female rats.(26) Much remains to be defined with regards to expression and localization of ER α and β in cardiac myocytes.

In contrast, more is known about VSMCs. In male SD rat mesenteric artery VSMCs ER α (the ER46 fragment) and ER β associated with caveolin-1.(13) Both ER α and ER β are found in adult human aortic VSMCs and both are predominantly nuclear.(13;14) VSMCs from the aortic arch of 8 week old rats expressed ER α and ER β based on RT-PCR.(15) Similarly,

primary adult human VSMCs contained both receptors, but less ER β than ER α was found. (16) VSMC cell lines had clearly different ER α and ER β expression compared primary cells and pathology specimens. Three different VSMCs cell lines expressed either ER α or ER β , but not both, which makes these cell lines less than desirable models for investigation of ER α and ER β in VSMCs. (14;15) Multiple studies have shown that ER α and ER β are both expressed in VSMCs and are primarily in the nucleus, however both ER α and ER β are also at the plasma membrane and co-localize with caveolin-1. The receptors' relative quantities and intracellular allocation remains to be conclusively defined.

ER Membrane Complex, Caveolae and Caveolin—As noted above, ER α and β demonstrate both genomic and nongenomic effects, also referred to as nuclear and non-nuclear events. ER α is found in caveolae, where it complexes with caveolin 1, HSP90, eNOS, c-Src, Akt, and PI3K in the plasma membrane (27–29), and this is regulated by palmitoylation. (30–32) ER46, an ER α splice variant, rather than the full length receptor, is present in the caveolar complex. (27;29;30;33) The co-localization of ER α , eNOS and HSP90 within the caveolae facilitates eNOS activation by E2. (30;32;34) The chaperone protein, HSP90, promotes interaction of ER with signaling enzymes, such as eNOS. In contrast, caveolin-1 inhibits eNOS activity. Caveolin-1 knockout leads to increased eNOS activity in mice. (35) The multiple proteins needed for the caveolar-ER α signaling complex have been reviewed in great detail. (36) As would be expected given caveolin-1's inhibitory effects, eNOS activation entails separation from caveolin-1; this separation is usually mediated by calcium/calmodulin, but in special circumstances, including shear stress, eNOS activation does not require calcium. Several excellent papers have reviewed in details the steps involved in regulation of eNOS activation. (37;38)

GPER—There persists some lingering controversy regarding whether GPER is an ER, and more work is necessary to fully delineate the role of GPER in estrogen signaling. (39) Nonetheless, there is increasing acceptance of GPER as key receptor for estrogen signaling in the cardiovascular system. (40) GPER, which is also known as G-protein coupled receptor 30 (GPR30) and membrane estrogen receptor (mER), is the most recently recognized membrane associated ER. (41) RT-PCR and western blotting were used to detect GPER expression in porcine coronary arteries and human coronary artery VSMCs. (42) In addition, GPER is present in all four cardiac chambers with greater expression in the ventricles than the atria. (43) GPER has not been specifically shown to be in cardiac myocytes, but is present in HL-1 cells, which are an atrial tumor derived cell line with many properties found in cardiac myocytes. (44) There is one report that male GPER deficient mice had impaired cardiac function with decreased +dP/dt and –dP/dt, as well as increased LVEDP. (45) Heart weight/tibia length and lung weight/tibia length did not differ from wild type, nor did systolic pressure. Female hearts were normal. As it has yet to be demonstrated conclusively that GPER is expressed in cardiac myocytes, the authors concluded, after ruling out metabolic abnormalities, that the impaired ventricular function in male mice was not a direct cardiac effect of GPER loss, but might reflect vascular changes secondary to GPER loss.

GPER, which has seven transmembrane domains similar to traditional membrane receptors, can mediate E2 driven vasodilation. (21;46) G1, a selective agonist for GPER, relaxed both

porcine aortic rings and human coronary artery VSMCs in endothelium independent manner. (42) G15, a selective inhibitor of GPER prevents G1 mediated relaxation. (42) Further work showed GPER activation of the large-conductance calcium and voltage activated potassium channel (BK_{ca}) in coronary artery VSMCs by GPER, but, interestingly G1 had no direct effect on BK_{ca} . (42) GPER inhibited endothelin-1 mediated vasoconstriction. (47) These findings provide new insights into GPER mediated vascular relaxation. Several groups have also reported activation of GPER by aldosterone. (48–51) Aldosterone increased cardiac vagal tone via activation of GPER. (51) These findings support nongenomic functions for aldosterone, and raise questions about aldosterone/estrogen interactions in regulating GPER activation. The role of aldosterone in GPER activation has recently been reviewed. (48) Thus GPER, unlike $ER\alpha$ and β which are controlled solely by estrogens, has two or possibly more hormones able to activate it.

Clinical Trials, Hormone Replacement Therapy (HRT) and the Timing

Hypothesis—Estrogen had been long viewed as cardioprotective based on anecdotal reports over many years. However, double blind, controlled trials of estrogen replacement post-menopause conducted in the 1990's, such as the Women's Health Initiative (WHI) and the Heart and Estrogen/progestin Replacement Study (HERS) demonstrated greater risk of cancer and cardiovascular disease (CHD) with estrogen replacement. (52–54) This was a surprising outcome as premenopausal women have a much lower incidence of CHD than males. Careful consideration of the unexpected results of these clinical trials led to the realization that on average the women enrolled were 10 years post-menopause, and thus were receiving late estrogen replacement long after their ovaries had ceased to make estrogen. This suggested that such a long gap before initiating estrogen could result in unexpected deleterious effects, and this was termed the **timing hypothesis**. (55–57) Recently we have demonstrated that delayed estrogen replacement in aged ovariectomized (OVX) female Norway Brown (NB) rats led to a substantial number of gene changes in the heart including increased expression of CD11b, MIP-1 β , STAT3, EMAP II, fibronectin, caspase 6 and MADD. (58) Although neither TNF α or iNOS had increased RNA levels in this model, both had marked increase in their protein levels with late replacement. A striking finding was increased expression of genes that stimulate leukocyte attraction and adhesion, initial steps in atherosclerosis. The most significant finding was the growth in the expression of pro-inflammatory and pro-apoptotic proteins with late estrogen replacement, exemplified by the increase in TNF α and iNOS protein levels with late replacement. Since TNF α enhances the expression of adhesion molecules this would only further promote atherosclerosis. (58)

HRT Controversies B—An important issue in HRT is actual type and delivery method of hormone replacement. In HRT estrogen is generally used in combination with progesterone, except in women who have had a hysterectomy, who given the absence of the uterus can receive estrogen alone without risk of endometrial cancer. Estrogen is a powerful steroid hormone with 17 β -estradiol (E2) the most active metabolite in humans. There are two important issues in estrogen replacement. Estrogen taken orally leads to high hepatic estrogen levels secondary to first pass metabolism. These high hepatic levels of estrogen can stimulate protein synthesis and are thought to increase the risk of thrombosis compared to transdermal delivery, but this interpretation remains contentious. (59–62) Increasingly studies

support transdermal estrogen as safer with regards to thrombotic events.(63) Estrogen delivery by transdermal patch precludes the first pass surge in hepatic estrogen and may have fewer complications. Indeed, the Kronos Early Estrogen Prevention Study (KEEPS) represents an ongoing clinical intervention trial, which aims to determine the protective efficacy of early transdermal HRT administration in younger menopausal women on cardiovascular outcomes to provide a more definitive test of this hypothesis.(64)

Differences Among Estrogens—The second issue in estrogen therapy is the type of estrogen replacement. The most frequently used estrogen for HRT is conjugated equine estrogen (CEE), which is extracted from the urine of pregnant mares (Premarin). CEE consists of sodium estrone sulfate, sodium equilin sulfate, and sodium 17 α -dihydroequilenin, but no 17 β -estradiol, the primary and most potent human estrogen.(65) Various estrogens have disparate binding affinities and specificity for the estrogen receptors (ERs); this will lead to distinct downstream effects. These differences in the estrogen composition of HRT could have a significant impact on the effects of estrogen replacement.

CEE, Estrogens and Thrombosis—Estrone has been connected with thrombin generation, a critical step in the coagulation cascade.(59) In a population based case-control study of HRT in postmenopausal women over six years, CEE were associated with greater risk of venous thromboembolism (odds ratio 2.08).(66) Neither the risk of MI nor stroke reached statistical significance, but the study was underpowered with only 68 subjects. Others have reported less thrombotic events with estradiol vs. CEE.(67) Hence, estrones have the possibility to significantly enhance the chance of thrombosis, and as CEE includes a significant amount of estrones, CEE would be anticipated to increase the probability of thrombotic events. However, there are only limited studies in this area and thus it is too early to make definitive conclusions.

Early Estrogen Replacemen—As a result of the clinical trials of the 1990's and the timing hypothesis, there was a move to initiate HRT early, at the first sign of menopause, rather than waiting. As noted above, the KEEPS trial was initiated in 2005 to investigate the effects of a low dose oral CEE or transdermal patch of combination estrogen and progestin in women with an average age of 52 years (64), and results should be forthcoming. Another approach has been low dose unopposed estrogen, and there is some very limited data to suggest that this is without harm.(68;69) Although the timing hypothesis may prove supportive for short-term cardioprotection with HRT in younger postmenopausal women, the evidence for diminished efficacy and possible detrimental effects of HRT on coronary heart disease (CHD) risk in older women, as well as concerns regarding increased breast and ovarian cancer risk with long-term HRT use remain problematic.

Estrogen and Myocardial Ischemia—Premenopausal women have decreased risk for CHD compared to age-matched men(70), as well as a decreased incidence of LV hypertrophy and cardiac remodeling following myocardial infarction (MI).(71) Post menopause the prevalence of CHD increases, such that aged women have both reduced ischemic tolerance(72;73) and increased mortality following MI(74) relative to age-matched men. In distinction from aging, independent effects of E₂ deficiency on cardiovascular risk

have also been observed. As early as 1953, Wuest *et al.* noted the increased prevalence of coronary artery disease in autopsy studies of premenopausal women who had undergone ovariectomy (OVX) (75), and numerous studies conducted throughout the ensuing five decades have demonstrated increased risk for CHD and MI in both postmenopausal and ovariectomized premenopausal women.(70;76–79) Indeed, E₂ deficiency alone reduces ischemic tolerance in hearts of both adult mice(80) and rats.(81) Epidemiological data also indicate the interaction of gender and aging, and influence of menopausal status on the determination of cardiovascular risk in aging women. Animal and human studies have each identified both functional and cellular alterations in ischemic tolerance and cardioprotection due to the independent and combined effects of aging and E₂ deficiency. A notable limitation in identifying specific mechanistic underpinnings in the adult and aged female heart has been differences in experimental models used to recapitulate postmenopausal E₂ deficiency.(82) In the paragraphs that follow, information regarding the influence of E₂ on cardioprotective signals is presented within the context of information gleaned from studies associated with the protective phenomenon of ischemic preconditioning (IPC). Particular emphasis is also placed on available experimental models of E₂ deficiency and aging.

Estrogen Receptors (ER) and Cardioprotection—As noted above, effects of E₂ in the heart are primarily mediated by ER α and ER β , although the precise subcellular distribution of cardiac ER receptors remains to be fully elucidated. Evidence linking ER α and ER β polymorphisms to adverse cardiac outcomes in women(83–85) suggest that ER α and ER β may each play distinct roles in cardioprotection. While the importance of cardiac ER subtypes in ischemic injury remains controversial, studies employing ER α and ER β deficient mice have each demonstrated reductions in ischemic tolerance.(86;87) However, it is important to note that ER deficiency in these models is not cardiac-specific, and some results are confounded by use of mice which encode a truncated ER α , as well as a metabolic phenotype which develops with age.(88;89) Nevertheless, in mice completely null for ER α , greater I/R (ischemia/reperfusion) injury and more impaired mitochondrial function(90;91) were observed vs. wild type mice. Further, activation of ER α with the specific agonist, propyl pyrazole triol (PPT), protects the *in vivo* rabbit heart from I/R injury, while the specific ER β activator, diarylpropionitrile (DPN) was without effect.(92) Recent studies also suggest a greater role for ER α vs. ER β in the modulation of endothelial progenitor cells and cardiac repair.(93;94) Taken together, these data support a dominant role for ER α as the cardioprotective ER involved in I/R injury.(92)

In contrast, acute ER β activation does not appear to impact functional recovery following I/R injury in either adult or aged female rats, and ER β mRNA was not detected.(26) The lack of measurable ER β in the F344 rat myocardium(26) was surprising given results gleaned from past studies utilizing the ER β knockout mouse model (86;95;96) mentioned heretofore. In this regard, ER β expression in the rodent myocardium remains controversial (10;97–102), and the protein signal produced by ER β antibodies in cardiac homogenates may be the result of cross-reactivity with ER α .(26) Combined with these previous findings, either ER β signaling varies substantially between rat, rabbit and murine models or, cardioprotection observed in mouse models may be mediated indirectly through extra-cardiac ER β signaling. For instance, DPN injection at the rostral ventrolateral medulla, an

area associated with autonomic cardiovascular control, has been shown to reduce systemic arterial pressure in rats.(103) That ER β activation can reduce systemic arterial pressure via autonomic influence indicates that additional autonomic cardioprotective mechanisms attributed to E₂ may be mediated through ER β . Indeed, E₂-linked cardioprotection has been associated with reduced sympathetic input to the heart and vasculature during ischemia in female rats, resulting in reduced heart rate, mean arterial pressure, arrhythmia frequency, and overall improved ischemic tolerance vs. males.(104;105) Therefore it is plausible that hypertension and vascular dysfunction observed in whole body ER β knockout mice as well as cardioprotection observed in chronic DPN treated mice may be explained by indirect ER β effects on autonomic cardiac control and not direct effects on the myocardium.(106;107) Definitive studies on ER subtype distribution in adult myocardium are needed.

GPER may also activate the nonnuclear protective response via a truncated 36 kDa ER α .(108) GPER activation by G1 reduced infarct size in both the isolated perfused male mouse heart and the isolated perfused male Wistar rat heart.(43;109) Opening of the mitochondrial permeability transition pore (mPTP) was also inhibited by GPER activation in the same study. Treatment with the ERK inhibitor, PD98059, eliminated the protective benefits of GPER activation. GPER activation can attenuate diastolic dysfunction and ventricular remodeling after OVX.(40) The demonstration that rapid ER α and/or GPER activation reduces I/R injury in the female heart supports a key role for non-genomic ER signaling in the maintenance of cardioprotection. In this regard, 17- β estradiol is the major physiological E₂, but it has a similar affinity for ER α and ER β . As noted, a number of selective ER α and ER β agonists have been created and described; however, only a minority of these compounds has been evaluated extensively *in vivo*. The discovery of the GPER has also reinforced the need for additional ER specific modulators. Selective estrogen receptor modulators (SERMs) may be of great utility and in understanding the role of ERs in ischemic tolerance with aging.

PI3K-Akt-GSK-3 β Signaling and Estrogen—Interestingly, many of the protective actions mediated by rapid ER signaling involve downstream effectors known to be associated with IPC, such as PI3K-Akt, eNOS and PKC ϵ (figure 2, for review see 110). Increased levels and/or activity of Akt have also been observed in female (vs male) animal and human myocardium.(111;112) ER α -mediated nuclear transcription is also affected by Akt, and nuclear accumulation of Akt in human cardiocytes is increased 5.8-fold in adult women over men and reduced in postmenopausal women.(111) Collectively, these data suggest that the PI3K-Akt pathway is acutely activated by E₂. Urata and colleagues(113) recently demonstrated that E₂ administration (18 hrs.) in H9c2 cells leads to a reduction in hydrogen peroxide (H₂O₂)-induced apoptosis through upregulation of glutaredoxin (GRX), which was abolished by the ER inhibitor ICI-182,780. Effects were presumed ER β -mediated since these cells do not express ER α .

A target of Akt which has been proposed as a convergence point for many cardioprotective signals is inactivation of mitochondrial GSK-3 β and associated apoptotic signaling. This model is supported by I/R- and OVX-dependent changes in mitochondrial pGSK-3 β which mirror changes in pAkt in adult but interestingly *not* aged rats (114), suggesting dysregulated Akt-GSK-3 β interactions in aged. Additional mechanisms by which rapid E₂

signaling may influence GSK-3 β and subsequent ischemic tolerance are worth noting. Recent studies suggest that GSK-3 β can enhance ER α -mediated transcription (115) implicating the nuclear compartment as an potentially important site of regulation in the female heart. If this is so, several cardioprotective or apoptotic proteins that are modulated by E₂, such as heat shock proteins (116), ANT-1(117), or Cx43 (118) may show altered expression or activity, thus contributing to reduced ischemic tolerance in aging. Future studies are indicated to determine the role, if any, of altered gene expression in relation to cell survival with age-associated E₂ deficiency.

Age, Estrogen, and Ischemic Tolerance

Ischemia/Reperfusion Injury in Aging—Reduced ischemic tolerance and increased susceptibility of the heart to I/R injury is a hallmark adaptation of both aged human and animal hearts.(119–129) The aged heart is also refractory to endogenous protection from interventions like IPC, verifying inadequate protective cellular reserves.(129–131) The precise cellular mechanisms underlying this dysfunction, however, are incompletely understood. The problem is further exacerbated by the **paucity** of studies using females, limiting extrapolation of results. Reversal of cardioprotection with senescence is likely to involve aberrations in both intrinsic (i.e. excitation-contraction coupling) and extrinsic (adrenergic) inotropic regulatory mechanisms (for review see 122;131). However, alterations in cell signaling pathways related to metabolic and oxidative stress may also shift the balance from cell survival to cell death regulating pathways.(132–136) Extensive and ongoing research has thus focused on the identification of effective treatments for the reduction of I/R injury (termed *cardioprotection*), which may be implemented in a clinical setting of acute MI to limit infarct size and minimize loss of cardiac function.

With regard to mechanistic underpinnings, increases in infarct size following I/R in aged relative to adult female rats have been associated with decreased Akt and mitochondrial PKC ϵ levels, as well as increased mitochondrial GSK-3 β .(114) Hunter *et al.* further found that aged OVX rats exhibited more severely impaired functional recovery and greater infarct size following I/R than was seen with aging or OVX alone, suggesting an additive detriment of aging and OVX in the female rat heart.(114) In support of a protective role for PKC ϵ targeting in the aged, E₂-deficient female rat heart, acute activation of PKC ϵ prior to ischemia by local delivery of ψ eRACK peptide has been associated with 1) improved functional recovery and reduced infarct size, 2) increased mitochondrial targeting of PKC ϵ , and 3) candidate downstream signaling targets suggesting a role for activation of antioxidant enzymes as a mechanism of PKC ϵ -mediated protection.(137) Specifically, mitochondrial HSP10, GPX, and SOD2 (MnSOD) abundance are significantly increased with ψ eRACK administration in aged OVX hearts (by ~10, 20, and 30%, respectively) and likely attributable to PKC ϵ -stimulated mitochondrial translocation or import of identified proteins.

With regard to the potential cardioprotective role of non-genomic ER activation in reducing I/R injury in aged female hearts, our work has also implicated a possible role for selective ER α activation as follows(25): 1) effectively reduced infarct size, 2) resulted in greater mitochondrial and particulate ER α localization coordinate with a protective pattern of PKC ϵ activation, and 3) enhanced gene expression of the PKC ϵ anchoring protein RACK2.

Collectively, these results demonstrate a protective role for non-genomic ER α signaling in the aged female rat heart, the cellular basis of which may involve two distinct PKC ϵ -dependent mechanisms. What are less clear are the mechanisms which underlie altered cardiac ER translocation. As noted above, PTMs such as phosphorylation, acetylation and sumoylation (for review see 17–20;138) are known to effect ER targeting, the effects of which are unstudied in aging. Since some non-genomic ER effects are specific to aged animals(25), it will be important that future studies incorporate true models of aging in conjunction with E₂ deficiency to fully characterize the non-genomic ER response.

Mitochondrial Mechanisms of Cell Death—Mitochondria are the main source of both ATP and reactive oxygen species (ROS) in the heart ideally positioning them as mediators of, and therapeutic targets for, ischemic CHD. Because of the pivotal role played by the mitochondria in the maintenance of cell survival and cardioprotection, it is logical that age-associated reductions in ischemic tolerance might arise from alterations in mitochondrial proteins. Given the estimate that 1000 to 2000 proteins are expressed in the mitochondria(139), it is likely that the adaptation of additional mitochondrial proteins in aging and/or E₂ deficiency may contribute to the reductions in ischemic tolerance and increased I/R injury associated with advancing age and menopause. While correlational relationships between age-dependent declines in ischemic tolerance and altered expression and localization of cardioprotective signaling proteins have been noted in the female heart, the breadth and extent of protein changes have only recently been addressed. Notably, the reader is referred to several recent complimentary reviews on mitochondrial aging and mechanisms of cell death.(6;140–142)

Using a high throughput proteomics approach targeting the cardiac mitochondrial subproteome in adult and aged F344 female rats, we observed significant directional changes in **67** proteins with aged and/or aged OVX, and **32** were unique to aged OVX.(143) Notably only **6** proteins were similarly altered in adult OVX (voltage-dependent ion channel 1, adenine nucleotide translocator 1, cytochrome c oxidase subunits VIIc and VIc, catalase, and myosin binding protein C), highlighting the specificity of the E₂ deficiency response in adult vs. aged female rats. Proteins affected by aging were primarily related to cellular metabolism, oxidative stress and cell death, with the largest change seen in monoamine oxidase-A, a potential source of oxidative stress. About 50% of the identified proteins altered in aged OVX were associated with mitochondrial ATP production.(143) In aged, E₂-deficient female hearts, reduced quantity of protein subunits of electron transport chain complex I (NADH dehydrogenase), II (succinate dehydrogenase), III (cytochrome *bc*₁ complex), IV (cytochrome *c* oxidase), and V (F₀F₁ ATPase), and bidirectional changes in proteins involved in fatty acid substrate metabolism (acyl Co-A synthetase subunits) have been observed. In contrast, increases were primarily observed for proteins involved in carbohydrate and amino acid metabolism (pyruvate dehydrogenase subunits) and enzymes of the tricarboxylic acid cycle.(143) Increased levels of HSP60 and mtHSP70 in aged OVX are consistent with previous studies in aged male hearts(144) and may be related to alterations in mitochondrial matrix protein import of nuclear-encoded enzymes, which may or may not be balanced by changes in proteolysis. Measurement of the activity and/or

phosphorylation status(145) of these proteins is indicated for a more comprehensive characterization of metabolic alterations and substrate utilization in the aged female heart.

Estrogen, Cardiac Myocytes and the Heat Shock Response

E2 indirectly regulates cardiac HSP 72 expression in both male and female adult rat cardiac myocytes.(116;146;147) Cardiac HSP72 levels in female Sprague Dawley (SD) rats are twice that of males.(147) OVX reduces HSP72 to the levels of males over a period of 9 weeks, supporting an indirect estrogen effect. Besides its facilitating function in protein folding, HSP72 has cardioprotective effects, and reduces apoptosis by stabilizing the mitochondrial membrane and preventing apoptosome formation.(148–150) Preventing the normal increase in HSP72 in response to hypoxia/reoxygenation with antisense increases cardiac myocyte injury.(151) In isolated adult SD male and female cardiac myocytes, pharmacologic concentrations of E2 increased HSP 72 expression through consecutive activation of NFκB and HSF-1.(figure 2;116;146) Pre-treatment with E2 reduced injury after hypoxia/reoxy-genation.(116;146) E2 had similar effects in human coronary artery endothelial cells (HCAEC) from young donors.(152)

In young adult cardiac myocytes and in HCAEC, E2 can increase HSP 72 expression. In HCAEC this occurs through simultaneous activation of P38, JNK and Akt leading to ERK 1/2 activation followed by activation of NFκB.(153) The SERMS, raloxifene and tamoxifen activated the same pathways, except they did not activate JNK.(153) However in cardiac myocytes from aged OVX (studied 9 weeks post-OVX) NB rats with and without immediate E2 replacement, the activation of NFκB and HSF1 by E2 was lost.(24) Similarly, treatment with hypoxia and reoxygenation, which readily induces expression of HSP72, had no effect on HSP72 in aged cardiac myocytes, but did increase HSP72 in cardiac myocytes from adult (5–6 mo.) rats with OVX and with/without E2 replacement.(24) Thus, two different activators of HSP expression had no effect in aged female myocytes, regardless of estrogen status. HSP expression is controlled primarily by HSF(heat shock factor)1. In male hearts and skeletal muscle there is a reduction in activation of HSF1 and a loss of the heat shock response (upregulation of HSPs).(154;155) Similarly, aged male hepatocytes have loss of activation of HSF1 with stress.(156) In skeletal muscle there was no difference in HSF1 binding by EMSA, but no increase in HSP72 with exercise.(157) Electromobility shift (EMSA) measures binding of HSF1 to the heat shock element, but this needs to be followed by increased transcription. Phosphorylation of serines 303/307 on HSF1, as we saw in the aged female heart, does not interfere with binding, but prevents initiation of transcription.(24;158) Different causes for the loss of the heat shock response have been found in male tissues including reduced binding of HSF1 to the heat shock element (HSE) and loss of stability of the HSF1 trimer, which forms prior to nuclear translocation and binding of HSF1 to the HSE.(156) As the normal heat shock response reduces inflammation, stabilizes protein structure and facilitates removing irreversibly denatured proteins, loss of this response with aging is quite detrimental.

Cardiac Myocyte Changes with Aging—Basic studies are important to understand differences that occur with aging and loss of estrogen, so that we can better comprehend the mechanisms and extent of cardiovascular changes in aging females. Studies of cardiac

myocytes in aging and estrogen loss are remarkably limited, as most studies of estrogen loss using very young rodent models. We have found that cardiac myocytes from aged (22 mo.) OVX NB rats with and without immediate E2 replacement had differences in the expression of inflammatory cytokines.(24) Although plasma levels of IL-6 and TNF α did not differ amongst the groups in the study, the aged OVX cultured cardiac myocytes had increased expression of IL-6 and TNF α mRNA compared to all other groups, indicating that cytokines tissue levels may be more informative than plasma levels.(24) IL-6 and TNF α levels were decreased by treating the cardiac myocytes with E2 for 6 h. The aged OVX-derived cardiac myocytes also had a greatly reduced ability to handle reactive oxygen species (ROS), which increase with age.(24;159) E2 treatment in culture did not reduce ROS levels, even though it had reduced inflammatory cytokine levels. In other work, Ross and Howlett found that cardiac myocytes isolated from adult (3 mo.) Fisher 344 female rats had better recovery after simulated I/R and better resistance to injury compared to cardiac myocytes from male Fisher 344 rats.(160) Cardiac myocytes from adult OVX (5 mo. age) and aged Fisher 344 rats (24 mo.) lost the gender benefits seen in cardiac myocytes from young intact females. Overall, aged OVX derived cardiac myocytes had loss of the heat shock response, increased inflammatory cytokines and markedly reduced ability to handle ROS. E2 replacement at the time of OVX improved many of the changes, but inactivation of HSF1 by phosphorylation of serines 303/307 persisted.

Summary/Conclusions

Much has been learned with regards to the ERs, the cardioprotective effects of estradiol, and the nongenomic, downstream signaling cascade. However, many questions remain unanswered and/or incomplete. Further knowledge of the distribution of the three receptors is needed, and to what degree these receptors mediate functional responses to different estrogen compounds in different cell types/cellular locations. There is also a substantial literature supporting the use of 17 β -estradiol to ameliorate traumatic injury.(161) A better understanding of the non-genomic actions of estrogen can potentially lead to improved clinical therapeutic interventions for treating acute coronary syndrome in aged women, specifically selective modulation of cardiac ERs and non-genomic ER signaling in an attempt to harness the protection associated with E₂ observed in adult women without increased cardiovascular risk observed from chronic HRT.

Acknowledgments

This work was supported by a Merit Award(5101BX000839) from the U.S. Department of Veterans' Affairs, Office of Research and Development, Biomedical Laboratory Research Program (AAK); NIH RO1 HL091907 (DHK) and AHA 12UFEL10340001 (DHK).

Abbreviation

ANT-1	Adenine Nucleotide Translocator 1
CEE	Conjugated Equine Estrogen
CHD	Coronary Heart Disease
Cx43	Connexin 43

EMSA	Electrophoretic Mobility Shift Assay
eNOS	Endothelial Nitric Oxide Synthase
ER	Estrogen Receptor
GPER	G-Protein-Coupled Estrogen Receptor
HSF	Heat Shock Factor
HSP	Heat Shock Protein
iNOS	Inducible Nitric Oxide Synthase
IPC	Ischemic Preconditioning
IR	Ischemia Reperfusion
MI	Myocardial Infarction
NB	Norway Brown
OVX	Ovariectomy
PTM	Post translational modification
VSMC	Vascular Smooth Muscle Cell

Reference List

1. Xing D, Nozell S, Chen YF, Hage F, Oparil S. Estrogen and Mechanisms of Vascular Protection. *Arterioscler Thromb Vasc Biol.* 2009; 29(3):289–295. [PubMed: 19221203]
2. Deschamps AM, Murphy E, Sun J. Estrogen Receptor Activation and Cardioprotection in Ischemia Reperfusion Injury. *Trends in Cardiovascular Medicine.* 2010; 20(3):73–78. [PubMed: 21130949]
3. Farsetti A, Grasselli A, Bacchetti S, Gaetano C, Capogrossi MC. The telomerase tale in vascular aging: Regulation by estrogens and nitric oxide signaling. *Journal of Applied Physiology.* 2008:91360.
4. Knowlton AA, Lee AR. Estrogen and the Cardiovascular System. *Pharmacology & Therapeutics.* 2012; 135(1):54–70. [PubMed: 22484805]
5. Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress: Mechanistic insights. *The Journal of Steroid Biochemistry and Molecular Biology.* 2013; 137(0):124–135. [PubMed: 23770428]
6. Korzick D, Lancaster T. Age-related differences in cardiac ischemia-reperfusion injury: effects of estrogen deficiency. *Pflugers Arch - Eur J Physiol.* 2013; 465(5):669–685. [PubMed: 23525672]
7. Deroo BJ, Korach KS. Estrogen receptors and human disease. *J Clin Invest.* 2006; 116(3):561–570. [PubMed: 16511588]
8. Driggers, PH., Segars, JH. Part II: the role of growth factors and phosphorylation in estrogen signaling. 13. 2002. *Estrogen action and cytoplasmic signaling pathways*; p. 422-427.
9. Alexaki VI, Charalampopoulos I, Kampa M, et al. Activation of membrane estrogen receptors induce pro-survival kinases. *The Journal of Steroid Biochemistry and Molecular Biology.* 2006; 98(2–3):97–110. [PubMed: 16414261]
10. Yang SH, Liu R, Perez EJ, et al. Mitochondrial localization of estrogen receptor beta. *Proc Natl Acad Sci U S A.* 2004; 101:4130–4135. [PubMed: 15024130]

11. Zhai P, Eurell TE, Cotthaus RP, Jeffery EH, Bahr JM, Gross DR. Effects of dietary phytoestrogen on global myocardial ischemia-reperfusion injury in isolated female rat hearts. *Am J Physiol Heart Circ Physiol*. 2001; 281(3):H1223–H1232. [PubMed: 11514291]
12. Meldrum DR. Estrogen increases protective proteins following trauma and hemorrhage. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*. 2006; 290(3):R809–R811.
13. Keung W, Chan MLY, Ho EYW, Vanhoutte PM, Man RYK. Non-genomic activation of adenylyl cyclase and protein kinase G by 17 β -estradiol in vascular smooth muscle of the rat superior mesenteric artery. *Pharmacological Research*. 2011; 64:509–516. [PubMed: 21641998]
14. Nakamura Y, Suzuki T, Miki Y, et al. Estrogen receptors in atherosclerotic human aorta: inhibition of human vascular smooth muscle cell proliferation by estrogens. *Molecular and Cellular Endocrinology*. 2004; 219:17–26. [PubMed: 15149723]
15. Watanabe T, Akishita M, Nakaoka T, et al. Estrogen receptor β mediates the inhibitory effect of estradiol on vascular smooth muscle cells proliferation. *Cardiovascular Research*. 2003; 59:734–744.
16. Takahashi K, Ohmichi M, Yoshida M, et al. Both estrogen and raloxifene cause G1 arrest of vascular smooth muscle cells. *Journal of Endocrinology*. 2003; 178:319–329. [PubMed: 12904179]
17. Chen D, Washbrook E, Sarwar N, et al. Phosphorylation of human estrogen receptor α at serine 118 by two distinct signal transduction pathways revealed by phosphorylation-specific antisera. *Oncogene*. 2002; 21:4921–4931. [PubMed: 12118371]
18. Faus H, Haendler B. Post-translational modifications of steroid receptors. *Biomedicine & Pharmacotherapy*. 2006; 60(9):520–528. [PubMed: 16949786]
19. Fu M, Wang C, Zhang X, Pestell RG. Acetylation of nuclear receptors in cellular growth and apoptosis. *Biochemical Pharmacology*. 2004; 68(6):1199–1208. [PubMed: 15313417]
20. Zhang D, Trudeau VL. Integration of membrane and nuclear estrogen receptor signaling. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2006; 144(3):306–315.
21. Revankar CM, Cimino DF, Sklar LA, Arterburn JB, Prossnitz ER. A Transmembrane Intracellular Estrogen Receptor Mediates Rapid Cell Signaling. *Science*. 2005; 1106943.
22. Dan P, Cheung JCY, Scriven DRL, Moore EDW. Epitope-dependent localization of estrogen receptor α , but not β in en face arterial endothelium. *American Journal of Physiology - Heart and Circulatory Physiology*. 2003; 284:H1295–H1306. [PubMed: 12531733]
23. Liao WX, Magness RR, Chen D. Expression of Estrogen Receptors- α and β in the Pregnant Ovine Uterine Artery. *Biol Reprod*. 2005; 72:530–537. [PubMed: 15564597]
24. Stice JP, Chen L, Kim SC, et al. 17 β -Estradiol, Aging, Inflammation and the Stress Response in the Female Heart. *Endocrinology*. 2011; 152:1589–1598. [PubMed: 21303943]
25. Novotny JL, Simpson AM, Tomicek NJ, Lancaster TS, Korzick DH. Rapid Estrogen Receptor- α Activation Improves Ischemic Tolerance in Aged Female Rats through a Novel Protein Kinase C ϵ -Dependent Mechanism. *Endocrinology*. 2009; 150(2):889–896. [PubMed: 19176323]
26. Tomicek NJ, Miller-Lee JL, Hunter JC, Korzick DH. Estrogen Receptor Beta Does Not Influence Ischemic Tolerance in the Aged Female Rat Heart. *Cardiovascular Therapeutics*. 2013; 31(1):32–37. [PubMed: 21884022]
27. Kim KH, Bender JR. Membrane-initiated actions of estrogen on the endothelium. *Molecular and Cellular Endocrinology*. 2009; 308(1–2):3–8. [PubMed: 19549586]
28. Haynes MP, Li L, Sinha D, et al. Src Kinase Mediates Phosphatidylinositol 3-Kinase/Akt-dependent Rapid Endothelial Nitric-oxide Synthase Activation by Estrogen. *Journal of Biological Chemistry*. 2003; 278(4):2118–2123. [PubMed: 12431978]
29. Li L, Haynes MP, Bender JR. Plasma membrane localization and function of the estrogen receptor α variant (ER46) in human endothelial cells. *PNAS*. 2003; 100(8):4807–4812. [PubMed: 12682286]
30. Levin ER. Plasma membrane estrogen receptors. *Trends in Endocrinology and Metabolism*. 2010; 20(10):477–482.

31. Acconcia F, Ascenzi P, Bocedi A, et al. Palmitoylation-dependent Estrogen Receptor α Membrane Localization: Regulation by 17β -Estradiol. *Molecular Biology of the Cell*. 2005; 16:231–237. [PubMed: 15496458]
32. Chambliss KL, Yuhanna IS, Mineo C, et al. Estrogen Receptor {alpha} and Endothelial Nitric Oxide Synthase Are Organized Into a Functional Signaling Module in Caveolae. *Circulation Research*. 2000; 87(11):44e–452.
33. Figtree GA, McDonald D, Watkins H, Channon KM. Truncated Estrogen Receptor {alpha} 46-kDa Isoform in Human Endothelial Cells: Relationship to Acute Activation of Nitric Oxide Synthase. *Circulation*. 2003; 107(1):120–126. [PubMed: 12515753]
34. Mineo C, Shaul PW. Circulating cardiovascular disease risk factors and signaling in endothelial cell caveolae. *Cardiovascular Research*. 2006; 70(1):31–41. [PubMed: 16540099]
35. Razani B, Engelman JA, Wang XB, et al. Caveolin-1 Null Mice Are Viable but Show Evidence of Hyperproliferative and Vascular Abnormalities. *Journal of Biological Chemistry*. 2001; 276(41):38121–38138. [PubMed: 11457855]
36. Boonyaratanakornkit V. Scaffolding proteins mediating membrane-initiated extra-nuclear actions of estrogen receptor. *Steroids*. 2011; 76:877–884. [PubMed: 21354435]
37. Fleming I. Molecular mechanisms underlying the activation of eNOS. *Pflugers Archives - European Journal of Physiology*. 2010; 459:793–806. [PubMed: 20012875]
38. Dessy C, Feron O, Balligand JL. The regulation of endothelial nitric oxide synthase by caveolin: a paradigm validated *in vivo* and share by the 'endothelium-derived hyperpolarizing factor'. *Pflugers Archives - European Journal of Physiology*. 2010; 459:817–827. [PubMed: 20339866]
39. Wu Q, Chambliss KL, Umetani M, Mineo C, Shaul PW. Non-nuclear Estrogen Receptor Signaling in the Endothelium. *Journal of Biological Chemistry*. 2011; 286(17):14737–14743. [PubMed: 21343284]
40. Wang H, Jessup JA, Lin MS, Chagas C, Lindsey SH, Groban L. Activation of GPR30 Attenuates Diastolic Dysfunction and Left Ventricle Remodeling in Oophorectomized mRen2.Lewis Rats. *Cardiovascular Research*. 2012 in press.
41. Deschamps AM, Murphy E. Activation of a novel estrogen receptor, GPER, is cardioprotective in male and female rats. *American Journal of Physiology - Heart and Circulatory Physiology*. 2009; 297:H1806–H1813. [PubMed: 19717735]
42. Yu X, Handong M, Barman SA, et al. Activation of G protein-coupled estrogen receptor induces endothelium-independent relaxation of coronary artery smooth muscle. *American Journal of Physiology - Endocrinology And Metabolism*. 2011; 301:E882–E888. [PubMed: 21791623]
43. Patel VH, Chen J, Ramanjaney A, et al. G-protein coupled estrogen receptor 1 expression in rat and human heart: Protective role during ischaemic stress. *International Journal of Molecular Medicine*. 2010; 26:193–199. [PubMed: 20596598]
44. Recchia AG, De Francesco EM, Vivacqua A, et al. The G Protein-coupled Receptor 30 Is Up-regulated by Hypoxia-inducible Factor-1 (HIF-1) in Breast Cancer Cells and Cardiomyocytes. *Journal of Biological Chemistry*. 2011; 286(12):10773–10782. [PubMed: 21266576]
45. Delbeck M, Golz S, Vonk R, et al. Impaired left-ventricular cardiac function in male GPR30-deficient mice. *Molecular Medicine Reports*. 2011; 4(1):37–40. [PubMed: 21461560]
46. Lindsey SH, Carver KA, Prossnitz ER, Chappell MC. Vasodilation in response to the GPR30 Agonist G-1 is Not Different from Estradiol in the mRen2.Lewis Female Rat. *Journal of Cardiovascular Pharmacology*. 2011; 57:598–603. [PubMed: 21326105]
47. Meyer MR, Field AS, Kanagy NL, Barton M, Prossnitz ER. GPER regulates endothelin-dependent vascular tone and intracellular calcium. *Life Sciences*. 2012; 91(13–14):623–627. [PubMed: 22326502]
48. Feldman RD, Gros R. Unraveling the mechanisms underlying the rapid vascular effects of steroids: sorting out the receptors and the pathways. *British Journal of Pharmacology*. 2011; 163(6):1163–1169. [PubMed: 21545416]
49. Batenburg WW, Jansen PM, van den Bogaerdt AJ, Danser J. Angiotensin II-aldosterone interaction in human coronary microarteries involves GPR30, EGFR, and endothelial NO synthase. *Cardiovascular Research*. 2012; 94(1):136–143. [PubMed: 22260839]

50. Gros R, Ding Q, Liu B, Chorazyczewski J, Feldman RD. Aldosterone mediates its rapid effects in vascular endothelial cells through GPER activation. *American Journal of Physiology - Cell Physiology*. 2013; 304(6):C532–C540. [PubMed: 23283935]
51. Brailoiu GC, Benamar K, Arterburn JB, et al. Aldosterone increases cardiac vagal tone via G protein-coupled oestrogen receptor activation. *The Journal of Physiology*. 2013; 591(17):4223–4235. [PubMed: 23878371]
52. Hulley S, Grady D, Bush T, et al. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women. Heart and Estrogen/progestin Replacement Study (HERS) Research Group. *JAMA*. 1998; 280:605–613. [PubMed: 9718051]
53. Grady D, Herrington D, Bittner V, et al. Cardiovascular disease outcomes during 6.8 years of hormone therapy: Heart and Estrogen/Progestin Replacement study follow-up (HERS II). *JAMA*. 2002; 288:49–57. [PubMed: 12090862]
54. Writing Group for the Women's Health Initiative Investigators. Risks and Benefits of Estrogen Plus Progestin in Healthy Postmenopausal Women: Principal Results From the Women's Health Initiative Randomized Controlled Trial. *JAMA*. 2002; 288(3):321–333. [PubMed: 12117397]
55. Barrett-Connor E. Hormones and heart disease in women: the timing hypothesis. *American Journal of Epidemiology*. 2007; 166:506–510. [PubMed: 17849510]
56. Dubey RK, Imthurn B, Barton M, Jackson EK. Vascular consequences of menopause and hormone therapy: Importance of timing of treatment and type of estrogen. *Cardiovascular Research*. 2005; 66:295–306. [PubMed: 15820198]
57. Grodstein F, Clarkson T, Manson JE. Understanding the Divergent Data on Postmenopausal Hormone Therapy. *New England Journal of Medicine*. 2003; 348(7):645–650. [PubMed: 12584376]
58. Pechenino AS, Lin L, Mbai FN, et al. Impact of Aging vs. Estrogen Loss on Cardiac Gene Expression: Late Estrogen Replacement and Inflammation. *Physiol Genomics*. 2011; 43:1065–1073. [PubMed: 21750230]
59. Bagot CN, Marsh MS, Whitehead M, et al. The effect of estrone on thrombin generation may explain the different thrombotic risk between oral and transdermal hormone replacement therapy. *Journal of Thrombosis and Haemostasis*. 2010; 8:1736–1744. [PubMed: 20553380]
60. Canonico M, Fournier A, Caracailon L, et al. Postmenopausal Hormone Therapy and Risk of Idiopathic Venous Thromboembolism: Results From E3N Cohort Study. *Arterioscler Thromb Vasc Biol*. 2010; 30:340–345. [PubMed: 19834106]
61. Tchaikovski SN, Rosing J. Mechanisms of Estrogen-Induced Venous Thromboembolism. *Thrombosis Research*. 2010; 126:5–11. [PubMed: 20163835]
62. Olié V, Canonico M, Scarabin PY. Postmenopausal hormone therapy and venous thromboembolism. *Thrombosis Research*. 2011; 127(Supplement 3)(0):S26–S29.
63. Roach, Rej, Lijfering, WM., Helmerhorst, FM., Cannegieter, Sc, Rosendaal, Fr, Van Hylckama Vlieg, A. The risk of venous thrombosis in women over 50-years old using oral contraception or postmenopausal hormone therapy. *Journal of Thrombosis and Haemostasis*. 2013; 11(1):124–131. [PubMed: 23136837]
64. Harman SM, Brinton EA, Cedars M, et al. KEEPS: The Kronos Early Estrogen Prevention Study. *Climacteric*. 2005; 8(1):3–12.
65. Booth EA, Lucchesi BR. Estrogen-Mediated Protection in Myocardial Ischemia-Reperfusion Injury. *Cardiovascular Toxicology*. 2008; 8:101–113. [PubMed: 18683081]
66. Smith NL, Blondon M, Wiggins KL. Lower risk of cardiovascular events in postmenopausal women taking oral estradiol compared with oral conjugated equine estrogens. *JAMA Internal Medicine*. 2013
67. Smith NL, Heckbert SR, Lemaitre RN, et al. Conjugated Equine Estrogen, Esterified Estrogen, Prothrombotic Variants, and the Risk of Venous Thrombosis in Postmenopausal Women. *Arterioscler Thromb Vasc Biol*. 2006; 26(12):2807–2812. [PubMed: 16973976]
68. Stevenson JC, Hodis H, Pickar JH, Lobo RA. Coronary heart disease and menopause management: The swinging pendulum of HRT. *Atherosclerosis*. 2009; 207:336–340. [PubMed: 19560146]

69. Grodstein F, Manson JE, Stampfer MJ. Hormone Therapy and Coronary Heart Disease: The Role of Time since Menopause and Age at Hormone Initiation. *Journal of Women's Health*. 2006; 15(1):35–44.
70. Barrett-Connor E. Sex Differences in Coronary Heart Disease: Why Are Women So Superior? The 1995 Ancel Keys Lecture. *Circulation*. 1997; 95(1):252–264. [PubMed: 8994444]
71. Hayward CS, Kelly RP, Collins P. The roles of gender, the menopause and hormone replacement on cardiovascular function. *Cardiovascular Research*. 2000; 46(1):28–49. [PubMed: 10727651]
72. Arnstein PM, Buselli EF, Rankin SH. Women and heart attacks: prevention, diagnosis and care. *Nurse Pract*. 1996; 21:57–58.
73. Robertson T, Kennard ED, Mehta S, et al. Influence of gender on in-hospital clinical and angiographic outcomes and on one-year follow-up in the New Approaches to Coronary Intervention (NACI) registry. *American Journal of Cardiology*. 1997; 80:26K–39K.
74. Wenger NK, Shaw LJ, Vaccarino V. Coronary Heart Disease in Women: Update 2008. *Clin Pharmacol Ther*. 2007; 83(1):37–51. [PubMed: 18043680]
75. WUEST JH, DRY TJ, EDWARDS JE. The Degree of Coronary Atherosclerosis in Bilaterally Oophorectomized Women. *Circulation*. 1953; 7(6):801–809. [PubMed: 13051821]
76. COCHRAN R, GWINUP G. Coronary artery disease in young females: Possibility of oophorectomy as an etiologic agent. *Archives of Internal Medicine*. 1962; 110(2):162–165. [PubMed: 13880078]
77. Colditz GA, Willett WC, Stampfer MJ, Rosner B, Speizer FE, Hennekens CH. Menopause and the Risk of Coronary Heart Disease in Women. *New England Journal of Medicine*. 1987; 316(18): 1105–1110. [PubMed: 3574358]
78. Falkeborn, M., Schairer, C., Naessén, T., Persson, I. Risk of myocardial infarction after oophorectomy and hysterectomy. 53. 2000. p. 832-837.
79. Rosenberg L, Hennekens CH, Rosner B, Belanger C, Rothman KJ, Speizer FE. Early menopause and the risk of myocardial infarction. *Am J Obstet Gynecol*. 1981; 139:47–51. [PubMed: 7457520]
80. Song X, Li G, Vaage J, Valen G. Effects of sex, gonadectomy, and oestrogen substitution on ischaemic preconditioning and ischaemia–reperfusion injury in mice. *Acta Physiologica Scandinavica*. 2003; 177(4):459–466. [PubMed: 12648163]
81. Kam KWL, Qi JS, Chen M, Wong TM. Estrogen Reduces Cardiac Injury and Expression of {beta}1-Adrenoceptor upon Ischemic Insult in the Rat Heart. *Journal of Pharmacology and Experimental Therapeutics*. 2004; 309(1):8–15. [PubMed: 14718598]
82. Miller V, Kaplan J, Schork N, et al. Strategies and methods to study sex differences in cardiovascular structure and function: a guide for basic scientists. *Biology of Sex Differences*. 2011; 2(1):14. [PubMed: 22152231]
83. Peter I, Shearman AM, Vasan RS, et al. Association of Estrogen Receptor β Gene Polymorphisms With Left Ventricular Mass and Wall Thickness in Women*. *American Journal of Hypertension*. 2005; 18(11):1388–1395. [PubMed: 16280269]
84. Phillips GB. Is Atherosclerotic Cardiovascular Disease an Endocrinological Disorder? The Estrogen-Androgen Paradox. *Journal of Clinical Endocrinology & Metabolism*. 2005; 90(5):2708–2711. [PubMed: 15687329]
85. Villablanca A, Lubahn D, Shelby L, Lloyd K, Barthold S. Susceptibility to Early Atherosclerosis in Male Mice Is Mediated by Estrogen Receptor {alpha}. *Arterioscler Thromb Vasc Biol*. 2004; 24(6):1055–1061. [PubMed: 15117737]
86. Gabel SA, Walker VR, London RE, Steenbergen C, Korach KS, Murphy E. Estrogen receptor beta mediates gender differences in ischemia/reperfusion injury. *Journal of Molecular and Cellular Cardiology*. 2005; 38(2):289–297. [PubMed: 15698835]
87. Wang M, Crisostomo P, Wairiuko GM, Meldrum DR. Estrogen receptor- α mediates acute myocardial protection in females. *American Journal of Physiology - Heart and Circulatory Physiology*. 2006; 290(6):H2204–H2209. [PubMed: 16415070]
88. Bryzgalova G, Gao H, Ahren B, et al. Evidence that oestrogen receptor- α plays an important role in the regulation of glucose homeostasis in mice: insulin sensitivity in the liver. *Diabetologia*. 2006; 49(3):588–597. [PubMed: 16463047]

89. Heine PA, Taylor JA, Iwamoto GA, Lubahn DB, Cooke PS. Increased adipose tissue in male and female estrogen receptor- α knockout mice. *PNAS*. 2000; 97(23):12729–12734. [PubMed: 11070086]
90. Wang M, Tsai BM, Reiger KM, Brown JW, Meldrum DR. 17- β -Estradiol decreases p38 MAPK-mediated myocardial inflammation and dysfunction following acute ischemia. *Journal of Molecular and Cellular Cardiology*. 2006; 40(2):205–212. [PubMed: 16427650]
91. Zhai P, Eurell TE, Cotthaus R, Jeffery EH, Bahr JM, Gross DR. Effect of estrogen on global myocardial ischemia-reperfusion injury in female rats. *American Journal of Physiology*. 2000; 279:H2766–H2775. [PubMed: 11087231]
92. Booth EA, Obeid NR, Lucchesi BR. Activation of estrogen receptor- $\{\alpha\}$ protects the in vivo rabbit heart from ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol*. 2005; 289(5):H2039–H2047. [PubMed: 15994857]
93. Dawn B, Bolli R. Increasing Evidence That Estrogen Is an Important Modulator of Bone Marrow Mediated Cardiac Repair After Acute Infarction. *Circulation*. 2006; 114(21):2203–2205. [PubMed: 17116779]
94. Hamada H, Kim MK, Iwakura A, et al. Estrogen Receptors $\{\alpha\}$ and $\{\beta\}$ Mediate Contribution of Bone Marrow-Derived Endothelial Progenitor Cells to Functional Recovery After Myocardial Infarction. *Circulation*. 2006; 114(21):2261–2270. [PubMed: 17088460]
95. Wang M, Crisostomo PR, Markel T, Wang Y, Lillemoe KD, Meldrum DR. Estrogen receptor beta mediates acute myocardial protection following ischemia. *Surgery*. 2008; 144(2):233–238. [PubMed: 18656630]
96. Wang M, Wang Y, Weil B, et al. Estrogen receptor β mediates increased activation of PI3K/Akt signaling and improved myocardial function in female hearts following acute ischemia. *Am J Physiol Regul Integr Comp Physiol*. 2009; 296:R972–R978. [PubMed: 19211725]
97. Forster C, Kietz S, Hultenby K, Warner M, Gustafsson JA. Characterization of the ER β –/– mouse heart. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(39):14234–14239. [PubMed: 15375213]
98. Jankowski M, Rachelska G, Donghao W, McCann SM, Gutkowska J. Estrogen receptors activate atrial natriuretic peptide in the rat heart. *Proceedings of the National Academy of Science, USA*. 2001; 98:11765–11770.
99. Kuiper GGJM, Carlsson B, Grandien K, et al. Comparison of the Ligand Binding Specificity and Transcript Tissue Distribution of Estrogen Receptors α and β . *Endocrinology*. 1997; 138(3):863–870. [PubMed: 9048584]
100. Saunders PT, Maguire SM, Gaughan J, Millar MR. Expression of oestrogen receptor beta (ER β) in multiple rat tissues visualised by immunohistochemistry. *Journal of Endocrinology*. 1997; 154(3):R13–R16. [PubMed: 9379111]
101. Schwend T, Gustafsson J+ False positives in MALDI-TOF detection of ER α in mitochondria. *Biochemical and Biophysical Research Communications*. 2006; 343(3):707–711. [PubMed: 16563354]
102. Xu Y, Arenas IA, Armstrong SJ, Davidge ST. Estrogen modulation of left ventricular remodeling in the aged heart. *Cardiovascular Research*. 2003; 57(2):388–394. [PubMed: 12566111]
103. Shih CD. Activation of estrogen receptor beta-dependent nitric oxide signaling mediates the hypotensive effects of estrogen in the rostral ventrolateral medulla of anesthetized rats. *Journal of Biomedical Science*. 2009; 16(1):60. [PubMed: 19583861]
104. Du XJ, Riemersma RA, Dart AM. Cardiovascular protection by oestrogen is partly mediated through modulation of autonomic nervous function. *Cardiovascular Research*. 1995; 30(2):161–165. [PubMed: 7585800]
105. Du XJ, Dart AM, Riemersma RA, Oliver MF. Sex difference in presynaptic adrenergic inhibition of norepinephrine release during normoxia and ischemia in the rat heart. *Circulation Research*. 1991; 68(3):827–835. [PubMed: 1683821]
106. Nikolic I, Liu D, Bell JA, Collins J, Steenbergen C, Murphy E. Treatment with an estrogen receptor-beta-selective agonist is cardioprotective. *Journal of Molecular and Cellular Cardiology*. 2007; 42(4):769–780. [PubMed: 17362982]

107. Pelzer T, Arias Loza PA, Hu K, et al. Increased Mortality and Aggravation of Heart Failure in Estrogen Receptor- β Knockout Mice After Myocardial Infarction. *Circulation*. 2005;01.
108. Kang L, Zhang X, Xie Y, et al. Involvement of Estrogen Receptor Variant ER- α 36, Not GPR30, in Nongenomic Signaling. *Molecular Endocrinology*. 2010; 24:709–721. [PubMed: 20197310]
109. Bopassa JC, Eghbali M, Toro L, Stefani E. A novel estrogen receptor GPER inhibits mitochondria permeability transition pore opening and protects the heart against ischemia-reperfusion injury. *American Journal of Physiology - Heart and Circulatory Physiology*. 2010; 298:H16–H23. [PubMed: 19880667]
110. Murphy E, Steenbergen C. Preconditioning: The Mitochondrial Connection. *Annual Review of Physiology*. 2007; 69(1):51–67.
111. Camper-Kirby D, Welch S, Walker A, et al. Myocardial Akt Activation and Gender: Increased Nuclear Activity in Females Versus Males. *Circulation Research*. 2001; 88(10):1020–1027. [PubMed: 11375271]
112. Zhao X, Eghbali-Webb M. Gender-related differences in basal and hypoxia-induced activation of signal transduction pathways controlling cell cycle progression and apoptosis, in cardiac fibroblasts. *Endocr*. 2002; 18(2):137–145.
113. Urata Y, Ihara Y, Murata H, et al. 17 β -Estradiol Protects against Oxidative Stress-induced Cell Death through the Glutathione/Glutaredoxin-dependent Redox Regulation of Akt in Myocardiac H9c2 Cells. *Journal of Biological Chemistry*. 2006; 281(19):13092–13102. [PubMed: 16549430]
114. Hunter JC, Kostyak JC, Novotny JL, Simpson AM, Korzick DH. Estrogen deficiency decreases ischemic tolerance in the aged rat heart: roles of PKC δ , PKC ϵ , Akt, and GSK3 β . *Am J Physiol Regul Integr Comp Physiol*. 2007; 292(2):R800–R809. [PubMed: 17008461]
115. Mendez P, Garcia-Segura LM. Phosphatidylinositol 3-Kinase and Glycogen Synthase Kinase 3 Regulate Estrogen Receptor-Mediated Transcription in Neuronal Cells. *Endocrinology*. 2006; 147(6):3027–3039. [PubMed: 16497810]
116. Hamilton KL, Gupta S, Knowlton AA. Estrogen and Regulation of Heat Shock Protein Expression in Female Cardiomyocytes: Cross-talk with NF κ B Signaling. *Journal of Molecular and Cellular Cardiology*. 2004; 36(4):579–586.
117. Too CKL, Giles A, Wilkinson M. Estrogen stimulates expression of adenine nucleotide translocator ANT1 messenger RNA in female rat hearts. *Molecular and Cellular Endocrinology*. 1999; 150(1–2):161–167. [PubMed: 10411310]
118. Yu W, Dahl G, Werner R. The Connexin43 Gene is Responsive to Oestrogen. *Proceedings of the Royal Society of London Series B: Biological Sciences*. 1994; 255(1343):125–132. [PubMed: 8165225]
119. Abete P, Calabrese C, Ferrara N, et al. Exercise training restores ischemic preconditioning in the aging heart. *Journal of the American College of Cardiology*. 2000; 36(2):643–650. [PubMed: 10933383]
120. Burns PG, Krukenkamp IB, Caldarone CA, Kirvaitis RJ, Gaudette GR, Levitsky S. Is the preconditioning response conserved in senescent myocardium? *The Annals of Thoracic Surgery*. 1996; 61(3):925–929. [PubMed: 8619719]
121. Isoyama S, Nitta-Komatsubara Y. Acute and chronic adaptation to hemodynamic overload and ischemia in the aged heart. *Heart Failure Reviews*. 2002; 7:63–69. [PubMed: 11790923]
122. Lakatta EG, SolloTT SJ. Perspectives on mammalian cardiovascular aging: humans to molecules. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2002; 132(4):699–721.
123. Liu P, Xu B, Cavalieri TA, Hock CE. Age-related difference in myocardial function and inflammation in a rat model of myocardial ischemia-reperfusion. *Cardiovascular Research*. 2002; 56(3):443–453. [PubMed: 12445885]
124. Mariani, J., Ou, R., Bailey, M., et al. Tolerance to ischemia and hypoxia is reduced in aged human myocardium. 120. 2000. p. 660-667.
125. Pepe S. Dysfunctional ischemic preconditioning mechanisms in aging. *Cardiovascular Research*. 2001; 49(1):11–14. [PubMed: 11121790]

126. Schulman D, Latchman DS, Yellon DM. Effect of aging on the ability of preconditioning to protect rat hearts from ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol*. 2001; 281(4):H1630–H1636. [PubMed: 11557553]
127. Tani M, Honma Y, Hasegawa H, Tamaki K. Direct activation of mitochondrial KATP channels mimics preconditioning but protein kinase C activation is less effective in middle-aged rat hearts. *Cardiovascular Research*. 2001; 49(1):56–68. [PubMed: 11121796]
128. Willems L, Zatta A, Holmgren K, Ashton KJ, Headrick JP. Age-related changes in ischemic tolerance in male and female mouse hearts. *Journal of Molecular and Cellular Cardiology*. 2005; 38(2):245–256. [PubMed: 15698831]
129. Fenton RA, Dickson EW, Dobson J. Inhibition of phosphatase activity enhances preconditioning and limits cell death in the ischemic/reperfused aged rat heart. *Life Sciences*. 2005; 77(26):3375–3388. [PubMed: 16098993]
130. Fenton RA, Dickson EW, Meyer TE, Dobson JG Jr. Aging Reduces the Cardioprotective Effect of Ischemic Preconditioning in the Rat Heart. *Journal of Molecular and Cellular Cardiology*. 2000; 32(7):1371–1375. [PubMed: 10860777]
131. Juhaszova M, Rabel C, Zorov DB, Lakatta EG, Sollott SJ. Protection in the aged heart: preventing the heart-break of old age? *Cardiovascular Research*. 2005; 66(2):233–244. [PubMed: 15820192]
132. Chimenti C, Kajstura J, Torella D, et al. Senescence and Death of Primitive Cells and Myocytes Lead to Premature Cardiac Aging and Heart Failure. *Circulation Research*. 2003; 93(7):604–613. [PubMed: 12958145]
133. Hansford RG. Lipid oxidation by heart mitochondria from young adult and senescent rats. *Biochemistry Journal*. 1978; 170:285–295.
134. Kajstura J, Cheng W, Sarangarajan R, et al. Necrotic and apoptotic myocyte cell death in aging fischer 344 rats. *American Journal of Physiology*. 1996; 271:H1215–H1228. [PubMed: 8853362]
135. Pepe S, Tsuchiya N, Lakatta EG, Hansford RG. PUFA and aging modulate cardiac mitochondrial membrane lipid composition and Ca²⁺ activation of PDH. *American Journal of Physiology - Heart and Circulatory Physiology*. 1999; 276(1):H149–H158.
136. Saeedi R, Wambolt R, Parsons H, et al. Gender and post-ischemic recovery of hypertrophied rat hearts. *BMC Cardiovascular Disorders*. 2006; 6(1):8. [PubMed: 16509993]
137. Lancaster TS, Jefferson SJ, Korzick DH. Local delivery of a PKC ϵ -activating peptide limits ischemia reperfusion injury in the aged female rat heart. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*. 2011; 301(5):R1242–R1249.
138. Cheskis BJ, Greger JG, Nagpal S, Freedman LP. Signaling by estrogens. *Journal of Cellular Physiology*. 2007; 213(3):610–617. [PubMed: 17886255]
139. McDonald TG, Van Eyk JE. Mitochondrial proteomics: undercover in the lipid bilayer. *Basic Res Cardiol*. 2003; 98:219–227. [PubMed: 12835951]
140. Dai DF, Rabinovitch PS, Ungvari Z. Mitochondria and Cardiovascular Aging. *Circulation Research*. 2012; 110(8):1109–1124. [PubMed: 22499901]
141. Dutta D, Calvani R, Bernabei R, Leeuwenburgh C, Marzetti E. Contribution of Impaired Mitochondrial Autophagy to Cardiac Aging: Mechanisms and Therapeutic Opportunities. *Circulation Research*. 2012; 110(8):1125–1138. [PubMed: 22499902]
142. Murphy E. Estrogen Signaling and Cardiovascular Disease. *Circulation Research*. 2011; 109(6):687–696. [PubMed: 21885836]
143. Lancaster TS, Jefferson SJ, Hunter JC, et al. Quantitative proteomic analysis reveals novel mitochondrial targets of estrogen deficiency in the aged female rat heart. *Physiol Genomics*. 2012; 44(20):957–969. [PubMed: 22930739]
144. Craig EE, Hood DA. Influence of aging on protein import into cardiac mitochondria. *American Journal of Physiology - Heart and Circulatory Physiology*. 1997; 272(6):H2983–H2988.
145. Deng N, Zhang J, Zong C, et al. Phosphoproteome Analysis Reveals Regulatory Sites in Major Pathways of Cardiac Mitochondria. *Molecular & Cellular Proteomics*. 2011; 10(2)
146. Knowlton AA, Sun L. Heat shock factor-1, steroid hormones, and regulation of heat shock protein expression in the heart. *American Journal of Physiology*. 2001; 280:H455–H464. [PubMed: 11123263]

147. Voss MR, Stallone JN, Li M, Cornelussen RNM, Knuefermann P, Knowlton AA. Gender Differences in the Expression of Heat Shock Proteins: The Effect of Estrogen. *Am J Physiol Heart Circ Physiol*. 2003; 285:H687–H692. [PubMed: 12714326]
148. Marber MS, Mestrl R, Chi SH, Sayen MR, Yellon DM, Dillmann WH. Overexpression of the rat inducible 70-kD heat stress protein in a transgenic mouse increases the resistance of the heart to ischemic injury. *Journal of Clinical Investigation*. 1995; 95:1854–1860. [PubMed: 7706492]
149. Trost SU, Omens JH, Karlson WJ, et al. Protection Against Myocardial Dysfunction After a Brief Ischemic Period in Transgenic Mice Expressing Inducible Heat Shock Protein 70. *Journal of Clinical Investigation*. 1998; 101(4):855–862. [PubMed: 9466981]
150. Suzuki K, Murtuza B, Sammut IA, et al. Heat Shock Protein 72 Enhances Manganese Superoxide Dismutase Activity During Myocardial Ischemia-Reperfusion Injury, Associated With Mitochondrial Protection and Apoptosis Reduction. *Circulation*. 2002; 106(90121):270I–276.
151. Nakano M, Mann DL, Knowlton AA. Blocking the endogenous increase in HSP72 increases susceptibility to hypoxia and reoxygenation in isolated adult feline cardiocytes. *Circulation*. 1997; 95(6):1523–1531. [PubMed: 9118521]
152. Hamilton KL, Mbai FN, Gupta S, Knowlton AA. Estrogen, Heat Shock Proteins, and NF{ κ }B in Human Vascular Endothelium. *Arterioscler Thromb Vasc Biol*. 2004; 24(9):1628–1633. [PubMed: 15231513]
153. Stice JP, Mbai FN, Chen L, Knowlton AA. Rapid Activation of Nuclear Factor- κ B by 17 β -Estradiol and Selective Estrogen Receptor Modulators: Pathways Mediating Cellular Protection. *Shock*. 2012; 38(2):128–136. [PubMed: 22683727]
154. Locke M. Heat shock transcription factor activation and hsp72 accumulation in aged skeletal muscle. *Cell Stress and Chaperones*. 2000; 5:45–51. [PubMed: 10701839]
155. Locke M, Tanguay RM. Diminished heat shock response in the aged myocardium. *Cell Stress and Chaperones*. 1996; 1:251–260. [PubMed: 9222610]
156. Heydari AR, You S, Takahashi R, Gutschmann-Conrad A, Sarger KD, Richardson A. Age-related alterations in the activation of heat shock transcription factor 1 in rat hepatocytes. *Experimental Cell Research*. 2000; 256:83–93. [PubMed: 10739655]
157. Demirel HA, Hamilton KL, Shanely RA, Tumer N, Koroly MJ, Powers SK. Age and attenuation of exercise-induced myocardial HSP72 accumulation. *American Journal of Physiology - Heart and Circulatory Physiology*. 2003; 285:H1609–H1615. [PubMed: 12791594]
158. Chu B, Soncin F, Price BD, Stevenson MA, Calderwood SK. Sequential Phosphorylation by Mitogen-activated Protein Kinase and Glycogen Synthase Kinase 3 Represses Transcriptional Activation by Heat Shock Factor-1. *Journal of Biological Chemistry*. 1996; 271(48):30847–30857. [PubMed: 8940068]
159. Podlutzky A, Ballabh P, Csiszar A. Oxidative stress and endothelial dysfunction in pulmonary arteries of aged rats. *Am J Physiol Heart Circ Physiol*. 2010; 298(2):H346–H351. [PubMed: 19966046]
160. Ross JL, Howlett SE. Age and Ovariectomy Abolish Beneficial Effects of Female Sex on Rat Ventricular Myocytes Exposed to Simulated Ischemia and Reperfusion. *PLOS One*. 2012; 7(6):e38425. [PubMed: 22701638]
161. Kawasaki T, Chaudry IH. The effects of estrogen on various organs: therapeutic approach for sepsis, trauma, and reperfusion injury. Part 1: central nervous system, lung and heart. *J Anesth*. 2012; 26:883–891. [PubMed: 22729227]
162. Stice JP, Knowlton AA. Estrogen, NF κ B and the Heat Shock Response. *Molecular Medicine*. 2008; 14(7–8):517–527. [PubMed: 18431462]

Highlights

- Cardiac non-genomic and genomic estrogen receptor signaling is reviewed.
- Hormone replacement controversies and cardiac outcome are identified.
- Experimental models of age-associated estrogen loss are described.
- Defective estrogen signaling contributes to cardiac ischemic injury with aging.
- Cardioprotective targets in the aged heart include heat shock proteins and PKC.

Figure 1.

Simplified schematic of cardioprotective cellular signaling mediated by genomic and non-genomic actions of estrogen receptor (ER) activation. Genomic actions mediated by nuclear ERs involve nuclear translocation and activation of protective gene transcription. Non-genomic ER actions include activation of downstream cardioprotective signaling pathways and possible mitochondrial localization. See text for specific details. Abbreviations: BAD, Bcl-2-associated death promoter; diacylglycerol, DAG; eNOS, endothelial nitric oxide synthase; ERE, estrogen response element; ERK, extracellular signal-regulated kinases; GSK-3β, glycogen synthase kinase-3β; HSP90, heat shock90; PI3K, phosphoinositide 3-kinase; PLC, phospholipase C; PKCε, protein kinase Cε; TF, transcription factor. Updated from Stice et al.(162)

Early Effects

Late Effects

Figure 2.

Diagram summarizes interactions of NFκB, estrogen and the heat shock proteins. Treatment of cardiac myocytes or endothelial cells with estrogen leads to activation of NFκB followed by HSF1 activation and increased expression of HSP72. HSPs and HSF1 have feedback loops leading to suppression of inflammation and NFκB. Longer term estrogen treatment inhibits NFκB and increases HSP72 expression in the heart. See text for more detail.

Updated from Stice et al.(162)