Published in final edited form as:

Exp Brain Res. 2017 September ; 235(9): 2883–2892. doi:10.1007/s00221-017-5013-5.

Decreased Spontaneous Activity and Altered Evoked Nociceptive Response of Rat Thalamic Submedial Neurons to Lumbar Vertebra Thrust

William R. Reed^{1,†}, Jamie T. Cranston¹, Stephen M. Onifer¹, Joshua W. Little², and Randall S. Sozio¹

¹Palmer Center for Chiropractic Research, Palmer College of Chiropractic, Davenport, IA, USA

²Center for Anatomical Science and Education, Department of Surgery, Saint Louis University School of Medicine, St. Louis, MO, USA

Abstract

Objectives—The thalamus is a central structure important to modulating and processing all mechanoreceptor input destined for the cortex. A large number of diverse mechanoreceptor endings are stimulated when a high velocity low amplitude thrust is delivered to the lumbar spine during spinal manipulation. The objective of this study was to determine if a lumbar thrust alters spontaneous and/or evoked nociceptive activity in medial thalamic submedial (Sm) neurons.

Methods—Extracellular recordings were obtained from 94 thalamic Sm neurons in 54 urethane-anesthetized adult Wistar rats. Spontaneous activity was recorded 5min before and after an L₅ control (no thrust) and thrust (85% rat body weight; 100ms) procedure. In a subset of responsive nociceptive specific neurons, mean changes in noxious evoked response (10s pinch with clip; 795g) at 3 sites (tail, contra- and ipsilateral hindpaw) were determined following an L₅ thrust. Mean changes in Sm spontaneous activity (60s bins) and evoked noxious response were compared using a mixed model repeated measures ANOVA with Bonferroni *post hoc* t-tests and paired t-tests, respectively.

Results—Compared to control, spontaneous Sm activity decreased 180–240s following the lumbar thrust ($p < 0.005$). Inhibitory evoked responses were attenuated in the contralateral hindpaw following an L₅ thrust compared to control ($p < 0.05$). No other changes in spontaneous or noxious evoked Sm activity were found.

Conclusions—A delayed, but prolonged suppression of spontaneous Sm activity along with changes in noxious evoked inhibitory responses in the contralateral hindpaw following lumbar vertebra thrust suggest that thalamic submedial neurons may play a role in central pain modulation related to manual therapy intervention.

Correspondence to: William R. Reed.

[†]Present Address: School of Health Professions, Department of Physical Therapy, UAB | The University of Alabama at Birmingham, Webb 318 | 1720 2nd Avenue South | Birmingham, AL 35294-1212, P: 205.934.3261 | wreed@uab.edu

Ethical Approval: All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted.

Keywords

thalamus; submedialis; neurophysiology; spinal manipulation; lumbar spine; pain

Introduction

The thalamus is a central structure consisting of a large collection of individual subnuclei that process and modulate all mechanoreceptor/nociceptor input destined for the cortex (Jones 2007; Yen and Lu 2013). The thalamic submedialis (Sm) nucleus is medially located and is frequently divided into rostral and caudal parts (Craig and Burton 1981; Dado and Giesler 1990; Yoshida et al. 1991; Yoshida et al. 1992). The Sm receives afferent projections from the lateral hypothalamus, orbital cortex, trigeminal sensory nucleus, medial parabrachial nucleus, raphe nuclei, and bilateral dorsal horn neurons throughout the entire spinal cord. Spinal projections primarily terminate in the rostral half of Sm in both the cat and rat, while spinal trigeminal nuclei projections terminate in the caudal half. (Craig and Burton 1981; Craig et al. 1982; Dado and Giesler 1990; Blomqvist et al. 1992; Yoshida et al. 1992). In the cat, spinal projections arise almost exclusively from the marginal zone (lamina I), whereas in the rat the trigeminal projections arise from lamina I while the vast majority of spinal projections arise from the deep dorsal horn and intermediate zone/ventral horn (lamina V–VII) (Dado and Giesler 1990; Yoshida et al. 1991; Yoshida et al. 1992). Electrical stimulation, lesions, and pharmacological microinjections of the Sm nucleus have been shown to modulate pain response in animals. (Roberts and Dong 1994; Zhang et al. 1995; Zhang et al. 1998; Zhang et al. 1999; Xiao et al. 2005; Feng et al. 2008; Wang et al. 2008; Erfanparast et al. 2015). Evidence indicates that the Sm nucleus is involved in opioid-receptor mediated nociception modulation (Wang et al. 2008; Tang et al. 2009; Erfanparast et al. 2015), and as part of this nociceptive modulatory pathway, Sm efferent fibers terminate most heavily in the ipsilateral ventrolateral orbital cortex (VLO), lateral orbital cortex, and to a lesser extent in the insular cortex (Craig and Burton 1981; Yoshida et al. 1992; Tang et al. 2009). In addition to reciprocal connections between the Sm and VLO, VLO neurons also project to the midbrain periaqueductal gray (PAG), (Hardy and Leichnetz 1981; Craig et al. 1982) a region extensively implicated in descending inhibition of nociceptive inputs at the spinal cord level. The submedialis nucleus subserves a number of physiological functions, but afferent and efferent connections along with behavioral and electrophysiological studies support a direct involvement of the Sm in mechanical/thermal nociceptive modulation.

Spinal manipulation is a non-pharmacological integrative healthcare intervention that is recommended by clinical guidelines and evidence reports for treating certain types of neck and low back pain (Chou et al. 2007; Bronfort et al. 2010; Chou et al. 2017). However, its efficacy and appropriate utilization is hindered due to the lack of current knowledge regarding its underlying neurophysiological mechanisms. Lumbar spinal manipulation is a mechanical stimulus that typically involves applying a single, short lever, high velocity, low amplitude posterior-to-anterior thrust of short duration (< 150ms) to a particular vertebra equating to 30–130% body weight of a 70kg individual (Herzog et al. 1993; Triano 2001; Bergmann 2005; Herzog 2010). Similar biomechanical thrust characteristics have been simulated and scaled for use in various animal models (Pickar 1999; Reed et al. 2013; Reed

et al. 2014a). Clinically, mechanical and thermal pain thresholds are increased (i.e. decrease tissue sensitivity) within 5 minutes of the manipulative thrust in areas local and remote to those being treated in both symptomatic and asymptomatic individuals (George et al. 2006; Bialosky et al. 2009b; Coronado et al. 2012; Bialosky et al. 2014). This widespread hypoalgesia has been attributed to alterations in central pain processing (Wright 1995; Boal and Gillette 2004; Bialosky et al. 2009a), but the central pathways, structures, and mechanisms responsible remain undefined and thereby provides the underlying scientific rationale for the current study.

In the rat, submedius neurons typically have large (often bilateral) receptive fields and respond predominately to noxious convergent sensory input from spatially separated cutaneous, muscle, joint, and visceral tissues which suggests a non-discriminatory role in nociception (Miletic and Coffield 1989; Dostrovsky and Guilbaud 1990; Fu et al. 2002). These attributes are particularly relevant to manual therapy because spinal manipulation has been hypothesized to stimulate as many as 40 types of peripheral receptors in both cutaneous and deep spinal tissues (Gillette 1987). Taken together, these neuronal characteristics make Sm nociceptive-specific neurons good candidates for investigating supraspinal neurophysiological changes related to lumbar vertebra thrusts. The purpose of this study was to determine if adult rat lumbar vertebrae high velocity low amplitude thrusts alter spontaneous and/or evoked noxious thalamic Sm activity.

Materials and Methods

All experimental methods were approved by the Institutional Animal Care and Use Committee. Animals were housed in pairs and exposed to a 12-hour light/dark cycle, environmental enrichment with food and water ad libitum. For electrophysiological recordings, 54 adult male Wistar rats (320–524g; Envigo, Indianapolis, IN) were initially anesthetized with isoflurane (2.5%) followed by an intraperitoneal injection of 50% urethane (1.2g/kg). The jugular vein and carotid artery was catheterized for intravenous (iv) infusion of fluids and blood pressure monitoring. The trachea was intubated for pCO₂ monitoring. Saturated oxygen concentration, heart rate, and respiration were monitored by a MouseOx system (Starr Life Sciences Corp., Oakmont, PA). Body temperature was monitored via a rectal thermometer and maintained at 37°C with a water circulating heat pad. To maintain an anesthetic state III-3 (Friedberg et al. 1999) throughout the duration of the experiment, depth of anesthesia was regularly assessed by monitoring hindpaw pinch withdrawal, corneal reflex, heart rate, respiration rate and vibrissae movements. Supplemental anesthesia (5% urethane; iv) was administered as indicated (Hubscher and Johnson 2003; Reed et al. 2009).

Electrophysiology

The rat's head was secured in a stereotaxic device (Kopf Instruments, Tujunga, CA) with its dorsal surface positioned horizontally. A small hole was made bilaterally in the skull just lateral to midline and then expanded carefully with a small bone rongeurs to avoid rupture of the superior sagittal sinus. The dura was opened and the recording electrode was advanced incrementally into the thalamic submedius nucleus. Warm mineral oil was used to prevent tissue desiccation. Activity of thalamic Sm neurons was recorded extracellularly with DiI-

(1,1'-diocetabecyl-3,3,3',3'-tetramethyl-indocarbocyanine perchlorate; Invitrogen, Carlsbad, CA) coated tungsten microelectrodes (FHC, Bowdoin, ME) having 6 to 8 M Ω impedance as previously described (Massey et al. 2006; Chadha and Hubscher 2008; Reed et al. 2009; Reed et al. 2014a). Thalamic submedial nuclei search coordinates were between -2.3 and -2.6 mm caudal to bregma, 0.5 and 0.9 mm lateral to midline, and 6.0 – 6.9 mm below the cortex surface (Fig. 1A) (Kawakita et al. 1993; Sumiya and Kawakita 1997; Yang and Follett 2003). The electrode was slowly advanced at a rate of 1 – 5 μ m per step using a motorized micromanipulator (Neurostar, Germany) (Fig. 1B) until spontaneous isolated single unit Sm activity was identified. Signals from single thalamic Sm neurons were passed through a high impedance probe (HIP511, Grass, West Warwick, RI), amplified (P511K, Grass), recorded and evaluated off-line using a PC based data acquisition system (Spike 2, Cambridge Electronic Design, UK). Neuron receptive fields were mapped and recorded. All Sm neurons included in this study were classified as nociceptive specific (high threshold) neurons due to a lack of response to low threshold stimulation (innocuous stroking with nylon brush). While not exclusive, nociceptive specific neurons are the predominate type of Sm neuron found in the normal rats. (Dostrovsky and Guilbaud 1988; Miletic and Coffield 1989; Kawakita et al. 1993).

Spontaneous Activity

The effect of high velocity low amplitude lumbar thrusts on spontaneous activity was determined as follows. Once single unit thalamic Sm activity was isolated, a timed-control protocol was performed. This consisted of recording spontaneous neural activity for 5 min before and after a no thrust procedure. Afterwards, spontaneous activity was again recorded for 5 min before and after a lumbar thrust protocol (peak amplitude of 85% rat body weight, 100ms duration) (Reed et al. 2014a; Reed et al. 2014b). Neural activity was quantified as impulses per period of time (10s or 60s intervals). A computer controlled electronic feedback system (Lever System Model 310; Aurora Scientific, Ontario, Canada) (Pickar 1999) was used to deliver a linearly increasing dorsal-ventral thrust force at the L₅ vertebra via toothed forceps attached to the spinous process (Fig. 1B). Prominent rat bony landmarks (ridge of iliac crests and large L₆ spinous process) were used to identify the L₅ spinous process for forceps attachment. The scaled L₅ thrust profile simulates the thrust magnitude, thrust duration and vertebrae displacement (0.5 – 1.5 mm) of that delivered in clinical settings (Herzog et al. 1993; Nathan and Keller 1994; Gal et al. 1997; Triano 2001; Ianuzzi and Khalsa 2005) albeit via forceps attached directly to spinous process so as to directly determine the amount of vertebra displacement. For spontaneous activity recordings, the control (no thrust) protocol always preceded the lumbar thrust protocol so as to prevent any potential long-lasting influence related to the thrust.

Noxious Evoked Response

To determine the effects of high velocity low amplitude lumbar thrusts on evoked Sm noxious response, a 10s noxious pinch stimulus (small clip; 795g; Roboz-RS 5452, Gaithersburg, MD) was applied in random order to three sites (tip of the tail, contra- and ipsilateral hindpaws) prior to- and following control and thrust procedures. To minimize the risk of potential peripheral sensitization, responses to evoked noxious stimulation (at three sites) was performed on one Sm neuron per hemisphere. An example of a thrust protocol

neural recording is shown in Figure 2. The order of control and thrust protocols for noxious evoked Sm activity were randomized to minimize any possible ordering effects in addition to a waiting period of 5 minutes between test protocols. The 10s duration of noxious pinch stimulus has been used in previous studies of thalamic neuron responsiveness (Zhao et al. 2006; Fischer et al. 2009). Spontaneous (resting) activity (10s) immediately preceding each pinch stimulus was subtracted from the 10s evoked responses to yield a net discharge for the noxious stimulus. Changes in evoked Sm response were determined by subtracting the net discharge of 1st pinch stimulus from the net discharge of the 2nd pinch stimulus.

Histology

Following experimental protocol completion, electrolytic lesions were made by applying 30 μ A current for 30s. Afterwards, transcardiac perfusions were performed with an oxygenated calcium-free Tyrodes buffer solution followed by 4% paraformaldehyde in 0.1M phosphate buffer, pH 7.4 (Onifer et al. 2005). The brains were removed and post-fixed overnight at 4° C, and then transferred to 30% sucrose at 4° C for a minimum of 36hrs prior to coronal tissue sectioning (30 μ m) using a cryostat. DiI-labeled electrode tracks were located using a Nikon microscope equipped with fluorescent filters. Select sections were then stained with cresyl violet and postmortem histological reconstructions were performed using a combination of stereotaxic coordinates records, DiI-labeled electrode track measurements (Hubscher 2006; Massey et al. 2006; Reed et al. 2009) and electrolytic lesions (Fig. 3). Neurons identified as being outside the confines of the Sm nucleus were excluded.

Statistical Analysis

Changes in mean spontaneous responses (impulses/time bin) from control and L₅ thrust protocols were compared using a mixed model repeated measures ANOVA followed by Bonferroni *post hoc* t-tests (IBM SPSS v22, Armonk, NY). For spontaneous activity analyses, the 5 min pre- and post-procedure recording periods were subdivided into five 60s bins (60, 120, 180, 240, 300s). To evaluate the possibility of a more immediate change in spontaneous activity due to lumbar thrust, changes in neural activity for the 10s immediately following each procedure were also compared. Outlier values ($\pm$ 3 standard deviations from the mean) were removed to prevent the undue influence of atypical thalamic neural activity (prolonged bursting or silence) during the 10 min protocol recording period. Outlier removal resulted in spontaneous activity from 86–91 Sm neurons being compared per time interval. For evoked response analysis, 55/94 Sm neurons were tested (noxious testing was limited to one neuron per thalamic hemisphere). Only Sm neurons exhibiting greater than 20% increase over initial pre-pinch baseline activity (10s) at individual test sites were considered to be responsive to noxious stimulation. This criteria ensured that the stimuli was applied within the neuron's receptive field and resulted in the analysis of 35–43 neurons at a given site for single factor paired t-tests comparisons. Data shown are mean $\pm$ SEM unless otherwise noted and statistical significance was set at $p < 0.05$.

Results

While the entire rostral-caudal extent of the submedius nucleus was searched, the majority (68%) of the 94 single unit recordings were located in the rostral half of the nucleus having a mean depth of 6.41mm (± 0.28 SD) from the cortical surface. All Sm neurons were classified as nociceptive-specific (high threshold) by their lack of response to innocuous stroking (brush) and exhibited an excitatory response to noxious pinch (forceps). Lumbar thrust displacement-time profiles mirrored those of the “inverted V” shape of clinical force-time profiles (Herzog et al. 1993; Triano 2001) with a mean L₅ displacement of 1.02mm (± 0.47 mm SD).

Spontaneous activity

All recorded Sm neurons were spontaneously active to some degree. Prior to peripheral stimulation, twenty two (23%) neurons had a low firing rate (< 1 Hz), 69 (73%) a moderate rate (1–10Hz) and 3 (3%) a high firing rate (> 10 Hz). Compared to the control procedure, there were no significant changes in mean Sm spontaneous activity during the first 180s following the L₅ vertebra thrust. However, spontaneous Sm activity was significantly decreased 180–240s after the thrust (ANOVA: $F = 4.8$, $df = 1,5$, $p < 0.05$, *post hoc*: $p < 0.005$) (Fig. 4, Table 1). Decreased spontaneous Sm activity continued into the subsequent 60s bin (from 240–300s) but failed to reach significance (Fig. 4).

Evoked noxious response

The mean change in Sm neuron response to a 10s noxious stimulus applied before and after control and lumbar thrust was determined. Compared to control, inhibitory responses to noxious contralateral hindpaw stimulation were significantly attenuated after the L₅ thrust (Fig. 5, Table 2). Mixed neural responses were seen with noxious stimulation of the tail (inhibitory) and ipsilateral hindpaw (excitatory) following the L₅ thrust but these changes did not reach significance (Fig. 5). Figure 6 provides an example of a Sm neuron exhibiting a mixed response to tail and ipsilateral hindpaw stimulation after an L₅ thrust. It is possible that random bursts and/or time-related changes in resting baseline activity could have contributed to a mixed response (since the 10s of resting baseline activity immediately preceding the pinch stimulus was subtracted from the evoked noxious response); however, no significant differences were found in pre-post mean baseline activity between the control and thrust protocols at these test sites.

Discussion

The vast majority of nociceptive specific Sm neurons in the current study had large (bilateral) receptive fields and exhibited a low to moderate spontaneous activity firing rate. This is in agreement with previous findings from a study involving 204 Sm neurons and urethane anesthesia (Kawakita et al. 1993). In this particular study, authors divided the Sm nucleus into 4 sub-regions and reported no significant differences in spontaneous activity between subregions nor between neuron responses recorded early or late in the experiment (Kawakita et al. 1993). We found that dynamic high velocity low amplitude thrusts similar to those applied in clinically delivered lumbar spinal manipulation produces a delayed and

prolonged suppression of thalamic nociceptive specific Sm neuron spontaneous activity. The physiological impact of this delayed but prolonged decrease of Sm activity is not currently known. Since widespread increases in thalamic spontaneous activity has been shown to occur in both acute and chronic pain conditions (Rinaldi et al. 1991; Hains et al. 2005; Fischer et al. 2009; Masri et al. 2009; Iwata et al. 2011) suppression of thalamic activity may play a role in spinal manipulation's clinical efficacy. Sustained changes in thalamic resting activity may in turn have neurophysiological consequences at the VLO or other efferent Sm connections as well. It is interesting to note that the delayed timing of this suppression of medial thalamic activity, ~3 min after the thrust delivery, coincides with the timing (<5min) of reported mechanical/thermal hypoalgesic effects following clinically delivered spinal manipulative thrusts (George et al. 2006; Fernandez-de-las-Penas et al. 2007; Fernandez-Carnero et al. 2008; Bishop et al. 2011; de Camargo et al. 2011; Srbely et al. 2013). Decreased neural activation in other central structures following manual therapy intervention has been reported in both animal and human neuroimaging studies (Malisza et al. 2003a; Malisza et al. 2003b; Sparks et al. 2013). Decreased activity at thalamic and/or other key spinal/supraspinal pain modulatory centers may serve as a key neurobiological mechanism of spinal manipulation.

Studies using bilateral Sm electrolytic lesions indicate that Sm neurons are involved in supraspinal and/or spinal mediated inhibition of nociceptive input (Roberts and Dong 1994; Zhang et al. 1995). We observed a decrease of Sm neuron activity evoked by noxious contralateral hindpaw stimulation after the control procedure. This inhibitory Sm activity was attenuated after the L₅ thrust which suggests that lumbar thrusts may act to modulate nociceptive activity in higher pain centers. Despite Sm neurons typically having large (often bilateral) receptive fields, the contralateral hindpaw was the only testing site to demonstrate a significant change following L₅ thrust. This is likely due to a greater preponderance (64%) of anatomically crossed ascending spinal projections to the Sm in the rat (Dado and Giesler 1990). While the changes did not reach significance, the L₅ thrust decreased Sm neuron response to tail pinch and increased Sm neuron response to ipsilateral hindpaw pinch. Sm neurons exhibiting inhibitory responses to noxious stimulation at one anatomical site and excitatory responses at another has been previously reported (Kawakita et al. 1993). These alternating nociceptive responses seen after L₅ thrust may be attributable to anatomical differences in peripheral nociceptor density, nociceptor sensitivity, and/or to the anatomical proximity of L₅ dorsal root ganglion cells to the lumbar thrust site compared to the sacral (S₁-S₄) dorsal root ganglion cells innervating the tail (Danneman et al. 1994; MacKenzie et al. 2015).

Despite both local and remote somatic hypoalgesic effects being associated with clinical spinal manipulative thrusts; the peripheral and/or central pathways, structures and mechanisms responsible remain unknown (Wright 1999; Bialosky et al. 2009a). Basic science (Vaillant et al. 2012; Reed et al. 2013; Reed et al. 2014b; Onifer et al. 2015; Reed et al. 2015a; Reed et al. 2015b; Song et al. 2016) and clinical (Bishop et al. 2011; Nougrou et al. 2013; Sparks et al. 2013; Bialosky et al. 2014; Nougrou et al. 2014; Page et al. 2014) studies have begun to focus much greater attention on potential physiological mechanisms associated with non-pharmacological manual therapy interventions. A functional magnetic resonance imaging study investigating supraspinal activation related to noxious mechanical

stimulation pre-and post-thrust spinal manipulation reported that while bilateral activation of the thalami, cerebellum, amygdala, periaqueductal gray, insular cortex, anterior cingulate cortex, somatosensory cortices, supplementary motor area, and premotor areas occurred with peripheral noxious stimulation; only the insular cortex demonstrated a significant relationship between pain-reduction and post-spinal manipulation (Sparks et al. 2013). While this small (n=10) clinical imaging study was unable to draw any definitive conclusions, it is interesting to note that the insula receives direct projections from medial thalamic nuclei (including the Sm) and that these nuclei are thought to be more involved with the affective and motivational aspects of pain which may in turn contribute to the efficacy of spinal manipulation and other integrative medicine interventions (Bushnell and Duncan 1989; Basbaum 2000; Jones 2007; Bialosky et al. 2009a).

Conclusion

To our knowledge, this study is the first to investigate the effects of lumbar vertebra high velocity low amplitude thrusts on central spontaneous neural activity and noxiously evoked responses in the medial thalamus. The results indicate that lumbar vertebra thrusts cause a delayed and prolonged decrease in nociceptive specific Sm neuron spontaneous activity. In addition, inhibitory responses to noxious contralateral hindpaw stimulation are attenuated following L₅ thrust. Additional acute and chronic pain studies need to be performed in the Sm as well as other thalamic subnuclei to determine if these neural changes occur more globally within the thalamus and/or impact nociceptive somatosensory processing at higher central centers.

Acknowledgments

This work was supported by grants from the Australian Spinal Research Foundation (LG2010-11) and NIH National Center for Complementary and Integrated Health (K01AT005935) to WRR and was conducted in a facility with support from NIH National Center for Research Resources under Research Facilities Improvement Grant Number C06RR15433. We thank Darlene Burke (University of Louisville-KSCIRC) of the Neuroscience Core for statistical analyses support.

References

- Basbaum, AJT. The Perception of Pain. In: Kandel, ERSJ., Jessell, TM., editors. Principles of Neural Science. McGraw-Hill; New York: 2000. p. 472-491.
- Bergmann, TF. High-velocity low amplitude manipulative techniques. In: Haldeman, S.Dagenais, S.Budgell, B.Grunnet-Nilsson, P.Hooper, W.Meeker, W., et al., editors. Principles and Practice of Chiropractic. McGraw-Hill; New York, NY, USA: 2005. p. 755-766.
- Bialosky JE, Bishop MD, Price DD, Robinson ME, George SZ. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man Ther.* 2009a; 14:531–538. DOI: 10.1016/j.math.2008.09.001 [PubMed: 19027342]
- Bialosky JE, Bishop MD, Robinson ME, Zeppieri G Jr, George SZ. Spinal manipulative therapy has an immediate effect on thermal pain sensitivity in people with low back pain: a randomized controlled trial. *Phys Ther.* 2009b; 89:1292–1303. DOI: 10.2522/ptj.20090058 [PubMed: 19797305]
- Bialosky JE, George SZ, Horn ME, Price DD, Staud R, Robinson ME. Spinal manipulative therapy-specific changes in pain sensitivity in individuals with low back pain (NCT01168999). *J Pain.* 2014; 15:136–148. DOI: 10.1016/j.jpain.2013.10.005 [PubMed: 24361109]
- Bishop MD, Beneciuk JM, George SZ. Immediate reduction in temporal sensory summation after thoracic spinal manipulation. *Spine J.* 2011; 11:440–446. DOI: 10.1016/j.spinee.2011.03.001 [PubMed: 21463970]

- Blomqvist A, Ericson AC, Broman J, Craig AD. Electron microscopic identification of lamina I axon terminations in the nucleus submedialis of the cat thalamus. *Brain Res.* 1992; 585:425–430. [PubMed: 1511331]
- Boal RW, Gillette RG. Central neuronal plasticity, low back pain and spinal manipulative therapy. *J Manipulative Physiol Ther.* 2004; 27:314–326. DOI: 10.1016/j.jmpt.2004.04.005 [PubMed: 15195039]
- Bronfort G, Haas M, Evans R, Leininger B, Triano J. Effectiveness of manual therapies: the UK evidence report. *Chiropr Osteopat.* 2010; 18:3. doi: 10.1186/1746-1340-18-3 [PubMed: 20184717]
- Bushnell MC, Duncan GH. Sensory and affective aspects of pain perception: is medial thalamus restricted to emotional issues? *Exp Brain Res.* 1989; 78:415–418. [PubMed: 2599050]
- Chadha HK, Hubscher CH. Convergence of nociceptive information in the forebrain of female rats: reproductive organ response variations with stage of estrus. *Exp Neurol.* 2008; 210:375–387. DOI: 10.1016/j.expneurol.2007.11.010 [PubMed: 18096159]
- Chou R, Deyo R, Friedly J, et al. Nonpharmacologic Therapies for Low Back Pain: A Systematic Review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med.* 2017; doi: 10.7326/m16-2459
- Chou R, Qaseem A, Snow V, Casey D, Cross JT Jr, Shekelle P, Owens DK. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med.* 2007; 147:478–491. [PubMed: 17909209]
- Coronado RA, Gay CW, Bialosky JE, Carnaby GD, Bishop MD, George SZ. Changes in pain sensitivity following spinal manipulation: a systematic review and meta-analysis. *J Electromyogr Kinesiol.* 2012; 22:752–767. DOI: 10.1016/j.jelekin.2011.12.013 [PubMed: 22296867]
- Craig AD Jr, Burton H. Spinal and medullary lamina I projection to nucleus submedialis in medial thalamus: a possible pain center. *J Neurophysiol.* 1981; 45:443–466. [PubMed: 6783739]
- Craig AD Jr, Wiegand SJ, Price JL. The thalamo-cortical projection of the nucleus submedialis in the cat. *J Comp Neurol.* 1982; 206:28–48. DOI: 10.1002/cne.902060105 [PubMed: 6284805]
- Dado RJ, Giesler GJ Jr. Afferent input to nucleus submedialis in rats: retrograde labeling of neurons in the spinal cord and caudal medulla. *J Neurosci.* 1990; 10:2672–2686. [PubMed: 2388083]
- Danneman PJ, Kiritsy-Roy JA, Morrow TJ, Casey KL. Central delay of the laser-activated rat tail-flick reflex. *Pain.* 1994; 58:39–44. [PubMed: 7970838]
- de Camargo VM, Albuquerque-Sendin F, Berzin F, Stefanelli VC, de Souza DP, Fernandez-de-las-Penas C. Immediate effects on electromyographic activity and pressure pain thresholds after a cervical manipulation in mechanical neck pain: a randomized controlled trial. *J Manipulative Physiol Ther.* 2011; 34:211–220. DOI: 10.1016/j.jmpt.2011.02.002 [PubMed: 21621722]
- Dostrovsky JO, Guilbaud G. Noxious stimuli excite neurons in nucleus submedialis of the normal and arthritic rat. *Brain Res.* 1988; 460:269–280. [PubMed: 3224262]
- Dostrovsky JO, Guilbaud G. Nociceptive responses in medial thalamus of the normal and arthritic rat. *Pain.* 1990; 40:93–104. [PubMed: 2339022]
- Erfanparast A, Tamaddonfard E, Taati M, Dabaghi M. Role of the thalamic submedialis nucleus histamine H1 and H2 and opioid receptors in modulation of formalin-induced orofacial pain in rats. *Naunyn-Schmiedeberg's Arch Pharmacol.* 2015; 388:1089–1096. DOI: 10.1007/s00210-015-1143-0 [PubMed: 26077661]
- Feng J, Jia N, Han LN, Huang FS, Xie YF, Liu J, Tang JS. Microinjection of morphine into thalamic nucleus submedialis depresses bee venom-induced inflammatory pain in the rat. *J Pharm Pharmacol.* 2008; 60:1355–1363. DOI: 10.1211/jpp/60.10.0012 [PubMed: 18812029]
- Fernandez-Carnero J, Fernandez-de-las-Penas C, Cleland JA. Immediate hypoalgesic and motor effects after a single cervical spine manipulation in subjects with lateral epicondylalgia. *J Manipulative Physiol Ther.* 2008; 31:675–681. DOI: 10.1016/j.jmpt.2008.10.005 [PubMed: 19028251]
- Fernandez-de-las-Penas C, Perez-de-Heredia M, Brea-Rivero M, Miangolarra-Page JC. Immediate effects on pressure pain threshold following a single cervical spine manipulation in healthy subjects. *J Orthop Sports Phys Ther.* 2007; 37:325–329. DOI: 10.2519/jospt.2007.2542 [PubMed: 17612359]

- Fischer TZ, Tan AM, Waxman SG. Thalamic neuron hyperexcitability and enlarged receptive fields in the STZ model of diabetic pain. *Brain Res.* 2009; 1268:154–161. DOI: 10.1016/j.brainres.2009.02.063 [PubMed: 19285053]
- Friedberg MH, Lee SM, Ebner FF. Modulation of receptive field properties of thalamic somatosensory neurons by the depth of anesthesia. *J Neurophysiol.* 1999; 81:2243–2252. [PubMed: 10322063]
- Fu JJ, Tang JS, Yuan B, Jia H. Response of neurons in the thalamic nucleus submedialis (Sm) to noxious stimulation and electrophysiological identification of on- and off-cells in rats. *Pain.* 2002; 99:243–251. [PubMed: 12237202]
- Gal J, Herzog W, Kawchuk G, Conway PJ, Zhang YT. Movements of vertebrae during manipulative thrusts to unembalmed human cadavers. *J Manipulative Physiol Ther.* 1997; 20:30–40. [PubMed: 9004120]
- George SZ, Bishop MD, Bialosky JE, Zeppieri G Jr, Robinson ME. Immediate effects of spinal manipulation on thermal pain sensitivity: an experimental study. *BMC Musculoskelet Disord.* 2006; 7:68.doi: 10.1186/1471-2474-7-68 [PubMed: 16911795]
- Gillette RG. A speculative argument for the co-activation of diverse somatic receptor populations by forceful chiropractic adjustments. *Manual Medicine.* 1987; 3:1–14.
- Hains BC, Saab CY, Waxman SG. Changes in electrophysiological properties and sodium channel Nav1.3 expression in thalamic neurons after spinal cord injury. *Brain.* 2005; 128:2359–2371. DOI: 10.1093/brain/awh623 [PubMed: 16109750]
- Hardy SG, Leichnetz GR. Frontal cortical projections to the periaqueductal gray in the rat: a retrograde and orthograde horseradish peroxidase study. *Neurosci Lett.* 1981; 23:13–17. [PubMed: 6164964]
- Herzog W. The biomechanics of spinal manipulation. *J Bodyw Mov Ther.* 2010; 14:280–286. DOI: 10.1016/j.jbmt.2010.03.004 [PubMed: 20538226]
- Herzog W, Conway PJ, Kawchuk GN, Zhang Y, Hasler EM. Forces exerted during spinal manipulative therapy. *Spine (Phila Pa 1976).* 1993; 18:1206–1212. [PubMed: 8362328]
- Hubscher CH. Estradiol-associated variation in responses of rostral medullary neurons to somatovisceral stimulation. *Exp Neurol.* 2006; 200:227–239. DOI: 10.1016/j.expneurol.2006.02.008 [PubMed: 16624305]
- Hubscher CH, Johnson RD. Responses of thalamic neurons to input from the male genitalia. *J Neurophysiol.* 2003; 89:2–11. DOI: 10.1152/jn.00294.2002 [PubMed: 12522154]
- Ianuzzi A, Khalsa PS. High loading rate during spinal manipulation produces unique facet joint capsule strain patterns compared with axial rotations. *J Manipulative Physiol Ther.* 2005; 28:673–687. DOI: 10.1016/j.jmpt.2005.08.020 [PubMed: 16326237]
- Iwata M, LeBlanc BW, Kadasi LM, Zerach ML, Cosgrove RG, Saab CY. High-frequency stimulation in the ventral posterolateral thalamus reverses electrophysiologic changes and hyperalgesia in a rat model of peripheral neuropathic pain. *Pain.* 2011; 152:2505–2513. DOI: 10.1016/j.pain.2011.07.011 [PubMed: 21906880]
- Jones, EG. *The Thalamus.* Cambridge University Press; New York: 2007.
- Kawakita K, Dostrovsky JO, Tang JS, Chiang CY. Responses of neurons in the rat thalamic nucleus submedialis to cutaneous, muscle and visceral nociceptive stimuli. *Pain.* 1993; 55:327–338. [PubMed: 8121694]
- MacKenzie SJ, Yi JL, Singla A, Russell TM, Calancie B. Innervation and function of rat tail muscles for modeling cauda equina injury and repair. *Muscle Nerve.* 2015; 52:94–102. DOI: 10.1002/mus.24498 [PubMed: 25346299]
- Malisza KL, Gregorash L, Turner A, et al. Functional MRI involving painful stimulation of the ankle and the effect of physiotherapy joint mobilization. *Magn Reson Imaging.* 2003a; 21:489–496. [PubMed: 12878258]
- Malisza KL, Stroman PW, Turner A, Gregorash L, Foniok T, Wright A. Functional MRI of the rat lumbar spinal cord involving painful stimulation and the effect of peripheral joint mobilization. *J Magn Reson Imaging.* 2003b; 18:152–159. DOI: 10.1002/jmri.10339 [PubMed: 12884326]
- Masri R, Quiton RL, Lucas JM, Murray PD, Thompson SM, Keller A. Zona incerta: a role in central pain. *J Neurophysiol.* 2009; 102:181–191. DOI: 10.1152/jn.00152.2009 [PubMed: 19403748]
- Massey JM, Hubscher CH, Wagoner MR, Decker JA, Amps J, Silver J, Onifer SM. Chondroitinase ABC digestion of the perineuronal net promotes functional collateral sprouting in the cuneate

- nucleus after cervical spinal cord injury. *J Neurosci*. 2006; 26:4406–4414. DOI: 10.1523/jneurosci.5467-05.2006 [PubMed: 16624960]
- Miletic V, Coffield JA. Responses of neurons in the rat nucleus submedius to noxious and innocuous mechanical cutaneous stimulation. *Somatosens Mot Res*. 1989; 6:567–587. [PubMed: 2816204]
- Nathan M, Keller TS. Measurement and analysis of the in vivo posteroanterior impulse response of the human thoracolumbar spine: a feasibility study. *J Manipulative Physiol Ther*. 1994; 17:431–441. [PubMed: 7989876]
- Nougarou F, Dugas C, Deslauriers C, Page I, Descarreaux M. Physiological responses to spinal manipulation therapy: investigation of the relationship between electromyographic responses and peak force. *J Manipulative Physiol Ther*. 2013; 36:557–563. DOI: 10.1016/j.jmpt.2013.08.006 [PubMed: 24161387]
- Nougarou F, Dugas C, Loranger M, Page I, Descarreaux M. The role of preload forces in spinal manipulation: experimental investigation of kinematic and electromyographic responses in healthy adults. *J Manipulative Physiol Ther*. 2014; 37:287–293. DOI: 10.1016/j.jmpt.2014.04.002 [PubMed: 24928637]
- Onifer SM, Reed WR, Sozio RS, Long CR. Antinociceptive Effects of Spinal Manipulative Therapy on Nociceptive Behavior of Adult Rats during the Formalin Test. *Evid Based Complement Alternat Med*. 2015; 2015:520454. doi: 10.1155/2015/520454 [PubMed: 26693243]
- Onifer SM, Zhang YP, Burke DA, et al. Adult rat forelimb dysfunction after dorsal cervical spinal cord injury. *Exp Neurol*. 2005; 192:25–38. DOI: 10.1016/j.expneurol.2004.10.016 [PubMed: 15698616]
- Page I, Nougarou F, Dugas C, Descarreaux M. The effect of spinal manipulation impulse duration on spine neuromechanical responses. *J Can Chiropr Assoc*. 2014; 58:141–148. [PubMed: 24932018]
- Pickar JG. An in vivo preparation for investigating neural responses to controlled loading of a lumbar vertebra in the anesthetized cat. *J Neurosci Methods*. 1999; 89:87–96. [PubMed: 10491938]
- Reed WR, Cao DY, Long CR, Kawchuk GN, Pickar JG. Relationship between Biomechanical Characteristics of Spinal Manipulation and Neural Responses in an Animal Model: Effect of Linear Control of Thrust Displacement versus Force, Thrust Amplitude, Thrust Duration, and Thrust Rate. *Evid Based Complement Alternat Med*. 2013; 2013:492039. doi: 10.1155/2013/492039 [PubMed: 23401713]
- Reed WR, Chadha HK, Hubscher CH. Effects of 17beta-estradiol on responses of viscerosomatic convergent thalamic neurons in the ovariectomized female rat. *J Neurophysiol*. 2009; 102:1062–1074. DOI: 10.1152/jn.00165.2009 [PubMed: 19553492]
- Reed WR, Long CR, Kawchuk GN, Pickar JG. Neural responses to the mechanical characteristics of high velocity, low amplitude spinal manipulation: Effect of specific contact site. *Man Ther*. 2015a; 20:797–804. DOI: 10.1016/j.math.2015.03.008 [PubMed: 25841562]
- Reed WR, Long CR, Kawchuk GN, Sozio RS, Pickar JG. Neural Responses to Physical Characteristics of a High Velocity, Low Amplitude Spinal Manipulation: Effect of Thrust Direction. *Spine (Phila Pa 1976)*. 2015b; doi: 10.1097/BRS.0000000000001344
- Reed WR, Pickar JG, Sozio RS, Long CR. Effect of spinal manipulation thrust magnitude on trunk mechanical activation thresholds of lateral thalamic neurons. *J Manipulative Physiol Ther*. 2014a; 37:277–286. DOI: 10.1016/j.jmpt.2014.04.001 [PubMed: 24928636]
- Reed WR, Sozio R, Pickar JG, Onifer SM. Effect of spinal manipulation thrust duration on trunk mechanical activation thresholds of nociceptive-specific lateral thalamic neurons. *J Manipulative Physiol Ther*. 2014b; 37:552–560. DOI: 10.1016/j.jmpt.2014.08.006 [PubMed: 25220757]
- Rinaldi PC, Young RF, Albe-Fessard D, Chodakiewitz J. Spontaneous neuronal hyperactivity in the medial and intralaminar thalamic nuclei of patients with deafferentation pain. *J Neurosurg*. 1991; 74:415–421. DOI: 10.3171/jns.1991.74.3.0415 [PubMed: 1993906]
- Roberts VJ, Dong WK. The effect of thalamic nucleus submedius lesions on nociceptive responding in rats. *Pain*. 1994; 57:341–349. [PubMed: 7936712]
- Song XJ, Huang ZJ, Song WB, et al. Attenuation Effect of Spinal Manipulation on Neuropathic and Postoperative Pain Through Activating Endogenous Anti-Inflammatory Cytokine Interleukin 10 in Rat Spinal Cord. *J Manipulative Physiol Ther*. 2016; 39:42–53. DOI: 10.1016/j.jmpt.2015.12.004 [PubMed: 26837229]

- Sparks C, Cleland JA, Elliott JM, Zagardo M, Liu WC. Using functional magnetic resonance imaging to determine if cerebral hemodynamic responses to pain change following thoracic spine thrust manipulation in healthy individuals. *J Orthop Sports Phys Ther.* 2013; 43:340–348. DOI: 10.2519/jospt.2013.4631 [PubMed: 23485766]
- Srbely JZ, Vernon H, Lee D, Polgar M. Immediate effects of spinal manipulative therapy on regional antinociceptive effects in myofascial tissues in healthy young adults. *J Manipulative Physiol Ther.* 2013; 36:333–341. DOI: 10.1016/j.jmpt.2013.01.005 [PubMed: 23830709]
- Sumiya E, Kawakita K. Inhibitory effects of acupuncture manipulation and focal electrical stimulation of the nucleus submedialis on a viscerosomatic reflex in anesthetized rats. *Jpn J Physiol.* 1997; 47:121–130. [PubMed: 9159651]
- Tang JS, Qu CL, Huo FQ. The thalamic nucleus submedialis and ventrolateral orbital cortex are involved in nociceptive modulation: a novel pain modulation pathway. *Prog Neurobiol.* 2009; 89:383–389. DOI: 10.1016/j.pneurobio.2009.10.002 [PubMed: 19819292]
- Triano JJ. Biomechanics of spinal manipulative therapy. *Spine J.* 2001; 1:121–130. [PubMed: 14588392]
- Vaillant M, Edgecombe T, Long CR, Pickar JG, Kawchuk GN. The effect of duration and amplitude of spinal manipulative therapy (SMT) on spinal stiffness. *Man Ther.* 2012; 17:577–583. DOI: 10.1016/j.math.2012.06.006 [PubMed: 22809745]
- Wang JY, Zhao M, Huang FS, Tang JS, Yuan YK. Mu-opioid receptor in the nucleus submedialis: involvement in opioid-induced inhibition of mirror-image allodynia in a rat model of neuropathic pain. *Neurochem Res.* 2008; 33:2134–2141. DOI: 10.1007/s11064-008-9733-6 [PubMed: 18473169]
- Wright A. Hypoalgesia post-manipulative therapy: a review of a potential neurophysiological mechanism. *Man Ther.* 1995; 1:11–16. DOI: 10.1054/math.1995.0244 [PubMed: 11327789]
- Wright A. Recent concepts in the neurophysiology of pain. *Man Ther.* 1999; 4:196–202. DOI: 10.1054/math.1999.0207 [PubMed: 10593108]
- Xiao DQ, Zhu JX, Tang JS, Jia H. GABAergic modulation mediates antinociception produced by serotonin applied into thalamic nucleus submedialis of the rat. *Brain Res.* 2005; 1057:161–167. DOI: 10.1016/j.brainres.2005.07.053 [PubMed: 16125153]
- Yang S, Follett KA. Electrical stimulation of thalamic Nucleus Submedialis inhibits responses of spinal dorsal horn neurons to colorectal distension in the rat. *Brain Res Bull.* 2003; 59:413–420. [PubMed: 12576136]
- Yen CT, Lu PL. Thalamus and pain. *Acta Anaesthesiol Taiwan.* 2013; 51:73–80. DOI: 10.1016/j.aat.2013.06.011 [PubMed: 23968658]
- Yoshida A, Dostrovsky JO, Chiang CY. The afferent and efferent connections of the nucleus submedialis in the rat. *J Comp Neurol.* 1992; 324:115–133. DOI: 10.1002/cne.903240109 [PubMed: 1383287]
- Yoshida A, Dostrovsky JO, Sessle BJ, Chiang CY. Trigeminal projections to the nucleus submedialis of the thalamus in the rat. *J Comp Neurol.* 1991; 307:609–625. DOI: 10.1002/cne.903070408 [PubMed: 1714465]
- Zhang S, Tang JS, Yuan B, Jia H. Inhibitory effects of glutamate-induced activation of thalamic nucleus submedialis are mediated by ventrolateral orbital cortex and periaqueductal gray in rats. *Eur J Pain.* 1998; 2:153–163. [PubMed: 10700311]
- Zhang S, Tang JS, Yuan B, Jia H. Electrically-evoked inhibitory effects of the nucleus submedialis on the jaw-opening reflex are mediated by ventrolateral orbital cortex and periaqueductal gray matter in the rat. *Neuroscience.* 1999; 92:867–875. [PubMed: 10426528]
- Zhang YQ, Tang JS, Yuan B, Jia H. Effects of thalamic nucleus submedialis lesions on the tail flick reflex inhibition evoked by hindlimb electrical stimulation in the rat. *Neuroreport.* 1995; 6:1237–1240. [PubMed: 7669977]
- Zhao P, Waxman SG, Hains BC. Sodium channel expression in the ventral posterolateral nucleus of the thalamus after peripheral nerve injury. *Mol Pain.* 2006; 2:27. doi: 10.1186/1744-8069-2-27 [PubMed: 16916452]

Fig. 1. Experimental setup

Schematic showing the recording location of the thalamic submedial (Sm) neurons (a); and the experimental set-up (b) including the microdrive (1), lever arm of feedback motor (2), toothed forceps attached to the L5 spinous process (3), and spinal stabilization clamp attached to the L2 spinous process (4), and hip pins (5)

Fig. 2. Thrust protocol example

Shows a histogram (top) of neural activity recorded from a submedius neuron during 10s noxious stimulation (clip) applied to the contralateral hindpaw (cHP pinch). Pinches were applied approximately 60s prior to- and 10–15s following an L₅ vertebra thrust (85% body weight [BW] 100ms duration). Insets show the waveform for the L₅ vertebra thrust and action potential of this thalamic neuron. Control protocols were the same only without the thrust

Fig. 3. Electrolytic lesion site

Photomicrograph showing recording and electrolytic lesion site (arrow) with Di-I in the thalamic nucleus submedius (Sm). Hippocampus (HP), Thalamus (TH), Scale bar, 0.5mm

Fig. 4. Spontaneous activity

Mean changes ($\pm$ SEM) in spontaneous activity of nociceptive specific submedius neurons before and after a control and high velocity low amplitude L₅ vertebra thrust procedures. Note a significant decrease of spontaneous activity 180–240s after the delivery of the L₅ thrust (* p <0.005)

Fig. 5. Evoked response

Mean changes ($\pm$ SEM) in neural responses to 10s pinch applied to three sites (tail, contralateral hindpaw [cHP], ipsilateral hindpaw [iHP]) before and after control (no thrust) or high velocity low amplitude L₅ thrust procedures. All submedius neurons exhibited greater than 20% increase over baseline activity to the first applied pinch stimulus at each site. (* $p < 0.05$)

Fig. 6. Example of Sm neuron response

Histogram of Sm neural responses (impulses/1s bin) to noxious pinch at 3 sites (tail, contralateral and ipsilateral hindpaw) in both control and L₅ thrust protocols. Gray horizontal bars indicate noxious pinch application and arrowhead indicate time of L₅ thrusts. Note a decrease in tail and contralateral hindpaw post-thrust noxious response and an increase in ipsilateral hindpaw noxious response following an L₅ thrust. This neuron was located in the rostral Sm nucleus at 6.24mm from the cortical surface

Table 1

Change in Spontaneous Activity (imps)

Mean changes and SEM in spontaneous discharge activity (imps/time bin) of nociceptive specific Sm neurons following control or L5 thrust protocols. Group: $F=4.8$, $df=1,5$, $p<.05$, *post hoc* $^{\dagger}p<0.005$. Time (sec): $F=.53$, $df=5.5$

Seconds	Control		L5 Thrust	
	Mean	SEM	Mean	SEM
10	0.39	1.23	-0.28	0.80
60	4.75	3.65	-0.93	4.31
120	6.36	4.42	-0.88	7.72
180	3.35	5.56	6.64	6.16
240	11.63	7.39	-18.39 [†]	7.13
300	0.98	9.03	-5.43	9.22

Change in Noxious Evoked Activity (imps)
 Mean changes and SEM of noxious evoked response at 3 sites following control or L₅ thrust protocols among nociceptive specific Sm neurons that demonstrated a minimum of 20% increase over baseline discharge on initial pinch stimulus at each site. There was a significant decrease in the contralateral hindpaw response after the L₅ thrust (*p<.05)

Table 2

	Tail (n=35)		Contra-HP (n=43)		Ipsi-HP (n=39)	
	Mean	SEM	Mean	SEM	Mean	SEM
Control	0.60	6.38	-11.2*	2.33	0.14	3.38
L ₅ thrust	-8.51	4.82	-4.85	2.36	1.86	5.31
t=-.99, df=34, p>.05 t=2.3, df=42, *p<.05 t=-.34, df=38, p>.05						