Published in final edited form as:

Chem Commun (Camb). 2017 May 04; 53(37): 5136–5138. doi:10.1039/c7cc02062g.

Aminoquinoline-directed, cobalt-catalyzed carbonylation of sulfonamide sp^2 C–H bonds†

Tung Thanh Nguyen, Liene Grigorjeva, and Olafs Daugulis

Department of Chemistry, University of Houston, Houston, TX 77204-5003, USA

Abstract

We report a method for cobalt-catalyzed, aminoquinoline-directed sp^2 C–H bond carbonylation of sulfonamides. The reactions proceed in a dichloroethane solvent, and employ diisopropyl azodicarboxylate as a carbon monoxide source, $\text{Mn}(\text{OAc})_2$ as a cooxidant and potassium pivalate as a base. Halogen, ester, and amide functionalities are compatible with the reaction conditions. This method allows for a short and efficient synthesis of saccharin derivatives.

2-Sulfobenzoic acid imide functionality is important in pharmaceutical and food chemistry (Fig. 1).¹ For example, Repinotan **1** acts as a highly selective 5-HT_{1A} receptor agonist and is effective in counteracting the respiratory depression produced by morphine.^{1a} Ipsapirone **2** has been investigated for generalized anxiety disorder treatment.^{1b} Another imide of sulfobenzoic acid, supidimide **3**, has been used as a non-teratogenic analogue of thalidomide.^{1c} Saccharin **4** is an artificial sweetener,^{1d} while 6-nitrosaccharin **5** is its bitter derivative that activates G protein-coupled receptors hT2R44 and hT2R61, which are thought to mediate a bitter taste response in humans.^{1d,e}

The most direct synthesis of 2-sulfobenzoic acid imides would involve the direct carbonylation of arylsulfonic acid amides possessing an appropriate directing group, preferably using a first-row transition metal catalyst.² While the second-row transition metal-catalyzed carbonylation of C–H bonds has emerged as an efficient synthetic methodology,³ the use of first-row metals in these reactions is rare.⁴ The cobalt-catalyzed carbonylation of azobenzene and Schiff bases disclosed by Murahashi is perhaps the earliest example of directed C–H bond functionalization.^{4a} Previously, we have shown that aminoquinoline benzamide carbonylation can be effected at room temperature under 1 atm of CO by employing oxygen from air as the terminal oxidant.^{4b} Following this report, several groups have used aminoquinoline to direct cobalt- or nickel-catalyzed sp^2 and sp^3 C–H bond carbonylation.^{4c–e} All of these reports describe aminoquinoline amide C–H bond functionalization that results in the synthesis of substituted phthalic acids. Sulfonamide C–H bond carbonylation under first-row transition metal catalysis has not been reported yet.^{5a–c} A single paper by the group of Yu describes two examples of palladium-catalyzed C–H carbonylation.^{5d} We report here a method for cobalt-catalyzed, aminoquinoline-directed sp^2

†Electronic supplementary information (ESI) available: Detailed experimental procedures and characterization data for new compounds. See DOI: 10.1039/c7cc02062g

Correspondence to: Olafs Daugulis.

C–H bond carbonylation of sulfonamides. The cleavage of the directing group affords saccharin derivatives.

Based on cobalt-catalyzed, aminoquinoline-directed C–H functionalization reactions developed previously by our group,^{4b,6} simple cobalt salts were used as catalysts and manganese(II) acetate was chosen as a co-oxidant. Reaction optimization is illustrated in Table 1. The use of carbon monoxide gas gives low conversion to **7** (entry 1). Previously, we have shown that aminoquinoline benzamide carboxylation proceeds efficiently with carbon monoxide at room temperature.^{4b} At higher temperatures, required for the corresponding sulfonamide carbonylation, the use of gaseous carbon monoxide is technically challenging. A less volatile source of CO is needed for an efficient reaction. It is known that azodicarboxylate esters decompose at elevated temperatures releasing carbon monoxide,^{7a} and dialkyl azodicarboxylates have been used in directed alkoxycarbonylation and carbonylation of sp² C–H bonds.^{7b,c} When diisopropyl azodicarboxylate (DIAD) was used as a CO source, 38% yield of **7** was obtained (entry 2). The batchwise addition of DIAD increases the yield to 71% (entry 3). Cobalt(II) and (III) acetylacetonate catalysts afford lower yields compared with those of cobalt(II) acetate (entries 4 and 5). When manganese(III) acetate was used as a cooxidant, the yield decreased to 18% (entry 6). The presence of cobalt is necessary (entry 7), while the reaction under air without any cooxidant gave less than 5% of **7** (entry 8). The reaction under nitrogen gave a substantially lowered product yield. The use of trifluoroethanol as a solvent, which was used for previous cobalt-catalyzed reactions, gave no product. The use of *tert*-butyl azodicarboxylate instead of DIAD gave the product in lower yield (entry 9), while Mo(CO)₆ was inefficient as a carbon monoxide source (entry 10).

The reaction scope with respect to substitution in sulfonamides is shown in Table 2. Both electron-rich (entries 1–3, 8 and 12) and electron-poor (entries 5–7 and 9–11) substrates afford products in moderate to good yields. Scaling up of the reaction is feasible without the loss of yields (entry 1). Substitution at the *ortho*-position is tolerated, as sterically hindered 2-methylbenzene-sulfonamide gives the product in 55% yield (entry 8). Similar to cobalt-catalyzed aminoquinoline carboxamide carbonylation,^{4b} *meta*-substituted substrates react selectively at the less hindered C–H bond (entries 9–14) due to preferential metalation at the less sterically hindered position. The reaction is functional group tolerant, with alkoxy (entry 2), iodo (entry 5), bromo (entry 6), trifluoromethoxy (entry 7), trifluoromethyl (entry 9), chloro (entry 10), fluoro (entry 11), naphthyl (entry 12), unsaturated ester (entry 13), and amide (entry 14) substituents compatible with the reaction conditions. The trifluoroacetylated 1,2,3,4-tetrahydroisoquinoline derivative is carbonylated in 50% yield (entry 14), showing the relevance of the methodology in drug-like molecule derivatization.⁸ Nitro-substituted sulfonamides did not give any carbonylation product. The reaction mechanism likely includes aminoquinoline-directed cobaltation, carbon monoxide coordination, and migratory insertion followed by reductive elimination from acylcobalt(III) species. Reoxidation to cobalt(III) would complete the catalytic cycle. The reaction mechanism is under investigation and will be published as a full paper.

Chen introduced an 8-amino-5-methoxyquinoline auxiliary that can be oxidatively cleaved.⁹ The carbonylation of sulfonamide **8** gave product **9** in 54% yield (Scheme 1). The cleavage

of the 5-methoxyaminoquinoline auxiliary under the original conditions employing ceric ammonium nitrate was unsuccessful. However, the removal of the methyl group with BBr₃ followed by oxidation with an iodine(III) reagent afforded methylsaccharin **10** in 63% yield.¹⁰

In conclusion, we have developed a method for cobalt-catalyzed, aminoquinoline-directed carbonylation of aryl sulfonamides. The reactions proceed in the presence of DIAD as a carbon monoxide source, Mn(OAc)₂ as a cooxidant, and employ KOⁱPiv as a base. Functional groups such as halogens, esters, and amides are compatible with the reaction conditions. The removal of the directing group affords saccharin derivatives.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Welch Foundation (Chair E-0044) and NIGMS (Grant No. R01GM077635) for supporting this research.

Notes and references

- (a) Guenther U, Theuerkauf NU, Huse D, Boettcher MF, Wensing G, Putensen C, Hoeft A. *Anesthesiology*. 2012; 116:56. [PubMed: 22082683] (b) Albert PR, Le Francois B. *Front Neurosci*. 2010; 4:35. [PubMed: 20661455] (c) Neubert D, Krowke R. *Prog Clin Biol Res*. 1983; 110:387. [PubMed: 6298820] (d) Rohse H, Belitz HD. *Z Lebensm-Unters Forsch*. 1988; 187:425. (e) Pronin AN, Tang H, Connor J, Keung W. *Chem Senses*. 2004; 29:583. [PubMed: 15337684]
- General reviews: Lyons TW, Sanford MS. *Chem Rev*. 2010; 110:1147. [PubMed: 20078038] Gutekunst WR, Baran PS. *Chem Soc Rev*. 2011; 40:1976. [PubMed: 21298176] Engle KM, Mei TS, Wang X, Yu JQ. *Angew Chem, Int Ed*. 2011; 50:1478. Ackermann L. *Acc Chem Res*. 2014; 47:281. [PubMed: 23379589] Ye B, Cramer N. *Acc Chem Res*. 2015; 48:1308. [PubMed: 25884306] Daugulis O, Roane J, Tran LD. *Acc Chem Res*. 2015; 48:1053. [PubMed: 25756616] First-row transition metal catalysis: Kulkarni AA, Daugulis O. *Synthesis*. 2009; 4087. Moselage M, Li J, Ackermann L. *ACS Catal*. 2016; 6:498.
- (a) Fujiwara Y, Kawauchi T, Taniguchi H. *J Chem Soc, Chem Commun*. 1980:220. (b) Ohashi S, Sakaguchi S, Ishii Y. *Chem Commun*. 2005:486. (c) Giri R, Yu JQ. *J Am Chem Soc*. 2008; 130:14082. [PubMed: 18834125] (d) Giri R, Lam JK, Yu JQ. *J Am Chem Soc*. 2010; 132:686. [PubMed: 20000840] (e) Yoo EJ, Wasa M, Yu JQ. *J Am Chem Soc*. 2010; 132:17378. [PubMed: 21082838] (f) Orito K, Horibata A, Nakamura T, Ushito H, Nagasaki H, Yuguchi M, Yamashita S, Tokuda M. *J Am Chem Soc*. 2004; 126:14342. [PubMed: 15521739] (g) Houlden CE, Hutchby M, Bailey CD, Ford JG, Tyler SNG, Gagné SNG, Lloyd-Jones GC, Booker-Milburn KI. *Angew Chem, Int Ed*. 2009; 48:1830. (h) Guan ZH, Chen M, Ren ZH. *J Am Chem Soc*. 2012; 134:17490. [PubMed: 23051010] (i) Lang R, Shi L, Li D, Xia C, Li F. *Org Lett*. 2012; 14:4130. [PubMed: 22839509] (j) Zhang H, Liu D, Chen C, Liu C, Lei A. *Chem – Eur J*. 2011; 17:9581. [PubMed: 21793064] (k) Guan ZG, Ren ZH, Spinella SM, Yu S, Liang YM, Zhang X. *J Am Chem Soc*. 2009; 131:729. [PubMed: 19099479] (l) Du Y, Hyster TK, Rovis T. *Chem Commun*. 2011; 47:12074. (m) Inoue S, Shiota H, Fukumoto Y, Chatani N. *J Am Chem Soc*. 2009; 131:6898. [PubMed: 19413322] (n) Hasegawa N, Charra V, Inoue S, Fukumoto Y, Chatani N. *J Am Chem Soc*. 2011; 133:8070. [PubMed: 21542614] (o) Moore EJ, Pretzer WR, O'Connell TJ, Harris J, LaBounty L, Chou L, Grimmer SS. *J Am Chem Soc*. 1992; 114:5888. (p) Wang C, Zhang L, Chen C, Han J, Yao Y, Zhao Y. *Chem Sci*. 2015; 6:4610.
- (a) Murahashi S. *J Am Chem Soc*. 1955; 77:6403. (b) Grigorjeva L, Daugulis O. *Org Lett*. 2014; 16:4688. [PubMed: 25146415] (c) Barsu N, Bolli SK, Sundararaju B. *Chem Sci*. 2017; 8:2431.

- [PubMed: 28451350] (d) Williamson P, Galván A, Gaunt M. Chem Sci. 2017; 8:2588. [PubMed: 28553492] (e) Wu X, Zhao Y, Ge H. J Am Chem Soc. 2015; 137:4924. [PubMed: 25815529] (f) Liu XG, Zhang SS, Jiang CY, Wu JQ, Li Q, Wang H. Org Lett. 2015; 17:5404. [PubMed: 26451846]
5. Aminoquinoline sulfonamides in cobalt-directed C–H bond functionalization: Planas O, Whiteoak C, Company A, Ribas X. Adv Synth Catal. 2015; 357:4003. Kalsi D, Sundararaju B. Org Lett. 2015; 17:6118. [PubMed: 26624001] Ran Y, Yang Y, Zhang L. Tetrahedron Lett. 2016; 57:3322. Palladium-catalyzed sulfonamide C–H carbonylation: Dai HX, Stepan AF, Plummer MS, Zhang YH, Yu JQ. J Am Chem Soc. 2011; 133:7222. [PubMed: 21488638]
6. (a) Grigorjeva L, Daugulis O. Angew Chem, Int Ed. 2014; 53:10209. (b) Grigorjeva L, Daugulis O. Org Lett. 2014; 16:4684. [PubMed: 25146300] (c) Grigorjeva L, Daugulis O. Org Lett. 2015; 17:1204. [PubMed: 25689275] (d) Nguyen TT, Grigorjeva L, Daugulis O. ACS Catal. 2016; 6:551. [PubMed: 27525171]
7. (a) Stolle R, Reichert W. J Prakt Chem. 1929; 123:82. (b) Yu WY, Sit WN, Lai KM, Zhou Z, Chan ASC. J Am Chem Soc. 2008; 130:3304. [PubMed: 18298118] (c) Ni J, Li J, Fan Z, Zhang A. Org Lett. 2016; 18:5960. [PubMed: 27934491]
8. Busujima T, Tanaka H, Shirasaki Y, Munetomo E, Saito M, Kitano K, Minagawa T, Yoshida K, Osaki N, Sato N. Bioorg Med Chem. 2015; 23:5922. [PubMed: 26210160]
9. (a) He G, Zhang SY, Nack WA, Li Q, Chen G. Angew Chem, Int Ed. 2013; 52:11124. (b) Yamamoto C, Takamatsu K, Hirano K, Miura M. J Org Chem. 2016; 81:7675. [PubMed: 27504671]
10. Nakazaki A, Mori A, Kobayashi S, Nishikawa T. Tetrahedron Lett. 2012; 53:7131.

Fig. 1.
2-Sulfolbenzoic acid imides.

Scheme 1.
Removal of directing group.

Table 1

Optimization of reaction conditions^a

Entry	Catalyst	Cooxidant	Yield of 7, %
1 ^b	Co(OAc) ₂	Mn(OAc) ₂	<5
2 ^{c,d}	Co(OAc) ₂	Mn(OAc) ₂	38
3 ^d	Co(OAc) ₂	Mn(OAc) ₂	71
4 ^d	Co(acac) ₂	Mn(OAc) ₂	31
5 ^d	Co(acac) ₃	Mn(OAc) ₂	25
6 ^d	Co(OAc) ₂	Mn(OAc) ₃ ·2H ₂ O	18
7 ^d	None	Mn(OAc) ₂	0
8	Co(OAc) ₂	None	<5
9 ^{d,e}	Co(OAc) ₂	Mn(OAc) ₂	62
10 ^{d,f}	Co(OAc) ₂	Mn(OAc) ₂	0

^a Sulfonamide **6** (0.5 mmol), DIAD (2.5 mmol, added 4 times, 0.625 mmol per time), catalyst (0.15 mmol), cooxidant (1 mmol), KOPiv (1 mmol), DCE (10 mL), 100 °C, 24 h. Yields are isolated yields. Please see the ESI for details. Abbreviations: DIAD = diisopropyl azodicarboxylate, OPiv = pivalate, DCE = 1,2-dichloroethane, acac = acetylacetonate, and Q = 8-quinoliny1.

^b Carbon monoxide was used instead of DIAD.

^c DIAD (2.5 mmol) was added at once at the beginning of the reaction.

^d Reaction vessel periodically opened to air.

^e *tert*-Butyl azodicarboxylate used instead of DIAD.

^f Molybdenum hexacarbonyl used instead of DIAD.

Table 2

Carbonylation of aminoquinoline sulfonamides^a

$ \begin{array}{c} \text{O}=\text{O} \\ \diagup \quad \diagdown \\ \text{Ar}-\text{S}-\text{NHQ} \end{array} \xrightarrow[\text{DCE, 100 }^{\circ}\text{C}]{\begin{array}{l} \text{DIAD (5 equiv)} \\ \text{Co(OAc)}_2 \text{ (30 mol\%)} \\ \text{Mn(OAc)}_2 \text{ (2 equiv)} \\ \text{KO}^i\text{Piv (2 equiv)} \end{array}} \begin{array}{c} \text{O}=\text{O} \\ \diagup \quad \diagdown \\ \text{R}-\text{C}_6\text{H}_3-\text{S}-\text{N-Q} \\ \diagup \quad \diagdown \\ \text{O} \quad \text{O} \end{array} $			
Entry	Ar	Product	Yield, %
1	4-MeC ₆ H ₄		71 68 ^b
2	4-MeOC ₆ H ₄		59
3	4- <i>t</i> BuC ₆ H ₄		72
4	4-PhC ₆ H ₄		57
5	4-IC ₆ H ₄		73
6	4-BrC ₆ H ₄		61

Entry	Ar	Product	Yield, %
7	4-CF ₃ OC ₆ H ₄		63
8	2-MeC ₆ H ₄		55
9	3-CF ₃ C ₆ H ₄		47
10	3-ClC ₆ H ₄		48
11	3,4-F ₂ C ₆ H ₃		69
12 ^c	2-Naphthyl		51

Entry	Ar	Product	Yield, %
13	2-Oxo-2H-1-benzo-pyran-5-yl		65
14	2-CF ₃ CO-tetra-hydroisoquinol-1-yl		50 ^d

^a Sulfonamide (0.5 mmol), DIAD (2.5 mmol, added 4 times, 0.625 mmol per time), Co(OAc)₂ (0.15 mmol), Mn(OAc)₂ (1 mmol), KOPiv (1 mmol), DCE (10 mL), 100 °C, 24 h. Yields are isolated yields. Reaction vessel periodically opened to air. Please see the ESI for details.

^b Scale: 2.5 mmol, 30 h.

^c Cobalt(II) chloride (20 mol%) catalyst, DIAD (1.5 mmol), 24 h at 120 °C.

^d Temperature: 85 °C, 30 h.