Published in final edited form as:

ACS Chem Biol. 2017 August 18; 12(8): 2157–2163. doi:10.1021/acscchembio.7b00281.

Progressive Stereo Locking (PSL): A Residual Dipolar Coupling Based Force Field Method for Determining the Relative Configuration of Natural Products and Other Small Molecules

Gabriel Cornilescu^{†,‡,*}, René F. Ramos Alvarenga[§], Thomas P. Wyche^{§,||}, Tim S. Bugni[§], Roberto R. Gil[⊥], Claudia C. Cornilescu[‡], William M. Westler^{†,‡}, John L. Markley^{†,‡}, and Charles D. Schwieters^{#,*}

[†]National Magnetic Resonance Facility at Madison

[‡]Biochemistry Department, University of Wisconsin—Madison, Madison, Wisconsin 53706, United States

[§]Pharmaceutical Sciences Division, School of Pharmacy, University of Wisconsin—Madison, Madison, Wisconsin 53705, United States

^{||}Biological Chemistry and Molecular Pharmacology Department, Harvard Medical School, Boston, Massachusetts 02115, United States

[⊥]Department of Chemistry, Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, United States

[#]Center for Information Technology, National Institutes of Health, Bethesda, Maryland 20892-5624, United States

Abstract

Establishing the relative configuration of a bioactive natural product represents the most challenging part in determining its structure. Residual dipolar couplings (RDCs) are sensitive probes of the relative spatial orientation of internuclear vectors. We adapted a force field structure calculation methodology to allow free sampling of both *R* and *S* configurations of the stereocenters of interest. The algorithm uses a floating alignment tensor in a simulated annealing protocol to identify the conformations and configurations that best fit experimental RDC and distance restraints (from NOE and *J*-coupling data). A unique configuration (for rigid molecules)

*Corresponding Authors: gabrielc@nmrfam.wisc.edu, charles.schwieters@nih.gov.

ORCID

Gabriel Cornilescu: 0000-0002-1204-8904

Thomas P. Wyche: 0000-0003-4703-741X

Tim S. Bugni: 0000-0002-4502-3084

John L. Markley: 0000-0003-1799-6134

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI:10.1021/acscchembio.7b00281.

Details of experimental and computational methods together with the molecular representations of the cases tested (PDF)

or a very small number of configurations (for less rigid molecules) of the structural models having the lowest chiral angle energies and reasonable magnitudes of the alignment tensor are provided as the best predictions of the unknown configuration. For highly flexible molecules, the progressive locking of their stereocenters into their statistically dominant *R* or *S* state dramatically reduces the number of possible relative configurations. The result is verified by checking that the same configuration is obtained by initiating the locking from different regions of the molecule. For all molecules tested having known configurations (with conformations ranging from mostly rigid to highly flexible), the method accurately determined the correct configuration.

Graphical Abstract

Enzymatic reactions leading to the biosynthesis of secondary metabolites overwhelmingly preserve unique regio- and stereospecificity. Therefore, knowledge of relative configuration is of the utmost importance for establishing configuration–bioactivity relationships.¹ Traditional methods, including *J*-couplings,^{2,3} NOEs,⁴ and ¹³C chemical shift predictions,⁵ often fail to unambiguously determine the configuration of such molecules. For a large number of molecules that cannot be crystallized, organic synthesis of all possible configurations compatible with the spectroscopic data is often unrealistic; thus, better ways to identify the configuration of the bioactive compound are required.

Motivated by the isolation of structurally novel natural products from marine invertebrate-associated bacteria,^{6,7} we explored the potential of NMR residual dipolar couplings (RDCs) to determine their relative configuration. RDCs are small nuclear spin couplings resulting from weak alignment of the molecule of interest in liquid-crystalline media or in stretched gels. The small degree of molecular orientation is described by the Saupe order matrix or alignment tensor.⁸ Although RDCs have been used extensively to extract structural and dynamic information on biomacromolecules,^{9–11} similar applications to small molecules became established relatively recently.^{12–16}

RDCs reflect the relative orientation of internuclear vectors in the molecular frame, which makes them a valuable long-range complement to NOEs. While vicinal relative configurations can be deduced from the NOEs and *J*-couplings involving the surrounding atoms, a very large number of relative configurations of more distant stereocenters remain compatible with the experimental NOE and *J*-coupling data. RDCs can dramatically reduce this remaining ambiguity.

To date, NMR could not be convincingly used to distinguish enantiomers in chiral nonracemic alignment conditions;¹⁷ therefore, the absolute configuration is obtained by chiroptical spectroscopy.¹⁸ Most protocols for RDCs used to date require the generation of a large number (up to 2^{N-1} , where N is the number of centers with unknown configuration) of 3D structural models of each enantiomer and then selecting the ones that best fit the experimental NOEs and RDCs. The models usually are calculated by density functional theory (DFT),^{5,19,20} and while computationally demanding, this method works well for mostly rigid molecules. In more dynamic cases, the models are generated by molecular dynamics (MD) simulations^{21,22} or simulated annealing (SA).²³ While software packages like MSpin-RDC²⁴ can tackle phenyl rotor averaging, the modeling of various conformers is performed manually (e.g., by generating models for discrete rotations around a particular covalent bond, calculating the alignment tensor of each conformer, and then selecting the one(s) that best fit(s) the RDCs).

We propose here a simpler method that uses restraints from NOEs and one-bond C–H RDC ($^1D_{CH}$) measurements, along with a simulated annealing protocol in the Xplor-NIH software package²⁵ modified to allow unbiased sampling of both *R* and *S* configurations for a user-selected set of stereocenters. The resulting structural models having the lowest angular energies at the stereocenters and a reasonable magnitude of the fitted alignment tensor are selected as predictors of the true configuration and conformation in solution. In our testing of rigid molecules of known relative configuration, if the best thus-selected 10 structures exhibited a unique configuration, that configuration was invariably the correct one. With less rigid molecules, only a small subset of the carbon atoms had a consistent configuration among these best 10 structures. Subsequent runs with this subset of stereocenters locked into their consistent configuration resulted in increased configuration consistency among the remaining centers. The procedure is repeated to maximize the number of stereocenters exhibiting a consistent relative configuration, so as to dramatically reduce the number of possible diastereomers compatible with the experimental data.

This algorithm represents a major improvement over established methods that use Xplor to calculate structures with restraints derived from NOEs and *J*-couplings and then filter them by the quality of the RDC fit²⁶ or by including the RDC restraints as a separate structural refinement step.²³

RESULTS AND DISCUSSION

For biomacromolecules, several NMR spectroscopic methods exist for establishing which part of the backbone represents a mostly rigid core and which parts are dynamic, allowing the exclusion of RDCs from the less rigid parts when calculating the alignment tensor. In this case, the number of measured RDCs is relatively high, and their directions usually sample sufficiently the 3D space so that the molecule's alignment tensor can be determined with high precision (e.g., through SVD of an overdetermined system of linear equations). For smaller molecules, the conformational freedom may be higher while the RDC directional sampling is sparser; therefore, either a large number of RDCs needs to be measured (e.g., by inclusion of long-range RDCs^{30,34,35}) or measurements need to be performed in more than one alignment medium to avoid ambiguities resulting from

underdetermination. The discovery rate of new alignment media compatible with organic solvents has been steady in the past decade, allowing relatively straightforward measurement of two or more independent RDC sets, which dramatically reduces such ambiguities.

Because measuring long-range RDCs accurately is more challenging, we focused on using the most common $^1D_{CH}$ RDC in a single alignment medium for all of our tests, while recording them in a second alignment medium only for the more complex case of ecteinamycin (14 stereocenters).

We used our method of calculating structural models that best fit the RDCs in a simulated annealing force field protocol in Xplor-NIH, which allows free sampling of all configurations for the carbon atoms of interest. The methods to date require either an *a priori* estimate of the magnitude and rhombicity of a single alignment tensor or that of various conformational dependent tensors, together with their contribution to a dynamic model. This requirement is eliminated in our approach, because the tensor is automatically the one that best fits the experimental RDCs for each molecular conformation calculated.

The accuracy of configurations predicted by our method can be affected by the quality of initial topology and parameter files generated by the PRODRG Web server from a reasonable set of initial coordinates (e.g., a single 3D model with a random configuration). The topology and parameter files generated by PRODRG need to be carefully inspected for possible errors of interpreting the chemical formula, number of double bonds, values of dihedral and improper angles, etc. For the test cases where the correct configuration was known (*i.e.*, literature cases), we deliberately altered it to avoid bias in the results. Furthermore, to avoid any artificial bias in predicted configuration, we thoroughly tested and confirmed the capacity of Xplor-NIH to sample equally and randomly all possible configurations of the carbon atoms of interest, when no NOE or RDC restraints were active.

In our tests, we noticed that structures with incorrect configurations or distortions had artificially high magnitudes for the best fitting alignment tensor or high chiral angle energies. In practice, an optimal degree of alignment corresponding to a Da of 25 Hz and a principal order parameter of 10^{-3} gives rise to $^1D_{CH}$ RDCs with a maximum absolute value of 50 Hz (when the internuclear vector is parallel to the longest axis of the alignment tensor). A Da of 45 Hz (giving rise to RDCs up to 90 Hz) corresponds to overalignment that would dramatically deteriorate the spectra due to the dipolar broadening (long-range interactions) and to the loss of INEPT transfer due to the large variability of the total couplings ($^1J_{CH} + ^1D_{CH}$). Therefore, we select the 10 structures with the lowest chiral angle energies and with best-fitted alignment tensor magnitudes smaller than 45 Hz. Invariably, some of these 10 structures were found to have the lowest total energies. When a single configuration (or its mirror image) is consistent among the majority of these 10 structures, we predict that it is the correct one. This was the case for enterocin, and we expect it to be the case with most small, rigid molecules for which sufficient NOE and RDC restraints are available (a single set of RDCs is usually sufficient).

However, for less rigid molecules with a large number of chiral centers, the inherent degeneracy of the orientational restraints provided by RDCs, even when measured in a

second, independent set of RDCs, cannot result in a single, consistent configuration as we observed for ecteinamycin. In such cases, smaller regions of the molecule that are either more rigid or have sufficient restraints (e.g., proton-dense regions with a relatively large number of measured NOEs and RDCs) frequently exhibit high consistency in the calculated configurations. Therefore, we tested a Progressive Stereo Locking (PSL) method, in which we lock these few centers in their dominant configuration (i.e., those similar in at least 7 out of the 10 best structures, selected as described above) by reactivating their bond angle terms. We then repeat the Xplor-NIH calculation with floating chirality for the rest of the chiral carbons, which invariably results in improvement of their chiral consistency. We iterate this process to maximize the number of locked stereocenters, which dramatically reduces the number of possible configurations compatible with the experimental restraints and, in some test cases, yields a unique dominant configuration that proves to be the correct one.

The lack of quantitative information on the dynamics of the covalent bonds corresponding to the measured RDCs implies that their measured magnitude represents the lower limit of the true RDCs. One might expect that the current approach of using a single structure and a quadratic energy function for the RDCs would fail in cases where some portions of the molecule of interest are much more rigid than others or in cases where large populations with radically differing alignment tensors are present. The fact that the current protocol is successful suggests that a single, time-averaged representative may minimize the former problem as a consequence of rapid tumbling of these small molecules and the domination of their populations. It is conceivable that particular time scales of dynamic parts of a molecule combined with a rather disperse distribution of its conformers may impede obtaining a unique configuration through the presented method. However, this will be obvious in our protocol since in these particular cases the generated structures will have a smaller number of centers with an unambiguous configuration, while fitting equally well the experimental RDCs and NOEs. If a sufficiently large number of RDCs are available (e.g., by measuring C–C and precise long-range C–H RDCs), examining separately the rigid parts in conjunction with jack-knifing or bootstrapping analyses to assess the precision of these alignment tensors would vastly improve the confidence in the configuration of these rigid fragments and of the global configuration.

Enterocin

We used this small molecule isolated from a marine *Streptomyces* sp. (WMMB 285) as an initial simple test case. Two DFT models were generated (using Spartan 10 software³⁶) having the opposite configuration at a single stereocenter (C5), and a single set of $^1D_{CH}$ RDCs was measured in PMMA gels. The lower limit of the experimental error (validated by a duplicate measurement), given by $LW[Hz]/(S/N)$ (the line width divided by the signal-to-noise ratio) when measured in the ^{13}C (F1) dimension,³⁰ was approximately ~0.2 Hz, which propagated into ~0.3 Hz in the RDC values, calculated as differences between the aligned and isotropic values of the H–C coupled resonances. When analyzing the SVD fit of the experimental RDCs to these two DFT models we observed that while the correct configuration clearly fitted the experimental RDCs better than the incorrect configuration (Figure 1B), the root-mean-square deviations (RMSD) between the experimental and back-calculated RDCs using the DFT models was about an order of magnitude larger than the

experimental errors (3.9 and 8.8 Hz for the correct and incorrect configurations, respectively).

This suggested that, even for such a simple test case, the time-averaged conformation of the ring rotations and other structural fluctuations in solution, reflected in the RDC values (measured over tens of milliseconds), might not be accurately represented by the lowest energy DFT models. This can be problematic for studies that employ such models to distinguish configurations that fit the RDCs only slightly better than others.

We therefore proceeded with our new Xplor-NIH protocol starting from a 3D model with the incorrect configuration at C5. For enterocin, the Xplor-NIH calculations with chirality unlocked at all of its seven stereocenters resulted in the correct configuration of the majority or all of the 10 best structures. Despite the fact that all starting models had the incorrect C5 configuration, and contained variations in the number of aromatic ring atoms locked in a planar conformation or in the relative force constants used to properly balance the various energy terms, the results exhibited only small variations in the calculated configurations. Other minor corrections in the dihedral or improper angle parameters created by PRODRG similarly resulted in only small variations in the number of chiral centers with dominant chirality among the best 10 structures calculated. In all cases where ambiguity was present at any chiral center, we applied the PSL method and invariably obtained the correct configuration after just a few iterations.

The best calculated Xplor-NIH enterocin structures fitted the RDCs with RMSDs of approximately 1 Hz. When we compared these structures with the DFT models, we saw most of the differences consisting of small reorientations of the aromatic groups. Such minor structural reorientation nevertheless resulted in RMSDs of their RDC fit to the DFT models much larger than 1 Hz (Figure S2).

Ecteinaamycin.³⁷

This recently discovered polyether antibiotic from marine *Actinomadura* sp. (WMMB 499) has 14 stereocenters. It represents a truly challenging system as all attempts to determine its configuration by crystallization, by derivatization with a chiral auxiliary, or by using ¹³C chemical shift predictions were unsuccessful. Traditional NMR information based on NOEs, *J*-couplings, and chemical shifts were used to partially solve the relative configurations of the carbon atoms within the cyclic regions (Figure 2), which reduced the number of possible relative configurations from 8192 (2¹³) to 256 (2⁸).

Ecteinaamycin appeared to align differently in PMMA gels and PELG (Figure S1), but the experimental RDCs did not fit any of the 256 DFT models generated based on NOEs, *J*-couplings, and chemical shifts (the RMSDs between the experimental and back calculated RDCs were in the 10–13 Hz range, while RDC's precision was better than 0.5 Hz, see Supporting Information). Our attempts to use only various subsets of the two sets of RDCs (and keep the unused ones for cross-validation) were inconclusive, as only a very sparse consensus configuration was obtained. We therefore used both RDC sets in our PSL method and initiated the locking of stereocenters showing a consistent configuration among 8 or 9 out of the 10 best structures. In cases when we locked chirality consistent within the

minimum 7 out of 10 best structures, we also checked its alternative (*i.e.*, by locking its opposite stereoisomer). In these tests, the centers with a highly dominant configuration (8, 9, or 10 out of 10) remained in that configuration. This allowed us to clearly identify the stereocenters with a truly consistent configuration in contrast to the more ambiguous ones. Most of the stereocenters that could be locked after the first few calculations (in which all or most of the 14 centers were unlocked) were in the cyclic C8–C14 and C24–C33 regions (Figure 2), as expected for those more rigid parts of the molecule. We also tested multiple locking pathways (initiated either in the C8–C14 or the C24–C33 regions) and obtained a consistent configuration for 13 (out of 14) stereocenters (8*S*, 9*S*, 12*S*, 14*R*, 15*S*, 17*S*, 18*S*, 22*S*, 24*R*, 27*R*, 30*S*, 33*R*, 34*R*). The presence of a quaternary carbon at C33 and unhindered rotation around the C33–C36 bond explains the ambiguity of C36 even when RDCs from two independent alignment media are used as restraints.

We also developed an exhaustive, more computationally intensive, method to calculate 10 structures for each of the 2^{14} configurations restrained by experimental NOEs and RDCs. We then applied the same filters used in the PSL method (lowest angular energies at the stereocenters and a reasonable magnitude of the fitted alignment tensors) to select a smaller number of structures (out of these 10×2^{14}) and compared their configuration to that calculated by PSL. We noticed that the configuration of the more rigid regions (the cyclic parts of the molecule) was consistent with the PSL one, while the configuration of regions with many degrees of freedom (*i.e.*, linker connecting the cyclic parts) varied considerably among the selected structures, as expected, while no significant energy difference could grant differentiation among them. This confirmed that PSL is needed to further reduce this still large number of possible configurations.

Deoxycumambrin.³⁰

We chose the 10-*epi*-8-deoxycumambrin B case from the literature because, even though the molecule has only five stereocenters, three MM/DFT isomers were hard to distinguish on the basis of one-bond RDCs alone. Small long-range couplings, for which highly precise measurements on the order of 0.1 Hz are required, were used to select the correct isomer in the original study. The use of long-range couplings is problematic in general due to the need for very high precision, unresolved peaks, potential sign ambiguity of vanishingly small couplings, and errors in the large internuclear distances involved. We therefore tested our method of calculating structures of this molecule using only the published one-bond ($^1D_{CH}$) RDCs. Furthermore, although in the original study one stereocenter was maintained in a known configuration (resulting in only 2^4 remaining possible configurations), we allowed floating chirality at all five centers (Figure S4). The structures with correct configuration (or its mirror image) among the best 10 calculated structures (by the criteria mentioned before) could clearly be distinguished on the basis of lowest total energy (Table S1) without the need to perform PSL. It is worth noting that the correct configurations also showed similar values for the alignment tensor magnitude and rhombicity. Predictably, when testing PSL on this system, the correct answer was obtained after just two iterations (three stereocenters were already 100% consistent among the 10 best structures calculated with all five centers unlocked), as shown in Table S1.

Vatiparol.³¹

Vatiparol has eight stereocenters and three rotating phenyl rings (Figure S5). In the original study, the conformation of two of these aromatic rings in the DFT refined structures, which agreed with the experimental NOEs, did not fit the experimental RDCs well. Since a large number of long-range NOEs helped establish the compound's relative configuration but not the conformation of the rings, the authors resorted to calculating populations of various ring conformations (with uncorrelated torsion angles) to obtain a proper fit to the experimental RDCs. Their method involved selecting 16 configurations out of the possible 128 based on *J*-couplings, followed by fitting the RDCs to the core structures with phenyls excluded to determine the configuration that agreed best with the NOEs. Subsequently, MM modeling was used to generate aromatic ring conformers, and DFT was used to refine them. Finally, the RDCs were fitted to the entire structures, which allowed selecting the conformer with a configuration that best fitted the NOEs.

Our protocol used the published NOEs and RDCs (no *J*-couplings) and an initial structural model (with the correct configuration, as received from the authors) in an XPLOR-NIH run with floating chirality (that took 10 min). All 10 best structures (selected by chiral angle energy and Da criteria) had the correct *RSRRSRRS* configuration (or its enantiomer). When starting with the opposite enantiomer, again all 10 best structures had the correct configuration. After scrambling the starting model's configuration to *RRRRSRRS*, the best 10 calculated structures (by the same criteria mentioned above) exhibited only two configurations that differed at a single stereocenter: the correct one was represented by three structures with the lowest total energies, and seven higher energy structures had the configuration *RSRRSRRS*. This suggests that, for favorable cases like vatiparol (in which the large number of long-range NOEs combined with RDCs provides considerable configurational discrimination power), running a second Xplor-NIH calculation starting with the lowest total energy structure from the initial calculation may result in all 10 best structures having a unique, consistent configuration, without needing to lock any stereocenter by PSL. In such cases, the PSL method can still be used for cross-validation. An overlay of the lowest total energy structure calculated with Xplor-NIH (that fits the RDCs with an RMSD of 0.9 Hz) and the DFT calculated structure provided by the authors showed differences mostly in the ring orientations (Figure S4B).

Fibrosterol.²²

Fibrosterol sulfate A is a pseudosymmetric structure with two disulfated steroidal tetracyclic moieties with known configurations joined by a flexible linker containing a cyclopentyl unit (Figure S6). Three of the five stereocenters in the linker had unknown configuration, and large-scale motions were expected to generate conformers with substantially different alignment tensors, making this case a challenge for any RDC-based method. The original study used multiple alignment tensor analysis combined with molecular dynamics in explicit solvent, but also found that the single tensor approximation gave the same result, presumably due to the dominance of an extended conformer. The analysis identified eight possible configurations that were subjected to an unrestrained MD simulation (100 ns per trajectory). The trajectories were then filtered by agreement with NOEs, $^3J_{HH}$, and $^3J_{CH}$ couplings. The resulting structures were clustered in four conformations per configuration,

and an average alignment tensor was determined for each conformation. Subsequently, the eight configurations were analyzed by fitting the RDCs to determine the contribution of each conformer to an average alignment tensor. The best fitting configuration found for C22, C25, and C24' was *SRS*, respectively. The original paper used a different convention in naming C25*S*, and thus the configuration is labeled *SSS* in that work. We use the IUPAC convention, which denotes C25 as *R*. Long-range NOEs were observed between the two steroid systems (corresponding to a collapsed conformation), but their weak intensities were interpreted as attesting to the presence of other more extended conformations. We noticed, however, that several strong linker NOEs calibrated at 2–3 Å, were incompatible with the extended conformation.

When we tested PSL on fibrosterol sulfate A with the three unknown centers unlocked using the published NOEs and RDCs, the lowest energy structure (and the fifth lowest total energy) already had the correct configuration (*SRS*). Upon locking the two dominant stereoconfigurations, C22*S* (80%) and C25*R* (90%), seven out of the 10 best structures (including the three lowest total energy structures) had the correct *SRS* configuration of the linker.

All our calculated structures had the collapsed conformation, because we used all experimental NOEs from the original paper (including the ~5 Å intersteroid NOEs and the 2–3 Å linker NOEs, which are incompatible with an extended conformation). The RDCs fitted the two lowest energy structures with RMSDs of 0.6–0.8 Hz, clearly within the experimental error, and agreed well with the experimental NOEs.

This test case illustrates a major advantage of our method: if a particular conformer is reflected in the experimental restraints (NOEs, *J*-couplings, RDCs), we simply calculate its structure to extract the correct configuration of the molecule, without the need to detect other conformers. The alignment tensor is fitted on-the-fly to each particular conformer calculated, and as long as the configuration of the lowest total energy conformers shows consistence, the PSL approach leads to the correct answer.

Conclusions

In summary, we have developed a simple method that uses RDCs and NOEs to calculate the 3D structures of natural products and other small molecules having the most likely configuration(s). The PSL method correctly identified the known relative configuration of all test cases. For rigid molecules, supplementing NOE (and optionally *J*-coupling) information with the most commonly measured one-bond $^1D_{CH}$ dipolar coupling restraints (in a single alignment medium) is sufficient for determining their relative configuration. For more flexible molecules with a large number of stereocenters, measuring $^1D_{CH}$ in a second alignment medium provides sufficient discriminating power to dramatically reduce the number of possible relative configurations from 2^{N-1} to just a few. Furthermore, for any chosen configuration, Xplor-NIH can generate ensembles of conformers best fitting the NOEs and RDCs, which can serve as starting models for absolute configuration determination by comparing calculated and experimental electronic circular dichroism spectra.³⁸

The PSL method including an example case presented in this paper is part of Xplor-NIH software distribution (starting with release 2.45). All other test cases and the associated Xplor-NIH input files and scripts are available on the Web at <http://www.nmrham.wisc.edu/software.htm>

METHODS

We recorded RDCs in CDCl_3 in PMMA compressed gels²⁷ on enterocin, a simple rigid model with seven stereocenters,²⁸ and ecteinamycin, a challenging natural product with 14 stereocenters (details in the Supporting Information). For ecteinamycin, an additional set of RDCs was recorded in PELG²⁹ liquid crystalline medium. The minimum amount of PELG needed to maintain a nematic state was used, as the molecule was slightly overaligned even at this concentration (the maximum measured $^1D_{\text{CH}}$ was 46.5 Hz). The two media resulted in uncorrelated RDC values (Figure S1), with a good degree of linear independence confirmed by a scalar product of the corresponding alignment tensors ranging between -0.23 and -0.45 for most structural models calculated.

Additionally, we tested our method on several published molecules: 10-epi-8-deoxycumambrin B, a tricyclic natural compound with five stereocenters for which three MM/DFT isomers were hard to distinguish from one-bond RDCs;³⁰ vatiparol, a minor resveratrol trimer with three rotating phenyl rings and eight stereocenters;³¹ and fibrosterol sulfate A, a polysulfated steroid with a flexible linker and large-scale motions.²²

We used 16 NOE restraints in the $0.5\text{--}4.0$ Å range for ecteinamycin, none for enterocin, and the published NOE and RDC values for the literature test cases. Occasionally, J -coupling information (*i.e.*, when indicating unambiguous *cis* vs. *trans* geometries) was used indirectly, being translated either into additional proton distance restraints or into additional improper angle restraints. For methylenes with nonmagnetically equivalent protons, we used initially the sum of their individual CH RDCs. After obtaining a structural model that fitted the other RDCs well, we tested each of the two possible stereospecific CH RDC assignments for each methylene group and chose the one that best fitted that model.

Topology and parameter files were generated using the PRODRG Web server³² from a reasonable set of initial coordinates (*e.g.*, a DFT model) with a configuration that was randomly altered from the correct one, when known. We emphasize that we used a DFT starting model for the molecules in the published studies mentioned above only as a matter of convenience (they were available), but such 3D models are not required for our method. Simply drawing the 2D covalent structure of the molecule of interest in the Web interface of PRODRG is sufficient to generate an energy minimized 3D model, which is not intended nor needs to represent the absolute energy minimum of a molecule's structure. The conformation and configuration of this starting model will subsequently be modified by our protocol, and refined structures will be generated to best fit the experimental restraints (NOEs and RDCs). RDC experimental errors were not used to weigh differently the RDC restraints in the structure calculations. All dihedral angles that maintain double bond or ring planarity must be replaced with improper angles (in the PRODRG generated topology file), as per Xplor convention. The force constants for dihedral and improper angles were adjusted

uniformly to 50 and 500 kcal mol⁻¹ rad⁻², respectively, which represent Xplor-NIH optimized values for balancing various energy terms in proteins.

In this protocol, for the stereocenters to be determined, we inactivated the improper dihedral and bond angle terms to allow their chirality to float. While the implementation is different, the idea is analogous to the floating stereospecific assignment of methylene or isopropyl groups under NOE restraints when computing NMR biomolecular structures.³³ The XPLOR bond angle terms at these centers were replaced with Python versions for which the force constants can be independently scaled. Importantly, while only torsion angle degrees of freedom were allowed away from the chiral centers, all degrees of freedom were freed about each stereocenter. The planar aromatic rings were rigidized to maintain the initial optimized geometry under RDC restraints. In a final refinement step after configuration was determined, full angle force constants are utilized along with all improper dihedral angle terms to restrain each center to its determined chirality. As covalent geometry is satisfied at the beginning of this refinement stage and structure calculation consists of dynamics and minimization in torsion angle space, there are no bond or angle violations away from ring regions. Six-membered rings are manually inspected to ensure they assume proper chair conformations.

Our modified Xplor-NIH method uses a slow cooling protocol to calculate 100 structures, which are then sorted by the chiral angle energy and filtered by the requirement that SVD fitted alignment tensor magnitude (Da) is less than 45 Hz. The 10 models best fitting these criteria are used as predictors of the true configuration. Invariably, some of these models are also among the lowest total energy structures calculated.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

NMRFAM is supported by National Institutes of Health grant P41GM103399. T.S.B. acknowledges funding from NIH grant R01GM104192. C.D.S. is supported by the Intramural Research Program of the Center for Information Technology at the NIH. We thank C.M. Thiele (Technische Universität Darmstadt) for useful information on the PELG alignment medium.

References

1. Carlomagno T. NMR in natural products: understanding conformation, configuration and receptor interactions. *Nat Prod Rep*. 2012; 29:536–554. [PubMed: 22456471]
2. Karplus M, Grant DM. A criterion for orbital hybridization and charge distribution in chemical bonds. *Proc Natl Acad Sci U S A*. 1959; 45:1269–1273. [PubMed: 16590503]
3. Haasnoot CAG, de Leeuw FAAM, Altona C. The relationship between proton-proton NMR coupling constants and substituent electronegativities—I. *Tetrahedron*. 1980; 36:2783–2792.
4. Neuhaus, D., Williamson, MP. *The Nuclear Overhauser Effect in Structural and Conformational Analysis*. VCH; New York: 1989.
5. Bifulco G, Dambruoso P, Gomez-Paloma L, Riccio R. Determination of relative configuration in organic compounds by NMR spectroscopy and computational methods. *Chem Rev*. 2007; 107:3744–3779. [PubMed: 17649982]

6. Zhang F, Adnani N, Vazquez-Rivera E, Braun DR, Tonelli M, Andes DR, Bugni TS. Application of 3D NMR for structure determination of peptide natural products. *J Org Chem.* 2015; 80:8713–8719. [PubMed: 26273993]
7. Wyche TP, Piotrowski JS, Hou Y, Braun D, Deshpande R, McIlwain S, Ong IM, Myers CL, Guzei IA, Westler WM, Andes DR, Bugni TS. Forazoline A: marine-derived polyketide with antifungal in vivo efficacy. *Angew Chem, Int Ed.* 2014; 53:11583–11586.
8. Saupe A, Englert G. High-resolution Nuclear Magnetic Resonance spectra of orientated molecules. *Phys Rev Lett.* 1963; 11:462–464.
9. Tolman JR, Flanagan JM, Kennedy MA, Prestegard JH. Nuclear magnetic dipole interactions in field-oriented proteins: information for structure determination in solution. *Proc Natl Acad Sci U S A.* 1995; 92:9279–9283. [PubMed: 7568117]
10. Tjandra N, Bax A. Direct measurement of distances and angles in biomolecules by NMR in a dilute liquid crystalline medium. *Science.* 1997; 278:1111–1114. [PubMed: 9353189]
11. Bax A, Grishaev A. Weak alignment NMR: a hawkeyed view of biomolecular structure. *Curr Opin Struct Biol.* 2005; 15:563–570. [PubMed: 16140525]
12. Yan J, Delaglio F, Kaerner A, Kline AD, Mo H, Shapiro MJ, Smitka TA, Stephenson GA, Zartler ER. Complete relative stereochemistry of multiple stereocenters using only residual dipolar couplings. *J Am Chem Soc.* 2004; 126:5008–5017. [PubMed: 15080707]
13. Thiele CM. Residual dipolar couplings (RDCs) in organic structure determination. *Eur J Org Chem.* 2008; 2008:5673–5685.
14. Kummerlöwe, G., Luy, B. CHAPTER 4 - Residual dipolar couplings for the configurational and conformational analysis of organic molecules. In: Graham, AW., editor. *Annu Rep NMR Spectrosc.* Academic Press; 2009. p. 193-232.
15. Yan J, Kline AD, Mo H, Shapiro MJ, Zartler ER. A novel method for the determination of stereochemistry in six-membered chairlike rings using residual dipolar couplings. *J Org Chem.* 2003; 68:1786–1795. [PubMed: 12608792]
16. Mangoni A, Esposito V, Randazzo A. Configuration assignment in small organic molecules via residual dipolar couplings. *Chem Commun (Cambridge, U K).* 2003:154–155.
17. Berger R, Courtieu J, Gil RR, Griesinger C, Kock M, Lesot P, Luy B, Merlet D, Navarro-Vazquez A, Reggelin M, Reinscheid UM, Thiele CM, Zweckstetter M. Is enantiomer assignment possible by NMR spectroscopy using residual dipolar couplings from chiral nonracemic alignment media?—A critical assessment. *Angew Chem, Int Ed.* 2012; 51:8388–8391.
18. Polavarapu P. Determination of the absolute configurations of chiral drugs using chiroptical spectroscopy. *Molecules.* 2016; 21:1056.
19. Smith SG, Goodman JM. Assigning stereochemistry to single diastereoisomers by GIAO NMR calculation: the DP4 probability. *J Am Chem Soc.* 2010; 132:12946–12959. [PubMed: 20795713]
20. Smith SG, Paton RS, Burton JW, Goodman JM. Stereostructure assignment of flexible five-membered rings by GIAO 13C NMR calculations: prediction of the stereochemistry of elatinyne. *J Org Chem.* 2008; 73:4053–4062. [PubMed: 18471017]
21. Barone G, Duca D, Silvestri A, Gomez-Paloma L, Riccio R, Bifulco G. Determination of the relative stereochemistry of flexible organic compounds by Ab initio methods: conformational analysis and Boltzmann-averaged GIAO 13C NMR chemical shifts. *Chem - Eur J.* 2002; 8:3240–3245. [PubMed: 12203354]
22. Sun H, Reinscheid UM, Whitson EL, d'Auvergne EJ, Ireland CM, Navarro-Vázquez A, Griesinger C. Challenge of large-scale motion for residual dipolar coupling based analysis of configuration: the case of fibrosterol sulfate A. *J Am Chem Soc.* 2011; 133:14629–14636. [PubMed: 21776994]
23. Fares C, Hassfeld J, Menche D, Carlomagno T. Simultaneous determination of the conformation and relative configuration of archazolide a by using nuclear overhauser effects, J couplings, and residual dipolar couplings. *Angew Chem, Int Ed.* 2008; 47:3722–3726.
24. Navarro-Vazquez A. MSpin-RDC. A program for the use of residual dipolar couplings for structure elucidation of small molecules. *Magn Reson Chem.* 2012; 50(Suppl 1):S73–S79. [PubMed: 23280663]
25. Schwieters CD, Kuszewski JJ, Tjandra N, Clore GM. The Xplor-NIH NMR molecular structure determination package. *J Magn Reson.* 2003; 160:65–73. [PubMed: 12565051]

26. Schuetz A, Murakami T, Takada N, Junker J, Hashimoto M, Griesinger C. RDC-Enhanced NMR Spectroscopy in Structure Elucidation of Sucro-Neolambertellin. *Angew Chem, Int Ed.* 2008; 47:2032–2034.
27. Gayathri C, Tsarevsky NV, Gil RR. Residual dipolar couplings (RDCs) analysis of small molecules made easy: fast and tuneable alignment by reversible compression/relaxation of reusable PMMA Gels. *Chem - Eur J.* 2010; 16:3622–3626. [PubMed: 20209530]
28. Cheng Q, Xiang L, Izumikawa M, Meluzzi D, Moore BS. Enzymatic total synthesis of enterocin polyketides. *Nat Chem Biol.* 2007; 3:557–558. [PubMed: 17704772]
29. Thiele CM. Simultaneous assignment of all diastereotopic protons in strychnine using RDCs: PELG as alignment medium for organic molecules. *J Org Chem.* 2004; 69:7403–7413. [PubMed: 15497963]
30. Trigo-Mourino P, Navarro-Vazquez A, Ying J, Gil RR, Bax A. Structural discrimination in small molecules by accurate measurement of long-range proton-carbon NMR residual dipolar couplings. *Angew Chem, Int Ed.* 2011; 50:7576–7580.
31. Ge HM, Sun H, Jiang N, Qin YH, Dou H, Yan T, Hou YY, Griesinger C, Tan RX. Relative and absolute configuration of vatiparol (1 mg): a novel anti-inflammatory polyphenol. *Chem - Eur J.* 2012; 18:5213–5221. [PubMed: 22434621]
32. Schuttelkopf AW, van Aalten DM. PRODRG: a tool for high-throughput crystallography of protein-ligand complexes. *Acta Crystallogr, Sect D: Biol Crystallogr.* 2004; 60:1355–1363. [PubMed: 15272157]
33. Folmer RH, Hilbers CW, Konings RN, Nilges M. Floating stereospecific assignment revisited: application to an 18 kDa protein and comparison with J-coupling data. *J Biomol NMR.* 1997; 9:245–258. [PubMed: 9229503]
34. Nath N, d'Auvergne EJ, Griesinger C. Longrange residual dipolar couplings: a tool for determining the configuration of small molecules. *Angew Chem, Int Ed.* 2015; 54:12706–12710.
35. Pham TN, Hinchley SL, Rankin DW, Liptaj T, Uhrin D. Determination of sugar structures in solution from residual dipolar coupling constants: methodology and application to methyl beta-D-xylopyranoside. *J Am Chem Soc.* 2004; 126:13100–13110. [PubMed: 15469309]
36. Spartan 10, v 102. Wavefunction Inc; 2011.
37. Wyche, TP., Ramos Alvarenga, RF., Piotrowski, JS., Duster, MN., Warrack, S., Cornilescu, G., De Wolfe, TJ., Hou, Y., Braun, D., Ellis, GA., Simpkins, SW., Nelson, J., Myers, CL., Mori, H., Safdar, N., Markley, JL., Rajski, SR., Bugni, TS. Chemical genomics, structure elucidation, and *in vivo* studies of the marine-derived anticlostridial ecteinamycin. Personal communication.
38. Liu LY, Sun H, Griesinger C, Liu JK. The use of a combination of RDC and chiroptical spectroscopy for determination of the absolute configuration of Fusariumin A from the fungus *Fusarium* sp. *Nat Prod Bioprospect.* 2016; 6:41–48. [PubMed: 26791752]

Figure 1.

(A) DFT 3D models of enterocin with opposite chirality at C5-H5. Green is the correct and red is the incorrect configuration. (B) Correlation plot between experimental $^1D_{CH}$ and their SVD back-calculated values for using the structures in panel A. The lower limit of the RDC experimental errors (~ 0.5 Hz) is approximately the size of the dots.

Figure 2.

(A) Structure of ecteinamycin showing configurational assignment based on NOEs and *J*-couplings. The configurations of stereocenters marked by asterisks and that of stereocenters 8–14 relative to stereocenters 24–34 were unknown. (B) Final structure with PSL-derived relative configuration.