Published in final edited form as:

Curr Treat Options Pediatr. 2017 September ; 3(3): 193–207. doi:10.1007/s40746-017-0092-x.

Neurologic Outcomes Following Care in the Pediatric Intensive Care Unit

Sherrill D. Caprarola, M.D., M.S.¹, Sapna R. Kudchadkar, M.D.², and Melania M. Bembea, M.D., M.P.H., Ph.D.²

¹Department of Pediatrics, Division of Pediatric Cardiology, Baylor College of Medicine/Texas Children's Hospital, 6621 Fannin St, Houston, TX, United States, 77030

²Departments of Anesthesiology and Critical Care Medicine, and Pediatrics, Johns Hopkins University School of Medicine, 1800 Orleans Street, Baltimore, MD, United States, 21287

Abstract

Purpose of review—With increasing survival of children requiring admission to pediatric intensive care units (PICU), neurodevelopmental outcomes of these patients are an area of increased attention. Our goal was to systematically review recently published literature on neurologic outcomes of PICU patients.

Recent Findings—Decline in neurofunctional status occurs in 3%–20% of children requiring PICU care. This proportion varies based on primary diagnosis and severity of illness, with children admitted for primary neurologic diagnosis, children who suffer cardiac arrest or who require invasive interventions during the PICU admission, having worse outcomes. Recent research focuses on early identification and treatment of modifiable risk factors for unfavorable outcomes, and on long-term follow-up that moves beyond global cognitive outcomes and is increasingly including tests assessing multidimensional aspects of neurodevelopment.

Summary—The pediatric critical care research community has shifted focus from survival to survival with favorable neurologic outcomes of children admitted to the PICU.

Keywords

neurologic outcomes; neurodevelopmental outcomes; neurocognitive outcomes; PICU; pediatrics; intensive care unit

Introduction

Advances in science, technology and systems of care in emergency medicine and pediatric critical care medicine over the last several decades have led to significantly improved

Corresponding Author: Melania M. Bembea, M.D., M.P.H., Ph.D., Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University School of Medicine, 1800 Orleans Street, Suite 6321, Baltimore, MD, United States, 21287, Tel: 410-955-6412, Fax: 410-502-5327, mbembea1@jhmi.edu.

Conflict of Interest

Sherrill D. Caprarola and Sapna R. Kudchadkar each declare no potential conflicts of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

survival of children with critical illness (1–3). This trend of improved survival has been widely described in the overall pediatric intensive care unit (PICU) population (1) as well as in children with specific diagnoses, including sepsis (4), cardiac arrest (5,6), acute respiratory distress syndrome (ARDS) (7), and oncologic disorders (8). However, increased survival has also translated to increased morbidity among survivors (2,9). Therefore, post-discharge morbidity, and specifically, new neurologic morbidity, has become a more relevant and meaningful outcome than mortality in the contemporary era of PICU care. In this systematic review of the literature, we discuss results of studies published in the last five years that focus on neurofunctional outcomes following PICU care, as well as some of the standardized outcome measures most commonly used in these studies.

Methods

Electronic searches of PubMed and EMBASE were conducted in January 2017 using heading terms and keywords to define concepts of pediatric critical care and neurologic outcomes (Supplemental Digital Content 1). The search was limited to publications between 2012–2016. Two reviewers (S.D.C., M.M.B) reviewed all citations independently, and disagreements were resolved by a third reviewer (S.R.K.). Eligible studies included children <18 years of age admitted to a PICU, and evaluated neurofunctional outcomes using standardized outcome measure tool(s). Studies were excluded if they met any of the following criteria: (1) subjects were cared for in the neonatal or pediatric cardiac intensive care unit exclusively; (2) included an adult-only population or a mixed population with the inability to separate data for children; (3) case reports or case series with <10 subjects; (4) animal studies; (5) non-original data; or (6) studies published in language other than English.

Results

Search results are presented in Figure 1. The patient populations and/or pathologies evaluated in the 66 studies included in the final review were: general PICU patients (n=15 studies) (9–23), seizures and/or status epilepticus (SE) (n=7) (24–30), traumatic brain injury (TBI) (n=17) (31–47), cardiac arrest (n=22) (48–69), and infectious diseases (n=5) (70–74).

The most commonly used outcome measure was the Pediatric Cerebral Performance Category (PCPC) (75,76). The PCPC was developed to evaluate the global cognitive impairment; it ranges from 1–6 with 1 being normal, 2 mild disability, 3 moderate disability, 4 severe disability, 5 coma or vegetative state and 6 brain death (75,76). In most studies, PCPC is assessed at baseline (preadmission) and at hospital discharge, via medical record review or caregiver interview, and has very good to excellent interrater reliability, making it amenable to research (76). The second most commonly used outcome measure was the Glasgow Outcome Scale (GOS) and its extensions, GOS – Extended (GOS-E) and GOS-E Pediatric Revision (GOS-E Peds) (77–79). The GOS describes a range of disability, ranging from good recovery to death, and is administered as a semi-structured interview (79). Its pediatric version, GOS-E Peds, has recently been validated in children with TBI (78). The full list of outcome measures identified in this systematic review are listed in Table 1. While some variability is to be expected given the wide range of age and pathology of critically-ill

children, the large number of measures used to evaluate short- and long-term outcomes of children after PICU discharge are also reflective of the lack of consensus on a core group of measures that should be used for outcome research in the PICU population.

Neurologic outcomes following PICU care

General PICU Population

A decline in neurofunctional status following pediatric critical illness has been described in multiple recent studies. New morbidity occurs in up to 4.8% PICU patients, with higher rates (7.3%) seen in children with critical neurologic conditions (3,20). In a retrospective analysis of 29,352 admissions to 24 PICUs submitting data to the Virtual PICU Performance System network, 3.4% and 10.3% of participants acquired global cognitive or functional disability, respectively, as measured by PCPC and Pediatric Overall Performance Category (POPC) at the time of hospital discharge (19). Risk factors for acquiring functional and cognitive disability were trauma, oncologic and neurologic primary diagnoses, unscheduled admission to the PICU, need for mechanical ventilation, renal replacement therapy, cardiopulmonary resuscitation or extracorporeal membrane oxygenation (19).

Children admitted to the PICU remain fragile for long periods of time following their critical illness. In a single center prospective observational study of 91 children who required urgent PICU admission and were followed up to 1 month after discharge, 22% of subjects had PCPC ≤ 3 and 28% had poor adaptive behavior functioning (defined as a Vineland Adaptive Behavior Scale – Second Edition [VABS-II] composite score ≤ 70) (16). A greater PCPC at baseline, neurologic illness and longer ECMO duration were associated with worse PCPC (16). A greater PCPC at baseline, longer chest compression duration, lower oxygen saturations, and circulatory compromise on admission were associated with worse VABS-II composite scores (16). In a single center prospective longitudinal cohort of PICU patients followed up to 3 years post-discharge, the Functional Status Scale (FSS) worsened in 37.7% of subjects, remained unchanged in 44.1%, and improved in only 9% of subjects (20). Patients with unfavorable 3-year neurofunctional outcomes were more likely to have required vasoactive medications and mechanical ventilation, and to have had longer duration of mechanical ventilation and PICU length of stay (LOS) (20). As a likely surrogate for severity of illness, LOS is consistently reported as a risk factor for unfavorable neurofunctional outcomes in PICU patients. Prolonged PICU admissions beyond 28 days are followed by moderate or severe disability as defined by a modified Glasgow Outcome Scale in as many as 67% of survivors (13).

Studies obtaining detailed neuropsychological testing in PICU survivors are somewhat limited by sample size, heterogeneity of primary diagnoses, cost, and loss to follow-up. Cohorts developed in Europe suggest that cognition, adaptive behavior, and verbal and visual memory in children who survive critical illness are within normal range, but tend to be significantly lower compared to healthy controls (22,70). Multiple contributing factors have been studied or are currently under study, including glucose control, sedation strategies, overall PICU LOS, and pre-PICU co-morbidities (22,70,72).

The impact of ICU care on development of short and long-term psychiatric symptoms has been well documented in adult and pediatric ICU patients. In the ICU, delirium affects up to 80% of critically ill adults, and a recent international point prevalence study of critically ill children revealed that delirium was present in 25% of all children (80). In children admitted to the PICU greater than six days, prevalence increased to 38% (80). Similar to adults, independent risk factors for delirium in the PICU include benzodiazepines, use of physical restraints and mechanical ventilation (80). PICU length of stay is increased in children with delirium, as is duration of mechanical ventilation (81). Despite the availability of validated screening tools for ICU setting in infants and children of all ages (82–84), however, routine delirium screening is not yet the PICU standard of care (85).

The link between ICU delirium and long-term neurologic outcomes is an emerging area of interest to both the adult and pediatric critical care community. While delirium affects patients during their ICU admission, a strong foundation of evidence on sequelae of ICU survivorship in adults has led to the definition of post-intensive care syndrome, or PICS (86). PICS is defined as “new or worsening impairments in physical, cognitive, or mental health status arising after critical illness and persisting beyond hospitalization.” A systematic review by Herrup et al. synthesizing the available pediatric evidence on morbidities within the construct of PICS found nineteen studies which encompassed the characteristics of PICS, including nine studies on psychological morbidity after PICU admission (87). Anxiety and depression, culminating in posttraumatic stress disorder, has been well documented in adult survivors of critical illness (88). In children, limited evidence suggests post-traumatic stress symptoms after PICU admission, including delusional memories, intrusive thoughts, and avoidance, occur in up to 32% of PICU survivors (18,23). Septic illness, increased numbers of invasive procedures, and benzodiazepines have been implicated as risk factors (18,89–92), and psychoeducational tools have been piloted with some success (12). Pediatric PICS is an important and rapidly growing area of research with results of ongoing multicenter studies expected to be published in the next few years (93). Additionally, nine U.S. PICUs have joined together as part the Society of Critical Care Medicine (SCCM) Pediatric ICU Liberation Collaborative to investigate approaches to multicomponent, interdisciplinary care using the “ABCDEF” bundle to improve outcomes for critically ill children through analgesia (A), ventilator management (B=breathing), sedation titration (C=coordination), delirium (D) prevention and treatment, early mobilization (E) and family engagement (F) (94,95).

Neurocritical Care

As many as 61% of children with neurocritical illness have unfavorable outcomes at hospital discharge, defined as PCPC 4–6 (28). Among children with neurocritical illness, outcomes tend to be worse in children with stroke, cardiac arrest, and SE, compared to TBI (24,27,28).

Seizures and clinical or electrographic SE are reported in 10–40% of children in the PICU [26–29]. An increasing and impressive body of literature on the prevalence and adverse outcomes associated with seizures and SE in the PICU has been published by multidisciplinary groups comprised of pediatric intensivists, neurologists, and neurophysiologists in the last years, as part of the pediatric Status Epilepticus Research

Group (pSERG), the Critical Care Electroencephalography (EEG) Monitoring Research Consortium, the Approaches and Decisions in Acute Pediatric TBI Trial (ADAPT), and others (96–99).

In a retrospective observational cohort of children admitted to the ICU at a single institution, Payne et al recently reported that as many as 93/259 (36%) experienced seizures and 23/259 (9%) experienced SE (26). Children who spent more time per hour with a seizure (increased seizure burden) were younger, had a longer ICU LOS, and had larger decline in PCPC score at discharge compared to baseline (26). Patients with a decline in PCPC at discharge compared to baseline were more likely to have a history of epilepsy, seizure on presentation, increased seizure burden during the ICU stay, unreactive EEG background and a discharge diagnosis of acute brain injury (26). In a distinct cohort of 200 children with acute encephalopathy admitted to the PICU and who underwent continuous EEG monitoring, Topjian et al determined that 84 (42%) had seizures, categorized as electrographic seizures in 41 (20.5%) and electrographic SE in 43 (21.5%) (30). After accounting for potential confounders, electrographic SE was significantly associated with increased risk of mortality and decline in PCPC at hospital discharge compared to baseline (30). Electrographic seizures without SE were not associated with worse outcomes (30).

Electrographic SE appears to be a significant risk factor for unfavorable outcomes beyond the time of hospital discharge. In a cohort of 137 children who were neurodevelopmentally normal prior to PICU admission and who underwent clinically indicated EEG monitoring while critically ill, unfavorable outcomes based on the GOS-E Peds obtained at median time of 2.7 years after PICU discharge were found in 21/60 (35%) of subjects seen in follow-up (25). Electrographic SE during the PICU stay was associated with unfavorable outcomes while electrographic seizures were not (25). Children who experienced electrographic SE during the PICU stay also had a significantly worse quality of life at follow-up (25). In this same cohort, Abend et al reported that, in multivariable analysis, both electrographic seizures and electrographic SE during the PICU stay were significantly associated with lower adaptive behavior global composite scores compared to subjects with no seizures (29).

Similarly, *TBI* represents a significant cause of morbidity and mortality in children admitted to the PICU (31,36,42,100). As shown in other fields, protocolized care and adherence to guidelines, even if of low evidence level, can be associated with improved outcomes. O’Lynnger et al evaluated the institution of protocolized therapy for TBI in a single center and demonstrated a significant decrease in mortality, but no significant difference in GOS scores among survivors, after vs before protocol institution (43). In a study evaluating adherence to the 2003 Pediatric TBI guidelines in children with severe TBI who survived past 48 hours of admission, for every 1% increase in adherence there was a 1% decrease in the chance of a poor outcome at discharge (42). Multiple predictors of unfavorable neurologic outcome following pediatric TBI have been described, including elevated intracranial pressure and decreased cerebral perfusion pressures (37,42,101), low Glasgow Coma Scale (GCS) on admission (40,41,44,46), subarachnoid hemorrhage and abnormal basal cisterns (35,44), and cerebral edema (35,41,46) on neuroimaging. However, several randomized controlled trials in pediatric TBI investigating therapies such as hypothermia (102), decompressive craniectomy (103), or immune-enhancing diet (104), have failed to

improve patient outcomes. Enrolling children in moderate to severe TBI trials has also been shown to be extremely challenging, due to heterogeneity of injury, early availability of guardians for informed consent, and the need for large number of centers requiring significant efforts in training and coordination (105). Using alternative approaches to randomized controlled trials, two large ongoing multicenter studies are starting to challenge paradigms of pediatric trauma systems and clinical care. Preliminary results have already been published, with more publications systematically addressing pediatric trauma guideline adherence (the Pediatric Guideline Adherence Outcomes [PEGASUS] study) and specific aspects of pediatric TBI care (ADAPT Trial; www.ADAPTTrial.org) are expected in the near future (106–110). ADAPT is an observational cohort study of 1000 children comparing the effectiveness of the following therapeutic strategies employed in severe TBI: first-line intracranial pressure management (cerebrospinal fluid diversion and hyperosmolar therapies), mitigation of secondary insults (hyperventilation without intracranial hypertension and hypoxia detection/treatment using interstitial brain oxygen monitoring), and metabolic support (nutritional support and glucose management) (106). Detailed neurodevelopmental follow-up data will be available in survivors from multiple international participating sites (106).

Cardiac Arrest

Major progress has been achieved in the contemporary cardiopulmonary resuscitation of children after cardiac arrest. Using data from the American Heart Association Get With The Guidelines – Resuscitation (AHA GWTG-R) registry, Berg et al demonstrated in a study published in 2013 that the ratio of PICU vs general ward cardiopulmonary resuscitation events in 315 U.S. hospitals has significantly increased over the study period of 2000–2010, a phenomenon thought to be linked to the successful implementation of rapid response teams in hospitals around the country, with earlier recognition and transfer of patients with deteriorating clinical status to the PICU (111,112). A parallel increase in survival and survival with good neurologic function has also been described (5,6,51,112). In a 1997 study conducted in 32 North-American PICUs, the incidence of cardiopulmonary resuscitation events >2 minutes in PICU patients was 205/11,165 (1.8%) – only 13.7% survived to hospital discharge (5). While the incidence of cardiac arrest in the PICU has not changed significantly since the 1990s, outcomes have certainly seen a remarkable improvement. In a recently published large prospective multicenter study that took place at 11 National Institute of Child Health and Human Development's Collaborative Pediatric Critical Care Research Network (NICHD/CPCCRN) academic children's hospitals, the incidence of cardiac arrest in the PICU was similar, 139/10078 (1.4%) (51). However, of the 139 children who required cardiopulmonary resuscitation 1 minute and/or defibrillation in the PICU, 78% had return of circulation, 45% survived to hospital discharge, and 89% of survivors had favorable neurologic outcome, defined as PCPC at hospital discharge 1–3 or no decline from baseline PCPC (51).

In contrast, out-of-hospital pediatric cardiac arrest is still fraught with poor outcomes. The results of a major randomized controlled trial of therapeutic hypothermia after out-of-hospital pediatric cardiac arrest (THAPCA-OH), were published in 2015 (67). There were 295 children who required cardiopulmonary resuscitation for 2 minutes, were comatose

and mechanically ventilated after return of circulation, and who were randomized to therapeutic hypothermia vs therapeutic normothermia (67). Compared to therapeutic normothermia, therapeutic hypothermia did not improve survival with good functional outcome at 1 year post-arrest (67). Favorable neurobehavioral outcome at 1 year post-arrest, defined as VABS-II composite score >70, was present in 27/138 (20%) subjects in the hypothermia group and 15/112 (12%) subject in the normothermia group (67). Risk factors for poor long-term neurobehavioral function were acute life threatening event/sudden unexpected infant death, respiratory causes of arrest other than drowning, unknown causes of arrest, and duration of chest compressions >30 minutes (68). There are significant ongoing efforts to optimize emergency medical systems of care and therapies aimed at improving outcomes after pediatric out-of-hospital cardiac arrest. Successes are starting to be reported in increased utilization of bystander cardiopulmonary resuscitation leading to improved overall survival and survival with favorable neurologic outcomes (69).

Drowning is the leading cause of accidental death among toddlers and continues carry high morbidity in some children who survive (66,113). Our systematic review identified four articles published in the last five years which examined outcomes after drowning. Overall mortality from drowning remains high, recently reported as 17–73% (65,66,113). Unfavorable outcome was associated with longer CPR time (64,66,113), CPR by emergency medical providers vs early initiation by bystander (65), more than 4 doses of epinephrine needed during cardiac arrest (64), lower base deficit and pH (65,113), longer duration of mechanical ventilation (65), longer PICU LOS (65), and multiple failing organs (66). Favorable outcomes have been associated with drowning in the winter (113).

Submersion for <10 minutes and drowning in water <6°C has been associated with a normal full scale intelligence quotient (65). Having a low intelligent quotient at median follow up of 8.5 years was associated with requiring CPR by emergency medical providers compared to those who received CPR only by a bystander (65). In the THAPCA-OH trial, children who suffered drowning and required CPR for >30 minutes vs 30 minutes had significantly lower VABS-II composite scores (median 41 vs 91) (64). VABS-II is a caregiver report measure developed to assess communication, daily living, social, and motor domains of adaptive behavior (114). The age-corrected composite score has a mean of 100 (standard deviation, 15) in the general population (114).

Infectious Diseases

A large point prevalence study conducted in 128 international PICUs and published in 2015 showed that pediatric sepsis remains a significant public health challenge around the world (74). In this study, 569 of 6925 PICU patients (8.2%), met diagnostic criteria for severe sepsis (74). Hospital mortality was 25% and 17% of survivors had acquired moderate disability at hospital discharge, defined as discharge POPC 3 and increased 1 from baseline POPC (74). Recent studies of long-term outcomes of children with severe sepsis showed that, at 3–9 months post-discharge, children with sepsis-associated encephalopathy had significantly decreased global and verbal intelligence quotient, compared to control sepsis patients not admitted to the PICU and to healthy controls (70–72). Evaluation of academic performance also revealed that children with sepsis who required PICU care had

significantly higher rates of difficulties in completing school work, deterioration in academic performance, dividing attention and considering a variety of possible outcomes (70).

Acute respiratory distress syndrome

Outcomes of pediatric ARDS survivors have been comprehensively and systematically summarized by the Pediatric Acute Lung Injury Consensus Conference (PALICC) group (7). In a multicenter randomized controlled trial of prone positioning, 16% of children who survived ARDS showed a decline in POPC and 11% showed a decline in PCPC at hospital discharge (115). However, long-term outcomes of pediatric ARDS are difficult to ascertain given that most PICU studies do not separate children who meet ARDS diagnostic criteria (7). The 2015 PALICC recommendations on long-term outcome assessment of children with ARDS of any etiology are to conduct physical, neurocognitive, emotional, family, and social function evaluation within 3 months of hospital discharge in children with moderate-to-severe pediatric ARDS, and that the same evaluation should be conducted prior to school start in children who developed pediatric ARDS as infants or toddlers (116). Future studies of neurocognitive outcomes after pediatric ARDS were recommended with strong agreement (116). An ongoing large multicenter study is aiming to determine the association between sedative exposure and neurocognitive function at 2.5 years post-ICU discharge, in 500 subjects that had been enrolled in the Randomized Evaluation of Sedation Titration for Respiratory Failure (RESTORE) study (117,118).

Conclusions

Decline in neurofunctional status occurs in up to a quarter of children requiring PICU care. This proportion varies based on primary diagnosis and severity of illness, with children admitted for primary neurologic diagnoses, children who suffer cardiac arrest or who require invasive interventions during the PICU admission, having worse outcomes. Recent research focuses on early identification and treatment of modifiable risk factors for unfavorable outcomes, and on long-term follow-up that moves beyond global cognitive outcomes and is increasingly relying on tests assessing multidimensional aspects of neurodevelopment.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Melania M. Bembea reports support for this work included funding from: the National Institute of Neurological Disorders and Stroke of the National Institutes of Health under Award Number K23NS076674 (MMB).

References

Important references •

1. Straney LD, Schlapbach LJ, Yong G, Bray JE, Millar J, Slater A, et al. Trends in PICU Admission and Survival Rates in Children in Australia and New Zealand Following Cardiac Arrest. *Pediatr Crit Care Med*. 2015 Sep; 16(7):613–620. [PubMed: 25901547]

2. Namachivayam P, Shann F, Shekerdemian L, Taylor A, van Sloten I, Delzoppo C, et al. Three decades of pediatric intensive care: Who was admitted, what happened in intensive care, and what happened afterward. *Pediatr Crit Care Med*. 2010 Sep; 11(5):549–555. [PubMed: 20124947]
3. Pollack MM, Holubkov R, Funai T, Clark A, Berger JT, Meert K, et al. PMC4221484; Pediatric intensive care outcomes: development of new morbidities during pediatric critical care. *Pediatr Crit Care Med*. 2014 Nov; 15(9):821–827. [PubMed: 25226501]
4. Ruth A, McCracken CE, Fortenberry JD, Hall M, Simon HK, Hebbar KB. Pediatric severe sepsis: current trends and outcomes from the Pediatric Health Information Systems database. *Pediatr Crit Care Med*. 2014 Nov; 15(9):828–838. [PubMed: 25226500]
5. Slonim AD, Patel KM, Ruttimann UE, Pollack MM. Cardiopulmonary resuscitation in pediatric intensive care units. *Crit Care Med*. 1997 Dec; 25(12):1951–1955. [PubMed: 9403741]
6. Girotra S, Spertus JA, Li Y, Berg RA, Nadkarni VM, Chan PS, et al. Survival trends in pediatric in-hospital cardiac arrests: an analysis from Get With the Guidelines-Resuscitation. *Circ Cardiovasc Qual Outcomes*. 2013 Jan 1; 6(1):42–49. [PubMed: 23250980]
7. Quasney MW, Lopez-Fernandez Y, Santschi M, Watson RS. The outcomes of children with pediatric acute respiratory distress syndrome: proceedings from the Pediatric Acute Lung Injury Consensus Conference. *Pediatr Crit Care Med*. 2015 Jun; 16(5):S118–31. [PubMed: 26035362]
8. Dalton HJ, Slonim AD, Pollack MM. MultiCenter outcome of pediatric oncology patients requiring intensive care. *Pediatr Hematol Oncol*. 2003 Dec; 20(8):643–649. [PubMed: 14578035]
9. Pollack MM, Holubkov R, Funai T, Clark A, Berger JT, Meert K, et al. Pediatric intensive care outcomes: Development of new morbidities during pediatric critical care. *Pediatric Critical Care Medicine*. 2014; 15(9):821–827. [PubMed: 25226501]
10. Colville GA, Pierce CM. Children's self-reported quality of life after intensive care treatment. *Pediatr Crit Care Med*. 2013 Feb; 14(2):e85–92. [PubMed: 23337805]
11. Cunha F, Mota T, Teixeira-Pinto A, Carvalho L, Estrada J, Marques A, et al. Factors associated with health-related quality of life changes in survivors to pediatric intensive care. *Pediatr Crit Care Med*. 2013 Jan; 14(1):e8–15. [PubMed: 23249784]
12. Als LC, Nadel S, Cooper M, Vickers B, Garralda ME. PMC4710808; A supported psychoeducational intervention to improve family mental health following discharge from paediatric intensive care: feasibility and pilot randomised controlled trial. *BMJ Open*. 2015 Dec 29; 5(12):e009581.
13. Namachivayam P, Taylor A, Montague T, Moran K, Barrie J, Delzoppo C, et al. Long-stay children in intensive care: long-term functional outcome and quality of life from a 20-yr institutional study. *Pediatr Crit Care Med*. 2012 Sep; 13(5):520–528. [PubMed: 22805156]
14. Als LC, Tennant A, Nadel S, Cooper M, Pierce CM, Garralda ME. Persistence of Neuropsychological Deficits Following Pediatric Critical Illness. *Crit Care Med*. 2015 Aug; 43(8):e312–5. [PubMed: 25962081]
15. Verstraete S, Vanhorebeek I, Covaci A, Guiza F, Malarvannan G, Jorens PG, et al. Circulating phthalates during critical illness in children are associated with long-term attention deficit: a study of a development and a validation cohort. *Intensive Care Med*. 2016 Mar; 42(3):379–392. [PubMed: 26667027]
16. Ebrahim S, Singh S, Hutchison JS, Kulkarni AV, Sananes R, Bowman KW, et al. Adaptive behavior, functional outcomes, and quality of life outcomes of children requiring urgent ICU admission. *Pediatr Crit Care Med*. 2013 Jan; 14(1):10–18. [PubMed: 23132399]
17. Polic B, Mestrovic J, Markic J, Mestrovic M, Capkun V, Utrobicic I, et al. Long-term quality of life of patients treated in paediatric intensive care unit. *Eur J Pediatr*. 2013 Jan; 172(1):85–90. [PubMed: 23052616]
18. Stowman S, Kearney CA, Daphtary K. Mediators of Initial Acute and Later Posttraumatic Stress in Youth in a PICU. *Pediatr Crit Care Med*. 2015 May; 16(4):e113–8. [PubMed: 21926659]
19. Bone MF, Feinglass JM, Goodman DM. Risk factors for acquiring functional and cognitive disabilities during admission to a PICU. *Pediatric Critical Care Medicine*. 2014; 15(7):640–648. [PubMed: 25072478]
20. Pinto NP, Rhinesmith EW, Kim TY, Ladner PH, Pollack MM. Long-Term Function After Pediatric Critical Illness: Results From the Survivor Outcomes Study. *Pediatr Crit Care Med*. 2017 Jan 19.

21. Pollack MM, Holubkov R, Funai T, Clark A, Moler F, Shanley T, et al. PMC4589215; Relationship between the functional status scale and the pediatric overall performance category and pediatric cerebral performance category scales. *JAMA Pediatr.* 2014 Jul; 168(7):671–676. [PubMed: 24862461]
22. Mesotten D, Gielen M, Sterken C, Claessens K, Hermans G, Vlasselaers D, et al. Neurocognitive development of children 4 years after critical illness and treatment with tight glucose control: a randomized controlled trial. *JAMA.* 2012 Oct 24; 308(16):1641–1650. [PubMed: 23101118]
23. Vet NJ, de Wildt SN, Verlaet CW, Mooij MG, Tibboel D, de Hoog M, et al. Short-Term Health-Related Quality of Life of Critically Ill Children Following Daily Sedation Interruption. *Pediatr Crit Care Med.* 2016 Nov; 17(11):e513–e520. [PubMed: 27662565]
24. Nishiyama M, Nagase H, Tanaka T, Fujita K, Maruyama A, Toyoshima D, et al. Demographics and outcomes of patients with pediatric febrile convulsive status epilepticus. *Pediatr Neurol.* 2015 May; 52(5):499–503. [PubMed: 25769239]
25. Wagenman KL, Blake TP, Sanchez SM, Schultheis MT, Radcliffe J, Berg RA, et al. PMC3917684; Electrographic status epilepticus and long-term outcome in critically ill children. *Neurology.* 2014 Feb 04; 82(5):396–404. [PubMed: 24384638]
26. Payne ET, Zhao XY, Frndova H, McBain K, Sharma R, Hutchison JS, et al. PMC3999716; Seizure burden is independently associated with short term outcome in critically ill children. *Brain.* 2014 May; 137:1429–1438. [PubMed: 24595203]
27. Nishiyama M, Nagase H, Tanaka T, Fujita K, Kusumoto M, Kajihara S, et al. Short and long-term outcomes in children with suspected acute encephalopathy. *Brain Dev.* 2016 Sep; 38(8):731–737. [PubMed: 26952815]
28. Nenadovic V, Perez Velazquez JL, Hutchison JS. PMC3994059; Phase synchronization in electroencephalographic recordings prognosticates outcome in paediatric coma. *PLoS One.* 2014; 9(4):e94942. [PubMed: 24752289]
29. Abend NS, Wagenman KL, Blake TP, Schultheis MT, Radcliffe J, Berg RA, et al. PMC4536172; Electrographic status epilepticus and neurobehavioral outcomes in critically ill children. *Epilepsy Behav.* 2015 Aug; 49:238–244. [PubMed: 25908325]
30. Topjian AA, Gutierrez-Colina A, Sanchez SM, Berg RA, Friess SH, Dlugos DJ, et al. Electrographic status epilepticus is associated with mortality and worse short-term outcome in critically ill children. *Crit Care Med.* 2013; 41(1):215–223. [PubMed: 23164815]
31. Emami P, Czorlich P, Fritzsche FS, Westphal M, Rueger JM, Lefering R, et al. Impact of Glasgow Coma Scale score and pupil parameters on mortality rate and outcome in pediatric and adult severe traumatic brain injury: a retrospective, multicenter cohort study. *J Neurosurg.* 2016 Apr 01; 1–8.
32. Mhanna MJ, Mallah WE, Verrees M, Shah R, Super DM. Outcome of children with severe traumatic brain injury who are treated with decompressive craniectomy. *J Neurosurg Pediatr.* 2015 Jul 31; 1–7.
33. Homolkova H, Prchlik M, Tomek P. The relationship between S100B protein serum levels, injury severity and Glasgow Outcome Scale values in children with CNS injuries. *Neuro Endocrinol Lett.* 2012; 33(2):207–211. [PubMed: 22592203]
34. Au AK, Aneja RK, Bell MJ, Bayir H, Feldman K, Adelson PD, et al. PMC3408241; Cerebrospinal fluid levels of high-mobility group box 1 and cytochrome C predict outcome after pediatric traumatic brain injury. *J Neurotrauma.* 2012 Jul 20; 29(11):2013–2021. [PubMed: 22540160]
35. Bennett TD, Dixon RR, Kartchner C, DeWitt PE, Sierra Y, Ladell D, et al. PMC5138132; Functional Status Scale in Children With Traumatic Brain Injury: A Prospective Cohort Study. *Pediatr Crit Care Med.* 2016 Dec; 17(12):1147–1156. [PubMed: 27753754]
36. Beca J, McSharry B, Erickson S, Yung M, Schibler A, Slater A, et al. Hypothermia for Traumatic Brain Injury in Children-A Phase II Randomized Controlled Trial. *Crit Care Med.* 2015 Jul; 43(7):1458–1466. [PubMed: 25803648]
37. Miller Ferguson N, Shein SL, Kochanek PM, Luther J, Wisniewski SR, Clark RS, et al. PMC4856573; Intracranial Hypertension and Cerebral Hypoperfusion in Children With Severe Traumatic Brain Injury: Thresholds and Burden in Accidental and Abusive Insults. *Pediatr Crit Care Med.* 2016 May; 17(5):444–450. [PubMed: 27028792]

38. Falk AC. Impact of elevated ICP on outcome after paediatric traumatic brain injury requiring intensive care. *Childs Nerv Syst.* 2012 Jul; 28(7):1069–1075. [PubMed: 22573139]
39. Sigurt AA, Zanaboni C, Canavesi K, Citerio G, Beretta L, Stocchetti N. Intensive care for pediatric traumatic brain injury. *Intensive Care Med.* 2013; 39(1):129–136. [PubMed: 23179331]
40. Chin KH, Bell MJ, Wisniewski SR, Balasubramani GK, Kochanek PM, Beers SR, et al. Effect of Administration of Neuromuscular Blocking Agents in Children with Severe Traumatic Brain Injury on Acute Complication Rates and Outcomes: A Secondary Analysis from a Randomized, Controlled Trial of Therapeutic Hypothermia. *Pediatric Critical Care Medicine.* 2015; 16(4):352–358. [PubMed: 25599147]
41. Lewis PM, Czosnyka M, Carter BG, Rosenfeld JV, Paul E, Singhal N, et al. Cerebrovascular pressure reactivity in children with traumatic brain injury. *Pediatric Critical Care Medicine.* 2015; 16(8):739–745. [PubMed: 26132743]
42. Vavilala MS, Kernic MA, Wang J, Kannan N, Mink RB, Wainwright MS, et al. PMC4167478; Acute care clinical indicators associated with discharge outcomes in children with severe traumatic brain injury. *Crit Care Med.* 2014 Oct; 42(10):2258–2266. [PubMed: 25083982]
43. O'Lynn TM, Shannon CN, Le TM, Greeno A, Chung D, Lamb FS, et al. Standardizing ICU management of pediatric traumatic brain injury is associated with improved outcomes at discharge. *J Neurosurg Pediatr.* 2016 Jan; 17(1):19–26. [PubMed: 26451717]
44. Lin C, Wang N, Shen ZP, Zhao ZY. Plasma copeptin concentration and outcome after pediatric traumatic brain injury. *Peptides.* 2013 Apr; 42:43–47. [PubMed: 23402790]
45. Young AM, Guilfoyle MR, Fernandes H, Garnett MR, Agrawal S, Hutchinson PJ. The application of adult traumatic brain injury models in a pediatric cohort. *J Neurosurg Pediatr.* 2016 Nov; 18(5): 558–564. [PubMed: 27564785]
46. Rhine T, Wade SL, Makoroff KL, Cassedy A, Michaud LJ. Clinical predictors of outcome following inflicted traumatic brain injury in children. *J Trauma Acute Care Surg.* 2012 Oct; 73(4):S248–53. [PubMed: 23026962]
47. Hasbani DM, Topjian AA, Friess SH, Kilbaugh TJ, Berg RA, Christian CW, et al. PMC4082326; Nonconvulsive electrographic seizures are common in children with abusive head trauma*. *Pediatr Crit Care Med.* 2013 Sep; 14(7):709–715. [PubMed: 23842589]
48. Topjian AA, French B, Sutton RM, Conlon T, Nadkarni VM, Moler FW, et al. PMC4085105; Early postresuscitation hypotension is associated with increased mortality following pediatric cardiac arrest. *Crit Care Med.* 2014 Jun; 42(6):1518–1523. [PubMed: 24561563]
49. Abend NS, Topjian AA, Kessler SK, Gutierrez-Colina A, Berg RA, Nadkarni V, et al. PMC3264396; Outcome prediction by motor and pupillary responses in children treated with therapeutic hypothermia after cardiac arrest. *Pediatr Crit Care Med.* 2012 Jan; 13(1):32–38. [PubMed: 21499174]
50. Starling RM, Shekdar K, Licht D, Nadkarni VM, Berg RA, Topjian AA. PMC4497876; Early Head CT Findings Are Associated With Outcomes After Pediatric Out-of-Hospital Cardiac Arrest. *Pediatr Crit Care Med.* 2015 Jul; 16(6):542–548. [PubMed: 25844694]
51. Berg RA, Nadkarni VM, Clark AE, Moler F, Meert K, Harrison RE, et al. Incidence and Outcomes of Cardiopulmonary Resuscitation in PICUs. *Crit Care Med.* 2016 Apr; 44(4):798–808. [PubMed: 26646466]
52. Michiels EA, Dumas F, Quan L, Selby L, Copass M, Rea T. Long-term outcomes following pediatric out-of-hospital cardiac arrest. *Pediatric Critical Care Medicine.* 2013; 14(8):755–760. [PubMed: 23925145]
53. Lee JK, Brady KM, Chung SE, Jennings JM, Whitaker EE, Aganga D, et al. PMC4164565; A pilot study of cerebrovascular reactivity autoregulation after pediatric cardiac arrest. *Resuscitation.* 2014 Oct; 85(10):1387–1393. [PubMed: 25046743]
54. Topjian AA, Sanchez SM, Shults J, Berg RA, Dlugos DJ, Abend NS. PMC5201170; Early Electroencephalographic Background Features Predict Outcomes in Children Resuscitated From Cardiac Arrest. *Pediatr Crit Care Med.* 2016 Jun; 17(6):547–557. [PubMed: 27097270]
55. Eltayeb AA, Monazea EM, Elsayeh KI. Cardiac Arrest in Children: Relation to Resuscitation and Outcome. *Indian J Pediatr.* 2015 Jul; 82(7):612–618. [PubMed: 25604246]

56. Akahane M, Tanabe S, Ogawa T, Koike S, Horiguchi H, Yasunaga H, et al. Characteristics and outcomes of pediatric out-of-hospital cardiac arrest by scholastic age category. *Pediatric Critical Care Medicine*. 2013; 14(2):130–136. [PubMed: 23314182]
57. van Zelle L, Utens EM, Legerstee JS, Cransberg K, Hulst JM, Tibboel D, et al. Cardiac Arrest in Children: Long-Term Health Status and Health-Related Quality of Life. *Pediatr Crit Care Med*. 2015 Oct; 16(8):693–702. [PubMed: 26020858]
58. van Zelle L, Utens EM, Madderom M, Legerstee JS, Aarsen F, Tibboel D, et al. PMC4908156; Cardiac arrest in infants, children, and adolescents: long-term emotional and behavioral functioning. *Eur J Pediatr*. 2016 Jul; 175(7):977–986. [PubMed: 27177532]
59. Fink EL, Berger RP, Clark RS, Watson RS, Angus DC, Richichi R, et al. PMC4478619; Serum biomarkers of brain injury to classify outcome after pediatric cardiac arrest*. *Crit Care Med*. 2014 Mar; 42(3):664–674. [PubMed: 24164954]
60. Ostendorf AP, Hartman ME, Friess SH. PMC5189632; Early Electroencephalographic Findings Correlate With Neurologic Outcome in Children Following Cardiac Arrest. *Pediatr Crit Care Med*. 2016 Jul; 17(7):667–676. [PubMed: 27164188]
61. Del Castillo J, Lopez-Herce J, Canadas S, Matamoros M, Rodriguez-Nunez A, Rodriguez-Calvo A, et al. Cardiac arrest and resuscitation in the pediatric intensive care unit: a prospective multicenter multinational study. *Resuscitation*. 2014 Oct; 85(10):1380–1386. [PubMed: 25008138]
62. Oualha M, Gatterre P, Boddaert N, Dupic L, De SB, Hubert P, et al. Early diffusion-weighted magnetic resonance imaging in children after cardiac arrest may provide valuable prognostic information on clinical outcome. *Intensive Care Med*. 2013 Jul; 39(7):1306–1312. [PubMed: 23632740]
63. Kieboom JK, Verkade HJ, Burgerhof JG, Bierens JJ, Van Rheenen PF, Kneyber MC, et al. Outcome after resuscitation beyond 30 minutes in drowned children with cardiac arrest and hypothermia: Dutch nationwide retrospective cohort study. *BMJ (Online)*. 2015:350.
64. Moler FW, Hutchison JS, Nadkarni VM, Silverstein FS, Meert KL, Holubkov R, et al. Targeted temperature management after pediatric cardiac arrest due to drowning: Outcomes and complications. *Pediatric Critical Care Medicine*. 2016; 17(8):712–720. [PubMed: 27362855]
65. Suominen PK, Sutinen N, Valle S, Olkkola KT, Lnnqvist T. Neurocognitive long term follow-up study on drowned children. *Resuscitation*. 2014; 85(8):1059–1064. [PubMed: 24709615]
66. Mtaweh H, Kochanek PM, Carcillo JA, Bell MJ, Fink EL. PMC4404206; Patterns of multiorgan dysfunction after pediatric drowning. *Resuscitation*. 2015 May; 90:91–96. [PubMed: 25703785]
67. Moler FW, Silverstein FS, Holubkov R, Slomine BS, Christensen JR, Nadkarni VM, et al. Therapeutic hypothermia after out-of-hospital cardiac arrest in children. *N Engl J Med*. 2015 May 14; 372(20):1898–1908. This was a major multicenter randomized controlled trial comparing therapeutic hypothermia vs therapeutic normothermia in children who were comatose following out-of-hospital cardiac arrest (THAPCA-OH). Therapeutic hypothermia did not confer a significant benefit in survival with a good functional outcome at 1 year, defined as a Vineland Adaptive Behavior Scale – Second Edition (VABS-II) composite score 70. Multiple secondary analyses are expected to improve understanding of risk factors for unfavorable short- and long-term neurocognitive outcomes in children who suffer out-of-hospital cardiac arrest. [PubMed: 25913022]
68. Meert KL, Telford R, Holubkov R, Slomine BS, Christensen JR, Dean JM, et al. Pediatric Out-of-Hospital Cardiac Arrest Characteristics and Their Association With Survival and Neurobehavioral Outcome. *Pediatr Crit Care Med*. 2016 Dec; 17(12):e543–e550. [PubMed: 27679965]
69. Naim MY, Burke RV, McNally BF, Song L, Griffis HM, Berg RA, et al. Association of Bystander Cardiopulmonary Resuscitation With Overall and Neurologically Favorable Survival After Pediatric Out-of-Hospital Cardiac Arrest in the United States: A Report From the Cardiac Arrest Registry to Enhance Survival Surveillance Registry. *JAMA Pediatr*. 2017 Feb 1; 171(2):133–141. [PubMed: 27837587]
70. Als LC, Nadel S, Cooper M, Pierce CM, Sahakian BJ, Garraida ME. Neuropsychologic function three to six months following admission to the PICU with meningoencephalitis, sepsis, and other disorders: a prospective study of school-aged children. *Crit Care Med*. 2013 Apr; 41(4):1094–1103. [PubMed: 23385103]

71. Kaur J, Singhi P, Singhi S, Malhi P, Saini AG. Neurodevelopmental and Behavioral Outcomes in Children With Sepsis-Associated Encephalopathy Admitted to Pediatric Intensive Care Unit: A Prospective Case Control Study. *J Child Neurol*. 2016 May; 31(6):683–690. [PubMed: 26500243]
72. van Zelle L, Utens EM, de Wildt SN, Vet NJ, Tibboel D, Buysse C. Analgesia-sedation in PICU and neurological outcome: a secondary analysis of long-term neuropsychological follow-up in meningococcal septic shock survivors*. *Pediatr Crit Care Med*. 2014 Mar; 15(3):189–196. [PubMed: 24366510]
73. Sasaki J, Chegondi M, Raszynski A, Totapally BR. Outcome of children with acute encephalitis and refractory status epilepticus. *J Child Neurol*. 2014 Dec; 29(12):1638–1644. [PubMed: 24413358]
74. Weiss SL, Fitzgerald JC, Pappachan J, Wheeler D, Jaramillo-Bustamante JC, Salloo A, et al. Global epidemiology of pediatric severe sepsis: the sepsis prevalence, outcomes, and therapies study. *Am J Respir Crit Care Med*. 2015 May 15; 191(10):1147–1157. [PubMed: 25734408]
75. Fiser DH. Assessing the outcome of pediatric intensive care. *J Pediatr*. 1992 Jul; 121(1):68–74. [PubMed: 1625096]
76. Fiser DH, Long N, Roberson PK, Hefley G, Zolten K, Brodie-Fowler M. Relationship of pediatric overall performance category and pediatric cerebral performance category scores at pediatric intensive care unit discharge with outcome measures collected at hospital discharge and 1- and 6-month follow-up assessments. *Crit Care Med*. 2000 Jul; 28(7):2616–2620. [PubMed: 10921604]
77. McCauley SR, Wilde EA, Anderson VA, Bedell G, Beers SR, Campbell TF, et al. Recommendations for the use of common outcome measures in pediatric traumatic brain injury research. *J Neurotrauma*. 2012 Mar 1; 29(4):678–705. [PubMed: 21644810]
78. Beers SR, Wisniewski SR, Garcia-Filion P, Tian Y, Hahner T, Berger RP, et al. Validity of a pediatric version of the Glasgow Outcome Scale-Extended. *J Neurotrauma*. 2012 Apr 10; 29(6):1126–1139. [PubMed: 22220819]
79. McMillan T, Wilson L, Ponsford J, Levin H, Teasdale G, Bond M. The Glasgow Outcome Scale - 40 years of application and refinement. *Nat Rev Neurol*. 2016 Aug; 12(8):477–485. [PubMed: 27418377]
80. Traube C, Silver G, Reeder RW, Doyle H, Hegel E, Wolfe HA, et al. Delirium in Critically Ill Children: An International Point Prevalence Study. *Crit Care Med*. 2017 Apr; 45(4):584–590. [PubMed: 28079605]
81. Traube C, Silver G, Gerber LM, Kaur S, Mauer EA, Kerson A, et al. Delirium and Mortality in Critically Ill Children: Epidemiology and Outcomes of Pediatric Delirium. *Crit Care Med*. 2017 May; 45(5):891–898. [PubMed: 28288026]
82. Smith HA, Boyd J, Fuchs DC, Melvin K, Berry P, Shintani A, et al. Diagnosing delirium in critically ill children: Validity and reliability of the Pediatric Confusion Assessment Method for the Intensive Care Unit. *Crit Care Med*. 2011 Jan; 39(1):150–157. [PubMed: 20959783]
83. Smith HA, Gangopadhyay M, Goben CM, Jacobowski NL, Chestnut MH, Savage S, et al. The Preschool Confusion Assessment Method for the ICU: Valid and Reliable Delirium Monitoring for Critically Ill Infants and Children. *Crit Care Med*. 2016 Mar; 44(3):592–600. [PubMed: 26565631]
84. Traube C, Silver G, Kearney J, Patel A, Atkinson TM, Yoon MJ, et al. Cornell Assessment of Pediatric Delirium: a valid, rapid, observational tool for screening delirium in the PICU*. *Crit Care Med*. 2014 Mar; 42(3):656–663. [PubMed: 24145848]
85. Kudchadkar SR, Yaster M, Punjabi NM. Sedation, sleep promotion, and delirium screening practices in the care of mechanically ventilated children: a wake-up call for the pediatric critical care community*. *Crit Care Med*. 2014 Jul; 42(7):1592–1600. [PubMed: 24717461]
86. Needham DM, Davidson J, Cohen H, Hopkins RO, Weinert C, Wunsch H, et al. Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit Care Med*. 2012 Feb; 40(2):502–509. [PubMed: 21946660]
87. Herrup EA, Wiecezorek B, Kudchadkar SR. Characteristics of postintensive care syndrome in survivors of pediatric critical illness: A systematic review. *World J Crit Care Med*. 2017 May 4; 6(2):124–134. [PubMed: 28529914]

88. Parker AM, Sricharoenchai T, Raparla S, Schneck KW, Bienvenu OJ, Needham DM. Posttraumatic stress disorder in critical illness survivors: a metaanalysis. *Crit Care Med*. 2015 May; 43(5):1121–1129. [PubMed: 25654178]
89. Als LC, Picouto MD, Hau SM, Nadel S, Cooper M, Pierce CM, et al. Mental and physical well-being following admission to pediatric intensive care. *Pediatr Crit Care Med*. 2015 Jun; 16(5):e141–9. [PubMed: 25901544]
90. Muranjan MN, Birajdar SB, Shah HR, Sundaraman P, Tullu MS. Psychological consequences in pediatric intensive care unit survivors: the neglected outcome. *Indian Pediatr*. 2008 Feb; 45(2):99–103. [PubMed: 18310787]
91. Rennick JE, Johnston CC, Dougherty G, Platt R, Ritchie JA. Children's psychological responses after critical illness and exposure to invasive technology. *J Dev Behav Pediatr*. 2002 Jun; 23(3):133–144. [PubMed: 12055495]
92. Rennick JE, Morin I, Kim D, Johnston CC, Dougherty G, Platt R. Identifying children at high risk for psychological sequelae after pediatric intensive care unit hospitalization. *Pediatr Crit Care Med*. 2004 Jul; 5(4):358–363. [PubMed: 15215006]
93. AfterPICU.com. Available at: <http://www.afterpicu.com/>. Accessed May 23, 2017.
94. Marra A, Ely EW, Pandharipande PP, Patel MB. The ABCDEF Bundle in Critical Care. *Crit Care Clin*. 2017 Apr; 33(2):225–243. [PubMed: 28284292]
95. Society of Critical Care Medicine. ICU Liberation. Available at: <http://www.iculiberation.org/About/collaborative/Pages/default.aspx> Accessed May 23, 2017.
96. Du Pont-Thibodeau G, Sanchez SM, Jawad AF, Nadkarni VM, Berg RA, Abend NS, et al. Seizure Detection by Critical Care Providers Using Amplitude-Integrated Electroencephalography and Color Density Spectral Array in Pediatric Cardiac Arrest Patients. *Pediatr Crit Care Med*. 2017 Feb 23.
97. Haider HA, Esteller R, Hahn CD, Westover MB, Halford JJ, Lee JW, et al. Sensitivity of quantitative EEG for seizure identification in the intensive care unit. *Neurology*. 2016 Aug 30; 87(9):935–944. [PubMed: 27466474]
98. Kurz JE, Poloyac SM, Abend NS, Fabio A, Bell MJ, Wainwright MS, et al. Variation in Anticonvulsant Selection and Electroencephalographic Monitoring Following Severe Traumatic Brain Injury in Children-Understanding Resource Availability in Sites Participating in a Comparative Effectiveness Study. *Pediatr Crit Care Med*. 2016 Jul; 17(7):649–657. [PubMed: 27243415]
99. Tasker RC, Goodkin HP, Sanchez Fernandez I, Chapman KE, Abend NS, Arya R, et al. Refractory Status Epilepticus in Children: Intention to Treat With Continuous Infusions of Midazolam and Pentobarbital. *Pediatr Crit Care Med*. 2016 Oct; 17(10):968–975. [PubMed: 27500721]
100. Mhanna MJ, El Mallah W, Verrees M, Shah R, Super DM. Outcome of children with severe traumatic brain injury who are treated with decompressive craniectomy. *Journal of Neurosurgery: Pediatrics*. 2015; 16(5):508–514.
101. Hutchison JS, Frndova H, Lo TY, Guerguerian AM, Hypothermia Pediatric Head Injury Trial Investigators, Canadian Critical Care Trials Group. Impact of hypotension and low cerebral perfusion pressure on outcomes in children treated with hypothermia therapy following severe traumatic brain injury: a post hoc analysis of the Hypothermia Pediatric Head Injury Trial. *Dev Neurosci*. 2010; 32(5–6):406–412. [PubMed: 21252486]
102. Hutchison JS, Ward RE, Lacroix J, Hebert PC, Barnes MA, Bohn DJ, et al. Hypothermia therapy after traumatic brain injury in children. *N Engl J Med*. 2008 Jun 5; 358(23):2447–2456. [PubMed: 18525042]
103. Taylor A, Butt W, Rosenfeld J, Shann F, Ditchfield M, Lewis E, et al. A randomized trial of very early decompressive craniectomy in children with traumatic brain injury and sustained intracranial hypertension. *Childs Nerv Syst*. 2001 Feb; 17(3):154–162. [PubMed: 11305769]
104. Briassoulis G, Filippou O, Kanariou M, Papassotiropoulos I, Hatzis T. Temporal nutritional and inflammatory changes in children with severe head injury fed a regular or an immune-enhancing diet: A randomized, controlled trial. *Pediatr Crit Care Med*. 2006 Jan; 7(1):56–62. [PubMed: 16395076]

105. Stanley RM, Johnson MD, Vance C, Bajaj L, Babcock L, Atabaki S, et al. Challenges Enrolling Children Into Traumatic Brain Injury Trials: An Observational Study. *Acad Emerg Med*. 2017 Jan; 24(1):31–39. [PubMed: 27618167]
106. Larsen GY, Schober M, Fabio A, Wisniewski SR, Grant MJ, Shafi N, et al. Structure, Process, and Culture Differences of Pediatric Trauma Centers Participating in an International Comparative Effectiveness Study of Children with Severe Traumatic Brain Injury. *Neurocrit Care*. 2016 Jun; 24(3):353–360. [PubMed: 26627225]
107. Kannan N, Wang J, Mink RB, Wainwright MS, Groner JJ, Bell MJ, et al. Timely Hemodynamic Resuscitation and Outcomes in Severe Pediatric Traumatic Brain Injury: Preliminary Findings. *Pediatr Emerg Care*. 2016 Jul 12.
108. Kernic MA, Rivara FP, Zatzick DF, Bell MJ, Wainwright MS, Groner JJ, et al. Triage of children with moderate and severe traumatic brain injury to trauma centers. *J Neurotrauma*. 2013 Jul 1; 30(13):1129–1136. [PubMed: 23343131]
109. Ostahowski PJ, Kannan N, Wainwright MS, Qiu Q, Mink RB, Groner JJ, et al. Variation in seizure prophylaxis in severe pediatric traumatic brain injury. *J Neurosurg Pediatr*. 2016 Oct; 18(4):499–506. [PubMed: 27258588]
110. Van Cleve W, Kernic MA, Ellenbogen RG, Wang J, Zatzick DF, Bell MJ, et al. National variability in intracranial pressure monitoring and craniotomy for children with moderate to severe traumatic brain injury. *Neurosurgery*. 2013 Nov; 73(5):746–52. discussion 752; quiz 752. [PubMed: 23863766]
111. Chan PS, Jain R, Nallmothu BK, Berg RA, Sasson C. Rapid Response Teams: A Systematic Review and Meta-analysis. *Arch Intern Med*. 2010 Jan 11; 170(1):18–26. [PubMed: 20065195]
112. Berg RA, Sutton RM, Holubkov R, Nicholson CE, Dean JM, Harrison R, et al. Ratio of PICU versus ward cardiopulmonary resuscitation events is increasing. *Crit Care Med*. 2013 Oct; 41(10):2292–2297. [PubMed: 23921270]
113. Kieboom JK, Verkade HJ, Burgerhof JG, Bierens JJ, Rheenen PF, Kneyber MC, et al. PMC4353310; Outcome after resuscitation beyond 30 minutes in drowned children with cardiac arrest and hypothermia: Dutch nationwide retrospective cohort study. *BMJ*. 2015 Feb 10; 350:h418. [PubMed: 25670715]
114. Sparrow, S.Cicchetti, D., Balla, D., editors. *Vineland Adaptive Behavior Scales*. Second. Circle Pines, MN: AGS Publishing; 2005.
115. Curley MA, Hibberd PL, Fineman LD, Wypij D, Shih MC, Thompson JE, et al. Effect of prone positioning on clinical outcomes in children with acute lung injury: a randomized controlled trial. *JAMA*. 2005 Jul 13; 294(2):229–237. [PubMed: 16014597]
116. Pediatric Acute Lung Injury Consensus Conference Group. Pediatric acute respiratory distress syndrome: consensus recommendations from the Pediatric Acute Lung Injury Consensus Conference. *Pediatr Crit Care Med*. 2015 Jun; 16(5):428–439. [PubMed: 25647235]
117. Sedation Strategy and Cognitive Outcome After Critical Illness in Early Childhood (RESTORE-cog). Jun 9. 2015 Available at: <https://clinicaltrials.gov/ct2/show/NCT02225041>. Accessed 03/03, 2017.
118. Curley MA, Wypij D, Watson RS, Grant MJ, Asaro LA, Cheifetz IM, et al. Protocolized sedation vs usual care in pediatric patients mechanically ventilated for acute respiratory failure: a randomized clinical trial. *JAMA*. 2015 Jan 27; 313(4):379–389. This is a multicenter cluster randomized trial comparing a goal-directed sedation protocol to usual care in children with acute respiratory failure on mechanical ventilation in the PICU. The primary outcome of duration of mechanical ventilation was similar in the two arms. The study led to significant advances in understanding the relationship between wakefulness, pain and agitation, and is being followed by long-term follow-up studies of functional outcomes that will help elucidate the relationship between exposure sedatives in the PICU and neurocognitive outcomes. [PubMed: 25602358]

Figure 1.
Study flowchart

Table 1**Outcome Measures Used in Pediatric Critical Care Studies**

Outcome Measure	No. of studies
PCPC	25
GOS*	16
WIQS	6
POPC	5
CBCL	5
FSS	4
CMS	4
BBDT	3
HUI	3
PedsQL	3
VABS-II	2
CHQ	2
WRIT	2
CNTAB	2
ANT	2
IES	2
Other ^b	20

PCPC: Pediatric Cerebral Performance Category, GOS: Glasgow Outcome Scale, WIQS: Wechsler Intelligence Quotient Scales, POPC: Pediatric Overall Performance Category, CBCL: Child Behavior Checklist, FSS: Functional Status Score, CMS: Children's Memory Scale, BBDT: Beery-Buktenica Development test of Visual-Motor Integration, HUI: Health Utilities Index, PedsQL: Pediatric Quality of Life Inventory, VABS-II: Vineland Adaptive Behavior Scale-II, CHQ - Child Health Questionnaire, WRIT: Wide Range Intelligence Test, CNTAB: Cambridge Neuropsychological Test Automated Battery, ANT: Amsterdam Neuropsychological Tasks, IES: Impact of Events

^a GOS includes GOS (n=12 studies), GOS-Extended for Pediatrics (n=2) and modified GOS (n=2).

^b Other outcome measures used in one study each. ABAS: Adaptive Behavior Assessment System, ABQ: A Behavioral Questionnaire, ASDS: Acute Stress Disorder Scale, BDI-II: Beck depression Inventory: II, BRIEF: Behavior Rating Inventory of Executive Function, CPMS: Childhood Psychopathology Measurement Schedule, CSDC: Child Stress Disorders Checklist, CSHQ: Child Sleep Habits Questionnaire, DP3: Development Profile 3, HADS: Hospital Anxiety and Depression Scale, HSUI: Health State Utilities Index, KOSCHI: King's Outcome Scale for Childhood head injury, MAQ: Multidimensional Anxiety Questionnaire, MISIC: Malin's Intelligence scale for Indian Children, PTSDC-C: PTSD Checklist-Civilian, RAH MOF: Royal Alexandra Hospital for Children Measure of Function, SCORE: Subtest 'SCORE' of the Test of Everyday Attention for Children, SDQ: Strength and Difficulties Questionnaire, TRF: Teacher Report Form, YSR: Youth Self Report