CORRESPONDENCE

The Diagnosis and Treatment of Behavioral Disorders in Dementia

by Prof. Dr. med. Torsten Kratz in issue 26/2017

Additional Comments

I have some brief additional comments to the overall very welcome review article (1), the subject matter of which is relevant for most practicing clinicians.

Risperidone is licensed for treating psychotic symptoms—such as increased psychomotor activity, and aggressiveness—only in Alzheimer's dementia, and only in limited dosages and for restricted periods. In off-label use, penalties are to be expected as prescribing is not permitted for doctors practicing under contract with a statutory health insurer. Other atypical antipsychotics are not licensed for this particular indication.

In Parkinsonian manifestations, quetiapine often leads to a deterioration over time, occasionally with the most severe symptoms, especially in Lewy body dementia. Unfortunately, the only antipsychotic that is licensed for this indication is clozapine, but this is promising and usually low in adverse effects. In all experience, the increased effort required in terms of obtaining blood counts can be managed in collaboration with general practitioners.

I think the recommendation to prefer atypical neuroleptic drugs to treat behavioral disorders in dementia is problematic. These drugs are by no means as low in adverse effects as had been hoped initially. There are even indications of comparatively common and, in particular, severe adverse effects for quetiapine (2).

Citalopram and escitalopram should continue not to be given in combination with other drugs that prolong the QT interval. The use of the latter is likely to be considered as the rule rather than the exception in behavioral disorders in dementia (for example, risperidone).

The review article by Wenzel-Seifert et al. can be consulted for the differential indication of psychotropic drugs in the relevant patient population (3). Accordingly, sertraline should be the preferred serotonin reuptake inhibitor.

In my opinion, the fact that doctors employed within the statutory healthcare system are not permitted to prescribe certain drugs because this would constitute off-label use, as well as important contraindications, should be taken more seriously into account in evidence based (S3) guidelines.

DOI: 10.3238/arztebl.2017.0753a

REFERENCES

1. Kratz T: The diagnosis and treatment of behavioral disorders in dementia. Dtsch Arztebl Int 2017; 114: 447–54.
2. Jin H, Shih PA, Golshan S, et al.: Comparison of longer-term safety and effectiveness of four atypical antipsychotics in patients over age 40: a trial using equipoise-stratified randomization. J Clinical Psychiatry 2013; 74: 10–8.
3. Wenzel-Seifert K, Wittmann M, Haen E: QTc prolongation by psychotropic drugs and the risk of torsade de pointes. Dtsch Arztebl Int 2011; 108: 687–93.

Dr. med. Wolfgang J. Stein
Koblenz
wjstein@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Paying Attention to Electrolytes

The author thankfully prioritized in his article non-medicinal treatment options for behavioral disorders in dementia (1).

The fact is that in older patients in particular, whose multimorbidity often results in polypharmacy, additional drugs to treat behavioral disorders are associated with substantial risks. Study data and the S3 guideline for dementia recommend treatment with citalopram in addition to risperidone, which is also licensed for this indication. The guideline describes indications of the efficacy of citalopram in dementia patients displaying agitation with an evidence level of Ib. Of note in this context is the fact that in a *Rote Hand Brief* (technical information disseminated from pharmaceutical companies directly to healthcare professionals) of 2011, simultaneous use of citalopram with drugs that are known to potentially prolong the QT interval is contraindicated. Because of the multitude of substances that prolong the QT interval, the use of citalopram in gerontopsychiatric patients substantially raises the probability that this contraindication is ignored and therefore the risk of developing torsade de pointes tachycardias increases substantially.

Because of its more favorable interaction profile in the cytochrome P-450 system compared with other SSRIs (fluoxetine, fluvoxamine, paroxetine), sertraline should be singled out as an alternative.

Especially in older patients, the risk of electrolyte imbalances in the sense of hyponatremia as a result of treatment with any SSRI should be emphasized, especially at the start of the treatment.

Furthermore, a critical cost-benefit analysis is recommended because of the potentially increased risk of hemorrhage—both gastrointestinal and intracerebral—when SSRIs and non-steroidal anti-inflammatories/acetylsalicylic acid are combined (2, 3). Such combination treatment should be undertaken extremely cautiously in patients at risk of falls, and all non-medicinal options for treatment should be exhausted completely.

DOI: 10.3238/arztebl.2017.0753b

REFERENCES

1. Kratz T: The diagnosis and treatment of behavioral disorders in dementia. Dtsch Arztebl Int 2017; 114: 447–54.
2. Strubel T, Birkhofer A, Mößner G, Förstl H: Blutungsrisiko unter SSRI-Behandlung. Nervenarzt 2010; 81: 549–55.
3. Shin JY, Park MJ, Lee SH, et al.: Risk of intracranial haemorrhage in antidepressant users with concurrent use of non-steroidal anti-inflammatory drugs: nationwide propensity score matched study. BMJ 2015; 351: h3517.

Dr. med. Sara Bientreau
Fachklinik Marienborn gGmbH
Zülpich
s.bientreau@marienborn-zuelpich.de

Conflict of interest statement

Dr Bientreau has received consultancy fees from Janssen-Cilag and Lundbeck. She has received funding for preparing scientific conferences from Janssen-Cilag.

In Reply:

I thank Dr. Bienentreu and Dr. Stein for their important comments regarding the drug treatment of behavioral disorders in dementia. In particular, their mention of the numerous adverse effects and interactions of drugs for behavioral disorders underlines the necessity of critically weighing up the risks and benefits and highlight the ensuring dilemma. In the present article, *Table 2* (Problematic psychoactive drugs in the treatment of behavioral disorders in demented patients) provides relevant details on this, while also taking into account the PRISCUS list (1). Even though the guidelines (2) recommend treatment attempts using drugs, it makes sense to investigate somatic or environmental causes and treat those specifically before starting any such symptomatic treatment. In many cases, high risk medicinal treatment is then no longer needed. For this reason, my article focused on non-drug treatment approaches in the context of behavioral disorders in dementia. The effectiveness of non-drug based approaches can be superior to solely drug-based treatments. Treating agitation and aggressiveness with risperidone achieves an effect size of $d = 0.33$ (2, 3), whereas the use of occupational therapy achieves a greater effect size when used in apathy, restlessness, and agitation—namely, $d = 0.72$ (2, 4). My article was intended to point out that the global effect of non-drug-based therapeutic approaches on behavioral disorders has been proven. Because of the heterogeneity of the interventions, however, it is currently not possible to say which specific approaches should be used in which specific behavioral disorder. More research is needed in this setting. It is undisputable, how-

ever, that non-drug-based therapeutic approaches should be implemented more full scale into daily practice in care/nursing homes and hospitals, so as to improve the quality of life of those affected and their carers, and to avoid polypharmacy and its known risks.

DOI: 10.3238/arztebl.2017.0754a

REFERENCES

1. Kratz T: The diagnosis and treatment of behavioral disorders in dementia. *Dtsch Arztebl Int* 2017; 114: 447–54.
2. Deuschl G, Maier W, Deutsche Gesellschaft für Psychiatrie, Psychotherapie und Nervenheilkunde (DGPPN), Deutsche Gesellschaft für Neurologie (DGN): Diagnose- und Behandlungsleitlinie Demenz – Interdisziplinäre S3 Praxisleitlinien 2016. www.dgn.org/images/red_leitlinien/LL_2016/PDFs_Download/038013_LL_Demenzen_2016.pdf (last accessed on 17 August 2017).
3. Chan WC, Lam LC, Choy CN, Leung VP, Li SW, Chiu HF: A double-blind randomised comparison of risperidone and haloperidol in the treatment of behavioural and psychological symptoms in Chinese dementia patients. *Int J Geriatr Psychiatry* 2001; 16: 1156–62.
4. Gitlin LN, Winter L, Burke J, et al.: Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduce caregiver burden: a randomized pilot study. *Am J Geriatr Psychiatry* 2008; 16: 229–39.

Prof. Dr. med. Torsten Kratz

Abteilung für Psychiatrie, Psychotherapie und Psychosomatik
Funktionsbereich Gerontopsychiatrie
Königin-Elisabeth-Herzberge Krankenhaus, Berlin
T.Kratz@keh-berlin.de

Conflict of interest statement

Prof. Kratz has received lecture honoraria from Lilly and Janssen-Cilag.

CLINICAL SNAPSHOT

Oriental Sore (Old-World Cutaneous Leishmaniasis)

A 23-year-old pregnant woman from Afghanistan complained of a nodule on the left cheek that had been present for six months and had become secondarily ulcerated. Skin biopsy revealed a granulomatous, plasma-cell-rich infiltrate of the entire dermis, with many non-flagellated parasitic inclusions within macrophages. *Leishmania* spp. was demonstrated with the aid of a specific polymerase chain reaction; subspeciation from paraffin-embedded tissue was unsuccessful (native tissue is the gold standard). Systemic pharmacotherapy was relatively contraindicated because of pregnancy, so cryotherapy was tried first. In the ensuing three weeks, marked enlargement of the lesion was seen. Treatment with fluconazole was then initiated but was only able to prevent further progression. The spontaneous course after delivery was benign, and the lesion healed with a scar. This disease, previously encountered in Germany as a travel-associated dermatosis, has become more common here as asylum-seekers from endemic areas may suffer from it. Single lesions on the limbs can easily be controlled with local treatment, but facial lesions are an indication for systemic treatment with azole antifungal agents or miltefosine. Larger and more treatment-resistant lesions are seen in pregnancy because of a lessened Th-1 immune response leading to an impaired ability to fend off parasites.

Lesion on the left cheek

Dr. med. Monica Schüürmann, Prof. Dr. med. Manfred Kunz, Klinik und Poliklinik für Dermatologie, Venerologie und Allergologie, Department für Innere Medizin, Neurologie und Dermatologie, Universitätsklinikum Leipzig

Prof. Dr. med. Christoph Lübbert, D.T.M. & H., Fachbereich Infektions- und Tropenmedizin, Klinik und Poliklinik für Gastroenterologie und Rheumatologie, Department für Innere Medizin, Neurologie und Dermatologie, Universitätsklinikum Leipzig; christoph.luebbert@medizin.uni-leipzig.de

Conflict of interest statement: The authors state that they have no conflict of interest.

Cite this as: Schüürmann M, Kunz M, Lübbert C: Oriental sore (old-world cutaneous leishmaniasis). *Dtsch Arztebl Int* 2017; 114: 754. DOI:10.3238/arztebl.2017.0754b

Translated from the original German by Ethan Taub, MD