Published in final edited form as:

Curr Opin Immunol. 2017 June ; 46: 75–81. doi:10.1016/j.coi.2017.04.009.

Unusual antigen presentation offers new insight into HIV vaccine design

Andrew J McMichael¹ and Louis J Picker²

¹Nuffield Department of Medicine, Oxford University, Old Road Campus, Oxford OX37FZ, UK

²Vaccine and Gene Therapy Institute, Oregon Health & Science University, 505 NW 185th Avenue, Beaverton, OR 97006-3448, United States

Abstract

Recent findings with a rhesus monkey cytomegalovirus based simian immunodeficiency virus vaccine have identified strong CD8+ T cell responses that are restricted by MHC-E. Also mycobacteria specific CD8+ T cells, that are MHC-E restricted, have been identified. MHC-E therefore can present a wide range of epitope peptides to CD8+ T cells, alongside its well defined role in presenting a conserved MHC-class I signal peptide to the NKG2A/C-CD94 receptor on natural killer cells. Here we explore the antigen processing pathways involved in these atypical T cell responses.

Introduction

The current well-established paradigm of T cell epitope targeting has CD4+ and CD8+ T cells recognizing diverse peptide antigens in the context of polymorphic MHC-II and classical MHC-Ia molecules, respectively. These CD4+ and CD8+ T cell responses have specifically evolved to survey the extra-cellular and intra-cellular compartments, respectively, for foreign antigens. Non-classical MHC-Ib molecules are much less polymorphic than MHC-II and MHC-Ia, and typically present non-peptide antigens to semi-innate T cells, often with invariant T cell receptors, or regulate the activity of natural killer (NK) cells [1]. Although these functions are clearly the default pathways in all mammalian immune systems examined in detail, accumulating data has shown important exceptions to these general rules, in particular, the description of CD8+ T cells restricted by MHC-II [2] and MHC-E [3]. Of these observations, the most dramatic is the finding that a modified RhCMV vector strain (68-1) exclusively elicits, in Rhesus macaques, broadly targeted CD8+ T cells that are restricted by either MHC-II or MHC-E, but not MHC-Ia [4,5••]. CD8+ T cell responses to SIVgag inserts in these vectors target an average of 20 diverse epitopes in MHC-E, and 25–30 in MHC-II. This contrasts with less than 20 MHC-Ia restricted T cell responses to SIVgag in rhesus monkeys vaccinated with conventional viral vectors or infected with SIV itself [4,5••]. Moreover, many of the MHC-E- and MHC-II-restricted epitopes recognized by the strain 68-1 RhCMV vector elicited responses are so-called

supertopes, epitopes targeted in 100% of vaccinated animals. This unconventional targeting is not restricted to SIVgag epitopes, but is observed to both endogenous RhCMV proteins and to any exogenous insert engineered into these vectors.

These unusual responses are dependent on genetic alterations of the RhCMV 68-1 vector that occurred during extensive *in vitro* passage of the original RhCMV obtained from animals. In particular, fibroblast adaptation led to loss of expression of Rh157.5 and Rh157.4, which are RhCMV orthologs of the human CMV UL128 and UL130 proteins and contribute to the pentameric receptor complex which is used by both RhCMV and HCMV to facilitate infection of non-fibroblast cells. Restoration of Rh157.5 and Rh157.4 expression in RhCMV 68-1.2 completely reverts the vector-elicited CD8⁺ T cells response to conventional MHC-Ia restriction, similar to the responses elicited by natural (colony circulating) strains of RhCMV [5••]. The mechanisms used by the RhCMV 68-1 to prime MHC-E- and MHC-II-restricted responses is under intense study, as is the role of these unconventional responses and the unique ‘control and clear’ efficacy manifested by 68-1 RhCMV/SIV vectors against highly pathogenic SIVmac239 challenge [6,7]. However, the mere existence of these striking unconventional responses suggests a need for re-evaluation of CD8⁺ T cell recognition paradigms. In particular, the ability of RhCMV 68-1 vectors to elicit CD8⁺ T cells to an average 4 distinct MHC-E-restricted epitopes per 100 amino acids of sequence for diverse proteins (likely any protein) raises the issue of the possible role of non-classical, relatively non-polymorphic, MHC-E molecule in other adaptive CD8⁺ T cell responses.

MHC-E

MHC-E molecules are non-classical class I molecules of limited genetic polymorphism. H-2 Qa1, HLA-E and Mamu-E are all closely related and function similarly. Originally, an alloreactive H-2 Qa1 specific CD8⁺ T cell clone was shown to have specificity for Qa1 plus a peptide derived from the signal sequence of classical H-2D or K molecules [8]. It was then found that HLA-E bound the same peptide derived from the signal sequence of HLA A, B, C and G molecules, but that the sequence was truncated in HLA-E itself [9]. The sequence, known as VL9, typically has the sequence VMAPRTLVL; there is some conservative polymorphism at peptide residues 7 and 8 in different HLA molecules. Also, more than half of the HLA-B molecules have a threonine for methionine substitution at position two, which greatly decreases HLA-E binding [10]. HLA-E tetramers, refolded with the VL9 peptide, were shown to bind to the CD94-NKG2 receptors found on natural killer cells and some T cells [10]. NKG2A delivers an inhibitory signal, binds with higher affinity than NKG2C, which is linked to DAP12 and delivers an activating signal [11]. Thus, HLA-E expression marks the presence or absence of classical HLA class I at the cell surface and makes the cells resistant or susceptible to NK cell attack, respectively. H-2Qa1 and Mamu-E serve the same function.

The crystal structure of HLA-E with the VL9 peptide showed a very good fit of the peptide to the groove with all pockets filled and multiple hydrogen bonds to the peptide backbone [12]. Thus, it appears that the two had co-evolved to control efficiently NK cell activity. Interestingly, there are exact matches to the VL9 peptide in the signal sequence of CMV UL40 and in Rhesus CMV Rh67. These peptides enhance MHC-E surface expression and

inhibit NK cell activity when CMV blocks the natural generation of the classical MHC-derived VL9 [5,13].

The VL9 peptide is present at positions 3–11 in the whole signal peptide (SP) (Figure 1). After translocation of the newly synthesized HLA class I heavy chain into the lumen of the endoplasmic reticulum (ER), the SP is cleaved by the signal peptidase. This releases a 24 amino acid fragment that is further cleaved by the signal peptide peptidase in the membrane of the ER, releasing a 14 amino acid fragment into the cytosol [14]. This peptide is processed by the proteasome, shortened at the carboxy terminus and an eleven amino acid fragment is transported back into the ER by the Transporter Associated with Antigen Processing (TAP) [14]. Finally the two amino terminal amino acids are cleaved by the TAP associated ERAP1 aminopeptidase and the exact VL9 nonamer is generated [15,16]. This binds to HLA-E, releasing the folded stable HLA-E protein from TAP and tapsin allowing it to move to the cell surface via the Golgi [17]. This whole process efficiently puts the VL9 peptide into the HLA-E molecules, where it is probably the most abundant peptide.

Infection with HCMV and RhCMV severely disrupts this process. There are five CMV gene products that directly interfere with the classical antigen processing pathway [18]. US2, US3 and US11 target the nascent heavy chain for degradation after aberrant translocation back into the cytosol. It is unclear how this affects the signal peptide, but US11 has been shown to be important in abrogating the classical canonical MHC class I restricted responses in the RhCMV vaccine experiments [4]. US6 efficiently blocks the TAP peptide transporter. Similar TAP blockade by the Herpes Simplex virus ICP47 protein completely alters the repertoire of peptides that bind to HLA-E [19]. Once transported, the two amino-terminal SP amino acids are removed by the endoplasmic amino-peptidase ERAP-1. Knock-out of ERAAP (the murine equivalent of ERAP-1) allowed alternative immunogenic self peptide(s) to bind to H-2 Qa-1 and elicit T cell responses in mice [16]. HCMV expresses a miRNA in US4 that blocks ERAP1 activity, so probably has the same effect [20].

HCMV and RhCMV compensate for loss of the natural VL9 by expressing UL40 or Rh67 respectively [5•]. Both have the VL9 peptide in their SPs, in proteins that are otherwise completely different. It is clear that both can put VL9, or possibly a longer version, into MHC-E so as to inhibit NK cells. This process is TAP independent, and occurs within the ER, allowing HLA-E to access the cell surface in CMV infected cells.

T cell responses restricted by MHC-E

Several HLA-E-restricted T cell responses have been described (Figure 2) [3,5•,21]. The peptides include self peptides, viral and bacterial peptides. However, when compared to the number of peptide epitopes restricted by classical MHC molecules described in the same time period, they are relatively few. This is consistent with the low level of surface expression of MHC-E, which is 5–10% of that of classical MHC-Ia molecules. Disruption of the classical VL9 pathway may contribute to these T cell responses by forcing HLA-E to utilize alternative antigen processing pathways, whilst other pathogens may enter these pathways directly. The RhCMV 68-1-SIV responses [5•] may be examples of the former, while the mycobacterial responses [3] may be the latter. At least some of these alternative

pathways can put enough peptide into MHC-E molecules to directly prime CD8+ T cell responses.

The HLA-E restricted mycobacterial specific T cell responses were originally identified as targets of a Mtb specific CD8+ T cell clone [22]. Then, using an epitope predictive algorithm, 67 peptide epitopes restricted by HLA-E were discovered and it was shown that nearly all tuberculin-reactive donors responded to several of these [3]. At the time of these writing, the relative proportion of these MHC-E-restricted CD8+ T cell responses relative to classically restricted MTb specific CD8+ T cells is unclear. Mycobacteria reside in the phagolysosomes of infected macrophages and this compartment, which expresses relatively high amounts of HLA-E, is thought to directly process mycobacterial proteins to prime CD8+ T cells [23]. This process is TAP dependent, but does not require newly synthesized HLA-E; instead this is probably recycling from the cell surface. This implies a pathway of low affinity peptide binding, dissociation, recycling and then acquisition of new peptides that can be immunogenic.

There may be lessons here for the RhCMV 68-1 SIV epitopes. These epitopes do not show any simple sequence motif. In addition, the carboxy terminal amino acids are very unusual for epitopes presented by class I MHC molecules. Nearly all classically presented epitopes end in L, I, V, K, R, F, Y or W, residues imposed by the proteasome specificity and the TAP preferences for transportable short peptides [24]. Q, G, E, S, N are very rare in well-characterized, classically processed epitopes (LANL HIV database), but here they appear common suggesting an alternative antigen processing pathway. That may not be dissimilar to the class II pathway where the newly synthesized class II molecule in the ER binds invariant chain, focused on the CLIP sequence, and is then relocated to endosomes. Here, the invariant chain is digested and the CLIP peptide removed, to be replaced with locally generated peptides derived from externally acquired proteins [25]. It is possible that UL40 and Rh67 serve a similar function, alongside their role in inhibiting NK cells. Invariant chain itself could also be involved as it has been shown to bind to class I MHC molecules [26].

Other described HLA-E restricted T cells were specific bacterial antigens and some self peptides. Particularly intriguing are the regulatory H-2 Qa1 restricted CD8T cells that control B cells and T helper cells in lymphoid follicles [27]. These are specific for HSP60 proteins, TCR sequences and possibly other self-antigens. They are lytic in function and regulate both T follicular helper cells and B cells. Similar cells have been described in humans [28] and in both species they appear to be important in preventing autoimmunity. They are of particular interest because T follicular helper cells are a known reservoir for long-term latent HIV infection. HIV is hard to eradicate from this site because there are few classical CD8+ T cells there [29••]. Recently CXCR5 expressing HIV specific T cells were described in lymphoid follicles and were capable of suppressing virus infection there, but they were probably not MHC-E restricted [30••].

Conclusions

In the original paradigm of MHC-E biology, the high affinity, almost 'lock and key', binding of MHC-Ia leader sequence VL9 peptides to MHC-E and the ability of the MHC-E-VL9

complex to regulate NK cell function via CD94-NKG2 receptor signaling suggested a highly specialized role for MHC-E as an innate immune ‘sensor’ of altered MHC-Ia expression [10]. This is an elegant system to distinguish cells in which MHC-Ia expression is compromised, and mark these cells, which are highly likely to be virally infected or neoplastically transformed, for innate immune destruction. Of course, this system would be rather easily compromised by pathogen adaptations that provide for MHC-E expression independent of MHC-Ia expression. However, as described in this review, the mammalian immune system had a second ‘trick up its sleeve’—the ability of non-polymorphic MHC-E to not only bind canonical VL9 peptides, but also to bind diverse pathogen-derived peptides and present them to CD8+ T cells. Thus, pathogens that up-regulate MHC-E to evade NK cells run the risk of priming adaptive MHC-E-restricted CD8+ T cells that would potentially be as, or more, destructive to the pathogen than the originally evaded NK cells!

Wildtype RhCMV, which markedly up-regulates MHC-E expression on infected cells, appears to handle this problem by specific evolutionary adaptations (Rh157.5 and Rh157.4, and likely other virus genes) that almost completely inhibit priming of MHC-E restricted CD8+ T cells, although allelic variants of the VL9 peptide itself can prime T cell responses during wildtype CMV infection [33]. Deletion of Rh157.5 and Rh157.4, and probably some other CMV genes, in RhCMV 68-1 enables the priming of CD8T cell responses to peptides bound to MHC-E, other than VL9. Thus CD8+ T cells elicited by wildtype CMV are only MHC-Ia restricted [5•,34] and probably elicited by both direct priming, before full downregulation of MHC-Ia in infected cells, and then cross priming. Indeed, the latter must also be impaired in RhCMV68-1 infection because no MHC-Ia restricted T cell responses are seen. Inhibition of programmed cell death in the restricted cell type infected by RhCMV 68-1 may limit access of dendritic cells to CMV protein debris [35•,36•]. Alternatively, direct priming of the MHC-E restricted responses may inhibit or simply out-compete cross-priming.

For pathogens that lack the sophisticated immune evasion potential of CMV, the solution may be to limit MHC-E expression to below the levels required for efficient CD8+ T cell priming, so as to achieve partial inhibition of NK cell recognition without awakening adaptive MHC-E restricted T cell responses. HIV and SIV likely belong to this later category of pathogen, down-regulating MHC-Ia on infected cells, modestly up-regulating MHC-E expression, but largely avoiding priming MHC-E-restricted CD8+ T cell responses [5•]. Importantly, the MHC-peptide expression threshold allowing recognition by pre-existing CD8+ T cells (<10 molecules) [31] is considerably less (likely 2–4 orders of magnitude) than that required for initial CD8+ T cell priming [32], and indeed, SIV-infected cells are efficiently recognized by the MHC-E-restricted SIV-specific CD8+ T cells elicited by 68-1 RhCMV/SIV vectors (Figure 3) [5•]. If, as we expect, the efficacy of these vectors is wholly or even partially dependent on their ability to prime MHC E-restricted, pathogen-specific CD 8+ T cells, this protection would essentially be exploiting a pathogen adaptation to evade NK cells.

Thus, for pathogens that evade NK cells by enforcing MHC-E expression, MHC-E expression constitutes a point of immunologic vulnerability that could be exploited by vaccines or other immunotherapeutic approaches. In this situation the non-polymorphism of

MHC-E becomes an asset as MHC-E-restricted CD8⁺ T cell responses are much more likely to be shared among individuals than MHC-Ia-restricted CD8⁺ T cell responses. The potential to exploit MHC E-restricted CD8⁺ T cell responses would extend beyond SIV/HIV to other infections as well as cancer, making the understanding of the mechanisms by which non-canonical peptides are processed and presented on MHC-E and by which MHC-E-restricted CD8⁺ T cells are primed a potentially very exciting research investment, perhaps opening an entirely new category of immunotherapy.

Acknowledgments

We are grateful to many colleagues for discussion of the papers discussed in this review including Persephone Borrow, Nilu Goonetilleke, Simon Brackenridge, Geraldine Gillespie, and Suzanne Campion. AJMcM is supported by Grants from the BMGF (OPP1133649), NIAID, NIH (UM1-AI100645-01), MRC (MR/K012037). LJ Picker is supported by grants from NIH, NIAID (P01-AI094417, R37-AI054292, U19 AI095985, U19 AI096109, and P51-OD011092), and from the Bill and Melinda Gates Foundation (OPP1107409 and OPP1108533).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Adams EJ, Luoma AM. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol.* 2013; 31:529–561. [PubMed: 23298204]
2. Ranasinghe S, Lamothe PA, Soghoian DZ, Kazer SW, Cole MB, Shalek AK, Yosef N, Jones RB, Donaghey F, Nwonu C, et al. Antiviral CD8⁺ T cells restricted by human leukocyte antigen class II exist during natural HIV infection and exhibit clonal expansion. *Immunity.* 2016; 45:917–930. [PubMed: 27760342]
3. Joosten SA, van Meijgaarden KE, van Weeren PC, Kazi F, Geluk A, Savage ND, Drijfhout JW, Flower DR, Hanekom WA, Klein MR, et al. Mycobacterium tuberculosis peptides presented by HLA-E molecules are targets for human CD8 T-cells with cytotoxic as well as regulatory activity. *PLoS Pathog.* 2010; 6:e1000782. [PubMed: 20195504]
4. Hansen SG, Sacha JB, Hughes CM, Ford JC, Burwitz BJ, Scholz I, Gilbride RM, Lewis MS, Gilliam AN, Ventura AB, et al. Cytomegalovirus vectors violate CD8⁺ T cell epitope recognition paradigms. *Science.* 2013; 340:1237874. [PubMed: 23704576]
- 5•• Hansen SG, Wu HL, Burwitz BJ, Hughes CM, Hammond KB, Ventura AB, Reed JS, Gilbride RM, Ainslie E, Morrow DW, et al. Broadly targeted CD8(+) T cell responses restricted by major histocompatibility complex E. *Science.* 2016; 351:714–720. The paper that identified the MHC-I restricted CD8⁺ T cells primed by RhCMV68-1 as being restricted by MHC-E. [PubMed: 26797147]
6. Hansen SG, Ford JC, Lewis MS, Ventura AB, Hughes CM, Coyne-Johnson L, Whizin N, Oswald K, Shoemaker R, Swanson T, et al. Profound early control of highly pathogenic SIV by an effector memory T-cell vaccine. *Nature.* 2011; 473:523–527. [PubMed: 21562493]
7. Hansen SG, Piatak M Jr, Ventura AB, Hughes CM, Gilbride RM, Ford JC, Oswald K, Shoemaker R, Li Y, Lewis MS, et al. Immune clearance of highly pathogenic SIV infection. *Nature.* 2013; 502:100–104. [PubMed: 24025770]
8. Aldrich CJ, DeCloux A, Woods AS, Cotter RJ, Soloski MJ, Forman J. Identification of a Tap-dependent leader peptide recognized by alloreactive T cells specific for a class Ib antigen. *Cell.* 1994; 79:649–658. [PubMed: 7525079]

9. Braud V, Jones EY, McMichael A. The human major histocompatibility complex class Ib molecule HLA-E binds signal sequence-derived peptides with primary anchor residues at positions 2 and 9. *Eur J Immunol.* 1997; 27:1164–1169. [PubMed: 9174606]
10. Braud VM, Allan DS, O'Callaghan CA, Soderstrom K, D'Andrea A, Ogg GS, Lazetic S, Young NT, Bell JI, Phillips JH, et al. HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C. *Nature.* 1998; 391:795–799. [PubMed: 9486650]
11. Kaiser BK, Pizarro JC, Kerns J, Strong RK. Structural basis for NKG2A/CD94 recognition of HLA-E. *Proc Natl Acad Sci U S A.* 2008; 105:6696–6701. [PubMed: 18448674]
12. O'Callaghan CA, Tormo J, Willcox BE, Braud VM, Jakobsen BK, Stuart DI, McMichael AJ, Bell JI, Jones EY. Structural features impose tight peptide binding specificity in the nonclassical MHC molecule HLA-E. *Mol Cell.* 1998; 1:531–541. [PubMed: 9660937]
13. Tomasec P, Braud VM, Rickards C, Powell MB, McSharry BP, Gadola S, Cerundolo V, Borysiewicz LK, McMichael AJ, Wilkinson GW. Surface expression of HLA-E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40. *Science.* 2000; 287:1031. [PubMed: 10669413]
14. Bland FA, Lemberg MK, McMichael AJ, Martoglio B, Braud VM. Requirement of the proteasome for the trimming of signal peptide-derived epitopes presented by the nonclassical major histocompatibility complex class I molecule HLA-E. *J Biol Chem.* 2003; 278:33747–33752. [PubMed: 12821659]
15. Hammer GE, Gonzalez F, Champsaur M, Cado D, Shastri N. The aminopeptidase ERAAP shapes the peptide repertoire displayed by major histocompatibility complex class I molecules. *Nat Immunol.* 2006; 7:103–112. [PubMed: 16299505]
16. Nagarajan NA, Gonzalez F, Shastri N. Nonclassical MHC class Ib-restricted cytotoxic T cells monitor antigen processing in the endoplasmic reticulum. *Nat Immunol.* 2012; 13:579–586. [PubMed: 22522492]
17. Braud VM, Allan DS, Wilson D, McMichael AJ. TAP- and tapasin-dependent HLA-E surface expression correlates with the binding of an MHC class I leader peptide. *Curr Biol.* 1998; 8:1–10. [PubMed: 9427624]
18. Reddehase MJ. Antigens and immunoevasins: opponents in cytomegalovirus immune surveillance. *Nat Rev Immunol.* 2002; 2:831–844. [PubMed: 12415307]
19. Lampen MH, Hassan C, Sluijter M, Geluk A, Dijkman K, Tjon JM, de Ru AH, van der Burg SH, van Veelen PA, van Hall T. Alternative peptide repertoire of HLA-E reveals a binding motif that is strikingly similar to HLA-A2. *Mol Immunol.* 2013; 53:126–131. [PubMed: 22898188]
20. Kim S, Lee S, Shin J, Kim Y, Evnouchidou I, Kim D, Kim YK, Kim YE, Ahn JH, Riddell SR, et al. Human cytomegalovirus microRNA miR-US4–1 inhibits CD8(+) T cell responses by targeting the aminopeptidase ERAP1. *Nat Immunol.* 2011; 12:984–991. [PubMed: 21892175]
21. Pietra G, Romagnani C, Manzini C, Moretta L, Mingari MC. The emerging role of HLA-E-restricted CD8+ T lymphocytes in the adaptive immune response to pathogens and tumors. *J Biomed Biotechnol.* 2010; 2010:907092. [PubMed: 20634877]
22. Heinzel AS, Grotzke JE, Lines RA, Lewinsohn DA, McNabb AL, Streblow DN, Braud VM, Grieser HJ, Belisle JT, Lewinsohn DM. HLA-E-dependent presentation of Mtb-derived antigen to human CD8+ T cells. *J Exp Med.* 2002; 196:1473–1481. [PubMed: 12461082]
23. Grotzke JE, Harriff MJ, Siler AC, Nolt D, Delepine J, Lewinsohn DA, Lewinsohn DM. The Mycobacterium tuberculosis phagosome is a HLA-I processing competent organelle. *PLoS Pathog.* 2009; 5:e1000374. [PubMed: 19360129]
24. Elliott T, Smith M, Driscoll P, McMichael A. Peptide selection by class I molecules of the major histocompatibility complex. *Curr Biol.* 1993; 3:854–866. [PubMed: 15335818]
25. Germain RN. MHC-dependent antigen processing and peptide presentation: providing ligands for T lymphocyte activation. *Cell.* 1994; 76:287–299. [PubMed: 8293464]
26. Cerundolo V, Elliott T, Elvin J, Bastin J, Townsend A. Association of the human invariant chain with H-2 Db class I molecules. *Eur J Immunol.* 1992; 22:2243–2248. [PubMed: 1516617]
27. Kim HJ, Verbrinnen B, Tang X, Lu L, Cantor H. Inhibition of follicular T-helper cells by CD8(+) regulatory T cells is essential for self tolerance. *Nature.* 2010; 467:328–332. [PubMed: 20844537]

28. Jiang H, Canfield SM, Gallagher MP, Jiang HH, Jiang Y, Zheng Z, Chess L. HLA-E-restricted regulatory CD8(+) T cells are involved in development and control of human autoimmune type 1 diabetes. *J Clin Invest*. 2010; 120:3641–3650. [PubMed: 20877010]
- 29••. Fukazawa Y, Lum R, Okoye AA, Park H, Matsuda K, Bae JY, Hagen SI, Shoemaker R, Deleage C, Lucero C, et al. B cell follicle sanctuary permits persistent productive simian immunodeficiency virus infection in elite controllers. *Nat Med*. 2015; 21:132–139. Demonstration that the lymphoid follicles are an immunologically privileged site for persistent SIV infection. [PubMed: 25599132]
- 30••. He R, Hou S, Liu C, Zhang A, Bai Q, Han M, Yang Y, Wei G, Shen T, Yang X, et al. Follicular CXCR5-expressing CD8+ T cells curtail chronic viral infection. *Nature*. 2016; 537:412–428. Demonstration that CXCR5 positive antiviral CD8+ T cells can limit chronic virus infection in this immunologically privileged site. [PubMed: 27501245]
31. Purbhoo MA, Irvine DJ, Huppa JB, Davis MM. T cell killing does not require the formation of a stable mature immunological synapse. *Nat Immunol*. 2004; 5:524–530. [PubMed: 15048111]
32. Met O, Buus S, Claesson MH. Peptide-loaded dendritic cells prime and activate MHC-class I-restricted T cells more efficiently than protein-loaded cross-presenting DC. *Cell Immunol*. 2003; 222:126–133. [PubMed: 12826082]
33. Pietra G, Romagnani C, Mazzarino P, Falco M, Millo E, Moretta A, Moretta L, Mingari MC. HLA-E-restricted recognition of cytomegalovirus-derived peptides by human CD8+ cytolytic T lymphocytes. *Proc Natl Acad Sci U S A*. 2003; 100:10896–10901. [PubMed: 12960383]
34. Sylwester AW, Mitchell BL, Edgar JB, Taormina C, Pelte C, Ruchti F, Sleath PR, Grabstein KH, Hosken NA, Kern F, et al. Broadly targeted human cytomegalovirus-specific CD4+ and CD8+ T cells dominate the memory compartments of exposed subjects. *J Exp Med*. 2005; 202:673–685. [PubMed: 16147978]
- 35•. Peppenelli MA, Arend KC, Cojohari O, Moorman NJ, Chan GC. Human cytomegalovirus stimulates the synthesis of select Akt-dependent antiapoptotic proteins during viral entry to promote survival of infected monocytes. *J Virol*. 2016; 90:3138–3147. Inhibition of programmed cell death by HCMV. [PubMed: 26739047]
- 36•. Yatim N, Jusforgues-Saklani H, Orozco S, Schulz O, Barreira da Silva R, Reis e Sousa C, Green DR, Oberst A, Albert ML. RIPK1 and NF-kappaB signaling in dying cells determines cross-priming of CD8(+) T cells. *Science*. 2015; 350:328–334. How programmed cell death is involved in cross priming. [PubMed: 26405229]

Figure 1.

Generation of the VL9 peptide from the HLA-A2 signal peptide. The signal peptide is translocated into the lumen of the endoplasmic reticulum (ER) where it is cleaved from the HLA heavy chain by the signal peptidase. This is further cleaved by the transmembrane signal peptide peptidase to generate a 14-mer fragment which is then digested by the proteasome. The fragments truncated at the carboxy terminus are conveyed by the transporter associated with antigen processing (TAP) back into the lumen of the ER where the amino terminal two amino acids are removed by the endoplasmic reticulum amino peptidase 1 (ERAP1) leaving the VL9 peptide which binds to the TAP and tapasin associated HLA-E. This is then released to travel to the cell surface. In parallel other peptides are being generated from self and foreign proteins, being transported and binding in a similar way to other HLA class I molecules.

Peptide Epitopes presented by MHC-E		
MHC-I Signal peptide: V MAP R TL V L		
L		
F		
M2 – L9 motif	Partial motif	No motif
CMV UL40 V MAP R TLL L	MRP7 ALALVRML I	SIVgag APLVPTG S E
Hsp60 Q MRPVSR V L	FAM49B FYAEATP M L	SIVgag SENLKSL Y N
GroEL G MKFDRG Y I	EBV BZLF1 SQAPLPC V L	SIVgag EKVKHTE E A
Prdx5 Δ 2 A MAPIKTH L	FLU M ILGFVFT L	SIVgag KTSRPTA S
SIV gag R MYNP T NIL	HCV core YLLPR R GP L	SIVgag QIIRDI E
	SIV gag GGNYPVQ Q I	SIVgag KPIKCWN C G
	SIV gag VGNIYRR I	SIVgag LGKQQRE K Q
		SIVgag EKQRESRE K
+ 35 <i>Mtb</i> peptides	+ 34 <i>Mtb</i> peptides eg RLP A KAP L L	

Figure 2.

Examples of peptides that have been shown to bind to MHC-E; most are also epitopes for MHC-E restricted T cells. The peptides are grouped according to whether or not they fit the (red) M2-L9 motif originally identified for the VL9 peptide. The unusual C terminal residue of the SIV peptides are indicated in green. The peptides are described in Refs [3,5•,15,20], the last review giving their original descriptions.

Figure 3.

A comparison of antigen processing in a RhCMV68-1-SIV infected cell with an SIV infected cell. On the left the RhCMV68-1-SIV vaccine infects a cell capable of direct T cell priming. RhCMV genes equivalent to HCMV US2,3 and 11 down-regulate classical MHC class I molecules by targeting them for degradation. Furthermore, the RhCMV gene equivalent to HCMV US6 blocks the TAP. MHC-E expression is enhanced probably because the RhCMV gene Rh67 (similar to HCMV UL40) provides the VL9 peptide from its own signal sequence in a TAP independent manner. Because the normal antigen presentation pathway is disturbed, other peptides can bind to MHC-E and prime CD8⁺ T cell responses. The Rh67 derived VL9 ensures inhibition of NK cell attack. Because there are no CD8⁺ T cell responses specific for peptides presented by classical MHC class Ia molecules, cross priming must be blocked.

In the SIV infected cells there are no equivalents to HCMV US2,3,11 or 6. There is partial MHC-Ia downregulation but this results from interference with post-ER trafficking so does not affect generation of the VL9 peptide. MHC-E then presents both the dominant VL9 peptide and some other peptides. These are sufficient for the cells to be targets for CD8⁺ T cells but are insufficient to prime T cell responses. Direct and cross priming generates MHC-Ia restricted CD8⁺ T cells.