Published in final edited form as:

Dev Psychopathol. 2017 December ; 29(5): 1619–1626. doi:10.1017/S0954579417001274.

Stress exposure and psychopathology alter methylation of the serotonin receptor 2A (*HTR2A*) gene in preschoolers

Stephanie H. Parade^{a,b}, Andrew M. Novick^{a,c}, Justin Parent^{a,b,d}, Ronald Seifer^{a,b}, Samantha J. Klaver^b, Carmen J. Marsit^e, Asi Polly Gobin^c, Bao-Zhu Yang^f, and Audrey R. Tyrka^{a,c}

^aDepartment of Psychiatry and Human Behavior, Alpert Medical School of Brown University

^bBradley/Hasbro Children's Research Center, E. P. Bradley Hospital

^cMood Disorders Research Program and Laboratory for Clinical and Translational Neuroscience, Butler Hospital

^dCenter for Children and Families, Department of Psychology, Florida International University

^eDepartment of Environmental Health, Rollins School of Public Health, Emory University

^fDepartment of Psychiatry, Yale University School of Medicine

Abstract

Serotonin signaling pathways play a key role in brain development, stress reactivity, and mental health. Epigenetic alterations in the serotonin system may underlie the effect of early life stress on psychopathology. The current study examined methylation of the serotonin receptor gene *HTR2A* in a sample of 228 children including 119 with child welfare documentation of moderate to severe maltreatment within the last 6 months. Child protection records, semi-structured interviews in the home, and parent reports were used to assess child stress exposure, psychiatric symptoms, and behavior. *HTR2A* genotype and methylation of *HTR2A* was measured at two CpG sites (–1420 and –1224) from saliva DNA. *HTR2A* genotype was associated with *HTR2A* methylation at both CpG sites. *HTR2A* genotype also moderated associations of contextual stress exposure and *HTR2A* methylation at site –1420. Contextual stress was positively associated with –1420 methylation among A homozygotes, but negatively associated with –1420 methylation among G homozygotes. PTSD and MDD symptoms were negatively associated with methylation at –1420, but positively associated with methylation at –1224. Results support the view that the serotonin system is sensitive to stress exposure and psychopathology, and *HTR2A* methylation may be a mechanism by which early adversity is biologically encoded.

Keywords

HTR2A; methylation; contextual stress; serotonin; children

Childhood stress exposure including maltreatment and exposure to other contextual stressors associated with poverty confer risk for physical and mental health problems across the

lifespan (Benjet, Borges, & Medina-Mora, 2010; Cohen, Janicki-Deverts, & Miller, 2007; Felitti et al., 1998). Yet the mechanisms by which childhood stress exposure undermines long term health are not fully understood. Epigenetic processes are now thought to be a major mechanism linking early life stress to physical and mental health (Conradt, 2017; Szfy, 2013). The majority of research with children has focused on epigenetic processes related to the hypothalamic-pituitary-adrenal (HPA) axis stress response system, particularly the glucocorticoid receptor gene *NR3C1*. Yet multiple other systems regulate brain function and health. It is well established that serotonin signaling pathways play a key role in brain development, stress reactivity, and mental health (Booij, Tremblay, Szyf, & Benkelfat, 2015; Lin, Lee, & Yang, 2014), suggesting that epigenetic processes within the serotonin system may also underlie the association of childhood stress exposure and mental health.

Serotonin is a neurotransmitter that acts in the gastrointestinal system, blood, and throughout the brain to influence behavior and physiology. The serendipitous discovery that many of our first psychoactive drugs acted on serotonergic targets has made serotonin a focus of research in psychiatric disorders. Serotonin-producing neurons in the raphe nucleus of the midbrain project to several key regions involved in emotion, behavior, and cognition. From there, released serotonin can bind to over a dozen unique serotonin receptors, each with their own distribution and function. The serotonin receptor 2A (5-HT_{2A}) has been implicated in multiple psychiatric disorders. Activation of HTR_{2A} receptors in the periaqueductal gray region inhibits panic-like responses, while their activation in the amygdala is related more to generalized anxiety disorder and post-traumatic stress disorder through promotion of inhibitory avoidance and conditioned fear, respectively (Graeff, Viana, & Mora, 1997; Zangrossi et al., 2001). In the pre-frontal cortex activation of 5-HT_{2A} receptors facilitates top-down control of the amygdala, regulating the stress response (Arnsten, 2009; Aznar & Klein, 2013). However over activation of cortical 5-HT_{2A} receptors (especially in the context of glutamate receptor hypoactivity) has been linked to psychosis and the effects of hallucinogens (González-Maeso et al., 2007; González-Maeso & Sealfon, 2009).

Multiple post-mortem and in-vivo binding studies have found changes in 5-HT_{2A} receptor binding and expression in the brains of individuals with various forms of psychopathology. Increases in cortical 5-HT_{2A} receptor binding have been found in suicide victims and in individuals with major depressive disorder who did not die by suicide, but results are conflicting (Stockmeier, 2003). Findings also exist for differences in cortical 5-HT_{2A} levels and/or binding affinity in schizophrenia (Hernandez & Sokolov, 2000; Selvaraj, Arnone, Cappai, & Howes, 2014), borderline personality disorder (Soloff et al., 2007), and eating disorders (Frank et al., 2002).

Genetic variation in the 5-HT_{2A} receptor gene (*HTR2A*) may be an important determinant of differences in expression and activity of the 5-HT_{2A} receptor. One of the most widely studied genetic variants of *HTR2A* is the single nucleotide polymorphism (SNP), rs6311 (−1438 A/G). This polymorphism, located in the promoter region of the gene, has been thought to influence gene expression, with initial evidence suggesting that the A allele increases promoter activity and subsequent expression of the 5-HT_{2A} receptor (Parsons, D'Souza, Arranz, Kerwin, & Makoff, 2004; Turecki et al., 1999). The AA genotype has been associated with seasonal affective disorder (Enoch et al., 1999; Lee, Sung, Lim, Paik,

& Leen Kim, 2006), bipolar disorder (Bonnier et al., 2002), reduced responses and decreased side effects from antidepressants (Kato & Serretti, 2010) and greater reduction in negative symptoms in response to olanzapine (Ellingrod et al., 2003). For a comprehensive review see (Serretti, Drago, & De Ronchi, 2007).

In addition to genetic variation, epigenetic modifications play an important role in *HTR2A* regulation. Epigenetic processes include alterations to DNA which allow for adaptations to the environment but do not change the DNA sequence. Epigenetic modulation of DNA has the potential to alter gene expression, rendering it more or less likely to be expressed, with either positive or negative long-term consequences (Moore et al., 2013; Szyf, 2007). DNA methylation is the most commonly studied epigenetic process, and occurs when a methyl group is added at sites in the DNA where a cytosine nucleotide occurs next to a guanine nucleotide (CpG dinucleotides). Methylation at CpG sites in gene promoter regions may alter transcription factor binding, which may result in transcriptional gene silencing. Consequently, increases in methylation are typically associated with reductions in gene expression, and decreases in methylation are usually associated with increases in gene expression (Egger, Liang, Aparicio, & Jones, 2004; Reik, 2007; Uddin et al., 2011), although the effects of methylation depend on the transcriptional mechanisms at work in the particular region of the gene in question.

Methylation at CpG sites in *HTR2A* can either block or enhance gene transcription by altering binding of transcription factors (Falkenberg et al., 2011). Two CpG sites within the *HTR2A* promoter include those located at –1420bp and –1224bp (Falkenberg, Gurbaxani, Unger, & Rajeevan, 2011; Zhu, Chen, & Shih, 1995). While studies investigating the role of *HTR2A* genotype and methylation in psychiatric disorders are limited, hypermethylation has been found in adult schizophrenic and bipolar patients at CpG sites, including –1420, and –1224 (Abdolmaleky et al., 2011; Ghadirivassfi et al., 2011), and it is hypothesized that this might be the mechanism behind findings of decreased *HTR2A* expression in the frontal cortex of patients with schizophrenia (Hurlemann et al., 2007).

Epigenetic changes through methylation are now considered to be a primary mechanism by which the early life environment influences the development of behavioral phenotype. Starting in embryogenesis, serotonin plays an essential role in neural development, with the 5-HT_{2A} receptor promoting neuron proliferation (Azmitia, 2001). Thus epigenetic changes to *HTR2A* expression via environmental stressors would be expected to influence brain development and subsequently, behavior. In the neonatal period, methylation of placental *HTR2A* at sites –1420 and –1224 was associated with higher levels of attention in infants (Paquette et al., 2013). Here it is likely that methylation induced differences in *HTR2A* receptor expression altered placental physiology, with downstream effects on fetal brain development (Paquette & Marsit, 2014). While this study did not investigate how *HTR2A* expression was influenced in offspring, rodent models have found that both prenatal and postnatal stressors can alter levels of *HTR2A* receptors in adulthood (Holloway & González-Maeso, 2015; Peters, 1988; Rentesi et al., 2013). As it is unknown how early life experience affects *HTR2A* epigenetics, the present study sought to investigate the relationship between *HTR2A* genotype and methylation as it relates to early life stress and psychiatric symptomatology in a sample of preschoolers with a range of stress exposure.

Method

Data were available from 228 families for the current report. One child from each family was included. Children ranged in age from 3 to 5 years ($M = 50.9$ months; $SD = 9$ months), 121 were female and 107 were male. The sample was racially and ethnically diverse. Eighty-nine children were white, 39 black, 51 biracial, and 49 other races. One hundred and two children were Hispanic. Most caregivers were biological mothers ($n = 214$). Forty-seven caregivers had less than a high school degree, 88 completed high school, 71 had some post-secondary education, 21 had a bachelor's degree, and one did not provide education information. One-hundred and twenty-five caregivers in the study were unemployed and 205 of the families qualified for public assistance. One hundred and nineteen children (52%) had substantiated cases of moderate to severe maltreatment within the past 6 months as described below.

Procedure

Families with a maltreated child were identified from the local child welfare agency and an emergency maltreatment assessment service via record review. Families of children with no indicated case of maltreatment were recruited from childcare centers and a pediatric medical clinic during a well-child visit. Based on review of available medical records and parent report, children with a chronic illness, consistent medication use, obesity, or failure-to-thrive were excluded. Those with acute illness or medication use were included no less than 2 weeks following resolution of illness and discontinuation of medication. Families completed a baseline set of assessments at the time of initial study enrollment and a follow-up set of assessments 6 months following enrollment. At each wave of assessment, families completed two home visits and questionnaires between the visits. The current report focuses on the first home visit during the baseline set of assessments, during which caregivers completed interviews on child stress exposure and behavior. A saliva sample for DNA isolation was also collected from the children during this visit.

Measures

Child maltreatment status—All families consented to examination of child welfare records to determine maltreatment status. Trained research staff coded the records using the System for Coding Subtype and Severity of Maltreatment in Child Protective Records (Barnett, Manly, & Cicchetti, 1993). Five maltreatment subtypes and severity scores ranging from 1 (least severe) to 5 (most severe) were derived. Children with an episode that met the criteria for moderate to severe maltreatment (score of 3–5) within the last 6 months were included in the maltreated group ($n = 119$). Twenty-two children had substantiated cases of physical abuse, 28 sexual abuse, 14 physical neglect/failure to provide, 33 physical neglect/lack of supervision, and 75 emotional maltreatment. The comparison group included children who had never had a substantiated case of maltreatment ($n = 109$).

Contextual stress—Caregivers completed a semi-structured interview developed in our laboratory to assess contextual stressors experienced in the child's lifetime. Categories were: death of a caregiver, separation from a caregiver, housing instability, inadequate food or clothing, and other stressful events which included witnessing neighborhood violence or

parental arrest. Interviews were conducted and scored by trained clinical social workers and Ph.D. level psychologists. The project coordinator reviewed each interview to ensure compliance with the scoring protocol. Each domain was scored positive if at least one episode occurred, and domains were summed to determine the number of contextual stressor categories the child experienced in their lifetime. Possible scores ranged from 0 (no stressors) to 5 (stressors in all five domains). In the current sample the number of stressor categories ranged from 0 to 5 ($M = 1.54$, $SD = 1.22$).

Traumatic life events—The Diagnostic Infant and Preschool Assessment (Scheeringa & Haslett, 2010) interview was conducted with caregivers to assess child experiences of traumatic life events. Traumatic events in each domain were dichotomized (no trauma versus 1 trauma), then summed to create a scale for number of types of traumas experienced in the child's lifetime. Physical and sexual abuse were not included because they were assessed as maltreatment (above). Possible scores ranged from 0 (no traumas) to 9 (traumas in all nine domains). In the current sample the number of traumatic life events ranged from 0 to 4 ($M = 1.12$, $SD = 1.06$).

PTSD and MDD symptoms—The Diagnostic Infant and Preschool Assessment (DIPA; Scheeringa & Haslett, 2010) interview was also conducted with caregivers to assess symptoms of Post-Traumatic Stress Disorder (PTSD) and Major Depressive Disorder (MDD). Interviews were conducted by trained clinical social workers and Ph.D. level psychologists, reviewed in a group supervision format, and scored based upon group consensus. Consistent with the low prevalence of MDD and PTSD diagnoses reported in other studies of children under the age of six recruited from low-income mental health clinics (Bufferd, Dougherty, Carlson, & Klein, 2011; Scheeringa & Haslett, 2010), none of the children met full DSM-IV criteria for MDD and only one met criteria for PTSD; therefore the variables for analysis were the number of symptoms of MDD and of PTSD experienced within the past month. In the current sample the number of PTSD symptoms ranged from 0 – 10 ($M = 1.44$, $SD = 2.46$) and the number of MDD symptoms ranged from 0 to 3 ($M = 0.41$, $SD = 0.78$).

Behavior problems—Caregivers completed the Child Behavior Checklist for Ages 1.5 to 5 (CBCL; Achenbach & Rescorla, 2000) to assess internalizing and externalizing behavior problems. For each of the 100 behaviors, parents assessed their children on a 3-point scale from 0 (Not True) to 2 (Very True). T scores were used for data analysis. The CBCL is a reliable and valid measure with strong test-retest reliability ($r = .90$ and $.87$ for internalizing and externalizing scales, respectively) as well as discriminant validity between children who were and were not referred for behavioral health services (Achenbach & Rescorla, 2000). In the current sample the range for internalizing problems was 29 to 79 ($M = 52.84$, $SD = 8.21$) and for externalizing problems was 28 to 78 ($M = 47.50$, $SD = 11.54$).

Methylation of *HTR2A*—Two CpGs (–1420 and –1224) were studied based on findings of Falkenberg and colleagues (2011) and Paquette and colleagues (2013). Saliva samples were obtained using the Oragene DISCOVER kits (OGR-575) for Assisted Collections (DNA Genotek, Kanata, Ontario, Canada), and DNA was isolated following the

manufacturer's instructions. A 276 bp. region of the *HTR2A* promoter region (Chr 7: 28415063-28414801) was amplified from bisulfite modified DNA using the pyromark PCR kit (Qiagen, 203443), and DNA methylation assessed through pyrosequencing following a serial pyrosequencing protocol using a PyroMark MD system (Qiagen) and two sequencing primers as described in Paquette et al (2013). Each pyrosequencing assay contained a bisulfite conversion control to assess conversion efficiency. All samples examined demonstrated >95% conversion.

***HTR2A* genotyping**—Genotyping of the *HTR2A* (rs6311) single nucleotide polymorphism (SNP), rs6311 (–1438 A/G) was determined visual inspection of the pyrogram resulting from the first pyrosequencing assay. Thirty-five children were homozygotes for the A allele, 121 children were AG heterozygotes, and 72 children were homozygotes for the G allele.

Modeling Ancestry Differences Using Principal Component Analysis—Allele frequency differences due to systematic ancestry differences could cause spurious associations. We used principal components analysis (PCA) to model ancestry differences in the current study using genome-wide SNP markers from saliva DNA genotyped using the Illumina Infinium PsychArray-24 beadchip (over 588,000 autosomal SNPs). Genotypes were cleaned using standard quality control procedures. We conducted the linkage disequilibrium (LD)-based pruning first, and followed by PCA using PLINK (Purcell et al., 2007). LD-based pruning reduces correlation among SNPs such that the principal components (PCs) of the genetic variation in the sample would not be over-weighted by the contribution of correlated SNPs. The first two PCs obtained using PLINK were used for controlling the potential population stratification (Price et al., 2006).

Statistical analysis

Mean differences in demographic characteristics, stress exposure, and child behavior based on genotype were examined using *t*-tests and Chi-square. Simple correlations between demographic characteristics and methylation were conducted to determine inclusion of covariates. General linear modeling was used for hypothesis testing. Two PCs to adjust for genetic ancestry were included in the general linear models as a-priori covariates. Child age was associated with genotype and methylation as described below, and therefore was also included in the models as a covariate. *HTR2A* genotype was associated with methylation as described below, and was included as a covariate in models testing main effects of stress exposures and child symptoms and behavior on methylation. Outliers, defined as values more than three standard deviations from the mean, were Winsorized by setting them to the next highest value within three standard deviations. Missing data were imputed using SPSS 22. Less than 3% of data was missing overall. Little's Missing Completely at Random test (Little, 1988) demonstrated that the data were missing completely at random. Ten imputed datasets were derived from the original dataset, analyses were run on each individual dataset, and parameter estimates from all ten datasets were pooled to derive the final results. Results derived using multiple imputation were consistent with those derived from the original dataset with missing values, with the exception that the effect of MDD symptoms on

methylation at -1420 was at trend level ($p < .10$) using the original dataset and reached significance using the imputed dataset.

Results

Sample characteristics

The minor allele frequency (MAF) of the *HTR2A* allelic variant in the sample was .42 and the distribution conformed to the Hardy-Weinberg equilibrium ($X^2 = 1.85$, $p = .17$). Table 1 shows sample characteristics in relation to *HTR2A* genotype. A homozygotes were significantly older than heterozygotes. Child sex, the PCs to adjust for genetic ancestry, stress exposure, and behavior did not differ according to genotype. Child age was not associated with methylation at -1420, but was negatively associated with methylation at -1224 ($r = -.14$, $p = .034$). Child sex and the PCs to adjust for genetic ancestry were not associated with methylation at -1420 or -1224.

HTR2A genotype effects on methylation

HTR2A genotype was associated with methylation at -1420 ($F(2, 225) = 8.33$, $p < .001$) and -1224 ($F(2, 225) = 43.77$, $p < .001$). As displayed in Figure 1, A homozygotes had lower methylation at -1420 than heterozygotes and G homozygotes. A homozygotes also had lower methylation at -1224 than heterozygotes and G homozygotes, and heterozygotes had lower methylation at -1224 than G homozygotes.

Stress exposure and methylation

Main effects of child maltreatment status, traumatic life events, and contextual stress on methylation at -1420 and -1224 were not significant. The interaction of contextual stress and genotype was a significant predictor of methylation at -1420 ($F(2, 219) = 3.52$, $p = .031$). To understand the interaction effect we examined partial correlations of contextual stress and -1420 methylation within each genotype, controlling for age and the PCs to adjust for genetic ancestry. As illustrated in Figure 2, contextual stress was positively associated with methylation among A homozygotes ($r = .21$), negatively associated with methylation among G homozygotes ($r = -.28$), and not associated with methylation among heterozygotes ($r = .02$). In contrast, the interaction of child maltreatment status and genotype was not significant, nor was the interaction of traumatic life events and genotype.

Child symptoms and methylation

Main effects of PTSD symptoms ($F(1, 221) = 5.35$, $p = .022$) and MDD symptoms ($F(1, 221) = 5.16$, $p = .024$) on methylation at -1420 were significant. Likewise, main effects of PTSD symptoms ($F(1, 221) = 11.06$, $p = .001$) and MDD symptoms ($F(1, 221) = 6.30$, $p = .013$) on methylation at -1224 were significant. Examination of partial correlations controlling for age, the PCs to adjust for genetic ancestry, and *HTR2A* genotype indicated that PTSD and MDD symptoms were negatively associated with methylation at -1420 ($r = -.15$, $p = .022$ for both PTSD and MDD symptoms), but positively associated with methylation at -1224 ($r = .22$, $p = .001$ and $r = .17$, $p = .014$ for PTSD and MDD symptoms, respectively). In contrast, interactions of PTSD and MDD symptoms with genotype were not significant predictors of methylation at -1420 or -1224.

Child behavior problems and methylation

Internalizing and externalizing behavior problems were not associated with methylation at –1420 or –1224. Likewise, interactions of *HTR2A* genotype and internalizing and externalizing behavior problems were not significant predictors of methylation at –1420 or –1224.

Discussion

The current study examined methylation of *HTR2A* with early life stress and psychiatric symptoms in a sample of preschoolers with early adversity. We demonstrated that *HTR2A* genotype is associated with *HTR2A* methylation at CpGs –1420 and –1224, such that A homozygotes had lower methylation than children with the G allele. *HTR2A* genotype also moderated associations of contextual stress exposure and *HTR2A* methylation at –1420. Contextual stress was positively associated with –1420 methylation among A homozygotes, but negatively associated with –1420 methylation among G homozygotes. PTSD and MDD symptoms were negatively associated with methylation at –1420, but positively associated with methylation at –1224. Collectively, these results suggest that *HTR2A* is sensitive to stress exposure and psychopathology, and that environmental factors work in conjunction with DNA variation to affect this epigenetic process.

In the present study, we found decreased methylation at both the –1420 and –1224 CpG sites in those with the AA genotype. Few studies have examined this issue, but initial evidence has suggested that the A allele results in increased *HTR2A* expression (Parsons et al., 2004; Turecki et al., 1999) (but also see Bray, Buckland, Hall, Owen, & O'Donovan, 2004; Smith et al., 2013 for alternate evidence). Given that methylation typically functions to silence gene expression (Egger et al., 2004), the decreased methylation in AA individuals found in the present study might facilitate increased gene expression associated with this allele. As the present study measured methylation of two CpG sites in *HTR2A* in saliva DNA and did not evaluate gene expression, it is unknown how such differences in methylation might affect receptor expression within the brain.

While the data on the relationship between *HTR2A* methylation and receptor expression is limited, Falkenberg et al. (2011) created a model of *HTR2A* regulation based on a combination of computational, animal and human evidence. In this model, decreased methylation in the AA genotype at –1420 and –1224 allows for binding of the glucocorticoid receptor and transcription factor SP1, respectively. These transcription factors have opposing actions on gene expression, with the glucocorticoid receptor acting as a repressor and SP1 as a promoter. This model suggests that our findings of decreased methylation at –1420 and –1224 in AA individuals could have mixed rather than unilateral effects on gene expression.

Contextual stress was associated with increases or decreases in methylation at –1420 dependent on genotype. In AA individuals, contextual stress was associated with increased methylation while the opposite was true in GG individuals. Based on the model of Falkenberg et al. (2011), by increasing methylation of the AA genotype (and blocking the repressive glucocorticoid response element), one would expect increased expression, while

the decreased methylation in the GG genotype reduces expression. That contextual stress resulted in a differing propensity for methylation based on genotype suggests a mechanism for potential gene $\times$ environment interaction. While the details and consequences of such an interaction would require further study, it is known that other serotonin genes demonstrate different methylation based on stress exposure and genotype. For example, early life and recent stress has been correlated with higher levels of methylation in individuals with the SS version of the serotonin transporter gene (*SLC6A4*) (Duman & Canli, 2015) with this genotype being linked to higher stress sensitivity (Karg, Burmeister, Shedden, & Sen, 2011).

Animal models support a role for stress in altering *HTR2A* expression, although results are conflicting. Prenatal stress has been found to result in both increased and decreased *htr2a* receptor expression in adult offspring (Holloway & González-Maeso, 2015; Peters, 1988; Rentesi et al., 2013). Postnatal maternal deprivation stress has been shown to result in decreased *HTR2A* expression the frontal cortex of rats (Rentesi et al., 2013), while repeated exposure to inescapable shock in adulthood increased *HTR2A* expression (Dwivedi, Mondal, Payappagoudar, & Rizavi, 2005). As these studies deal with stress exposure occurring during discrete periods of development, they have limited applicability to the contextual stress measurement in the present study. That child maltreatment and traumatic life events were not associated with *HTR2A* methylation, but contextual stress exposure was associated with methylation in conjunction with genotype, suggests that developmental timing and other stressor characteristics are possible determinants of *HTR2A* methylation and gene expression. Further research is needed to understand these processes.

Although overall symptom levels of MDD and PTSD were low in our sample, we did find significant associations with *HTR2A* methylation. To our knowledge, we are the first to show a relationship with PTSD and MDD symptoms and methylation of *HTR2A* –1420 and –1224 CpG sites. Specifically, symptom levels were positively correlated with methylation at the –1420 site and negatively correlated with symptoms at the –1224 site. This pattern of methylation would enhance binding of the promoter transcription factor Sp1 (Zhu et al., 1995) while decreasing that of the repressive glucocorticoid receptor (Falkenberg et al., 2011) and therefore might be expected to increase overall expression. This is in line with a recent study demonstrating that expression of *HTR2A* mRNA in peripheral blood cells was associated with greater depression severity (Amidfar et al., 2017). The current study did not find a relationship between genotype and MDD or PTSD symptoms. While there is a lack of data on PTSD, recent meta-analyses have found both no relationship between –1438 A/G polymorphism and MDD (Jin, Xu, Yuan, Wang, & Cheng, 2013) and increased risk with AA genotype (Zhao et al., 2014). These conflicting results might be reconciled with additional studies that take into account methylation and contextual stressors, as our results suggest these play a role in genotype-dependent regulation of the 5-HT_{2A} receptor.

The current study is the first to examine associations of early life stress and psychopathology with *HTR2A* methylation. Nonetheless there are limitations that warrant future research in this domain. First, our comparison group of children with no maltreatment history were exposed to a range of other contextual stressors associated with poverty which may have reduced the strength of associations among maltreatment and methylation. Future work should draw upon a comparison group of children with no maltreatment history or other

contextual stressors to determine if the non-significant associations among maltreatment and *HTR2A* methylation are unique to our sample with a range of other stress exposures. Likewise, although child protection records were reviewed for all children in the study, it is possible that children within the comparison group had undocumented maltreatment, further reducing the strength of associations. Second, the current study focused on contextual risk factors and psychopathology as they relate to methylation and is an important first step in understanding environmental effects on *HTR2A* methylation. Examination of protective factors including parenting sensitivity and engagement in therapeutic interventions that potentially enhance resilience among children exposed to early adversity is an important next step. Finally, and as discussed above, data on gene expression is not currently available from the sample and would be useful in understanding the mechanisms underlying our results. Despite these limitations, the current study provides novel data linking stress exposure and psychiatric symptoms to *HTR2A* methylation in a sample of children with early adversity. Results support the perspective that DNA methylation is a mechanism by which early life stress impacts child mental health, and implicates serotonin signaling pathways as a relevant epigenetic process.

Acknowledgments

This research was supported by grant R01 MH083704 (ART) and R25 MH101076 (AMN) from the National Institute of Mental Health. The content is solely the responsibility of the authors and does not necessarily reflect the official views of the NIMH. We are grateful to the children and families who participated in this study, and we thank Hasbro Children's Hospital, Rhode Island Head Start, and the Rhode Island Department of Children, Youth, and Families for assisting in recruitment of study participants. We also thank Brittney Josefson and the numerous other research assistants who contributed to this project. Isolation of DNA and the genotyping array were done in the laboratory of Joel Gelernter, M.D., and we are grateful to Dr. Gelernter and his staff for their contribution.

References

- Abdolmaleky HM, Yaqubi S, Papageorgis P, Lambert AW, Ozturk S, Sivaraman V, Thiagalingam S. Epigenetic dysregulation of *HTR2A* in the brain of patients with schizophrenia and bipolar disorder. *Schizophrenia Research*. 2011; 129:183–190. DOI: 10.1016/j.schres.2011.04.007 [PubMed: 21550210]
- Achenbach, TM., Rescorla, LA. Manual for the ASEBA preschool forms & profiles: An integrated system of multi-informant assessment; Child behavior checklist for ages 11/2–5; Language development survey; Caregiver-teacher report form. University of Vermont: 2000.
- Amidfah M, Kim Y-K, Colic L, Arbabi M, Mobaraki G, Hassanzadeh G, Walter M. Increased levels of 5HT2A receptor mRNA expression in peripheral blood mononuclear cells of patients with major depression: correlations with severity and duration of illness. *Nordic Journal of Psychiatry*. 2017; 71:282–288. DOI: 10.1080/08039488.2016.1276624 [PubMed: 28125323]
- Arnsten AFT. Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews. Neuroscience*. 2009; 10:410–422. DOI: 10.1038/nrn2648 [PubMed: 19455173]
- Azmitia EC. Modern views on an ancient chemical: Serotonin effects on cell proliferation, maturation, and apoptosis. *Brain Research Bulletin*. 2001; 56:413–424. DOI: 10.1016/S0361-9230(01)00614-1 [PubMed: 11750787]
- Aznar S, Klein AB. Regulating Prefrontal Cortex Activation: An Emerging Role for the 5-HT2A Serotonin Receptor in the Modulation of Emotion-Based Actions? *Molecular Neurobiology*. 2013; 48:841–853. DOI: 10.1007/s12035-013-8472 [PubMed: 23696058]
- Barnett, D., Manly, JT., Cicchetti, D. Defining child maltreatment: The interface between policy and research. In: Cicchetti, D., Toth, SL., editors. *Child abuse, child development, and social policy*. Norwood, NJ: Ablex; 1993. p. 7-73.

- Benjet C, Borges G, Medina-Mora ME. Chronic childhood adversity and onset of psychopathology during three life stages: childhood, adolescence and adulthood. *Journal of Psychiatric Research*. 2010; 44:732–740. DOI: 10.1016/j.jpsychires.2010.01.004 [PubMed: 20144464]
- Bonnier B, Gorwood P, Hamon M, Sarfati Y, Boni C, Hardy-Bayle M-C. Association of 5-HT(2A) receptor gene polymorphism with major affective disorders: the case of a subgroup of bipolar disorder with low suicide risk. *Biological Psychiatry*. 2002; 51:762–765. DOI: 10.1016/S0006-3223(01)01228-8 [PubMed: 11983190]
- Booij L, Tremblay R, Szyf M, Benkelfat C. Genetic and early environmental influences on the serotonin system: consequences for brain development and risk for psychopathology. *Journal of Psychiatry & Neuroscience*. 2015; 40:5–18. DOI: 10.1503/jpn.140099 [PubMed: 25285876]
- Bray NJ, Buckland PR, Hall H, Owen MJ, O'Donovan MC. The serotonin 2A receptor gene locus does not contain common polymorphism affecting mRNA levels in adult brain. *Molecular Psychiatry*. 2004; 9:109–114. DOI: 10.1038/sj.mp.4001366 [PubMed: 14699448]
- Bufferd SJ, Dougherty LR, Carlson GA, Klein DN. Parent-reported mental health in preschoolers: findings using a diagnostic interview. *Comprehensive Psychiatry*. 2011; 52:359–369. DOI: 10.1016/j.comppsy.2010.08.006 [PubMed: 21683173]
- Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *Journal of the American Medical Association*. 2007; 298:1685–1687. DOI: 10.1001/jama.298.14.1685 [PubMed: 17925521]
- Conradt E. Using Principles of Behavioral Epigenetics to Advance Research on Early Life Stress. *Child Development Perspectives*. 2017; 11:107–112. DOI: 10.1111/cdep.12219 [PubMed: 29081832]
- Duman EA, Canli T. Influence of life stress, 5-HTTLPR genotype, and SLC6A4 methylation on gene expression and stress response in healthy Caucasian males. *Biology of Mood & Anxiety Disorders*. 2015; 5:2.doi: 10.1186/s13587-015-0017 [PubMed: 25995833]
- Dwivedi Y, Mondal AC, Payappagoudar GV, Rizavi HS. Differential regulation of serotonin (5HT)2A receptor mRNA and protein levels after single and repeated stress in rat brain: role in learned helplessness behavior. *Neuropharmacology*. 2005; 48:204–214. DOI: 10.1016/j.neuropharm.2004.10.004 [PubMed: 15695159]
- Egger G, Liang G, Aparicio A, Jones PA. Epigenetics in human disease and prospects for epigenetic therapy. *Nature*. 2004; 429:457–463. DOI: 10.1038/nature02625 [PubMed: 15164071]
- Ellingrod VL, Lund BC, Miller D, Fleming F, Perry P, Holman TL, Bever-Stille K. 5-HT2A receptor promoter polymorphism, –438G/A and negative symptom response to olanzapine in schizophrenia. *Psychopharmacology Bulletin*. 2003; 37:109–112. [PubMed: 14566219]
- Enoch MA, Goldman D, Barnett R, Sher L, Mazzanti CM, Rosenthal NE. Association between seasonal affective disorder and the 5-HT2A promoter polymorphism, –1438G/A. *Molecular Psychiatry*. 1999; 4:89–92. DOI: 10.1038/sj.mp.4000439 [PubMed: 10089016]
- Falkenberg VR, Gurbaxani BM, Unger ER, Rajeevan MS. Functional genomics of serotonin receptor 2A (HTR2A): interaction of polymorphism, methylation, expression and disease association. *Neuromolecular Medicine*. 2011; 13:66–76. DOI: 10.1007/s12017-010-8138-2 [PubMed: 20941551]
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, Marks JS. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*. 1998; 14:245–258. DOI: 10.1016/S0749-3797(98)00017-8 [PubMed: 9635069]
- Frank GK, Kaye WH, Meltzer CC, Price JC, Greer P, McConaha C, Skovira K. Reduced 5-HT2A receptor binding after recovery from anorexia nervosa. *Biological Psychiatry*. 2002; 52:896–906. DOI: 10.1016/S0006-3223(02)01378-1 [PubMed: 12399143]
- Ghadirivasfi M, Nohesara S, Ahmadkhaniha H-R, Eskandari M-R, Mostafavi S, Thiagalingam S, Abdolmaleky HM. Hypomethylation of the serotonin receptor type-2A Gene (HTR2A) at T102C polymorphic site in DNA derived from the saliva of patients with schizophrenia and bipolar disorder. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics: The Official Publication of the International Society of Psychiatric Genetics*. 2011; 156B:536–545. DOI: 10.1002/ajmg.b.31192

- González-Maeso J, Sealfon SC. Psychedelics and schizophrenia. *Trends in Neurosciences*. 2009; 32:225–232. DOI: 10.1016/j.tins.2008.12.005 [PubMed: 19269047]
- González-Maeso J, Weisstaub NV, Zhou M, Chan P, Ivic L, Ang R, Gingrich JA. Hallucinogens Recruit Specific Cortical 5-HT_{2A} Receptor-Mediated Signaling Pathways to Affect Behavior. *Neuron*. 2007; 53:439–452. DOI: 10.1016/j.neuron.2007.01.008 [PubMed: 17270739]
- Graeff FG, Viana MB, Mora PO. Dual role of 5-HT in defense and anxiety. *Neuroscience and Biobehavioral Reviews*. 1997; 21:791–9. DOI: 10.1016/S0149-7634(96)00059-0 [PubMed: 9415904]
- Hernandez I, Sokolov BP. Abnormalities in 5-HT_{2A} receptor mRNA expression in frontal cortex of chronic elderly schizophrenics with varying histories of neuroleptic treatment. *Journal of Neuroscience Research*. 2000; 59:218–225. DOI: 10.1002/(SICI)1097-4547(20000115)59:2<218::AID-JNR8>3.0.CO;2-H [PubMed: 10650880]
- Holloway T, González-Maeso J. Epigenetic Mechanisms of Serotonin Signaling. *ACS Chemical Neuroscience*. 2015; 6:1099–1109. DOI: 10.1021/acschemneuro.5b00033 [PubMed: 25734378]
- Hurlmann R, Matusch A, Kuhn K-U, Berning J, Elmenhorst D, Winz O, Kolsch H, Zilles K, Wagner M, Bauer A. 5-HT_{2A} receptor density is decreased in the at-risk mental state. *Psychopharmacology*. 2007; 195:579–590. DOI: 10.1007/s00213-007-0921-x [PubMed: 17899021]
- Jin C, Xu W, Yuan J, Wang G, Cheng Z. Meta-analysis of association between the –1438A/G (rs6311) polymorphism of the serotonin 2A receptor gene and major depressive disorder. *Neurological Research*. 2013; 35:7–14. DOI: 10.1179/1743132812Y.0000000111 [PubMed: 23317793]
- Karg K, Burmeister M, Shedden K, Sen S. The Serotonin Transporter Promoter Variant (5-HTTLPR), Stress, and Depression Meta-analysis Revisited: Evidence of Genetic Moderation. *Archives of General Psychiatry*. 2011; 68:444. doi: 10.1001/archgenpsychiatry.2010.189 [PubMed: 21199959]
- Kato M, Serretti A. Review and meta-analysis of antidepressant pharmacogenetic findings in major depressive disorder. *Molecular Psychiatry*. 2010; 15:473–500. DOI: 10.1038/mp.2008.116 [PubMed: 18982004]
- Lee H-J, Sung S-M, Lim S-W, Paik J-W, Kim L. Seasonality associated with the serotonin 2A receptor-1438 A/G polymorphism. *Journal of Affective Disorders*. 2006; 95:145–148. DOI: 10.1016/j.jad.2006.05.001 [PubMed: 16762419]
- Lin SH, Lee LT, Yang YK. Serotonin and mental disorders: a concise review on molecular neuroimaging evidence. *Clinical Psychopharmacology and Neuroscience*. 2014; 12:196. doi: 10.9758/cpn.2014.12.3.196 [PubMed: 25598822]
- Little RJ. A test of missing completely at random for multivariate data with missing values. *Journal of the American Statistical Association*. 1988; 83:1198–1202. DOI: 10.2307/2290157
- Moore LD, Le T, Fan G. DNA methylation and its basic function. *Neuropsychopharmacology*. 2013; 38:23–38. DOI: 10.1038/npp.2012.112 [PubMed: 22781841]
- Paquette AG, Lesseur C, Armstrong DA, Koestler DC, Appleton AA, Lester BM, Marsit CJ. Placental HTR_{2A} methylation is associated with infant neurobehavioral outcomes. *Epigenetics*. 2013; 8:796–801. DOI: 10.4161/epi.25358 [PubMed: 23880519]
- Paquette AG, Marsit CJ. The Developmental Basis of Epigenetic Regulation of HTR_{2A} and Psychiatric Outcomes. *Journal of Cellular Biochemistry*. 2014; 115:2065–2072. DOI: 10.1002/jcb.24883 [PubMed: 25043477]
- Parsons MJ, D'Souza UM, Arranz M-J, Kerwin RW, Makoff AJ. The –1438A/G polymorphism in the 5-hydroxytryptamine type 2A receptor gene affects promoter activity. *Biological Psychiatry*. 2004; 56:406–410. DOI: 10.1016/j.biopsych.2004.06.020 [PubMed: 15364038]
- Peters DA. Both prenatal and postnatal factors contribute to the effects of maternal stress on offspring behavior and central 5-hydroxytryptamine receptors in the rat. *Pharmacology, Biochemistry, and Behavior*. 1988; 30:669–673. DOI: 10.1016/0091-3057(88)90081-0
- Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics*. 2006; 38:904–909. DOI: 10.1038/ng1847 [PubMed: 16862161]
- Reik W. Stability and flexibility of epigenetic gene regulation in mammalian development. *Nature*. 2007; 447:425–432. DOI: 10.1038/nature05918 [PubMed: 17522676]

- Rentesi G, Antoniou K, Marselos M, Syrrou M, Papadopoulou-Daifoti Z, Konstandi M. Early maternal deprivation-induced modifications in the neurobiological, neurochemical and behavioral profile of adult rats. *Behavioural Brain Research*. 2013; 244:29–37. DOI: 10.1016/j.bbr.2013.01.040 [PubMed: 23395600]
- Scheeringa MS, Haslett N. The reliability and criterion validity of the Diagnostic Infant and Preschool Assessment: a new diagnostic instrument for young children. *Child Psychiatry & Human Development*. 2010; 41:299–312. DOI: 10.1007/s10578-009-0169-2 [PubMed: 20052532]
- Selvaraj S, Arnone D, Cappai A, Howes O. Alterations in the serotonin system in schizophrenia: A systematic review and meta-analysis of postmortem and molecular imaging studies. *Neuroscience & Biobehavioral Reviews*. 2014; 45:233–245. DOI: 10.1016/j.neubiorev.2014.06.005 [PubMed: 24971825]
- Serretti A, Drago A, De Ronchi D. HTR2A gene variants and psychiatric disorders: a review of current literature and selection of SNPs for future studies. *Current Medicinal Chemistry*. 2007; 14:2053–2069. DOI: 10.2174/092986707781368450 [PubMed: 17691947]
- Smith RM, Papp AC, Webb A, Ruble CL, Munsie LM, Nisenbaum LK, Kleinman JE, Lipska BK, Sadee W. Multiple Regulatory Variants Modulate Expression of 5-Hydroxytryptamine 2A Receptors in Human Cortex. *Biological Psychiatry*. 2013; 73:546–554. DOI: 10.1016/j.biopsych.2012.09.028 [PubMed: 23158458]
- Soloff PH, Price JC, Meltzer CC, Fabio A, Frank GK, Kaye WH. 5HT2A receptor binding is increased in borderline personality disorder. *Biological Psychiatry*. 2007; 62:580–587. DOI: 10.1016/j.biopsych.2006.10.022 [PubMed: 17448449]
- Stockmeier CA. Involvement of serotonin in depression: evidence from postmortem and imaging studies of serotonin receptors and the serotonin transporter. *Journal of Psychiatric Research*. 2003; 37:357–373. DOI: 10.1016/S0022-3956(03)00050-5 [PubMed: 12849929]
- Szyf M. The dynamic epigenome and its implications in toxicology. *Toxicological Sciences*. 2007; 100:7–23. DOI: 10.1093/toxsci/kfm177 [PubMed: 17675334]
- Turecki G, Brière R, Dewar K, Antonetti T, Lesage AD, Séguin M, Chawky N, Vanier C, Alda M, Joobert R, Benkelfat C. Prediction of Level of Serotonin 2A Receptor Binding by Serotonin Receptor 2A Genetic Variation in Postmortem Brain Samples From Subjects Who Did or Did Not Commit Suicide. *American Journal of Psychiatry*. 1999; 156:1456–1458. DOI: 10.1176/ajp.156.9.1456 [PubMed: 10484964]
- Uddin M, Koenen KC, Aiello AE, Wildman DE, de Los Santos R, Galea S. Epigenetic and inflammatory marker profiles associated with depression in a community based epidemiologic sample. *Psychological Medicine*. 2011; 41:997. doi: 10.1017/S0033291710001674 [PubMed: 20836906]
- Zangrossi H, Viana MB, Zanoveli J, Bueno C, Nogueira RL, Graeff FG. Serotonergic regulation of inhibitory avoidance and one-way escape in the rat elevated T-maze. *Neuroscience and Biobehavioral Reviews*. 2001; 25:637–45. DOI: 10.1016/S0149-7634(01)00047-1 [PubMed: 11801289]
- Zhao X, Sun L, Sun Y-H, Ren C, Chen J, Wu Z-Q, Jiang YH, Lv X-L. Association of HTR2A T102C and A-1438G polymorphisms with susceptibility to major depressive disorder: a meta-analysis. *Neurological Sciences: Official Journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*. 2014; 35:1857–1866. DOI: 10.1007/s10072-014-1970-7
- Zhu QS, Chen K, Shih JC. Characterization of the human 5-HT2A receptor gene promoter. *Journal of Neuroscience*. 1995; 15:4885–4895. [PubMed: 7623119]

Figure 1.

Mean differences in methylation by *HTR2A* genotype

Note: Different letters indicate significant differences within the CpG site based on genotype at $p < .01$.

Figure 2.
HTR2A genotype moderates the association of contextual stress and *HTR2A* -1420 methylation

Table 1Descriptive statistics and mean differences by *HTR2A* genotype

	AA (<i>n</i> = 35)	AG (<i>n</i> = 121)	GG (<i>n</i> = 72)	<i>p</i>
Sex, <i>N</i> (%) female	20 (57.1)	63 (52.0)	38 (52.7)	.87
Age, <i>M</i> (<i>SD</i>)	4.5 (0.7) ^a	4.1 (0.7) ^a	4.3 (0.8)	.02
PC for Genetic Ancestry 1, <i>M</i> (<i>SD</i>)	-.02 (0.1)	.00 (0.1)	.01 (0.1)	.11
PC for Genetic Ancestry 2, <i>M</i> (<i>SD</i>)	.02 (0.1)	-.01 (0.1)	.00 (0.1)	.16
Maltreatment Status, <i>N</i> (%)	20 (57.1)	65 (53.7)	34 (47.2)	.56
Number of Stressors, <i>M</i> (<i>SD</i>)	1.7 (1.2)	1.5 (1.2)	1.6 (1.2)	.60
Traumatic Life Events, <i>M</i> (<i>SD</i>)	1.06 (1.0)	1.17 (1.1)	1.07 (1.1)	.74
PTSD Symptoms, <i>M</i> (<i>SD</i>)	1.8 (2.8)	1.3 (2.3)	1.5 (2.6)	.57
MDD Symptoms, <i>M</i> (<i>SD</i>)	0.6 (1.1)	0.4 (0.7)	0.4 (0.7)	.31
Internalizing Behavior, <i>M</i> (<i>SD</i>)	52.2 (7.3)	53.3 (8.5)	52.4 (8.2)	.71
Externalizing Behavior, <i>M</i> (<i>SD</i>)	48.8 (11.5)	47.1 (11.4)	47.6 (11.8)	.74

Note: *p* values indicate *t*-test or chi-square significance level.^aIndicates values are significantly different at $p < .01$.