Published in final edited form as:

Psychol Med. 2014 October ; 44(14): 2965–2974. doi:10.1017/S0033291714000786.

Neuropsychological Deficits in Past Suicide Attempters with Varying Levels of Depression Severity

John G. Keilp, Sue R. Beers, Ainsley K. Burke, Nadine M. Melhem, Maria A. Oquendo, David A. Brent, and J. John Mann

Department of Molecular Imaging and Neuropathology, New York State Psychiatric Institute, and the Department of Psychiatry, Columbia University College of Physicians and Surgeons (JGK, AKB, MAO, JJM), and the Department of Psychiatry, University of Pittsburgh School of Medicine, and the Western Psychiatric Institute and Clinic (SRB, NMM, DAB)

Abstract

Background—Our previous work identified deficits in interference processing and learning/memory in past suicide attempters who were currently depressed and medication-free. In this study, we extend this work to an independent sample studied at various stages of illness and treatment (mild symptoms, on average) to determine if these deficits in past suicide attempters are evident during a less severe clinical state and to identify potential intermediate phenotypes for familial aggregation studies.

Methods—Eighty individuals with a past history of major depression and suicide attempt were compared to eighty-one individuals with a history of major depression and no life-time suicide attempts on a battery of neurocognitive measures assessing attention, memory, abstract/contingent learning, working memory, language fluency, and impulse control.

Results—Past attempters performed more poorly in attention, memory, and working memory domains, but also in an estimate of premorbid intelligence. After correction for this estimate, tests that had previously distinguished past attempters – a computerized Stroop task and the Buschke Selective Reminding Test – remained significantly worse in attempters. In a secondary analysis, similar differences were found among those with the lowest levels of depression (HDRS<10) suggesting that these deficits may be trait markers independent of current symptomatology.

Conclusions—Deficits in interference processing and learning/memory constitute an enduring defect in information processing that may contribute to poor adaptation, other higher-order cognitive impairments, and risk for suicidal behavior.

Neurocognitive impairment plays a role in suicidal behavior, but the nature of that role and the specific contributions that various impairments make to the risk for suicidal behavior remain unknown. We have identified deficits in interference processing, memory, and working memory in past suicide attempters across two cohorts of currently depressed, medication-free individuals (Keilp et al., 2001; Keilp et al., 2013). Other recent studies have

*Address reprint requests to Dr. Keilp at Box 42, NYSPI, 1051 Riverside Drive, New York, New York 10032, Tel. (212) 543-5866, jgk13@columbia.edu, New York State Psychiatric Institute and Department of Psychiatry, Columbia University College of Physicians and Surgeons.

The authors have no other financial involvements that would create a conflict of interest based on results reported in this manuscript.

identified deficits in decision processes (Jollant et al., 2005; Jollant et al., 2011; Clark et al., 2011) reversal learning (Dombrovski et al., 2010) and other facets of executive performance (Marzuk, et al., 2005; Westheide et al., 2008; Dombrovski et al., 2011; McGirr et al., 2012; Bridge et al., 2012). Questions remain about the comparability of samples and the generalizability of these findings across samples. In addition, the nature of the attempts made by the population under study (i.e. attempt severity and choice of violent vs. non-violent methods) may play a critical role in the nature of the neurocognitive deficits observed (Keilp et al., 2013), further confounding attempts to characterize general risk factors for every type of suicidal behavior.

The present study extends our strategy of using a comprehensive neuropsychological assessment to evaluate depressed individuals with and without a past history of suicidal behavior, to an independent sample of individuals in the community with varying levels of depression severity. Our goal was to examine basic cognitive functions that may be related to suicidal behavior, and that may underlie the more complex dysfunction in reasoning and decision-making reported in other studies. Unlike our previous studies where patient participants were medication-free and met criteria for a current major depressive episode, participants in this study are a separate sample drawn from two sites, in various stages of depressive illness – on average with mild residual symptoms. The range of symptomatology, then, includes individuals who meet current criteria for a depressive episode, as well as individuals who meet rating-scale criteria for remission. There is no sample overlap with our previous studies.

Based on our previous findings (Keilp et al., 2001; Keilp et al., 2008; Keilp et al., 2013), we hypothesized that individuals with a past history of suicidal behavior would exhibit deficits in interference processing and learning/memory. There were two aims to this study. First, we sought to replicate our earlier findings and to determine how past attempter/non-attempter differences might be affected by the variable severity of depression in this sample. Our previous studies have found neurocognitive deficits in past suicide attempters while depressed; are these same deficits apparent when symptoms have been reduced by treatment and/or the passage of time? As a corollary to this analysis, we also examined performance differences between past attempters who had made higher versus lower lethality attempts, and those whose worst attempt involved a violent (e.g. hanging, cutting, jumping from a height) versus non-violent (overdose, substance ingestion) method. We (Keilp et al., 2013) and others (Jollant et al., 2005) had previously found differences between these subgroups that may or may not be evident in a less severely depressed sample.

Second, we sought to determine if deficits could be found among those with the fewest symptoms, in the subgroup of participants who met rating-scale criteria for remission. A number of previous studies of neurocognitive impairment in past suicide attempters had identified deficits while participants were remitted (Jollant et al., 2005; Malloy-Diniz et al., 2009; Jollant et al., 2011). If impairments can be identified among those whose depressive symptoms have been reduced to inter-episode levels, then these impairments are more likely to be trait markers of risk. The overall goal of these analyses was to identify neurocognitive intermediate phenotypes for familial transmission, given that all subjects were parents whose offspring have been evaluated on the same neurocognitive battery. If deficits are evident both

within as well as outside of a depressive episode, they may be characteristics that run within families as well, possibly with a genetic basis.

METHODS

Subjects were 161 individuals with a past history of major depressive episode (unipolar or bipolar), drawn from two sites at medical centers in Pittsburgh, Pennsylvania and New York, New York. Subjects were recruited by advertisement for participation in a family study, requiring their offspring to be evaluated as well. This study was approved by local Institutional Review Boards at each site, and all participants signed informed consent (offspring under the age of 18 signed assent, with parents signing consent).

Participants were free of any major medical or neurological illnesses and current substance abuse/dependence in the last two months (though past history was allowed). All participants had an estimated IQ above 80 (estimated using the Peabody Picture Vocabulary Test, Third Revision [PPVT]; Dunn et al., 1997) and were fluent in English (had spoken English as a primary language for at least 60% of their lives). 80 subjects had a past history of suicide attempt; 81 had no past history of suicide attempt.

Clinical Assessment

Lifetime diagnoses were established via SCID-I (Spitzer et al., 1990) and SCID-II interview (First et al., 1996), and past history of suicidal behavior was determined via structured interview (Oquendo et al., 2003). Participants also received measures of depression severity (Hamilton Depression Rating Scale [Hamilton, 1960, 1967], Beck Depression Scale [Beck et al., 1961]), hopelessness and suicidal thinking (Beck Hopelessness Scale [Beck et al., 1974], Beck Scale for Suicidal Ideation [Beck et al., 1979]), and traits associated with suicidal behavior (Barratt Impulsiveness Scale [Barratt 1985, 1994], Buss-Durkee Hostility Inventory [Buss & Durkee, 1961]). Past attempters also received an assessment of the severity (degree of medical damage) of past suicidal behavior (Beck Lethality Scale; Scale [Beck et al., 1975]), as in our previous studies (Mann et al., 1999; Brent et al., 2002; Keilp et al., op cit.).

All clinical assessments were completed by master's level clinicians or trained psychiatric nurses, with reliability of ratings regularly evaluated across sites, as reported in previous descriptions of our study (Brent, et al. 2002; Melhem et al., 2007).

Neuropsychological Assessment

Neuropsychological measures assessed six domains of functioning: **Attention** (computerized Continuous Performance Test – Identical Pairs; computerized Stroop Color/Word Test), **Memory** (Buschke Selective Reminding Test; Benton Visual Retention Test), **Abstract/Contingent Learning** (Wisconsin Card Sort, 64-card version), **Working Memory** (computerized A, Not B Reasoning Test; computerized N-Back Test), **Language Fluency** (Letter and Category Fluency), and **Impulse Control** (computerized Go-No Go Test; computerized Time Production Task). Tasks have been described in previous publications (Keilp et al., 2001; Keilp et al., 2005; Keilp et al., 2013). All assessors were trained by the

primary author (JGK) and supervised by each of the first two authors at each site (JGK in New York, SRB in Pittsburgh).

All test scores were adjusted for age, education, and/or sex effects based on available norms (Keilp et al., 2005; Spreen & Strauss, 2006). The principal measure from each task (see Table 2.) was averaged with others within each domain to produce six aggregate domain scores for analysis, as in our previous work (see Keilp et al., 2013).

Statistical Analyses

Demographic and clinical features of the samples were compared using univariate t-tests for continuous variables and chi-squared tests for categorical variables. Neuropsychological Domain scores were then analyzed in an initial omnibus analysis via a General Linear Model, treating domains as repeated measures, as in our previous work (Keilp et al., 2013). Covariates were selected based on univariate differences between the groups, and only those with a significant effect on test performance retained for the final model. Following omnibus analyses, individual test scores were compared between groups using t-tests and univariate Analysis of Covariance. Alpha level was set at $p = .05$.

Two secondary analyses were completed to address prior findings from the literature. Past attempters who had made higher vs. lower lethality attempts (with high lethality defined as a score of 4 or greater on the Beck Lethality Scale) were compared, as were those attempters who had used a violent (hanging, cutting, jumping from height) vs. non-violent (overdose, substance ingestion) method in their worst attempt. These groups had differed in our earlier work (Keilp et al., 2001; Keilp et al., 2013) as well as in other studies (Jollant et al., 2005). Finally, initial attempter vs. non-attempter comparisons were repeated in the subgroup of individuals who met criterion for remission at the time of their assessment, with remission defined as a 24-item Hamilton Depression Rating Scale score less than 10 (comparable to a suggested 21-item score <8 ; Riedel et al., 2010). As noted above, other studies had found deficits in remitted past attempters, but these studies have not included measures used here.

RESULTS

Demographic and Clinical Features

Samples were overwhelmingly female (88.8% of all participants), and both groups had a very high rate of reported past abuse (55.2% of overall samples report physical abuse; 60.1% report sexual abuse). However, past suicide attempters differed from non-attempters on a number of demographic and clinical measures (Table 1.). Past attempters had a larger proportion of females even though both groups were predominantly female. Past attempters had a year less education, lower overall estimated verbal intelligence, and higher levels of suicidal ideation. Both groups were mildly, but equivalently, depressed. Comorbidities, however, were much more prevalent in past suicide attempters, including Borderline Personality Disorder (BPD), past substance abuse, Post-Traumatic Stress Disorder (PTSD), and a history of past abuse.

Attempters were more likely to be on psychotropic medication (see Table 1., $\chi^2[1]=4.69$, $p=.03$). Percentage on antidepressants (42.5% vs. 35.8%; $\chi^2[1]=.36$, $p=.38$),

benzodiazepines/hypnotics (10.0% vs. 6.2%; $\chi^2[1]=.79$, $p=.37$), or stimulants (0.0% vs. 2.5%; $\chi^2[1]=2.00$, $p=.16$) did not differ significantly, but percentages on mood stabilizers or neuroleptics (13.8% vs. 4.9%; $\chi^2[1]=3.70$, $p=.05$) did. Remaining medications included cardiac, antihypertensive, or antihistamines (2.9% of sample), analgesic or other medications (12.9% of sample), and hormonal medications such as birth control (4.3% of sample).

Past attempters had higher hostility scores, but groups were equivalent in impulsiveness. Past attempters had made an average of just over two lifetime suicide attempts, with 56.8% ($n=46$) reporting multiple attempts.

Neuropsychological Performance, Aggregate Differences

In the omnibus comparison across all six neuropsychological domains, there was a significant difference between past attempters and non-attempters ($F[1,159]=11.16$, $p=.001$; see Figure 1.). The group by domain interaction was not significant ($F[5,795]=.97$, $p=.43$). At the level of individual domains (see Table 2.), differences were found in Attention, Memory, and Working Memory, with the difference in Impulse Control approaching significance ($p<.10$).

At the level of individual test scores (Table 2.), poorer performance by past attempters was found on the CPT, Stroop interference, Buschke SRT, Wisconsin Card Sort error score, and A, Not B Timed Reasoning task. Difference in N-Back was marginal and difference in Go-No Go approached significance. In all cases, past attempters performed more poorly than non-attempters.

Neuropsychological Performance, Covariates

The following covariates were tested for their potential effect on past attempter/non-attempter differences: PPVT score, gender, current suicidal ideation, history of BPD, history of substance abuse, history of PTSD, history of abuse, use of psychotropic medication, and research site (Pittsburgh vs. New York). Age and education were not included because all test scores were adjusted for their effects prior to analyses. The only significant covariate was PPVT ($F[1,134]=41.85$, $p<.001$) – and past attempt status remained significant even if all of these covariates were included in a single model ($F[1,134]=11.00$, $p=.001$).

In the final trimmed model including PPVT as the only covariate, PPVT remained significant ($F[1,158]=58.12$, $p<.001$), as did the attempter vs. non-attempter effect ($F[1,158]=6.01$, $p=.015$).

At the level of individual domains with PPVT as a covariate, however, only the difference in the Attention domain remained significant ($F[1,154]=7.65$, $p=.006$). Differences in the aggregate Memory ($F[1,158]=1.91$, $p=.17$), Working Memory ($F[1,158]=2.86$, $p=.09$), and Impulse Control ($F[1,158]=1.75$, $p=.19$) domains were non-significant. Differences in Abstract/Contingent Learning ($F[1,157]=.58$, $p=.45$) and Language Fluency ($F[1,157]=.50$, $p=.48$) remained non-significant.

At the level of individual tests with PPVT as a covariate, Stroop interference ($F[1,152]=7.28$, $p=.008$) remained significant. Although the aggregate Memory domain did

not differ with adjustment for PPVT score, the Buschke SRT score ($F[1,149]=3.88$, $p=.05$), one that has repeatedly distinguished attempters in our previous studies, did. WCST Error ($F[1,158]=3.39$, $p=.07$) approached significance. CPT ($F[1,153]=2.60$, $p=.11$) and A, Not B Timed Reasoning ($F[1,157]=2.52$, $p=.12$) were now non-significant, with Go-No Go in the same range ($F[1,158]=2.02$, $p=.16$). All other test score differences were non-significant.

Lethality and Violence of Past Attempts

In this sample, there were no aggregate differences in neuropsychological performance between higher lethality past attempters (lethality rating of worst attempt >3 ; $n=54$) and lower lethality past attempters (lethality ≤ 3 ; $n=26$). Neither the main effect for lethality grouping ($F[1,78]=1.72$, $p=.19$) nor the interaction with domain ($F[5,390]=1.97$, $p=.08$) were significant (PPVT did not differ between the groups). Both had equivalent levels of depression severity ($t[76]=.47$, $p=.64$).

Similarly, there were no differences between the small subsample of those who used a violent method during their worst attempt ($n=11$) and those who used non-violent means ($n=69$). Violent means included cutting ($n=6$), attempted hanging ($n=3$), jumping from a height ($n=1$) and attempted self-immolation ($n=1$). Main effect for violence grouping ($F[1,78]=.01$, $p=.94$) and the interaction with domain ($F[5,390]=.35$, $p=.88$) were non-significant (again with no difference in PPVT). Violent and non-violent attempters in this sample were equivalent in depression severity ($t[76]=.03$, $p=.98$). Of note, violent attempters had significantly *lower* lethality for their worst attempts than non-violent attempters (1.5 ± 1.6 vs. 2.8 ± 1.7 , $t[78]=2.44$, $p=.02$), lower than in our previous studies.

Remitted Participants

There were 34 past suicide attempters and 36 non-attempters with an HDRS <10 . Demographic and clinical differences were comparable to those in the total sample, with attempters having lower PPVT scores (99.4 ± 13.1 vs. 108.2 ± 12.8 , $t[68]=2.84$, $p=.006$) and less education (13.6 ± 2.5 vs. 15.2 ± 2.2 , $t[68]=2.92$, $p=.005$), and more likely to have Borderline Personality Disorder (23.5% vs. 0.0%; $\chi^2[1]=11.58$, $p=.003$), past substance abuse (60.6% vs. 38.9%; $\chi^2[1]=3.25$, $p=.07$), Post Traumatic Stress Disorder (26.5% vs. 8.3%; $\chi^2[1]=8.71$, $p=.01$), and reported past history of physical or sexual abuse (85.3% vs. 58.3%; $\chi^2[1]=6.22$, $p=.01$). In this subgroup, however, suicidal ideation was comparable in attempters and non-attempters and almost negligible ($.2 \pm 1.2$ vs. $.03 \pm .17$, $t[68]=.88$, $p=.38$), while impulsiveness (46.7 ± 16.0 vs. 38.0 ± 13.8 , $t[52]=2.09$, $p=.04$) and hostility (31.7 ± 10.1 vs. 23.4 ± 9.9 , $t[56]=3.11$, $p=.003$) were greater in attempters.

Neuropsychological domain scores in the remitted sample alone are presented in Figure 2. Without covariates, attempter group differences were significant in all domains (Attention, $t[67]=3.22$, $p=.002$; Memory, $t[68]=2.72$, $p=.008$; Abstract/Contingent Learning, $t[68]=2.53$, $p=.01$; Working Memory, $t[68]=2.49$, $p=.02$; Language Fluency, $t[67]=2.38$, $p=.02$; Impulse Control, $t[68]=2.27$, $p=.03$). On individual tests, attempters differed on CPT d' ($t[67]=2.88$, $p=.005$), Stroop interference ($t[67]=2.45$, $p=.02$), Buschke SRT ($t[66]=2.69$, $p=.009$), WCST error score ($t[68]=2.70$, $p=.009$), A, Not B Timed Reasoning ($t[68]=2.60$, $p=.01$) and Letter Fluency ($t[67]=2.50$, $p=.02$).

Covarying PPVT score, univariate differences were significant in the Attention domain ($F[1,66]=7.44$, $p=.008$) and Abstract/Contingent Learning ($F[1,67]=3.85$, $p=.05$), but non-significant in all other domains. On individual tests, Stroop interference ($F[1,66]=6.18$, $p=.02$), and WCST error score ($F[1,67]=4.97$, $p=.03$) remained significant. Differences on the CPT ($F[1,66]=3.63$, $p=.06$), Buschke SRT ($F[1,65]=3.76$, $p=.06$) and A, Not B Timed Reasoning ($F[1,67]=3.07$, $p=.08$) were marginal.

DISCUSSION

Past suicide attempters in this sample of variably depressed subjects with previous episodes of major depression, performed more poorly on many of the same tests that distinguished past attempters in our previous studies (Keilp et al., 2001; Keilp et al., 2008; Keilp et al., 2013). Before adjustment for differences in estimate intelligence, these included measures of interference processing, memory, and working memory, as well the efficiency of abstract/contingent learning. Even after adjustment for group differences in estimated intelligence, two core measures that have consistently distinguished past attempters in our earlier studies - Stroop interference and Buschke SRT recall - continued to differentiate past attempters here. This replication of our earlier findings in a sample with varying levels of depression severity confirms the importance of these measures as potential markers of suicide risk. Though the magnitude of differences is moderate, these neuropsychological characteristics provide a window into the type of cognitive impairment that may play a role in suicidal behavior. As we have noted in our previous work, the executive control of attention is a fundamental feature of both mindfulness-based therapies for suicidal behavior (Baer, 2003; Bishop et al., 2004) and dialectical behavior therapy (Linehan & Dexter-Mazza, 2008), each of which attempts to enhance the capacity to direct attention away from distressing feeling states. Memory impairments, in the form of over-general autobiographical recall have been linked to suicidal behavior as well (Sinclair et al., 2007; Arie et al., 2008); our data suggest that poor initial learning may play a role in these recall deficits.

It is also noteworthy that the estimated intelligence of the samples in this study is close to population normative levels, and lower than in our previous studies where depressed patients as well as comparison subjects were nearly a standard deviation above normative levels (Keilp et al., 2001; Keilp et al., 2008; Keilp et al., 2013). Thus, our earlier findings are not restricted to comparison groups of exceptional performers. Previous studies have found lower estimated IQ's in groups at risk for suicidal behavior (Gunnell et al., 2005; Sörberg et al., 2013), although it is unclear if IQ itself or life circumstances and other social pathologies that are related to lower IQ are more important as risk factors (see Sörberg et al., 2013). Nonetheless, controlling for intelligence suggests that there are more specific cognitive functions, ones that may covary with overall intelligence, that are more sensitive indicators of suicidal behavior risk.

Our findings with regard to interference processing and memory extended to the subsample of individuals whose depressive symptoms had largely remitted. These results can be compared directly to those of other studies of remitted subjects that have found impairments in decision-making (see Jollant et al., 2011). Our data suggest there are more fundamental deficits in information processing - that may underlie other higher order impairments - and

these persist outside of depressive episodes. After covariation for PPVT in the remitted subsample, the difference in Stroop interference was significant and difference in Buschke SRT recall marginal ($p=.06$), though the effect size of these differences was comparable in both the full and remitted samples (for Stroop, effect size .50 in full sample, .63 in remitted; for Buschke SRT, effect size .42 in full sample, .57 in remitted). Wisconsin Card Sort error score had been marginal in the full sample, but was significantly different between attempters and non-attempters in the remitted subsample, contrary to at least one previous study suggesting that deficits are most likely state-dependent (Marzuk et al., 2005). Of note, however, the Wisconsin Card Sort error score was the only subscore from the WCST that differed between groups. All others (categories, failure to maintain, perseverative error) were comparable, suggesting that the deficit observed on the WCST reflected an inefficiency of learning rather than a failure of abstraction or reasoning abilities.

High and low lethality attempters did not differ significantly in neuropsychological performance as they had in our earlier studies of individuals in a current depressive episode (Keilp et al., 2001; Keilp et al., 2008), although not our most recent study (Keilp et al., 2013). Mean neuropsychological performance levels tended to be poorer in high lethality attempters (data available upon request), but these differences did not reach statistical significance in this mildly depressed sample.

Additionally, those who had used a violent method in their most serious attempt did not differ from those who used a non-violent method. While we and others (Jollant et al., 2005; Keilp et al., 2013) had previously found differences in performance in violent attempters, the small size of the violent attempt sample here and the mild severity of their attempts (requiring only routine medical intervention) made finding any differences unlikely.

Overall, findings here suggest that deficits in interference processing and memory reflect a trait in past attempters. Although performance may be modulated to some degree by current clinical state, past attempters appear to perform consistently more poorly than non-attempters when both groups are in the same clinical state. The persistence of deficits in past attempters may also represent a degree of treatment resistance, at least in the cognitive realm. In the remitted sample, for example, non-attempters' performance falls approximately at normative levels (near a z-score of zero; see Figure 2.) while attempters' performance is at approximately the same level as attempters in the entire sample (Figure 1.). It is difficult to draw any definitive conclusions about the effects of clinical state without examining the same subjects at different points in treatment. Nonetheless, it is clear that relative deficits in past attempters persist outside of depressive episodes.

Limitations of this study include the variability in clinical state and treatment status of the sample. Though this study was designed as an evaluation of the robustness of our prior findings in a community sample reflecting outpatient clinical conditions and treatment practices, the variety of treatments and comorbidities make it difficult to characterize all possible influences on neurocognitive performance. The clinical course of these participants is also unknown, such that some may be improving with/without treatment, some stable, and some worsening in ways that may have impacted their performance. Participants in this study were also predominantly female, and in the vast majority of cases had made their

attempts by drug overdose. Data from a representative sample of male attempters as well as those using more violent methods are needed.

In the future, follow-up studies are needed to track the course of the deficits observed here, as well as their relationship to other measures that have distinguished suicide attempters in other studies, such as the Iowa Gambling Task (Jollant et al., 2010; Bridge et al., 2012) or reversal learning (Dombrovski, et al., 2010). Ideally, these studies might be undertaken in the context of a controlled clinical trial. Mechanistic studies of the role of these deficits in suicidal behavior are also needed. As noted in the Introduction above, the larger study of which this report is a part involved simultaneous evaluation of participants' offspring. In future analyses, we will be able to determine if the deficits observed here run in families, if they aggregate with other risk factors, and if they are markers of prospective risk in offspring.

Acknowledgments

This work was supported by grants from the National Institute of Mental Health (MH56612, MH56390, MH55123, MH66371, MH62185 and MH018951).

REFERENCES

- Arie M, Apter A, Orbach I, Yefet Y, Zalzman G. Autobiographical memory, interpersonal problem solving, and suicidal behavior in adolescent inpatients. *Comprehensive Psychiatry*. 2008; 49(1):22–29. [PubMed: 18063037]
- Baer RA. Mindfulness training as a clinical intervention: A conceptual and empirical review. *Clinical Psychology: Science and Practice*. 2003; 10(2):125–143.
- Barratt, ES. Impulsiveness defined within a system model of personality. In: Spielberger, CD., Butcher, JN., editors. *Advances in Personality Assessment*. Vol. 5. New York: Earlbaum; 1985. p. 113-132.
- Barratt, ES. Impulsivity and aggression. (2003). In: Monahan, J., Steadman, HJ., editors. *Violence and Mental Disorder : Developments in Risk Assessment*. Chicago: University of Chicago Press; 1994. p. 61-80.
- Beck AT, Beck R, Kovacs M. Classification of suicidal behaviors : I. Quantifying intent and medical lethality. *American Journal of Psychiatry*. 1975; 132:285–287. [PubMed: 1115273]
- Beck AT, Kovacs M, Weissman A. Assessment of suicidal ideation: the scale for suicidal ideation. *Journal of Consulting and Clinical Psychology*. 1979; 47:343–352. [PubMed: 469082]
- Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Archives of General Psychiatry*. 1961; 4:53–63.
- Beck AT, Weissman A, Lester D, Trexler L. The measurement of pessimism: the hopelessness scale. *Journal of Consulting and Clinical Psychology*. 1974; 46:861–865.
- Bishop SR, Lau M, Shapiro S, Carlson L, Anderson ND, Carmody J, Segal ZV, Abbey S, Speca M, Velting D, Devins G. Mindfulness: A proposed operational definition. *Clinical Psychology: Science and Practice*. 2004; 11(3):230–241.
- Brent DA, Oquendo M, Birmaher B, Greenhill L, Kolko D, Stanley B, Zelazny J, Brodsky B, Bridge J, Ellis S, Salazar JO, Mann JJ. Familial pathways to early-onset suicide attempt: risk for suicidal behavior in offspring of mood-disordered suicide attempters. *Archives of General Psychiatry*. 2002; 59(9):801–807. [PubMed: 12215079]
- Bridge JA, McBee-Strayer SM, Cannon EA, Sheftall AH, Reynolds B, Campo JV, Pajer KA, Barbe RP, Brent DA. Impaired decision making in adolescent suicide attempters. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2012; 51(4):394–403. [PubMed: 22449645]
- Buss AR, Durkee A. An inventory for assessing different kinds of hostility. *Journal of Consulting Psychology*. 1961; 21:343–349.

- Clark L, Dombrovski AY, Siegle GJ, Butters MA, Shollenberger CL, Sahakian BJ, Szanto K. Impairment in risk-sensitive decision-making in older suicide attempters with depression. *Psychology and Aging*. 2011; 26(2):321–330. [PubMed: 21443349]
- Dombrovski AY, Clark L, Siegle GJ, Butters MA, Ichikawa N, Sahakian BJ, Szanto K. Reward/Punishment reversal learning in older suicide attempters. *American Journal of Psychiatry*. 2010; 167(6):699–707. [PubMed: 20231320]
- Dombrovski AY, Szanto K, Siegle GJ, Wallace ML, Forman SD, Sahakian B, Reynolds CF 3rd, Clark L. Lethal forethought: delayed reward discounting differentiates high- and low-lethality suicide attempts in old age. *Biological Psychiatry*. 2011; 70(2):138–144. [PubMed: 21329911]
- Dunn, LM., Dunn, LM., Dunn, DM. Peabody Picture Vocabulary Test, Third Revision. Circle Pines, MN: American Guidance Service; 1997.
- First, MB., Spitzer, RL., Gibbon, M., Williams, JBW., Loma, B. Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II), (Version 2.0). New York, NY: New York State Psychiatric Institute, Biometric Research Department; 1996.
- Gunnell D, Magnusson PKE, Rasmussen F. Low intelligence test scores in 18 year old men and risk of suicide : cohort study. *British Medical Journal*. 2005; 330:167–171. [PubMed: 15615767]
- Hamilton M. A rating scale for depression. *Journal of Neurology, Neurosurgery and Psychiatry*. 1960; 23:56–62.
- Hamilton M. Development of a rating scale for primary depressive illness. *British Journal of Social and Clinical Psychology*. 1967; 6(4):278–296. [PubMed: 6080235]
- Jollant F, Bellivier F, Leboyer M, Astruc B, Torres S, Verdier R, Castelnau D, Malafosse A, Courtet P. Impaired decision making in suicide attempters. *American Journal of Psychiatry*. 2005; 162(2):304–310. [PubMed: 15677595]
- Jollant F, Lawrence NL, Olié E, Guillaume S, Courtet P. The suicidal mind and brain: A review of neuropsychological and neuroimaging studies. *World Journal of Biological Psychiatry*. 2011 Mar 8. [Epub ahead of print].
- Keilp JG, Gorlyn M, Oquendo MA, Burke AK, Mann JJ. Attention deficit in depressed suicide attempters. *Psychiatry Research*. 2008; 159(1–2):7–17. [PubMed: 18329724]
- Keilp JG, Gorlyn M, Russell M, Oquendo MA, Burke AK, Harkavy-Friedman J, Mann JJ. Neuropsychological function and suicidal behavior: attention control, memory and executive dysfunction in suicide attempt. *Psychological Medicine*. 2013; 43(3):539–551. [PubMed: 22781400]
- Keilp JG, Sackeim HA, Brodsky BS, Oquendo MA, Malone KM, Mann JJ. Neuropsychological dysfunction in depressed suicide attempters. *American Journal of Psychiatry*. 2001; 158(5):735–741. [PubMed: 11329395]
- Keilp JG, Sackeim HA, Mann JJ. Correlates of trait impulsiveness in performance measures and neuropsychological tests. *Psychiatry Research*. 2005; 135(3):191–201. [PubMed: 15996748]
- Linehan, MM., Dexter-Mazza, ET. Dialectical behavior therapy for borderline personality disorder. In: Barlow, DH., editor. *Clinical Handbook of Psychological Disorders: A Step-by-Step Treatment Manual*. 4th. New York, NY, US: Guilford Press; 2008. p. 365–420.
- Malloy-Diniz LF, Neves FS, Abrantes SS, Fuentes D, Corrêa H. Suicide behavior and neuropsychological assessment of type I bipolar patients. *Journal of Affective Disorders*. 2009; 112(1–3):231–236. [PubMed: 18485487]
- Mann JJ, Waternaux C, Haas GL, Malone KM. Toward a clinical model of suicidal behavior in psychiatric patients. *American Journal of Psychiatry*. 1999; 156(2):181–189. [PubMed: 9989552]
- Marzuk PM, Hartwell N, Leon AC, Portera L. Executive functioning in depressed patients with suicidal ideation. *Acta Psychiatrica Scandinavica*. 2005; 112(4):294–301. [PubMed: 16156837]
- McGirr A, Dombrovski AY, Butters MA, Clark L, Szanto K. Deterministic learning and attempted suicide among older depressed individuals: cognitive assessment using the Wisconsin Card Sorting Task. *Journal of Psychiatric Research*. 2012; 46(2):226–232. [PubMed: 22024486]
- Melhem NM, Brent DA, Ziegler M, Iyengar S, Kolko D, Oquendo M, Birmaher B, Burke A, Zelazny J, Stanley B, Mann JJ. Familial pathways to early-onset suicidal behavior: familial and individual antecedents of suicidal behavior. *American Journal of Psychiatry*. 2007; 164(9):1364–1370. [PubMed: 17728421]

- Oquendo, MA., Halberstam, B., Mann, JJ. Risk factors of suicidal behavior: The utility and limitations of research instruments. In: First, M., editor. *Standardized Evaluation in Clinical Practice: Review of Psychiatry*. Vol. 22. Washington, DC: American Psychiatric Publishing; 2003. p. 103-130.
- Riedel M, Möller HJ, Obermeier M, Schennach-Wolff R, Bauer M, Adli M, Kronmüller K, Nickel T, Brieger P, Laux G, Bender W, Heuser I, Zeiler J, Gaebel W, Seemüller F. Response and remission criteria in major depression--a validation of current practice. *Journal of Psychiatric Research*. 2010; 44(15):1063–1068. [PubMed: 20447651]
- Sinclair JM, Crane C, Hawton K, Williams JM. The role of autobiographical memory specificity in deliberate self-harm: correlates and consequences. *Journal of Affective Disorders*. 2007; 102(1–3): 11–18. [PubMed: 17258815]
- Sörberg A, Allebeck P, Melin B, Gunnell D, Hemmingsson T. Cognitive ability in early adulthood is associated with later suicide and suicide attempt: the role of risk factors over the life course. *Psychological Medicine*. 2013; 43(1):49–60. [PubMed: 22617391]
- Spitzer, RL., Williams, JBW., Gibbon, M., First, MB. *Instruction manual for the structured clinical interview for the DSM-IV (SCID-P)*. Washington, DC: American Psychiatric Press; 1990.
- Spren, O., Strauss, E. *A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary*. 3rd. New York, NY: Oxford University Press; 2006.
- Westheide J, Quednow BB, Kuhn KU, Hoppe C, Cooper-Mahkorn D, Hawellek B, Eichler P, Maier W, Wagner M. Executive performance of depressed suicide attempters: the role of suicidal ideation. *European Archives of Psychiatry and Clinical Neuroscience*. 2008; 258(7):414–421. [PubMed: 18330667]

Figure 1. Neuropsychological performance in adults with a lifetime history of depression, with and without a past history of suicidal behavior.

Figure 1. Neuropsychological performance in adults with remitted depression (current HDRS < 10), with and without a past history of suicidal behavior.

Table 1

Demographic and Clinical Rating Data

Variable	Non-Attempters	Suicide Attempters	p-value ^I
N	81	80	--
Age (yrs.)	44.6 ± 7.7	42.0 ± 9.1	.06
Sex (% Female)	84.0% (68)	93.8% (75)	.05*
Education (yrs.)	14.4 ± 2.3	13.5 ± 2.3	.01*
Peabody Picture Vocabulary	102.6 ± 13.2	98.0 ± 12.1	.02*
Hamilton Depression Scale	11.7 ± 8.5	13.4 ± 10.3	.25
Beck Depression Inventory	14.3 ± 10.8	16.5 ± 11.5	.23
Beck Hopelessness Scale	7.2 ± 5.6	8.7 ± 6.6	.13
Scale for Suicidal Ideation (current)	0.2 ± 1.4	2.5 ± 5.5	.001*
Scale for Suicidal Ideation (prior to enrollment)	0.5 ± 2.3	1.8 ± 5.2	.06
Bipolar Diagnosis	17.3% (14)	21.3% (17)	.48
Comorbid Borderline Personality Disorder	4.9% (4)	30.0% (24)	<.001*
Substance Abuse/Dependence	40.7% (33)	59.5% (47)	.02*
Post Traumatic Stress Disorder (Lifetime)	22.2% (18)	36.3% (29)	.001*
Past History of Abuse	64.2% (52)	82.3% (65)	.01*
Psychotropic Medication	49.4% (40)	66.3% (53)	.03*
Any Medication	90.1% (73)	86.3% (69)	.45
Number Previous Suicide Attempts (Attempters only)	--	2.4 ± 1.7	--
Maximum Lethality of Attempt (Attempters only)	--	2.6 ± 1.7	--
Violent Method of Attempt	--	13.8% (11)	--
Barratt Impulsiveness Scale	50.3 ± 19.2	51.6 ± 17.3	.71
Buss Durkee Hostility Inventory	31.0 ± 13.89	34.9 ± 10.8	.07

^I t-test for continuous variables, chi-squared for categorical variables.

Table 2

Neuropsychological Measures (Z-Scores)

Variable	Non-Attempters	Suicide Attempters	p-value ^I
<u>Attention</u>	.08 ± .68	-.32 ± .92	.002 *
CPT (d')	.07 ± .84	-.25 ± 1.04	.04 *
Stroop Interference	.11 ± .93	-.39 ± 1.31	.008 *
<u>Memory</u>	-.02 ± 1.01	-.39 ± .97	.02 *
Buschke SRT (total)	-.20 ± 1.21	-.73 ± 1.27	.009 *
Benton VRT (error)	.16 ± 1.26	-.04 ± 1.15	.28
<u>Abstract/Contingent Learning</u>	-.25 ± .64	-.38 ± .63	.20
Wisconsin Card Sort (categories)	-.28 ± .61	-.39 ± .66	.27
Wisconsin Card Sort (fail to maintain)	-.23 ± 1.12	-.20 ± .98	.88
Wisconsin Card Sort (error)	-.24 ± .77	-.55 ± .87	.02 *
Wisconsin Card Sort (perseverative error)	-.50 ± .65	-.57 ± .91	.62
<u>Working Memory</u>	-.25 ± .99	-.67 ± 1.12	.01 *
N-Back (d')	-.27 ± 1.11	-.62 ± 1.12	.05 *
A, Not B Timed Reasoning	-.18 ± 1.29	-.72 ± 1.58	.02 *
<u>Language Fluency</u>	.13 ± .97	-.12 ± .96	.10
Letter Fluency	.15 ± .98	-.11 ± 1.01	.11
Category Fluency	.11 ± 1.17	-.14 ± 1.16	.19
<u>Impulse Control</u>	.14 ± .76	-.07 ± .69	.07
Go-No Go Commission Error (Log)	.09 ± 1.17	-.23 ± .99	.06
Time Production (Deviation)	.18 ± .87	.10 ± .99	.59

^I t-test for continuous variables, chi-squared for categorical variables.

* p .05