Is current preoperative frailty assessment adequate?

Gilgamesh Eamer, MD, MSc
 Jennifer A. Gibson, MSN
 Chelsia Gillis, RD, MSc
 Amy T. Hsu, PhD
 Marian Krawczyk, PhD
 Emily MacDonald, RN, MN
 Reid Whitlock, MSc
 Rachel G. Khadaroo, MD, PhD

Accepted Apr. 5, 2017; Early-released
 Oct. 1, 2017

Correspondence to:

R. Khadaroo
 Department of Surgery
 University of Alberta
 8440-112 St
 2D Walter MacKenzie Centre
 Edmonton AB T6G 2B7
 khadaroo@ualberta.ca

DOI: 10.1503/cjs.001417

SUMMARY

Preoperative frailty predicts adverse postoperative outcomes. Recommendations for preoperative assessment of elderly patients include performing a frailty assessment. Despite the advantages of incorporating frailty assessment into surgical settings, there is limited research on surgical health care professionals' perception and use of frailty assessment for perioperative care. We surveyed local health care employees to assess their attitudes toward and practices for frail patients. Nurses and allied health professionals were more likely than surgeons to agree frailty should play a role in planning a patient's care. Lack of knowledge about frailty issues was a prominent barrier to the use of frailty assessments in practice, despite clinicians understanding that frailty affects their patients' outcomes. Results of this survey suggest further training in frailty issues and the use of frailty assessment instruments is necessary and could improve the uptake of such tools for perioperative care planning.

As the baby boom population ages, increasing numbers of patients aged 65 years and older are presenting with surgical disease. They are at increased risk for prolonged hospital admission.¹ Although age is commonly used to assess surgical risk, it has been well established that preoperative frailty is more accurate at predicting postoperative outcomes.¹ Frailty is a multisystem syndrome of low physiologic reserves resulting in increased risk for adverse events.² It is increasingly recognized as an important determinant of postoperative complications and recovery. Frailty screening and the implementation of early interventions has been associated with preserved autonomy and reduced adverse events. Well-designed studies examining comprehensive geriatric assessment among surgical patients have shown improved outcomes, mostly in orthogeriatric populations. A systematic review by the Cochrane collaboration is ongoing to assess the robustness of these findings.³ Recommendations developed by the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society⁴ suggest the use of multiple preoperative assessments, including preoperative assessment of each patient for frailty. Despite mounting evidence, frailty assessment is not routine surgical practice,⁵ and it remains unclear why.

Frailty assessment is not currently well taught to surgeons or surgical nursing staff, and little is known about interdisciplinary surgical health care providers' perception of frailty or its role in clinical assessment. We have recently undertaken a survey of our surgical staff at the University of Alberta Hospital to assess their beliefs about frailty and the barriers to frailty assessment, and we compared the perspectives across health care professions. The survey was distributed to all health care providers involved in the care of general surgery patients at our institution. It assessed the attitudes of 3 subgroups: surgeons, nurses and allied health professionals.⁶

Previous research has found that frailty assessment and management improves patient outcomes in both medical and surgical patients; however, it continues to have low uptake in most surgical settings. We found the highest

uptake among allied health practitioners; frailty assessment is less frequent outside of allied health professionals. Specifically, surgeons were less likely than both allied health professionals and nurses to use frailty in guiding patient care. A qualitative investigation by Age UK⁷ found frailty is viewed as something surgeons “know when they see,” yet numerous studies have reported that perceived frailty varies individually and is an inadequate proxy for measured frailty. This suggests that although health care professionals, particularly surgeons, acknowledge that frailty is an important factor in patients’ outcomes, they are overly reliant on their “gut” impression of a patient’s frailty and do not screen for or manage patients based on their frailty. Furthermore, we identified 4 key barriers to surgical care for frail patients: hospital-specific/institutional, health care system, professional knowledge, and patient/family members. These barriers lead to lack of confidence in conducting frailty assessments and inadequate delivery of elder-friendly surgical care.

A number of authors have identified similar knowledge gaps.⁵ Successfully addressing these gaps will require awareness of how health care professionals navigate relevant system complexities and constraints in their provision of care. Given that most health care delivery is based on a single problem-oriented diagnostic model and that health care professionals may not be trained to focus on the holistic care of patients, system reorganization around frailty is challenging. Furthermore, frailty is an evolving area of inquiry, and consensus has not yet identified a single optimal tool to identify frailty. That said, the availability of validated and rapidly administered tools permits the use of quick, reliable and easily interpreted frailty assessments in fast-paced surgical environments.

Our survey was limited by a low response rate, which raises the risk of response bias, and was limited by its single-centre design. There was strong agreement, however, for most items across the health professions, and the results were consistent with predicted attitudes of each profession.

It is clear that the use of formal frailty assessment tools has not been widely adopted in practice. Much of this may be due to inadequate education surrounding the effect of frailty on outcomes and tools to permit rapid assessment, particularly for surgeons. Creating a program to educate surgeons about the importance of frailty assessment — such as how it can improve their patients’ care and, most importantly, how to perform a rapid validated assessment — is key to improving uptake. Further research of comprehensive geriatric

assessment for surgical patients should also be performed to determine if it is effective outside of orthogeriatric patient populations. Addressing barriers to frailty assessment and high-quality care for frail patients could substantially improve care and postoperative outcomes for this vulnerable population.

Acknowledgements: The authors acknowledge the support of the Canadian Frailty Network Interdisciplinary Fellowship Program.

Affiliations: From the Department of Surgery, University of Alberta, Edmonton, Alta. (Eamer, Khadaroo); the School of Public Health, University of Alberta, Edmonton, Alta. (Eamer); the School of Nursing, University of British Columbia, Vancouver, BC (Gibson); the Cumming School of Medicine, University of Calgary, Calgary, Alta. (Gillis); the Clinical Epidemiology Program, Ottawa Hospital Research Institute, Ottawa, Ont. (Hsu); the Department of Epidemiology and Community Medicine, University of Ottawa, Ottawa, Ont. (Hsu); the Centre for Health Evaluation and Outcome Sciences, St. Paul’s Hospital, Vancouver, BC (Krawczyk); the Trinity Western University, Langley, BC (Krawczyk); the Department of Nursing and Health Sciences, University of New Brunswick, Saint John, NB (MacDonald); and the Department of Community Health Sciences, University of Manitoba, Winnipeg, Man. (Whitlock).

Competing interests: J. Gibson received training funding from the Canadian Frailty Network and is supported by the Government of Canada through the Networks of Centres of Excellence (NCE) Program; she has received travel assistance from these programs to attend academic conferences. No other competing interests declared.

Contributors: All authors contributed substantially to the conception, writing and revision of this article and approved the final version for publication.

References

1. Harari D, Hopper A, Dhese J, et al. Proactive care of older people undergoing surgery (‘POPS’): designing, embedding, evaluating and funding a comprehensive geriatric assessment service for older elective surgical patients. *Age Ageing* 2007;36:190-6.
2. Vermeiren S, Vella-Azzopardi R, Beckwée D, et al. Frailty and the prediction of negative health outcomes: a meta-analysis. *J Am Med Dir Assoc* 2016;17:1163.e1-e17.
3. Eamer G, Taheri A, Chen SS, et al. Comprehensive geriatric assessment for improving outcomes in elderly patients admitted to a surgical service. In: Khadaroo RG, editor. *Cochrane Database of Systematic Reviews*. Chichester (UK): John Wiley & Sons, Ltd; 2017.
4. Chow WB, Rosenthal RA, Merkow RP, et al. Optimal preoperative assessment of the geriatric surgical patient. A best practices guideline from the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society. *J Am Coll Surg* 2012;215:453-66.
5. Partridge JSL, Collingridge G, Gordon AL, et al. Where are we in perioperative medicine for older surgical patients? A UK survey of geriatric medicine delivered services in surgery. *Age Ageing* 2014;43:721-4.
6. Eamer G, Gibson JA, Gillis C, et al. Surgical frailty assessment: a missed opportunity. *BMC Anesthesiology* 2017;17:99.
7. Nicholson C, Gordon AL, Tinker A. Changing the way “we” view and talk about frailty. *Age Ageing* 2017;46:349-51.