Antiangiogenic effects of decorin restored by unfractionated, low molecular weight, and nonanticoagulant heparins

Amy K. L. Chui,¹ Tilini N. Gunatillake,¹ Vera Ignjatovic,^{2,3} Paul T. Monagle,^{2,3} Padma Murthi,⁴⁻⁶ Shaun P. Brennecke,^{6,7} John M. Whitelock,⁸ and Joanne M. Said^{1,9}

¹Department of Obstetrics and Gynaecology, Sunshine Hospital, The University of Melbourne, St Albans, VIC, Australia; ²Department of Clinical Haematology and

³Department of Paediatrics, Murdoch Childrens Research Institute, The Royal Children's Hospital, The University of Melbourne, Parkville, VIC, Australia; ⁴Department of

Medicine, School of Clinical Sciences, Monash University, Clayton, VIC, Australia; ⁵The Ritchie Centre, Hudson Institute of Medical Research, Clayton, VIC, Australia;

⁶Department of Maternal-Fetal Medicine Pregnancy Research Centre, The Royal Women's Hospital, Parkville, VIC, Australia; ⁷Department of Obstetrics and Gynaecology, The

Royal Women's Hospital, The University of Melbourne, Parkville, VIC, Australia; ⁸Graduate School of Biomedical Engineering, University of New South Wales, Kensington,

NSW, Australia; and ⁹Maternal Fetal Medicine, Sunshine Hospital, St Albans, VIC, Australia

Key Points

- UFH, LMWH, and NAC restored angiogenesis in decorin-reduced endothelial cells.
- NAC treatment was similar to, or better than, UFH or LMWH at improving endothelial angiogenesis without increasing anticoagulant activity.

Pregnancies affected by preeclampsia (PE) or fetal growth restriction (FGR) display increases in thrombin generation and reductions in angiogenesis and cell growth. There is significant interest in the potential for low molecular weight heparins (LMWHs) to reduce the recurrence of PE and FGR. However, LMWH is associated with an increased risk of bleeding. Therefore, it is of vital importance to determine the exact molecular function of heparins in pregnancy if they are used as therapy for pregnant women. We aimed to determine this using our model for PE/FGR in microvascular endothelial cells. The expression of decorin, a proteoglycan, was reduced to mimic PE/FGR in these cells compared with controls. Four concentrations of unfractionated heparin (UFH), LMWH, and nonanticoagulant heparin (NAC) were added to determine the effect on thrombin generation, angiogenesis, and cell growth. Treatment with UFH and LMWH reduced thrombin generation and restored angiogenesis but decreased cell growth. Treatment with NAC did not affect thrombin generation, restored angiogenesis, and showed a trend toward cell growth. In conclusion, treatment with NAC produced the same, if not better, results as treatment with UFH or LMWH, without the same impact on coagulation. Therefore, NAC could potentially be a better therapeutic option for prevention of PE/FGR in high-risk women, without the risk of the adverse effects of traditional anticoagulants.

Introduction

A successful pregnancy requires the establishment and proper development of an adequate placental circulation. Placenta-mediated pregnancy complications include preeclampsia (PE), fetal growth restriction (FGR), and late pregnancy loss. It has been documented that these complications are associated with abnormal placental development, with underdeveloped placental vasculature or placental inflammation.^{1,2} In addition, normal pregnancy is considered a hypercoagulable state and a period of increased risk of thrombotic complications. This is accompanied by all the elements of Virchow's triad: hypercoagulability, venous stasis, and vascular damage.³ Therefore, thrombosis in the placental bed is also at least partially responsible for placenta-mediated complications.⁴⁻⁶

Apart from the devastating effects of these pregnancy complications on both maternal and fetal health, the risk of recurrent placenta-mediated pregnancy complications is substantial. For example, women affected by prior severe PE have a 25% to 65% risk of recurrent PE, a 3% risk of placental abruption, and a 10% risk of small-for-gestation-age babies.^{6,7} There are currently no highly effective preventive

strategies that can be used for the prevention of these complications. Aspirin has been shown to offer a small relative risk reduction in patients with prior PE and small-for-gestation-age babies; however, meta-analyses suggest it is only effective if started within the first 16 weeks of pregnancy.^{8,9}

In recent years, anticoagulants such as heparin have been used increasingly to try to prevent recurrent pregnancy complications by reduction of thrombosis or utilization of the anti-inflammatory and proangiogenic functions of heparin. Although several randomized controlled trials have been undertaken to determine whether low molecular weight heparin (LMWH) can prevent recurrent pregnancy complications, the results have not been universal, and the studies have been underpowered and had inconsistent patient criteria.¹⁰⁻¹² Therefore, although it seems that LMWH could be a promising therapeutic treatment for recurrent pregnancy complications, there is still insufficient evidence proving the efficacy and impact of LMWH treatment during pregnancy. Furthermore, adoption of this intervention without sufficient information of its potential benefits and harms could expose women to the risk of undesirable and potentially fatal adverse effects, such as major bleeding, heparin-induced thrombocytopenia, osteoporotic fractures, and withholding of epidural analgesia.^{13,14} Therapy is also associated with regular injections and substantial costs.

Heparin is a glycosaminoglycan composed of chains of alternating residues of d-glucosamine and uronic acid. Its unique pentasaccharide structure has a high binding affinity to antithrombin,¹⁵ and this mediates the majority of the anticoagulant effect. Unfractionated heparin (UFH) has been used for many indications during pregnancy. It is a large molecule that does not cross the placenta and therefore does not cause teratogenic effects, unlike coumarin (Warfarin). The main adverse effects of UFH are inconvenience, with a minimum of twice daily injections, and the potential for osteoporosis and heparin-induced thrombocytopenia. LMWHs, in contrast, have become the preferred anticoagulant because they are equivalent or superior to UFH in efficacy and safety in the treatment of thrombotic problems outside of pregnancy.¹⁶⁻¹⁹ Several studies have suggested the safety and efficacy of LMWH during pregnancy.^{13,20,21} In addition, the risk of adverse heparin-induced thrombocytopenia and osteoporosis after LMWH treatment is greatly reduced compared with UFH.²⁰⁻²⁵

In our previous work, we showed that the messenger RNA and protein expression of the proteoglycan decorin (DCN) was reduced in PE- and FGR-affected placentae compared with controls.²⁶⁻²⁸ Furthermore, we also showed that downregulation of *DCN* in human microvascular endothelial cells resulted in decreased cell growth and proliferation, decreased network formation (angiogenesis), and a modest increase in thrombin generation.²⁶ We also demonstrated that DCN was downregulated in the first trimester in women who went on to develop growth-restricted infants, suggesting that there is both a temporal relationship and biologic plausibility for the association between reduced DCN expression and subsequent development of PE/FGR.²⁹ The results from these studies have led to the opportunity to investigate whether the addition of heparin to these *DCN*-reduced microvascular endothelial cells may result in a reversal of adverse cell growth and angiogenesis, as well as an expected anticoagulation effect. In addition, we also aimed to investigate the effect of adding a nonanticoagulant (NAC), de-N-desulfated heparin, to the *DCN*-reduced cells to

determine whether this heparin could also reverse the decrease in cell growth and angiogenesis observed, but without the detrimental anticoagulant effects during pregnancy. We used a wide range of heparin concentrations from the therapeutic doses according to the American College of Chest Physicians guidelines,³⁰ with one concentration above and below these values. This study will be the first to determine the effect of heparin (UFH, LMWH, and NAC) on cultured microvascular endothelial cells and determine whether this addition may help to reverse some of the adverse effects of *DCN* downregulation.

Methods

Cell lines

The telomerase-immortalized human microvascular endothelial cells from neonatal foreskin (TIME) were purchased from American Type Culture Collection (CRL-4025; Manassas, VA). TIME cells were cultured in Microvascular Endothelial Cell Growth Medium-2 (EGM-2 MV; Single Quot Kit; catalog number CC-4147; Lonza/Clonetics, Mt Waverley, VIC, Australia) containing 10% fetal bovine serum (Murdoch Childrens Research Institute Tissue Culture Supplies, Parkville, VIC, Australia) at 20% oxygen culture conditions. These cells have been shown to represent the functional and morphological characteristics of human placental endothelial cells.³¹

Reduction of *DCN* expression by siRNA

Four independent *DCN* small interference (siRNA) oligonucleotides were obtained as 4-For-Silencing siRNA Duplexes (Qiagen, Chadstone, VIC, Australia). The *DCN* siRNAs showed no significant DNA sequence similarity to other genes in GenBank complementary DNA databases (data not shown).²⁶

TIME cells were grown in EGM-2 MV and transfected with *DCN* siRNAs using HiPerfect transfection reagent (Qiagen). Negative control (NC) siRNA consisted of a pool of enzyme-generated siRNA oligonucleotides that were not specific for any known human genes (AllStars Negative siRNA; Qiagen).²⁶ The Mock control represented the cells in media only. The efficacies of the 4 siRNAs in downregulating *DCN*, and the subsequent selection of the 2 best siRNA as DCNS2 and DCNS3, are depicted in our previous published work.²⁶

Heparins and concentrations used

The following heparin concentrations were used: UFH, heparin sodium 5000 IU/mL (Pfizer, West Ryde, NSW, Australia); LMWH, dalteparin 10 000 IU/mL; and NAC, de-N-sulfated heparin sodium salt 5 mg, purchased from Sigma (Castle Hill, NSW, Australia) as prepared from porcine mucosal heparin by a modification of the method to render it completely without anticoagulant activity.³²

The actual concentration of heparins used was adjusted for tissue culture experiments to correspond to those used in an adult at prophylactic and therapeutic ranges. Therefore, because the therapeutic range of UFH in adults is usually aimed at 0.35 to 0.7 IU/mL as measured by anti-Xa activity, we used UFH at 0.15, 0.35, 0.5, and 1.0 IU/mL to cover this range. Similarly, the therapeutic range of LMWH in adults is between 0.5 and 1.0 IU/mL, also measured by anti-Xa activity; our LMWH concentrations were 0.15, 0.5, 0.7, and 1.2 IU/mL. Because NAC has not been used therapeutically, we tried to use concentrations similar to those used for UFH and LMWH as milligrams per milliliter. The concentrations we used for NAC were 0.010, 0.026, 0.040, and 0.073 mg/mL.

Stock solutions of UFH, LMWH, and NAC were made up in complete EGM-2 MV media and used throughout all experiments that contained the heparins.

TIME cell growth using the xCELLigence system with addition of UFH, LMWH, and NAC

TIME cell growth was assessed using the xCELLigence SP real-time system (Roche Diagnostics, Melbourne, VIC, Australia) according to the manufacturer's instructions.²⁶ Briefly, cells were prepared and added to the E-Plate 96 at a density of 5000 cells per well and transfected with *DCN* siRNAs or Mock/NC for 48 hours in 4 concentrations of each of UFH, LMWH, or NAC (Roche Diagnostics). The xCELLigence system recorded the background electrical impedance for 72 hours. The cell index was calculated using the RTCA-integrated software (version 1.2; John Morris Scientific, Melbourne, VIC, Australia), and the data were analyzed using GraphPad Prism 5 (GraphPad Software, San Diego, CA).

TIME cell network formation assays with addition of UFH, LMWH, and NAC

TIME cell network formation was assessed using the μ -Slide Angiogenesis system (IBIDI, Hallam, VIC, Australia).²⁶ Briefly, TIME cells were transfected with *DCN* siRNA or Mock/NC in 24-well plates with 4 concentrations of each of UFH, LMWH, or NAC and incubated for 24 hours. After the treatment, the cells were dissociated using enzyme-free cell dissociation solution (Millipore, Billerica, MA). The μ -Slide Angiogenesis wells were coated with 10 μ l of neat Growth-Factor Reduced Matrigel (BD, Scoresby, VIC, Australia) and allowed to polymerize for 1 hour at room temperature. The cells were seeded into the wells of the slide at a density of 8000 cells per well and returned to the incubator for an additional 24 hours, for a total siRNA/Mock/NC plus heparin incubation time of 48 hours. The media was then removed, stained with calcein-AM (Millipore), and visualized under the Olympus BX53 fluorescence microscope (Olympus, Tokyo, Japan). Photomicrographs at a magnification of $\times 10$ of entire wells were taken in triplicate, and the branching points between the TIME cells were counted by Wimasis image analysis (Munich, Germany).

Thrombin-generation assays with addition of UFH, LMWH, and NAC

TIME cells were plated into 96-well plates at a density of 5000 cells per well and transfected with *DCN* siRNAs or Mock/NC with 4 concentrations of each of UFH, LMWH, or NAC for 48 hours. Lyophilized Coag-Norm plasma (Diagnostica Stago, Doncaster, VIC, Australia) was obtained and resuspended as per manufacturer's instructions. This was then spiked with 4 concentrations of UFH, LMWH, or NAC. Measurement of endogenous thrombin potential (ETP) by calibrated automated thrombogram (CAT; Thromboscope; Diagnostica Stago) was performed according to the manufacturer's instructions.²⁶ All experiments were conducted in triplicate wells. The ETP represents the total enzymatic activity performed by thrombin, which is considered the most predictive parameter of bleeding/thrombosis risk.^{33,34} The ETP (nanomolar per minute) was calculated using the Thromboscope software (version 3.0.0.29; Diagnostica Stago) and represented as the area under the thrombin-generation curve.

Data analysis

All data in this study are described as mean $\pm$ standard error of the mean and were analyzed by the GraphPad Prism 6 statistical software (GraphPad Software). One-way analysis of variance (ANOVA) was used to assess the differences in *DCN* function between siRNA-treated and Mock/NC groups with or without added heparins. A probability value of $< .05$ was considered statistically significant.

Results

Thrombin-generation potential

Treatment with all 4 concentrations of UFH and LMWH decreased thrombin-generation levels of control and *DCN*-reduced TIME cells, with the exception of NAC, which had no statistically significant effects on thrombin generation at lower concentrations of NAC but a modest effect at the highest dose of NAC.

In Figure 1, the panels on the left column are representative images of the output from the CAT machine showing the thrombin-generation curve and how the ETP, or area under the curve, is calculated. The y-axis is in nanomolars and the x-axis is the time in minutes. Using the ETP calculated by the CAT program, treatment with all 4 concentrations of UFH or LMWH resulted in a significant dose-dependent decrease in the thrombin-generation potential of *DCN*-reduced TIME cells compared within each treatment group ($P < .00001$; $n = 9$; 1-way ANOVA; Figure 1A-B). However, treatment with NAC did not result in a decrease in ETP when compared within each individual treatment group ($P > .05$; $n = 9$; 1-way ANOVA), except at the higher concentrations of 0.6 IU (Mock vs Mock 0.6 IU: $P < .01$; $n = 9$; 1-way ANOVA) or 1.2 IU (NC vs NC 1.2 IU, D2 vs D2 1.2 IU, and D3 vs D3 1.2 IU: $P < .00001$; $n = 9$; 1-way ANOVA; Figure 1C).

Network formation

Treatment with all 4 concentrations of UFH, LMWH, or NAC restored the network-formation abilities of *DCN*-reduced TIME cells. Figures 2-4 show representative images of the network-formation potential of TIME cells, quantitated by the number of branch points.

UFH. Treatment with the midconcentrations of UFH 0.35 and 0.5 IU, but not the highest, resulted in a significant increase in branch points in Mock (0.35 IU: $P < .0001$; 0.5 IU: $P < .05$; $n = 9$; 1-way ANOVA) or NC (0.15 IU: $P < .05$; 0.5 IU: $P < .00001$; $n = 9$; 1-way ANOVA). Importantly, treatment with almost all 4 concentrations of UFH resulted in an increase in branch points in *DCN*-reduced cells back to non-heparin-treated Mock/NC levels ($P < .05$ to $P < .00001$; $n = 9$; 1-way ANOVA; Figure 2A-B).

LMWH. Unlike UFH, treatment with the low concentrations to midconcentrations of LMWH, but not the higher concentrations, significantly increased branch points back to non-heparin-treated Mock/NC levels in *DCN*-reduced cells ($P < .001$ to $P < .00001$; $n = 9$; 1-way ANOVA), without affecting Mock- or NC-treated cells (Figure 3A-B).

NAC. Similar to LMWH, only treatment with the low concentrations to midconcentrations of NAC significantly restored branch points to non-heparin-treated Mock/NC levels in *DCN*-reduced cells ($P < .05$ to $P < .001$; $n = 9$; 1-way ANOVA), without affecting heparin-treated Mock or NC cells (Figure 4A-B).

Figure 1. Treatment with all 4 concentrations of UFH and LMWH decreased thrombin generation levels of control and *DCN*-reduced TIME cells, with the exception of NAC. The ETP of the TIME cells after reduction of *DCN* and incubated with heparins was determined using the CAT system. Representative images from the CAT output are also shown for each heparin used, where the y-axis represents the ETP in nanomolars and the x-axis shows the time in minutes. (A) Treatment without heparin and treatment with all 4 concentrations of UFH (A), LMWH (B), and NAC (C) in a dose-dependent manner. * $P < .01$, **** $P < .00001$ (both significant); $n = 9$; 1-way ANOVA. The y-axis represents the ETP (nanomolars per minute).

Cell growth

Treatment with all 4 concentrations of UFH or LMWH, but not NAC, reduced cell growth of *DCN*-reduced TIME cells at 48 hours, as shown in Figure 5. Graphical representation of cell index from time point 0 to 48 hours is also shown.

UFH. Figure 5A is representative graph showing the cell index of TIME cells treated with *DCN* siRNA or Mock/NC incubated with 0.15, 0.35, 0.5, and 1.0 IU of UFH over 48 hours in culture. All 4 concentrations of UFH resulted in a significant decrease in cell

growth in cells treated with Mock, NC, or *DCN* siRNA ($P < .00001$; $n = 9$; 1-way ANOVA).

LMWH. The results after incubation with 0.15, 0.5, 0.7, and 1.2 IU of LMWH are shown in Figure 5B. Once again, all 4 concentrations of LMWH resulted in a significant decrease in cell growth in all cells treated ($P < .0001$ to $P < .00001$; $n = 9$; 1-way ANOVA).

NAC. Treatment with 0.15, 0.4, 0.6, and 1.1 IU of NAC resulted in a significant decrease in cell growth in Mock- and NC-treated cells ($P < .01$; $n = 9$; 1-way ANOVA), but not in *DCN*-reduced cells, and

Figure 2. Treatment with all 4 concentrations of UFH restored the network-formation abilities of DCN-reduced TIME cells. The ability of TIME cells to form networks after *DCN* siRNA or Mock/NC was determined using the μ -Slide Angiogenesis system by IBIDI. (A) Representative image of the networks stained with calcein-AM is shown for each concentration. Images were acquired using the Olympus BX53 fluorescence microscope and represent 10 $\times$ magnification to obtain the whole field. (B) The network-formation potential of TIME cells after treatment with 4 concentrations of UFH calculated by the Wimasys software. * $P < .05$, ** $P < .001$, *** $P < .0001$, **** $P < .00001$ (all significant); $n = 9$; 1-way ANOVA. The y-axis represents the number of branch points.

did not restore cell growth to non-heparin-treated Mock/NC levels ($P > .05$ to $P < .0001$; $n = 9$; 1-way ANOVA; Figure 5C).

Discussion

Over the recent years, an increasing number of pregnant women have been given heparins to prevent a wide range of pregnancy-related complications, such as recurrent miscarriages,³⁵ pregnancy-related thromboembolism,³⁶ and recurrent PE/FGR. The most common heparin being used is LMWH, which has a lower risk of serious adverse effects when used over a short time during pregnancy. Heparins are complex macromolecules with diverse actions that extend beyond purely anticoagulation.³⁷ Therefore, because pregnancy disorders such as PE and FGR are also complex

pathologies, it makes sense to further elucidate the mechanisms by which heparins affect placental development.

Previous randomized controlled trials have reported a lack of efficacy of LMWH in the prevention of recurrent miscarriages in thrombophilia⁺ and thrombophilia⁻ women.³⁸⁻⁴⁰ However, smaller trials have demonstrated that subcutaneous LMWH improved perinatal outcomes in thrombophilia⁻ women with previous severe PE.⁴¹ In addition, over recent years, the activity and functional role of NAC heparins in different forms have been widely investigated, including in angiogenesis and pulmonary diseases.⁴²⁻⁴⁴ The anti-inflammatory role, via inhibition of leukocyte infiltration,⁴⁵ and anti-metastatic role in lung and colon⁴⁶ and pancreatic cancer cells⁴⁷ of these NAC heparins has also been established. However, their efficacy and safety in pregnancy have not been investigated.

Figure 3. Treatment with all 4 concentrations of LMWH restored the network-formation abilities of *DCN*-reduced TIME cells. (A)

Representative image of the networks stained with calcein-AM is shown for each concentration. Images were acquired using the Olympus BX53 fluorescence microscope and represent 10× magnification to obtain the whole field. (B) The network-formation potential of TIME cells after treatment with LMWH calculated by the Wimasis software. $**P < .001$, $***P < .0001$, $****P < .00001$ (all significant); $n = 9$; 1-way ANOVA. The y-axis represents the number of branch points.

In this study, we used UFH, LMWH, and NAC to determine whether addition of these agents would potentially reverse the effects of *DCN* downregulation, specifically network formation and cell growth.²⁶ As expected, UFH and LMWH decreased the thrombin generation of the cells. In contrast, NAC had no effect on thrombin generation. Thrombin generation, with resultant placental infarcts, does not seem to be the main culprit in the development of pregnancy complications such as FGR and PE.⁴⁸ Therefore, this result is significant because the use of NAC could potentially eliminate the clinical bleeding risks involved with using either UFH or LMWH in high-risk pregnant women. Moreover, the reduction of anticoagulant activity means that women are not prevented from accessing epidurals during labor and birth.

The addition of UFH, LMWH, and NAC to *DCN*-downregulated cells significantly reversed the reduction in network formation to at

least Mock control or NC levels. However, UFH, the strongest anticoagulant used, also significantly increased network formation abilities of the cells in Mock controls and NCs, causing us to speculate that if this were used in women without pregnancy complications, the angiogenic potential of the microvascular endothelial cells in the placenta and potentially throughout the system would be increased. The effect and/or consequence of this increase have not been previously considered. However, the use of the lowest concentration of UFH, at 0.15 IU/mL, restored network formation to control level without affecting the controls themselves. Therefore, there is potential for the use of a lower concentration of UFH than what is currently used therapeutically. In contrast, this increase did not occur in controls treated with low doses to mid-doses of LMWH or NAC. The ability for LMWH to improve in vitro angiogenesis in serum of high-risk women to the equivalent of serum from low-risk

Figure 4. Treatment with all 4 concentrations of NAC restored the network-formation abilities of *DCN*-reduced TIME cells. (A)

Representative image of the networks stained with calcein-AM is shown for each concentration. Images were acquired using the Olympus BX53 fluorescence microscope and represent 10× magnification to obtain the whole field. (B) Network formation of TIME cells after treatment with NAC calculated by the Wimasis software. * $P < .05$, ** $P < .001$ (both significant); $n = 9$; 1-way ANOVA. The y-axis represents the number of branch points.

women in human umbilical vein endothelial cells has been recently documented,⁴⁹ substantiating the findings observed in the current experiments. In addition, the restoration of network formation occurred at the lowest doses of LMWH, therefore suggesting that at least in vitro, low doses have the potential to induce the beneficial effects in the placenta. Whether this translates to improvements in vivo at low doses is less certain, because various factors including maternal weight, body mass index, and renal function can alter the pharmacokinetics of heparins. The addition of low-dose NAC resulted in the same restorative effect in *DCN*-reduced cells as both UFH and LMWH. This is clinically significant because NAC could potentially be used in place of LMWH and still result in the same rescue effect. This could eliminate the anticoagulant risks posed by

the heparins, leading to potentially increased therapeutic safety with the benefit of improving pregnancy outcomes.

Cell growth, or proliferation, is an integral part of any developed or developing organ system, including the placenta. *DCN* reduction resulted in a significant decrease in cell growth compared with controls.²⁶ Treatment with all 4 doses of UFH or LMWH resulted in a significant further decrease in cell growth and thus did not rescue the effect of *DCN* reduction. Although there is limited information regarding the antiproliferative role of UFH or LMWH specifically in endothelial cells, it is well known that glycosaminoglycans play a role in endothelial cell function, where heparin and heparan sulfate can modulate the angiogenic and/or proliferative activities of growth factors such as fibroblast growth factor 2⁵⁰ by facilitating receptor

Figure 5. Treatment with all 4 concentrations of UFH or LMWH, but not NAC, reduced cell growth of DCN-reduced TIME cells. The effect on TIME-cell proliferation after 48 hours of incubation with DCN siRNA or Mock/NC incubated in 4 concentrations of different heparins was determined using the xCELLigence system. Representative images from the xCELLigence output are also shown for each heparin used, where the y-axis represents the cell index and the x-axis shows the time in minutes. (A) A representative graph showing the cell index of TIME cells treated with DCN siRNA or Mock/NC incubated with 0.15, 0.35, 0.5, and 1.0 IU of UFH over 48 hours in culture. **** $P < .00001$ (significant); $n = 9$; 1-way ANOVA. The y-axis represents the cell index at 48 hours. (B) The results after incubation with 0.15, 0.5, 0.7, and 1.2 IU of LMWH. *** $P < .0001$, **** $P < .00001$ (both significant); $n = 9$; 1-way ANOVA. The y-axis represents the cell index at 48 hours. (C) Treatment with 0.15, 0.4, 0.6, and 1.1 IU of NAC and cell growth in Mock/NC and DCN-reduced cells. * $P < .05$, ** $P < .001$, *** $P < .0001$ (all significant); $n = 9$; 1-way ANOVA. The y-axis represents the cell index at 48 hours.

interaction and activation.^{51,52} However, although heparins increased the angiogenic potential of TIME cells, the differential effects of fibroblast growth factor 2 signaling and function, including its separate role in proliferation, depend on specific structural

variations (ie, variations in the molecular weight of heparin and heparan sulfate).^{53,54} Khorana et al⁵⁵ reported an antiproliferative effect in cultured human umbilical vein endothelial cells of 94% and 58%, observed in 6kDa and 3kDa LMWH, respectively. In contrast,

no data regarding the antiproliferative effect of UFH or LMWH in microvascular endothelial cells have been reported. The LMWH (Dalteparin) used in this study is a 3-6kDa heparin and is the same LMWH that is currently used during pregnancy. Therefore, a significant antiproliferative effect was observed even at the lowest dose of this LMWH, indicating that there may actually be some further detrimental potential of using this LMWH to prevent adverse pregnancy outcomes in terms of a reduction in endothelial cell growth. This is especially relevant if a decrease in cell growth is already a problem in cells affected by downregulation of *DCN*, as observed in first-trimester samples from women who went on to develop pregnancies complicated by FGR.²⁹ In addition, women who do not have complicated pregnancies and are treated with the same LMWH may have detrimental effects in terms of endothelial cell growth. Alternatively, although a decrease in cell growth was also observed in control cells treated with doses of NAC, this decrease was not as significant as those observed for UFH and LMWH, and the *DCN*-reduced cells did not have an additional decrease in their cell-growth potential that was not already present in the non-heparin-treated cells. Therefore, if using our *DCN*-reduced cells as a model for PE/FGR, this treatment with NAC produced an overall reduction in cell growth but did not potentiate further reduction in the *DCN*-reduced cells. In other words, treatment with NAC may not exacerbate cell growth in women with PE and/or FGR compared with UFH or LMWH. Whether these in vitro effects translate into in vivo effects remains to be determined.

This study was based on our previous findings of the importance of *DCN* in the development of the human placenta and the potential contribution its downregulation has on the development of PE and/or FGR,^{26,29} together with the knowledge that heparins are currently used to prevent adverse pregnancy outcomes. We aimed to investigate the mechanism of heparin with respect to *DCN* downregulation in microvascular endothelial cells as our model for PE/FGR. In addition, the comparison between treatment with a NAC heparin and traditional anticoagulant heparins revealed that the NAC resulted in similar effects at a cellular level to LMWH or

UFH, with additional benefits including improved cellular proliferation and reduced anticoagulant activity. This then raises the possibility that NAC could potentially be a better candidate for the prevention of PE/FGR without the risk of significant adverse effects associated with the currently used LMWH. As such, future research into the extended use of NAC and its interaction and implications in the control of cellular functions in other cell types and tissues may improve our understanding of this drug, as well as pave the way for potentially extending this into clinical trials.

Acknowledgments

This work was supported by National Health and Medical Research Council Project Grants, Australia, awarded to J.M.S. (Chief Investigator A) (Application 10042239).

Authorship

Contribution: A.K.L.C. contributed to the study concept, performed all experiments, data analysis, and interpretation, and wrote the manuscript; T.N.G. contributed to the study concept, performed some experiments, and performed critical review of manuscript draft; V.I. contributed to the study concept, performed some of the thrombin generation assays, and performed critical review of manuscript drafts and final approval of manuscript for submission; P.T.M., P.M., S.P.B., and J.M.W. contributed to the study concept and performed critical review of manuscript drafts and final approval of manuscript for submission; and J.M.S. participated in the study concept and design, obtained funding to undertake this work, and performed interpretation of data, critical review of manuscript drafts, and final approval of manuscript for submission.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Amy K. L. Chui, Department of Obstetrics and Gynaecology, The University of Melbourne, Sunshine Hospital, PO Box 294, 176 Furlong Rd, St Albans, VIC 3021, Australia; e-mail: akl.chui83@gmail.com.

References

1. Sood R, Kalloway S, Mast AE, Hillard CJ, Weiler H. Fetomaternal cross talk in the placental vascular bed: control of coagulation by trophoblast cells. *Blood*. 2006;107(8):3173-3180.
2. Steegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. *Lancet*. 2010;376(9741):631-644.
3. Greer IA. Thrombosis in pregnancy: maternal and fetal issues. *Lancet*. 1999;353(9160):1258-1265.
4. Arias F, Romero R, Joist H, Kraus FT. Thrombophilia: a mechanism of disease in women with adverse pregnancy outcome and thrombotic lesions in the placenta. *J Matern Fetal Med*. 1998;7(6):277-286.
5. van der Molen EF, Verbruggen B, Nováková I, Eskes TK, Monnens LA, Blom HJ. Hyperhomocysteinemia and other thrombotic risk factors in women with placental vasculopathy. *BJOG*. 2000;107(6):785-791.
6. Weiner Z, Younis JS, Blumenfeld Z, Shalev E. Assessment of uterine placental circulation in thrombophilic women. *Semin Thromb Hemost*. 2003;29(2):213-218.
7. van Rijn BB, Hoeks LB, Bots ML, Franx A, Bruinse HW. Outcomes of subsequent pregnancy after first pregnancy with early-onset preeclampsia. *Am J Obstet Gynecol*. 2006;195(3):723-728.
8. Askie LM, Duley L, Henderson-Smart DJ, Stewart LA; PARIS Collaborative Group. Antiplatelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet*. 2007;369(9575):1791-1798.
9. Bujold E, Roberge S, Lacasse Y, et al. Prevention of preeclampsia and intrauterine growth restriction with aspirin started in early pregnancy: a meta-analysis. *Obstet Gynecol*. 2010;116(2 Pt 1):402-414.

10. Martinelli I, Ruggerenti P, Cetin I, et al; HAPPY Study Group. Heparin in pregnant women with previous placenta-mediated pregnancy complications: a prospective, randomized, multicenter, controlled clinical trial. *Blood*. 2012;119(14):3269-3275.
11. Rodger MA, Langlois NJ, de Vries JI, et al. Low-molecular-weight heparin for prevention of placenta-mediated pregnancy complications: protocol for a systematic review and individual patient data meta-analysis (AFFIRM). *Syst Rev*. 2014;3:69.
12. de Vries JI, van Pampus MG, Hague WM, Bezemer PD, Joosten JH; FRUIT Investigators. Low-molecular-weight heparin added to aspirin in the prevention of recurrent early-onset pre-eclampsia in women with inheritable thrombophilia: the FRUIT-RCT. *J Thromb Haemost*. 2012;10(1):64-72.
13. Greer IA, Nelson-Piercy C. Low-molecular-weight heparins for thromboprophylaxis and treatment of venous thromboembolism in pregnancy: a systematic review of safety and efficacy. *Blood*. 2005;106(2):401-407.
14. Huhle G, Geberth M, Hoffmann U, Heene DL, Harenberg J. Management of heparin-associated thrombocytopenia in pregnancy with subcutaneous r-hirudin. *Gynecol Obstet Invest*. 2000;49(1):67-69.
15. Quaranta M, Erez O, Mastrolia SA, et al. The physiologic and therapeutic role of heparin in implantation and placentation. *PeerJ*. 2015;3:e691.
16. Hull RD, Raskob GE, Brant RF, et al; American-Canadian Thrombosis Study Group. Low-molecular-weight heparin vs heparin in the treatment of patients with pulmonary embolism. *Arch Intern Med*. 2000;160(2):229-236.
17. Koopman MM, Prandoni P, Piovella F, et al; The Tasman Study Group. Treatment of venous thrombosis with intravenous unfractionated heparin administered in the hospital as compared with subcutaneous low-molecular-weight heparin administered at home. *N Engl J Med*. 1996;334(11):682-687.
18. Levine M, Gent M, Hirsh J, et al. A comparison of low-molecular-weight heparin administered primarily at home with unfractionated heparin administered in the hospital for proximal deep-vein thrombosis. *N Engl J Med*. 1996;334(11):677-681.
19. Simonneau G, Sors H, Charbonnier B, et al. A comparison of low-molecular-weight heparin with unfractionated heparin for acute pulmonary embolism. The THESEE Study Group. Tinzaparine ou Heparine Standard: Evaluations dans l'Embolie Pulmonaire. *N Engl J Med*. 1997;337(10):663-669.
20. Melissari E, Parker CJ, Wilson NV, et al. Use of low molecular weight heparin in pregnancy. *Thromb Haemost*. 1992;68(6):652-656.
21. Sanson BJ, Lensing AW, Prins MH, et al. Safety of low-molecular-weight heparin in pregnancy: a systematic review. *Thromb Haemost*. 1999;81(5):668-672.
22. Carlin AJ, Farquharson RG, Quenby SM, Topping J, Fraser WD. Prospective observational study of bone mineral density during pregnancy: low molecular weight heparin versus control. *Hum Reprod*. 2004;19(5):1211-1214.
23. Casele H, Haney EI, James A, Rosene-Montella K, Carson M. Bone density changes in women who receive thromboprophylaxis in pregnancy. *Am J Obstet Gynecol*. 2006;195(4):1109-1113.
24. Casele HL, Laifer SA. Prospective evaluation of bone density in pregnant women receiving the low molecular weight heparin enoxaparin sodium. *J Matern Fetal Med*. 2000;9(2):122-125.
25. Pettilä V, Leinonen P, Markkola A, Hiilesmaa V, Kaaja R. Postpartum bone mineral density in women treated for thromboprophylaxis with unfractionated heparin or LMW heparin. *Thromb Haemost*. 2002;87(2):182-186.
26. Chui A, Murthi P, Gunatillake T, et al. Altered decorin leads to disrupted endothelial cell function: a possible mechanism in the pathogenesis of fetal growth restriction? *Placenta*. 2014;35(8):596-605.
27. Swan BC, Murthi P, Rajaraman G, et al. Decorin expression is decreased in human idiopathic fetal growth restriction. *Reprod Fertil Dev*. 2010;22(6):949-955.
28. Chui A, Murthi P, Brennecke SP, Ignjatovic V, Monagle PT, Said JM. The expression of placental proteoglycans in pre-eclampsia. *Gynecol Obstet Invest*. 2012;73(4):277-284.
29. Murthi P, van Zanten DE, Eijssink JJ, et al. Decorin expression is decreased in first trimester placental tissue from pregnancies with small for gestation age infants at birth. *Placenta*. 2016;45:58-62.
30. Garcia DA, Baglin TP, Weitz JI, Samama MM; American College of Chest Physicians. Parenteral anticoagulants: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(2 suppl):e24S-43S.
31. Venetsanakos E, Mirza A, Fanton C, Romanov SR, Tlsty T, McMahon M. Induction of tubulogenesis in telomerase-immortalized human microvascular endothelial cells by glioblastoma cells. *Exp Cell Res*. 2002;273(1):21-33.
32. Ogamo A, Uchiyama H, Nagasawa K. Separation of heparin into fractions with different anticoagulant activity by hydrophobic interaction chromatography. *Biochim Biophys Acta*. 1980;626(2):477-485.
33. Williams KJ. Arterial wall chondroitin sulfate proteoglycans: diverse molecules with distinct roles in lipoprotein retention and atherogenesis. *Curr Opin Lipidol*. 2001;12(5):477-487.
34. Duchemin J, Pan-Petes B, Arnaud B, Blouch MT, Abgrall JF. Influence of coagulation factors and tissue factor concentration on the thrombin generation test in plasma. *Thromb Haemost*. 2008;99(4):767-773.
35. Rai R, Cohen H, Dave M, Regan L. Randomised controlled trial of aspirin and aspirin plus heparin in pregnant women with recurrent miscarriage associated with phospholipid antibodies (or antiphospholipid antibodies). *BMJ*. 1997;314(7076):253-257.
36. Gris JC, Mercier E, Quéré I, et al. Low-molecular-weight heparin versus low-dose aspirin in women with one fetal loss and a constitutional thrombophilic disorder. *Blood*. 2004;103(10):3695-3699.
37. Casu B, Vlodavsky I, Sanderson RD. Non-anticoagulant heparins and inhibition of cancer. *Pathophysiol Haemost Thromb*. 2008;36(3-4):195-203.

38. Clark P, Walker ID, Langhorne P, et al; Scottish Pregnancy Intervention Study (SPIN) collaborators. SPIN (Scottish Pregnancy Intervention) study: a multicenter, randomized controlled trial of low-molecular-weight heparin and low-dose aspirin in women with recurrent miscarriage. *Blood*. 2010; 115(21):4162-4167.
39. Kaandorp SP, Goddijn M, van der Post JA, et al. Aspirin plus heparin or aspirin alone in women with recurrent miscarriage. *N Engl J Med*. 2010;362(17):1586-1596.
40. Laskin CA, Spitzer KA, Clark CA, et al. Low molecular weight heparin and aspirin for recurrent pregnancy loss: results from the randomized, controlled HepASA Trial. *J Rheumatol*. 2009;36(2):279-287.
41. Dodd JM, McLeod A, Windrim RC, Kingdom J. Antithrombotic therapy for improving maternal or infant health outcomes in women considered at risk of placental dysfunction. *Cochrane Database Syst Rev*. 2010;(6):CD006780.
42. Cassinelli G, Naggi A. Old and new applications of non-anticoagulant heparin. *Int J Cardiol*. 2016;212(suppl 1):S14-S21.
43. Lever R, Page CP. Non-anticoagulant effects of heparin: an overview. *Handb Exp Pharmacol*. 2012;(207):281-305.
44. Yu L, Garg HG, Li B, Linhardt RJ, Hales CA. Antitumor effect of butanoylated heparin with low anticoagulant activity on lung cancer growth in mice and rats. *Curr Cancer Drug Targets*. 2010;10(2):229-241.
45. Riffo-Vasquez Y, Somani A, Man F, Amison R, Pitchford S, Page CP. A non-anticoagulant fraction of heparin inhibits leukocyte diapedesis into the lung by an effect on platelets. *Am J Respir Cell Mol Biol*. 2016;55(4):554-563.
46. Duckworth CA, Guimond SE, Sindrewicz P, et al. Chemically modified, non-anticoagulant heparin derivatives are potent galectin-3 binding inhibitors and inhibit circulating galectin-3-promoted metastasis. *Oncotarget*. 2015;6(27):23671-23687.
47. Alyahya R, Sudha T, Racz M, Stain SC, Mousa SA. Anti-metastasis efficacy and safety of non-anticoagulant heparin derivative versus low molecular weight heparin in surgical pancreatic cancer models. *Int J Oncol*. 2015;46(3):1225-1231.
48. Kingdom JC, Walker M, Proctor LK, et al. Unfractionated heparin for second trimester placental insufficiency: a pilot randomized trial. *J Thromb Haemost*. 2011;9(8):1483-1492.
49. McLaughlin K, Baczyk D, Potts A, Hladunewich M, Parker JD, Kingdom JC. Low molecular weight heparin improves endothelial function in pregnant women at high risk of preeclampsia. *Hypertension*. 2017;69(1):180-188.
50. Bikfalvi A, Klein S, Pintucci G, Rifkin DB. Biological roles of fibroblast growth factor-2. *Endocr Rev*. 1997;18(1):26-45.
51. DiGabriele AD, Lax I, Chen DI, et al. Structure of a heparin-linked biologically active dimer of fibroblast growth factor. *Nature*. 1998;393(6687):812-817.
52. Spivak-Kroizman T, Lemmon MA, Dikic I, et al. Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. *Cell*. 1994;79(6):1015-1024.
53. Guimond SE, Turnbull JE. Fibroblast growth factor receptor signalling is dictated by specific heparan sulphate saccharides. *Curr Biol*. 1999;9(22):1343-1346.
54. Lundin L, Larsson H, Kreuger J, et al. Selectively desulfated heparin inhibits fibroblast growth factor-induced mitogenicity and angiogenesis. *J Biol Chem*. 2000;275(32):24653-24660.
55. Khorana AA, Sahni A, Altland OD, Francis CW. Heparin inhibition of endothelial cell proliferation and organization is dependent on molecular weight. *Arterioscler Thromb Vasc Biol*. 2003;23(11):2110-2115.