Published in final edited form as:

Tetrahedron Lett. 2017 August 30; 58(35): 3406–3409. doi:10.1016/j.tetlet.2017.07.045.

Iodine-mediated rearrangements of diallylsilanes

Gregory W. O'Neil* and Elizabeth J. Cummins

Department of Chemistry, Western Washington University, Bellingham, WA, 98225, USA

Abstract

Diallylsilanes can be made to rearrange upon treatment with I_2 . Of the silanes tested, diallyldiphenylsilane showed the greatest propensity to undergo this intramolecular carbocation allylation process. After etherification of the initially formed iodosilane, the products from this transformation represent useful synthetic intermediates, suitable for alkylation and cross-metathesis/annulation reactions.

Graphical abstract

Keywords

Silacyclobutane; Diallylsilane; Rearrangement; Cross-metathesis; Amine annulation

During the course of ongoing investigations into activatable silanes for nucleophilic additions to carbonyls,¹ we became interested in exploring reactions of allylsilacyclobutane **1** (Scheme 1).² This compound has been reported to perform non-catalyzed and diastereoselective additions to α -hydroxyketones,² proceeding first through alkoxy ligand exchange followed by carbonyl activation by the Lewis acidic silacyclobutane³ and intramolecular allylation. In that original report,² compound **1** was prepared in 30% yield by adding an allyl Grignard reagent (1 equiv.) to 1,1-dichlorosilacyclobutane followed by etherification with cyclohexanol (CyOH) in the presence of triethylamine (Et_3N). When attempting this chemistry, we routinely obtained very low yields of **1** rather than the expected statistical mixture. Switching the order of addition (i.e. CyOH/ Et_3N followed by allylmagnesium bromide) gave similar results. Others have commented on the challenges of mono-substitution reactions of dichlorosilanes with Grignard reagents as a rationale for the development of more selective methods.^{4,5}

*Corresponding author. Tel.: +01-360-650-6283; fax: +01-360-650-2826; oneilg@wwwu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary Material

Analytical data and experimental procedures for compounds **8**, **10–13**.

As part of these efforts to synthesize **1**, significant quantities of diallylsilacyclobutane **2** were recovered (*ref.* Scheme 1). In an attempt to convert compound **2** to **1**, we turned to the iodine-promoted etherification procedure of Hosomi and Sakurai.⁶ In the event, treatment of **2** with catalytic iodine (10 mol%) in the presence of CyOH (1 equiv.) at 50°C gave a mixture of **1**, **2**, and the diether **3** from which **1** could be consistently isolated in ca. 30% yield (Scheme 2). Currently this is our method of choice (i.e. purposely synthesizing **2** followed by iodoetherification) for obtaining useful quantities of **1**.

Careful analysis of the ¹H NMR spectrum from the reaction of **2** with catalytic iodine and cyclohexanol revealed small amounts of a side product containing what appeared to be a diastereotopic methylene group (Figure 1). Drawn suggestively, one can envision how a compound of this type (**4**) might arise from an intramolecular allylation of the initially formed intermediate cation **5**.

A few other reports of related electrophile-promoted rearrangements of diallylsilanes can be found.^{7,8} For instance, Akiyama et al. reported a domino allylation of diallylsilanes to α,β-unsaturated ketones using BF₃·Et₂O (Scheme 3).⁷ For their reaction, the use of diisopropyldiallylsilane proved optimal, affording fluorosilanes of type **6** in yields between 32–63%. The group of Shainyan has reported a Brønsted-acid promoted rearrangement of diallylsilanes.⁸ For instance, treatment of diallyldiphenylsilane with BF₃·2AcOH gave fluorosilane **7** in 63% yield.

Despite a rich history of allylsilane research describing both inter- and intramolecular reactions with electrophiles,⁹ the electrophile-promoted unique chemistry available to diallylsilanes remains relatively underdeveloped. Combined with the potential synthetic utility of products of type **4**, we were inspired to further investigate this apparent iodine-mediated rearrangement. It was surmised that increasing the amount of iodine should favor the process. Indeed, treatment of **2** with 1.0 equivalent of I₂ followed by treating with CyOH/Et₃N¹⁰ allowed for the isolation of ether **8** in 54% yield (Scheme 4).

Substituents on silicon can have a major influence on reaction outcome.^{1b,11} We therefore set out to examine the impact of substitution at silicon on the selectivity of this iodination/rearrangement. As shown in Figure 2, this was conveniently accomplished by performing the reactions in dry CDCl₃ allowing for direct monitoring of the course of the reaction by NMR. From these studies, we uncovered that diallyldiphenylsilane gave the highest ratio of rearranged compound **9** to non-rearranged products (7.86:1, with the formation of non-rearranged products indicated by the presence of allyl iodide). The reaction with diallyldimethylsilane gave the lowest extent of rearrangement, presumably as a result of decreased steric bulk of the silyl group. Comparing results for diallyldiphenyl- to diallyldiisopropyl silanes, however, suggests that there is also an electronic component to the selectivity for these compounds to rearrange (7.86:1 vs. 6.63:1 respectively) which may involve the ability of neighboring Si-C bonds to stabilize the initially formed cation upon alkene iodination through hyperconjugation effects (*ref.* Scheme 3).¹²

Reactions at 60°C and room temperature gave identical ratios of rearrangement to non-rearrangement, however cooling the reaction to 0 °C resulted in an erosion in selectivity for

the rearranged product. The reaction proved favorable in chloroform (or dichloromethane), with toluene, tetrahydrofuran, DMF and DMSO all giving at best trace amounts of the corresponding rearranged product. Using our optimal conditions (diallyldiphenylsilane, 1 equiv. I₂, CDCl₃, room temp.), ether **10** was isolated in 72% yield (Scheme 5).

We argue that the versatility of the products provided by our iodine-mediated transformation makes this a useful compliment those previously reported related methods (*ref.* Scheme 3).^{7,8} For instance, iodide **10** could be alkylated with dimethylmalonate providing silyl diester **11** in 76% yield (Scheme 6).¹³ Cross-metathesis of **10** with methyl acrylate using the 2nd generation Grubbs' catalyst (**Ru-II**) gave **12** in very high yield.¹⁴ Compound **12** was then taken on to the silylmethyl-functionalized pyrrolidine **13** upon treatment with benzylamine which was isolated in 64% yield (85% based on recovered starting material).¹⁵ Overall, this 3-step conversion of diallyldiphenylsilane to pyrrolidine **13** represents an efficient increase in molecular complexity, producing a privileged structure¹⁶ containing yet additional reactive sites.

To summarize, during the course of investigating an iodine-catalyzed monoetherification of diallylsilanes we observed small amounts of what was assigned as a rearrangement product. This was thought to arise by intramolecular allylation of an intermediate β -silyl carbocation formed upon initial alkene iodination. Ultimately, it was found that good yields of a rearranged product could be obtained by treating diallyldiphenylsilane with stoichiometric iodine in dichloromethane (or chloroform) at room temperature. The initially formed iodosilane was then treated with cyclohexanol¹⁰ allowing for isolation of the corresponding silyl ether **10** in 72% yield. The products from this iodine-mediated process represent useful synthetic intermediates. For instance, we demonstrated that the alkyl iodide portion allows for alkylation with dimethyl malonate (and presumably other carbon nucleophiles). Additionally we were able to transform **10** into a complex silylmethyl-functionalized pyrrolidine **13** via cross-metathesis of the alkene followed by amine annulation. Future work will be aimed at further exploring the synthetic utility of this iodination/rearrangement as well as the development of stereoselective variants.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support from the National Institutes of Health (R15 GM119034) is gratefully acknowledged.

References and notes

1. Medina CR, Carter KP, Miller MM, Clark TB, O'Neil GW. *J Org Chem.* 2013; 78:9093–9101. [PubMed: 23968232] (b) O'Neil GW, Miller MM, Carter KP. *Org Lett.* 2010; 12:5350–5353. [PubMed: 20979374]
2. Matsumoto K, Oshima K, Utimoto K. *J Org Chem.* 1994; 59:7152–7155.
3. Myers AG, Kephart SE, Chen H. *J Am Chem Soc.* 1992; 114:7922–7923.(b) Denmark SE, Griedel BD, Coe DM, Schnute ME. *J Am Chem Soc.* 1994; 116:7026–7043.
4. Li Z, Yang C, Zheng H, Qiu H, Lai G. *J Organomet Chem.* 2008; 693:3771–3779.

5. Li ZF, Cao GQ, Lai JH, Liu Y, Ni JR, Wu HY, Qiu J. *J Organomet Chem.* 2006; 691:4740–4746.
6. Hosomi A, Sakurai H. *Chem Lett.* 1981:85–88.
7. Akiyama T, Asayama K, Fujiyoshi S. *J Chem Soc, Perkin Trans 1.* 1998:3655–3656.
8. Suslova EN, Albanov AI, Shainyan BA. *Russ J Gen Chem.* 2008; 78:1016–1017.(b) Suslova EN, Albanov AI, Shainyan BA. *J Organomet Chem.* 2009; 694:420–426.
9. Sakurai H. *Pure Appl Chem.* 1982; 54:1.(b) Schinzer D. *Synthesis.* 1988:263.(c) Hosomi H. *Acc Chem Res.* 1988; 21:200.(d) Majetich G, Song JS, Leigh AJ, Condon SM. *J Org Chem.* 1993; 58:1030–1037.(e) Majetich, G. *Allylsilanes in Organic Synthesis in Organic Synthesis, Theory, and Applications.* Hudlicky, T., editor. Jai Press Inc.; Greenwich, CT: 1989.
10. Other alcohols can also been used, however we generally opted for cyclohexanol to balance reactivity and stability of the resulting products.
11. Denmark SE, Winter SBD. *Synlett.* 1997:1087–1089.(b) Morkin TL, Leigh WJ. *Acc Chem Res.* 2001; 34:129–136. [PubMed: 11263871] (c) Denmark SE, Neuville L, Christy ME, Tymonko SA. *J Org Chem.* 2006; 71:8500–8509. [PubMed: 17064026]
12. Lambert JB, Wang G-t, Finzel RB, Teramura DH. *J Am Chem Soc.* 1987; 109:7838–7845.
13. Gualandi A, Canestrari P, Emer E, Cozzi PG. *Adv Synth Catal.* 2014; 356:528–536.
14. O'leary, DJ., O'Neil, GW. *Cross-Metathesis.* In: O'Leary, DJ., Grubbs, RH., editors. *Handbook of Metathesis.* 2nd. Wiley-VCH Verlag GmbH & Co. KGaA; Weinheim, Germany: 2015.
15. Bunce RA, Peebles CJ, Jones PB. *J Org Chem.* 1992; 57:1727–1733.
16. Barreiro, EJ. *Privileged Scaffolds in Medicinal Chemistry: Design, Synthesis, Evaluation.* The Royal Society of Chemistry; London, UK: 2015. *Privileged Scaffolds in Medicinal Chemistry: An Introduction.* RSC Drug Discovery Series(b) Zhang, S., Wang, W. *Proline Derivatives.* In: Zhou, QL., editor. *Privileged Chiral Ligands and Catalysts.* Wiley-VCH Verlag GmbH & Co. KGaA; Weinheim, Germany: 2011.

Highlights for: Iodine-mediated rearrangements of diallylsilanes

- A novel iodine-promoted rearrangement of diallylsilanes has been discovered.
- The reaction is proposed to proceed via intramolecular allylation of a carbocation intermediate.
- Selectivity for rearrangement to occur is sensitive to substituents on silicon.
- Products were suitable for alkylation and cross-metathesis/annulation reactions.

Figure 1.

^1H NMR analysis of iodine-catalyzed etherification of **2** showing small amounts of putative compound **4**.

Figure 2. NMR analysis of iodination/rearrangement for different substituted diallylsilanes. Numbers given are integration values for the signals indicated. Increased amounts of allyl iodide formation correspond to lower amounts of rearranged products **9**.

Scheme 1.
Reactivity and initial synthesis of **1**.

Scheme 2.
Synthesis of **1** via diallylsilane **2**.

J. Chem. Soc., Perkin Trans 1, **1998**, 3655

Russ. J. Gen. Chem. **2008**, 1016

Scheme 3.

Previously reported electrophilic rearrangements of diallylsilanes.

Scheme 4.
Iodine-mediated rearrangement of diallylsilane **2**.

Scheme 5.
Synthesis of **10** from diallyldiphenylsilane.

Scheme 6. Further synthetic manipulations of product **10**.