Divergent Responses of Forest Soil Microbial Communities under Elevated CO₂ in Different Depths of Upper Soil Layers

Hao Yu,^{a,b} Zhili He,^c Aijie Wang,^{a,d} Jianping Xie,^e Liyou Wu,^c Joy D. Van Nostrand,^c Decai Jin,^a Zhimin Shao,^b Christopher W. Schadt,^f Jizhong Zhou,^{c,g} Ye Deng^{a,c,h}

^aCAS Key Laboratory of Environmental Biotechnology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences (CAS), Beijing, China

^bCollege of Environmental Science and Engineering, Liaoning Technical University, Fuxin, China

^cDepartment of Microbiology and Plant Biology, Institute for Environmental Genomics, the University of Oklahoma, Norman, Oklahoma, USA

^dState Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology, Harbin, China

^eSchool of Minerals Processing and Bioengineering, Central South University, Changsha, Changsha, China

^fBiosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

^gState Key Joint Laboratory of Environment Simulation and Pollution Control, School of Environment, Tsinghua University, Beijing, China

^hCollege of Resources and Environment, University of Chinese Academy of Sciences, Beijing, China

ABSTRACT Numerous studies have shown that the continuous increase of atmosphere CO₂ concentrations may have profound effects on the forest ecosystem and its functions. However, little is known about the response of belowground soil microbial communities under elevated atmospheric CO₂ (eCO₂) at different soil depth profiles in forest ecosystems. Here, we examined soil microbial communities at two soil depths (0 to 5 cm and 5 to 15 cm) after a 10-year eCO₂ exposure using a high-throughput functional gene microarray (GeoChip). The results showed that eCO₂ significantly shifted the compositions, including phylogenetic and functional gene structures, of soil microbial communities at both soil depths. Key functional genes, including those involved in carbon degradation and fixation, methane metabolism, denitrification, ammonification, and nitrogen fixation, were stimulated under eCO₂ at both soil depths, although the stimulation effect of eCO₂ on these functional markers was greater at the soil depth of 0 to 5 cm than of 5 to 15 cm. Moreover, a canonical correspondence analysis suggested that NO₃-N, total nitrogen (TN), total carbon (TC), and leaf litter were significantly correlated with the composition of the whole microbial community. This study revealed a positive feedback of eCO₂ in forest soil microbial communities, which may provide new insight for a further understanding of forest ecosystem responses to global CO₂ increases.

IMPORTANCE The concentration of atmospheric carbon dioxide (CO₂) has continuously been increasing since the industrial revolution. Understanding the response of soil microbial communities to elevated atmospheric CO₂ (eCO₂) is important for predicting the contribution of the forest ecosystem to global atmospheric change. This study analyzed the effect of eCO₂ on microbial communities at two soil depths (0 to 5 cm and 5 to 15 cm) in a forest ecosystem. Our findings suggest that the compositional and functional structures of microbial communities shifted under eCO₂ at both soil depths. More functional genes involved in carbon, nitrogen, and phosphorus cycling were stimulated under eCO₂ at the soil depth of 0 to 5 cm than at the depth of 5 to 15 cm.

KEYWORDS microbial responses, elevated carbon dioxide, soil microbial community, free-air CO₂ enrichment, functional genes, forest ecosystem

Received 3 August 2017 Accepted 15 October 2017

Accepted manuscript posted online 27 October 2017

Citation Yu H, He Z, Wang A, Xie J, Wu L, Van Nostrand JD, Jin D, Shao Z, Schadt CW, Zhou J, Deng Y. 2018. Divergent responses of forest soil microbial communities under elevated CO₂ in different depths of upper soil layers. *Appl Environ Microbiol* 84:e01694-17. <https://doi.org/10.1128/AEM.01694-17>.

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Zhili He, zhili.he@ou.edu, or Ye Deng, yedeng@rcees.ac.cn.

The atmospheric carbon dioxide (CO₂) concentration is closely linked to global carbon (C) cycling and has significant effects on terrestrial ecosystems (1). The concentration of global atmosphere CO₂ has increased substantially since the preindustrial period, from 280 ppm to currently over 400 ppm (2). Elevated atmospheric CO₂ (eCO₂) could strongly influence ecosystem functioning through various processes, including the soil microbial communities that mediate C sequestration and nutrient cycling (3–6). Thus, understanding the response of microbial communities to eCO₂ is crucial for understanding multiple ecosystem functions and for predicting the changes in terrestrial ecosystems in response to future climates (7, 8).

Forests currently tie up more than 85% of all terrestrial biomass carbon (C) and provide the largest terrestrial C sinks for global atmospheric CO₂ (9, 10). Numerous studies have shown that eCO₂ dramatically influenced the productivity, transpiration, nitrogen (N) use efficiency, leaf litter productivity, fine root exudation, and carbon (C) assimilation rates of trees (11–16). Such responses of forest ecosystems may further influence the belowground microbial composition and function by changing soil geochemical properties and nutrient inputs (17, 18). However, to date, accurate observations of the effect of eCO₂ on microbial communities have been largely complicated by the extremely high diversity and the complexity of measuring belowground bioprocesses (3). Some studies have shown significant changes in the composition, structure, activity, and functional capacity of soil microbial communities (3, 6, 18–22) under eCO₂, while other studies have shown no significant differences of microbial community compositions between eCO₂ and ambient CO₂ (aCO₂) conditions (21–26). Therefore, further studies at additional sites and using new analysis approaches are necessary to understand whether the structures of soil microbial communities shift under eCO₂ and how soil microbial communities respond to complex interactions of biotic and abiotic factors in forests.

Soil microbial communities are expected to play important roles in mediating many belowground processes (e.g., organic matter decomposition and nutrient cycling) in forest ecosystems (27, 28). Since nearly all soil physiochemical parameters change with soil depth (e.g., C and N availability, oxygen content, soil moisture, and temperature) (17, 29), distinct responses of soil microbial communities to eCO₂ are expected to occur at different soil depths. Previous reports showed that numerous genes for nutrient (e.g., C and N) cycling significantly increased at soil depths of 0 to 5 cm with no significant changes at depths of 5 to 15 cm in a farmland ecosystem in response to eCO₂ (17, 30). However, most studies of forest ecosystems have evaluated the effect of eCO₂ on soil microbial communities at only one depth, which could mask finer-scale differences (e.g., at different depths) (23). Plant litter and root exudation inputs to soil vary with soil depth (29, 31), which may affect soil microbial communities associated with ecological processes at different depths. Thus, it is important to further understand whether eCO₂ shifts the structure and function of microbial communities at depth-resolved scales.

Free-air CO₂ enrichment (FACE) experiments provide an opportunity to understand many aspects of eCO₂ effects on ecosystems under natural field conditions (32). It was demonstrated in several FACE experiments that the photosynthesis, net primary production (NPP), fine-root production, and turnover of trees were enhanced by eCO₂ (16, 33–36). Consequently, an increase in soil C stock and a high C/N ratio were observed under eCO₂ at a soil depth of 0 to 5 cm in a forest ecosystem, where inputs from roots and aboveground litter were greatest (37, 38). These changes may directly or indirectly affect the structure of the soil microbial community and the functional capacity associated with nutrient cycling (e.g., C and N cycling). However, how microbial communities respond to eCO₂ at different depths in forest FACE experiments remains unknown. In this study, we evaluated the soil microbial community response to eCO₂ in a deciduous forest FACE experiment site (<http://face.ornl.gov/index.html>), which had been exposed to eCO₂ for more than 10 years, using a functional gene array (GeoChip). As a powerful high-throughput metagenomics technology, GeoChip 3.0 assesses 292 genes of key enzymes involved in C, N, S, and P cycling, key energy metabolism, metal resistance, organic contaminant degradation, antibiotic resistance, and a phylogenetic

TABLE 1 Soil properties at different depths under eCO₂ and aCO₂^a

Treatment	NH ₄ -N (mg/kg)	NO ₃ -N (mg/kg)	TC (%) ^b	TN (%) ^c	C/N
0- to 5-cm depth					
eCO ₂	34.2 ± 2.30	3.67 ± 0.254	2.75 ± 0.145	0.203 ± 0.007	13.78 ± 0.44
aCO ₂	39.4 ± 6.19	2.62 ± 0.623	2.14 ± 0.43	0.178 ± 0.018	12.47 ± 0.95
<i>P</i> value	0.189	0.023	0.037	0.070	0.071
5- to 15-cm depth					
eCO ₂	31.26 ± 3.42	1.34 ± 0.069	2.39 ± 0.105	0.178 ± 0.011	13.67 ± 0.31
aCO ₂	30.25 ± 3.39	1.00 ± 0.106	2.02 ± 0.068	0.138 ± 0.005	14.64 ± 0.32
<i>P</i> value	0.851	0.035	0.023	0.012	0.075

^aThe variables of soil properties at each depth were analyzed separately, and significances between different treatments (aCO₂ and eCO₂) were calculated by *t* tests. Significant differences (*P* < 0.05) are indicated by bold type.

^bTC, total carbon content.

^cTN, total nitrogen content.

marker (*gyrB*) (39). Recently, many studies demonstrated that GeoChip is an ideal tool for analyzing microbial communities from complex environments, such as soil (17, 19, 30), contaminated water (40, 41), oil fields (19, 42), bioreactor systems (43, 44), and other habitats (45). We hypothesized that the eCO₂ treatment would have significant effects on the functional diversity, composition, structure, and metabolic potential of soil microbial communities at different depths (0 to 5 cm and 5 to 15 cm) and that the plant and soil properties would have different impacts on soil microbial communities at different depths.

RESULTS

Effects of eCO₂ on plant and soil properties at two depths. The ORNL FACE experiment was constructed in 1998 in the Oak Ridge National Environmental Research Park within a sweetgum (*Liquidambar styraciflua* L.) plantation. After 10 years of elevated CO₂ (eCO₂) treatment, the plant net primary productivity (NPP) increased 9% compared with that at ambient CO₂ (aCO₂), although the enhancement of NPP was less significant (46). For soil properties, five important soil variables were measured or calculated, including soil nitrate (NO₃⁻-N), ammonium (NH₄⁺-N), total nitrogen (TN), total carbon (TC), and C:N ratio. Elevated CO₂ greatly increased the amount of soil NO₃⁻-N (*P* < 0.05) and TC (*P* < 0.05) in 6 replicate soil samples at both soil depths (0 to 5 cm and 5 to 15 cm) (Table 1). However, the effects of eCO₂ on soil C:N ratios were different in these two soil layers, with an increase in the 0- to 5-cm-depth layer but a decrease at a depth 5 to 15 cm, which were marginally significant (*P* < 0.1). There was no significant difference in the NH₄⁺-N contents between eCO₂ and aCO₂ at either soil depth.

Overall differences of soil microbial functional genes and phylogenetic markers. A total of 7,090 microbial function genes were detected by GeoChip 3.0 across 24 samples. The gene numbers detected under eCO₂ (3,502 ± 381) were significantly greater (*P* = 0.026) than that under aCO₂ (2,386 ± 85) at the soil depth of 0 to 5 cm. However, this difference was not significant (*P* > 0.1) at the depth of 5 to 15 cm. Similar patterns were also observed in alpha-diversity indexes. The Shannon index (*H'*) and the Simpson's reciprocal index (1/*D*) were significantly higher (*P* < 0.05) in eCO₂ at the depth of 0 to 5 cm, but no significant differences were found at the depth of 5 to 15 cm. The overall taxonomic compositions of the microbial communities were analyzed at phylum level based on GeoChip data (see Fig. S1 in the supplemental material). All detected genes were taxonomically derived from 2 archaeal phyla, 21 bacterial phyla, and 5 eukaryotic phyla. Among these phyla, functional genes derived from *Proteobacteria*, *Actinobacteria*, and *Firmicutes* had relative higher abundances than those from other phyla. eCO₂ highly increased the abundance of key genes derived from *Crenarchaeota*, *Euryarchaeota*, *Bacteroidetes*, *Chloroflexi*, *Cyanobacteria*, *Firmicutes*, and *Proteobacteria* at the soil depth of 0 to 5 cm and from *Actinobacteria* and *Proteobacteria* at the soil depth of 5 to 15 cm (see Fig. S2).

FIG 1 Detrended correspondence analysis (DCA) of all detected functional genes at soil depths of 0 to 5 cm and 5 to 15 cm under ambient CO₂ (aCO₂) and elevated CO₂ (eCO₂) conditions.

A detrended correspondence analysis (DCA) of all functional genes illustrated that eCO₂ and aCO₂ were clearly separated (Fig. 1), and similar separation by using the phylogenetic marker gene (*gyrB*) only (see Fig. S3) was observed at both soil active layers (0 to 5 cm and 5 to 15 cm), indicating that microbial functional gene structures and phylogenetic structures were both changed under eCO₂. In addition, the permutational multivariate analysis of variance (Adonis) test on all detected functional genes and *gyrB* phylogenetic markers showed that not only CO₂ treatment but also depth and their interaction had significant effects on both functional and phylogenetic structures of soil microbial communities ($P < 0.05$) (Table 2). The main factor contributing to the total variation of microbial functional gene structures was eCO₂ (12.9%), followed by depth (7.7%) and their interaction (7.7%), while the phylogenetic structures were mainly shaped by eCO₂ (12.1%), eCO₂ plus depth (9.5%), and depth (8.3%). Other statistical tests, namely, analysis of similarities (ANOSIM) and a multiresponse permutation procedure (MRPP), also detected significant changes according to both CO₂ treatment and depth (see Table S1). Thus, the compositions and phylogenetic and functional gene structures of soil microbial communities had been dramatically altered by eCO₂ at both soil depths.

Effect of eCO₂ on microbial functional genes involved in C, N, and P cycling. A total of 589 ± 74 and 524 ± 41 genes involved in C cycling were detected at the depths of 0 to 5 cm and 5 to 15 cm, respectively, under eCO₂, which were higher than those under aCO₂ (399 ± 13 and 428 ± 21 , respectively) (Table 3). A significant difference was observed at the depth of 0 to 5 cm ($P = 0.044$), while no significant difference was observed at the depth of 5 to 15 cm ($P = 0.087$) between aCO₂ and eCO₂. These results indicated that microbial C cycling might be altered by eCO₂.

Several important C fixation genes were further investigated, including the propionyl-CoA carboxylase (PCC) gene and the gene encoding ribulose-1,5-bisphosphate carboxylase/oxygenase (RubisCO). The total signal intensities for PCC and RubisCO genes were substantially higher ($P < 0.05$) under eCO₂ than under aCO₂ at the soil depth of 0 to

TABLE 2 Adonis analysis of the effect of eCO₂, soil depth, and their interaction on the functional and phylogenetic structures of microbial communities based on all detected genes and *gyrB*

Gene(s)	CO ₂		Depth		CO ₂ -depth interaction	
	R ²	P value	R ²	P value	R ²	P value
All functional genes	0.129	0.001 ^a	0.077	0.007 ^a	0.077	0.012 ^a
Phylogenetic marker (<i>gyrB</i>)	0.121	0.001 ^a	0.083	0.013 ^a	0.095	0.005 ^a

^a $P < 0.05$.

TABLE 3 Distribution of key gene categories involved in carbon, nitrogen, phosphorus, and sulfur cycling

Gene category	No. $\pm$ SEM of genes at 0- to 5-cm depth			No. $\pm$ SEM of genes at 5- to 15-cm depth		
	eCO ₂	aCO ₂	<i>P</i> value ^a	eCO ₂	aCO ₂	<i>P</i> value ^a
C cycling	589 $\pm$ 74	399 $\pm$ 13	0.044	524 $\pm$ 41	428 $\pm$ 21	0.087
Acetogenesis	7 $\pm$ 1	3 $\pm$ 0	0.043	5 $\pm$ 0	7 $\pm$ 1	0.178
C degradation	436 $\pm$ 56	296 $\pm$ 9	0.048	380 $\pm$ 30	317 $\pm$ 15	0.119
C fixation	132 $\pm$ 16	90 $\pm$ 5	0.045	121 $\pm$ 10	94 $\pm$ 5	0.06
Methane	15 $\pm$ 1	11 $\pm$ 1	0.011	18 $\pm$ 1	11 $\pm$ 1	<0.001
N cycling	390 $\pm$ 43	255 $\pm$ 11	0.019	327 $\pm$ 29	294 $\pm$ 22	0.42
Ammonification	39 $\pm$ 5	23 $\pm$ 1	0.021	29 $\pm$ 3	23 $\pm$ 2	0.132
Anammox	4 $\pm$ 0	3 $\pm$ 0	0.183	4 $\pm$ 0	3 $\pm$ 0	0.065
Assimilatory N reduction	27 $\pm$ 2	19 $\pm$ 1	0.021	25 $\pm$ 2	19 $\pm$ 1	0.038
Denitrification	135 $\pm$ 19	87 $\pm$ 6	0.049	113 $\pm$ 10	97 $\pm$ 7	0.25
Dissimilatory N reduction	25 $\pm$ 3	16 $\pm$ 1	0.033	20 $\pm$ 3	18 $\pm$ 2	0.557
Nitrification	2 $\pm$ 0	1 $\pm$ 0	0.038	2 $\pm$ 0	2 $\pm$ 0	0.734
N fixation	158 $\pm$ 13	107 $\pm$ 7	0.008	134 $\pm$ 12	132 $\pm$ 11	0.901
P utilization	68 $\pm$ 9	49 $\pm$ 2	0.091	54 $\pm$ 5	47 $\pm$ 4	0.4
S cycling	177 $\pm$ 23	119 $\pm$ 5	0.047	135 $\pm$ 12	126 $\pm$ 7	0.601
Total	1,224 $\pm$ 148	822 $\pm$ 29	0.036	1,039 $\pm$ 86	895 $\pm$ 54	0.223

^aThe effects of eCO₂ on distribution of key gene categories were calculated by *t* tests. Significant differences (*P* < 0.05) indicated by bold type.

5 cm, while the significant difference was only detected for the PCC gene at the soil depth of 5 to 15 cm (see Fig. S4).

Three functional genes involved in methane production and oxidation showed distinguishable responses to eCO₂ at different depths, including *mcrA* for CH₄ production and *mmoX* and *pmoA* for CH₄ oxidation. Among these genes, a total of 4 $\pm$ 0.7 genes derived from *Archaea* for methane production and 11 $\pm$ 0.5 genes derived from *Bacteria* for methane oxidation were detected in eCO₂ samples at a soil depth of 0 to 5 cm, while 3 $\pm$ 0.4 and 8 $\pm$ 0.7 genes, respectively, were detected in aCO₂ samples. An obviously higher number of genes (*P* = 0.004) derived from *Bacteria* was observed in eCO₂ samples than that in aCO₂ samples. The difference between eCO₂ and aCO₂ was even larger at a soil depth of 5 to 15 cm. Totals of 6 $\pm$ 0.6 (*Archaea*) and 12 $\pm$ 0.7 (*Bacteria*) genes were detected in eCO₂ samples, and 3 $\pm$ 0.3 (*Archaea*) and 8 $\pm$ 0.7 (*Bacteria*) genes were detected in aCO₂ samples. eCO₂ greatly increased the numbers of detected genes derived from *Archaea* (*P* = 0.002) and *Bacteria* (*P* = 0.005) at a soil depth of 5 to 15 cm. At a soil depth of 0 to 5 cm, 12 and 3 unique genes involved in methane metabolism, including those derived from uncultured bacterium, uncultured archaeon, uncultured methanogenic archaeon, uncultured *Methanobacteriales* archaeon, uncultured euryarchaeota, and *Methanobrevibacter smithii*, were detected in eCO₂ and aCO₂, respectively, while 22 and 4 unique genes derived from uncultured bacterium, uncultured methanotrophic bacterium, *Frankia* sp., *Azoarcus* sp., *Methylosinus trichosporium*, uncultured methanogenic archaeon, uncultured *Methanobacteriales* archaeon, uncultured archaeon, *Methanofollis liminatans*, *Methanomethylovorans thermophile*, *Methanosarcina lacustris*, and uncultured *Methanosarcinales* archaeon were detected under eCO₂ and aCO₂, respectively, at a soil depth of 5 to 15 cm (see Fig. S5). The total signal intensities of these genes were increased under eCO₂ at both soil depths. Significantly higher signal intensities were observed for *mcrA* (*P* = 0.001), *mmoX* (*P* = 0.033), and *pmoA* (*P* = 0.049) under eCO₂ at the soil depth of 5 to 15 cm (see Fig. S6). However, only *pmoA* showed a dramatic difference (*P* = 0.005) between eCO₂ and aCO₂ at the soil depth of 0 to 5 cm.

In C degradation processes, eCO₂ treatment also showed positive effects in both soil layers (Fig. 2), and most of the genes had significantly higher signal intensities

FIG 2 Normalized signal intensities of the detected genes involved in carbon degradation at the soil depths of 0 to 5 cm (top) and 5 to 15 cm (bottom). All data are presented as means $\pm$ standard errors (SEs). **, $P < 0.01$; *, $P < 0.05$ based on Student's t tests.

($P < 0.05$) under eCO₂ than under aCO₂. In the 0- to 5-cm-depth layer, those included genes for isocitrate lyase (*aceA*) and malate synthase (*aceB*) for the tricarboxylic acid (TCA) cycle, α -amylase (*amyA*), glucoamylase, pullulanase, and neopullulanase (*npIT*) for starch decomposition, arabinofuranosidase for hemicellulose decomposition, cellobiase and exoglucanase for cellulose decomposition, exochitinase for chitin decomposition, and glyoxalase (*glx*) and phenol oxidase for lignin decomposition (Fig. 2). At the soil depth of 5 to 15 cm, seven of the above genes were not significantly altered, including both TCA cycle genes (*aceA* and *aceB*). However, six other functional genes were dramatically changed, including two other hemicellulose degradation genes (mannanase and xylanase), one aromatics degradation gene (limonene hydrolase), one cellulose gene (endoglucanase), and two chitin degradation genes (endochitinase and acetylglucosaminidase). These results indicated that, although eCO₂ enhanced carbon degradation genes at both soil depths, the genes enriched at these two depths differed.

For N cycling processes at the soil depths of 0 to 5 cm and 5 to 15 cm, 390 ± 43 and 327 ± 29 genes, respectively were detected under eCO₂ and 255 ± 11 and 294 ± 22 genes, respectively, were detected under aCO₂. The difference is significant in the upper 0- to 5-cm-depth layer ($P = 0.019$) but not significant in the 5- to 15-cm-depth layer ($P = 0.42$) (Table 3). Under eCO₂, the total signal intensities of nine genes were substantially higher ($P < 0.05$) than those under aCO₂ in the 0- to 5-cm-depth soil layer, including *gdh* and *ureC* that are involved in ammonification, *nasA* and *nirB* involved in assimilatory N reduction, *narG* and *nirS* involved in denitrification, *nrfA* involved in dissimilatory N reduction, *amoA* involved in nitrification, and *nifH* involved in N₂ fixation (Fig. 3). However, at the soil depth of 5 to 15 cm, only the total signal intensities of three genes involved in ammonification (*gdh*), assimilatory N reduction (*nasA*), and denitrification (*narG*) were significantly ($P < 0.05$) increased under eCO₂. These results

FIG 3 Normalized signal intensities of detected genes involved in the N cycle. (A) ammonification, (B) anammox, (C) assimilatory N reduction, (D) denitrification, (E) dissimilatory N reduction, (F) nitrification, and (G) N₂ fixation. All data are presented as means $\pm$ SEs. **, $P < 0.01$; *, $P < 0.05$; •, $P < 0.1$ based on t tests.

suggested that soil N cycling was more active at the depth of 0 to 5 cm than at 5 to 15 cm under eCO₂.

The functional gene encoding phytase for P utilization was substantially higher ($P < 0.05$) under eCO₂ at the soil depth of 0 to 5 cm, and it was detected only in eCO₂ not in aCO₂ at the soil depth of 5 to 15 cm. For other genes, the differences were not significant between eCO₂ and aCO₂ at either depth, including for *Ppx* for inorganic polyphosphate degradation and for *Ppk* for polyphosphate biosynthesis (see Fig. S7).

Associations between the microbial functional gene structures and environmental variables. To determine how major environmental variables influence the microbial functional structure, a canonical correspondence analysis (CCA) was performed. Seven environmental variables were selected: NO₃-N, NH₄-N, TN, and TC for soil properties, and leaf litter and fine root production for plant properties (47). Totals of 13.8% and 11.4% of the constrained variations in microbial communities were explained by the first and second canonical axes, respectively (Fig. 4a). The communities under eCO₂ and aCO₂ separated clearly along the first canonical axis, while communities at the two soil depths separated along the second canonical axis (Fig. 4a). The results of the CCA showed that the microbial functional structures had significant correlations with selected soil properties ($P = 0.023$), plant properties ($P = 0.004$), and all environmental variables ($P = 0.003$) (Fig. 4b). Among these variables, NO₃-N, TN, TC, and leaf litter were significantly correlated with all detected functional genes, suggesting that they were important environmental variables in shaping the soil microbial community structure at this FACE site. Relationships between functional genes involved in C and N cycling and environmental variables at two soil depths were further analyzed by a Mantel test (see Table S2). The results showed that 9 and 8 genes involved in C and N cycling significantly correlated ($P < 0.05$) with soil or plant properties at the soil depths of 0 to 5 cm and 5 to 15 cm, respectively.

To further assess the contributions of soil and plant properties to the microbial functional structure at each soil depth, a CCA-based variation partitioning analysis (VPA) was conducted (Fig. 4c). At the soil depth of 0 to 5 cm, soil properties accounted for 35.1% of microbial community variation and plant properties accounted for 20.8%, while their

FIG 4 Canonical correspondence analysis (CCA) of GeoChip data under eCO₂ and aCO₂ with selected environmental variables (a). (b) Model significances. (c) Variation partitioning analysis (VPA) based on partial CCA in the soil depths 0 to 5 cm and 5 to 15 cm. Environmental variables, including NO₃-N, NH₄-N, TN, and TC for soil and leaf litter and fine roots for plants, were selected based on variance inflation factor (VIF) analysis.

interaction only accounted for 1.4%, with 42.7% unexplained. In comparison to those at a soil depth of 0 to 5 cm, lower amounts of variation were accounted for by soil properties (30.5%) and plant properties (16.1%) at the soil depth of 5 to 15 cm. However, the interaction of soil and plant properties accounted for a substantially higher percentage of microbial community variations (13.6%) than the 1.4% at the soil depth of 0 to 5 cm.

DISCUSSION

As soil microorganisms mediate several important aspects of nutrient (e.g., C, N, and P) cycling, understanding the responses of soil microbial communities in forest ecosystems to eCO₂ is critical to fully predict the effects of future global change. In this study, we demonstrated the long-term (>10 years) effect of eCO₂ on soil properties, functional diversity, structure and metabolic potential of microbial communities in this sweetgum forest ecosystem.

Since the response of soil microbial communities to eCO₂ remains inconclusive for FACE sites (18, 20, 23), more studies are needed to understand whether the compositions and structures of soil microbial communities are altered at eCO₂, especially across multiple ecosystems. The effects of eCO₂ treatment on soil microbial communities were observed in forest, grassland, and agriculture ecosystems (3, 8, 48), but some research also has shown a lack of significant changes in soil microbial community structures and compositions in FACE sites after 2 or 8 years of eCO₂ exposure by assaying enzymes, substrate utilization, and 16S rRNA gene clone libraries (23, 49). However, these conclusions may not be the same if eCO₂ treatment time was extended or a more comprehensive microbial technique was used for soil microbial communities and they were examined at finer scales (23, 49). It is not surprising that no significant responses

of microbial communities were detected, as the availability of soil nutrient pools were often not altered in the short-term treatment (49) and changes in the community structures would be diluted by combining different depths of soil together (23). In addition, the methods previously used for examining microbial community structure were at the phylum level, which can be imprecise for detecting the specific differences within a phylum (23, 50). As a high-resolution and high-throughput metagenomics technology (39, 50), GeoChip was employed in this study to examine the effect of eCO₂ on soil microbial communities. Our results show that soil microbial communities have significant responses ($P < 0.05$) to eCO₂ at both soil depths, which are associated with the change of soil properties, on the basis of dissimilarity tests (ANONISM and MRPP) (see Table S1 in the supplemental material). It had been substantiated by previous studies of the ORNL FACE site that eCO₂ largely increased the production of plant litter (e.g., leaves and roots) (13, 23, 37, 38), which could alter C and N storage and cycling in the soil (38). In accordance with the previous study, our research shows that total carbon content (TC) and total nitrogen content (TN) were significantly or marginally increased under eCO₂ treatment at two depths (Table 1) (38). Consequentially, these changes may result in shifts of microbial community functional diversity, composition, and structure.

As we expected, the microbial functional diversity, composition, and phylogenetic and functional structures of soil microbial communities shifted under eCO₂ at two soil depths (0 to 5 cm and 5 to 15 cm) (Fig. 1; see also Fig. S3). The results agreed well with a recent report of soil microbial communities at a FACE experimental site in a corn-soy agroecosystem (SoyFACE), which showed that after an 8-year exposure to eCO₂, the functional and phylogenetic structures of soil microbial communities had shifted at soil depths of 0 to 5 cm and 5 to 15 cm (17). Moreover, the relative abundance of functional genes derived from some key phyla was significantly enhanced, potentially indicating that the abundances of bacteria and fungi increased under eCO₂ at the two soil depths. The effects of eCO₂ on soil fungal and bacterial communities possibly occur via increased soil C input in the form of plant litter and root exudates (51, 52). This stimulatory effect of eCO₂ was also observed in other FACE experiments. The microbial and fungal abundances increased under eCO₂ treatment at a wheat field and in a biodiversity, CO₂, and N (BioCON) experiment (53, 54).

Whether the functional processes (e.g., C and N cycling) of soil microbial communities were stimulated by long-term eCO₂ is another critical issue for FACE site research. In this study, we found that a large number of the key functional genes responsible for C, N, and P cycling had significantly higher signal intensities under eCO₂ than under aCO₂, although some patterns for individual genes were different between the two depths. First, different stimulation effects of eCO₂ were observed for the detected key genes involved in C cycling between two depths. Elevated CO₂ significantly enhanced the signal intensities of 57.9% (11 of 19) of detected functional genes for C degradation at the soil depth of 0 to 5 cm and of only 31.6% (6 of 19) at the soil depth of 5 to 15 cm (Fig. 2). Moreover, eCO₂ significantly increased all functional genes for glyoxylate/TCA cycling and lignin decomposition but only one for starch decomposition at the soil depth of 0 to 5 cm, while we did not find significant differences of these genes at the soil depth of 5 to 15 cm. These significantly enriched genes are linked closely with decomposing plant litter and soil organic matter. The physiological activity of soil microorganisms is greatly driven by the input of organic substrates (27), and it has been shown that the effects of eCO₂ on plant litter production could change the substrate input to soil (36). Increases in plant litter productivity and soil C content were observed at the soil depth of 0 to 5 cm at the ORNL FACE site (Table 1) (13, 37, 38), which could stimulate plant-induced microbially mediated C decomposition in soil (17). Such effects of eCO₂ coincided with those found by previous research on soil microbial communities (3, 17, 30). However, this effect was weakened at the soil depth of 5 to 15 cm, and the total signal intensities of almost all of these genes increased only slightly under eCO₂ (Fig. 2). For C fixation, the total signal intensities of RubisCO and PCC genes increased significantly at the soil depth of 0 to 5 cm, and that for only the PCC gene increased significantly at the soil depth of 5 to 15 cm (Fig. S4), suggesting that C fixation processes

for microorganisms may be more abundant at the soil depth of 0 to 5 cm than at the depth of 5 to 15 cm. The total signal intensities of *mcrA* for methane production were enhanced under eCO_2 at two depths, and the difference was significant in the soil depth of 5 to 15 cm. In addition, the numbers of these genes derived from *Archaea* and *Bacteria* increased under eCO_2 at both depths, especially at the lower soil depth of 5 to 15 cm, where the difference was significant ($P < 0.05$). Previous studies showed that the abundance of methanogenic archaea significantly increased in rice-cultivated soil under eCO_2 (55, 56) because of the increase in the soil carbon input (e.g., root exudates) with elevated CO_2 . Our results revealed that soil methane emission was potentially stimulated in response to eCO_2 in a forest ecosystem, which is in agreement with a previous report as well (30).

Second, microbially mediated soil N cycling provided a positive feedback to eCO_2 . The soil C and N cycles have strong linkages in all terrestrial ecosystems, and N supply is well documented as an important constraint to limit the productivity of terrestrial ecosystems under eCO_2 at these FACE sites (36, 46, 57). Due to the high concentrations of N and carbohydrates, decaying plant materials are important for the flux of biological monomers C and N to the soil (58, 59). The production of plant materials was reported to be increased at an ORNL site under eCO_2 , leading to an increasing input of C and N to the soil profile (13, 38). It remains unclear how soil microbial communities modify their functional processes (N cycling) in response to eCO_2 . GeoChip data showed that the total signal intensities of *nifH* involved in N_2 fixation and *ureC* involved in ammonification significantly increased at eCO_2 at the soil depth of 0 to 5 cm and remained unchanged at the soil depth of 5 to 15 cm (Fig. 3). Such responses of these genes may increase N fixation and ammonification processes at the soil depth of 0 to 5 cm under eCO_2 , which could relieve progressive N limitation at FACE sites (4, 57, 60). The total signal intensities of key genes involved in denitrification were generally increased under eCO_2 , and significantly higher signal intensities were observed for *narG* and *nirS*. The results suggest that the denitrification processes could also be enhanced under eCO_2 , which may result in an increase of N_2O emission as previously observed (61, 62). In addition, the key genes involved in assimilatory N reduction (*nasA* and *nirB*), dissimilatory N reduction (*nrfA*), and nitrification (*amoA*) were also significantly increased. It appears that N cycling could be entirely stimulated under eCO_2 , especially at the soil depth of 0 to 5 cm, where microbial communities were more sensitive to environmental change. This could be well explained by the change of plant litter productivity. Plants may influence microbial communities directly by the changes in plant litter inputs and root exudates or indirectly by their capacity to influence soil properties (e.g., organic matter and nutrient availability) (63, 64). At this site, the C and N input to soil may be altered by the changes in leaf or fine root quantity at two depths (0 to 5 cm and 5 to 15 cm) (13, 23), which may have considerable effect on microbial communities. Previous research also reported that the increase of C and N input to the soil from fine roots under eCO_2 was mainly in deep soil (below 30 cm), which were not assessed in our study (13).

As soil microbial communities may shift with the changing of environmental variables (e.g., soil C and N concentrations) (65), it is imperative to establish links between environmental variables and microbial communities for FACE sites. Plant and soil properties were generally considered main factors that largely determine the soil microbial community structure (63, 66). The whole microbial functional composition significantly correlated with selected soil, plant, and environmental variables ($P < 0.05$) (Fig. 4b). Variations in microbial communities were explained by soil and plant properties at two depths (0 to 5 cm and 5 to 15 cm), indicating that at the soil depth of 0 to 5 cm, the microbial community is more sensitive to plant and soil properties than at the soil depth of 5 to 15 cm.

In summary, this study highlights that functional microbial communities may be altered under eCO_2 treatments in forest ecosystems. We found that eCO_2 significantly altered the microbial functional diversity, composition, and structure and increased the abundance of a large number of key functional genes involved in C and N cycling at the soil depths of 0 to 5 cm and 5 to 15 cm, which may ultimately feed back to the

ecosystem level responses to elevated CO₂ at the ORNL FACE site. In addition, the effect of eCO₂ was more pronounced at the soil depth of 0 to 5 cm, with increases of leaf litter productivity and C input to the soil. Since fine roots are critical to soil C input, further studies are needed to understand how microbial community structure and function shift in response to eCO₂ at deeper soil depths (below 15 cm) in forest ecosystems.

MATERIALS AND METHODS

Site description and experimental design. The ORNL FACE experiment was established on a sweetgum (*Liquidambar styraciflua* L.) plantation in the Oak Ridge National Environmental Research Park, TN, USA (35°54'N, 84°20'W). Five 25-m-diameter plots (46) were constructed at the FACE site, with two plots for elevated CO₂ (~560 ppm in 2008) treatment and three plots for ambient CO₂ (~405 ppm in 2008) treatment. The mean annual temperature and precipitation are 13.9°C and 1,322 mm, respectively. A more complete site description has been documented previously (33, 67). The soil at the FACE site is slightly acidic (pH 5.5 to 6.0) and is an alluvial aquic hapludult with a silty clay loam texture (68). Soil samples were collected from four plots, including two of three aCO₂ plots (plot 1 and plot 2) and two eCO₂ plots (plot 4 and plot 5). Triplicate soil samples were collected from each plot at the soil depths of 0 to 5 cm and 5 to 15 cm in July 2008 after more than 10 years of fumigation treatment. After the removal of plant roots and large stones, 24 soil samples were immediately stored at –20 or 4°C for DNA extraction or soil property analysis, respectively. Plant property data were collected from the FACE data management system (<https://facedata.ornl.gov/>) (69).

Soil property analysis. The total C and total N of soil samples were measured by a Leco Truspec dry combustion carbon analyzer (70), and NO₃-N and NH₄-N of were extracted from soil samples by the use of a 1.0 M KCl solution and were quantified by a flow injection autoanalyzer.

DNA extraction, amplification, labeling, and hybridization. Microbial community DNA was extracted from 5-g soil samples with the freeze-grinding method as described previously (71). DNA quality was assessed by an ND-1000 spectrophotometer (NanoDrop Technologies Inc., Wilmington, NC) using the ratios of 260/280 nm and 260/230 nm, and DNA concentration was quantified with Quant-It PicoGreen (Invitrogen, Carlsbad, CA). Approximately 100 ng of DNA from each sample was amplified with a TempliPhi amplification kit (Amersham Biosciences, Piscataway, NJ) and labeled with Cy5 (72). The fluorescently labeled DNA was hybridized with the GeoChip 3.0 on a MAUI hybridization system (Biomicro Systems, Salt Lake City, UT) at 42°C for 12 h (39).

GeoChip data processing and statistical analysis. Gene- and group-specific probes for GeoChip 3.0 were designed by CommOligo 2.0, and the oligonucleotide probes were synthesized by Invitrogen (Carlsbad, CA). These probes were arrayed onto Corning UltraGAPS (Corning, NY) slides by a Microgrid II arrayer (Genomic Solutions, Ann Arbor, MI) as described previously (39). The microarray slides were scanned by a Pro Scan array microarray scanner (PerkinElmer, Boston, MA) with 95% laser power and 75% photomultiplier tube (PMT) gain, and the images were analyzed by ImaGene 6.0 (Biodiscovery, El Segundo, CA). Raw data were upload to the IEG microarray processing pipeline (<http://ieg.ou.edu/microarray/>) after the removal of poor spots with a signal-to-noise ratio (SNR) (SNR = [signal mean – background mean]/background standard deviation) of >2.0 as previously reported (73). Positive genes were those that were detected with at least two probes from 6 replicates samples. These positive genes were left for further statistical analyses.

The differences between soil properties and the total signal intensities of individual genes at eCO₂ and aCO₂ at different depths (0 to 5 cm and 5 to 15 cm) were calculated by unpaired *t* tests. The differences of microbial functional gene structures and phylogenetic structures between eCO₂ and aCO₂ at each depth were analyzed by detrended correspondence analysis (DCA), permutational multivariate analysis of variance (Adonis), analysis of similarities (ANOSIM), and a multiresponse permutation procedure (MRPP). The correlation between the microbial functional structures and environmental variables was evaluated by canonical correspondence analysis (CCA) and variation partitioning analysis (VPA). All statistical analyses were performed with R project v.3.2.1 (www.R-project.org).

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/AEM.01694-17>.

SUPPLEMENTAL FILE 1, PDF file, 0.3 MB.

ACKNOWLEDGMENTS

We thank Richard J. Norby for providing plant response data. We also thank James Walter Voordeckers for carefully editing the grammar of the manuscript and for some valuable suggestions for the final version.

This project was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences ([CAS] grant XDB15010302), the National Natural Science Foundation of China (grant no. 31540071), the National Key Research and Development Program (grant no. 2016YFC0500702), the China Postdoctoral Science Foundation (2016M601145), and the Natural Science Foundation of Liaoning Province of China (201602361).

REFERENCES

- Beerling DJ, Taylor LL, Bradshaw CD, Lunt DJ, Valdes PJ, Banwart SA, Pagani M, Leake JR. 2012. Ecosystem CO₂ starvation and terrestrial silicate weathering: mechanisms and global-scale quantification during the late Miocene. *J Ecol* 100:31–41. <https://doi.org/10.1111/j.1365-2745.2011.01905.x>.
- Schröder P, Herzog R, Bojinov B, Ruttens A, Nehnevajova E, Stamatiadis S, Memon A, Vassilev A, Caviezel M, Vangronsveld J. 2008. Bioenergy to save the world. Producing novel energy plants for growth on abandoned land. *Environ Sci Pollut Res Int* 15:196–204. <https://doi.org/10.1065/espr2008.03.481>.
- He Z, Xu M, Deng Y, Kang S, Kellogg L, Wu L, Van Nostrand JD, Hobbie SE, Reich PB, Zhou J. 2010. Metagenomic analysis reveals a marked divergence in the structure of belowground microbial communities at elevated CO₂. *Ecol Lett* 13:564–575. <https://doi.org/10.1111/j.1461-0248.2010.01453.x>.
- Xu M, He Z, Deng Y, Wu L, Van Nostrand JD, Hobbie SE, Reich PB, Zhou J. 2013. Elevated CO₂ influences microbial carbon and nitrogen cycling. *BMC Microbiol* 13:124. <https://doi.org/10.1186/1471-2180-13-124>.
- Luo Y, Hui D, Zhang D. 2006. Elevated CO₂ stimulates net accumulations of carbon and nitrogen in land ecosystems: a meta-analysis. *Ecology* 87:53–63. <https://doi.org/10.1890/04-1724>.
- Bardgett RD, Freeman C, Ostle NJ. 2008. Microbial contributions to climate change through carbon cycle feedbacks. *ISME J* 2:805–814. <https://doi.org/10.1038/ismej.2008.58>.
- Delgado-Baquerizo M, Maestre FT, Reich PB, Jeffries TC, Gaitan JJ, Encinar D, Berdugo M, Campbell CD, Singh BK. 2016. Microbial diversity drives multifunctionality in terrestrial ecosystems. *Nat Commun* 7:10541. <https://doi.org/10.1038/ncomms10541>.
- Carney KM, Hungate BA, Drake BG, Megonigal JP. 2007. Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proc Natl Acad Sci U S A* 104:4990–4995. <https://doi.org/10.1073/pnas.0610045104>.
- ESF-Forest FACE Group, Calfapietra C, Ainsworth EA, Beier C, De Angelis P, Ellsworth DS, Godbold DL, Hendrey GR, Hickler T, Hoosbeek MR, Karnosky DF, King J, Körner C, Leakey AD, Lewin KF, Liberloo M, Long SP, Lukac M, Matyssek R, Miglietta F, Nagy J, Norby RJ, Oren R, Percy KE, Rogers A, Mugnozza GS, Stitt M, Taylor G, Ceulemans R. 2010. Challenges in elevated CO₂ experiments on forests. *Trends Plant Sci* 15:5–10. <https://doi.org/10.1016/j.tplants.2009.11.001>.
- Phillips RP, Finzi AC, Bernhardt ES. 2011. Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO₂ fumigation. *Ecol Lett* 14:187–194. <https://doi.org/10.1111/j.1461-0248.2010.01570.x>.
- Norby RJ, DeLucia EH, Gielen B, Calfapietra C, Giardina CP, King JS, Ledford J, McCarthy HR, Moore DJ, Ceulemans R. 2005. Forest response to elevated CO₂ is conserved across a broad range of productivity. *Proc Natl Acad Sci U S A* 102:18052–18056. <https://doi.org/10.1073/pnas.0509478102>.
- Ainsworth EA, Long SP. 2005. What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytol* 165:351–371. <https://doi.org/10.1111/j.1469-8137.2004.01224.x>.
- Iversen CM, Ledford J, Norby RJ. 2008. CO₂ enrichment increases carbon and nitrogen input from fine roots in a deciduous forest. *New Phytol* 179:837–847. <https://doi.org/10.1111/j.1469-8137.2008.02516.x>.
- Tausz-Posch S, Seneweera S, Norton RM, Fitzgerald GJ, Tausz M. 2012. Can a wheat cultivar with high transpiration efficiency maintain its yield advantage over a near-isogenic cultivar under elevated CO₂? *Field Crops Res* 133:160–166. <https://doi.org/10.1016/j.fcr.2012.04.007>.
- Iversen CM. 2010. Digging deeper: fine-root responses to rising atmospheric CO₂ concentration in forested ecosystems. *New Phytol* 186:346–357. <https://doi.org/10.1111/j.1469-8137.2009.03122.x>.
- Körner C, Aschhoff R, Bignucolo O, Hättenschwiler S, Keel SG, Peláez-Riedl S, Pepin S, Siegwolf RT, Zotz G. 2005. Carbon flux and growth in mature deciduous forest trees exposed to elevated CO₂. *Science* 309:1360–1362. <https://doi.org/10.1126/science.1113977>.
- Xiong J, He Z, Shi S, Kent A, Deng Y, Wu L, Van Nostrand JD, Zhou J. 2015. Elevated CO₂ shifts the functional structure and metabolic potentials of soil microbial communities in a C4 agroecosystem. *Sci Rep* 5:9316. <https://doi.org/10.1038/srep09316>.
- He Z, Piceno Y, Deng Y, Xu M, Lu Z, DeSantis T, Andersen G, Hobbie SE, Reich PB, Zhou J. 2012. The phylogenetic composition and structure of soil microbial communities shifts in response to elevated carbon dioxide. *ISME J* 6:259–272. <https://doi.org/10.1038/ismej.2011.99>.
- Deng Y, He Z, Xiong J, Yu H, Xu M, Hobbie SE, Reich PB, Schadt CW, Kent A, Pendall E. 2016. Elevated carbon dioxide accelerates the spatial turnover of soil microbial communities. *Glob Chang Biol* 22:957–964. <https://doi.org/10.1111/gcb.13098>.
- Deng Y, He Z, Xu M, Qin Y, Van Nostrand JD, Wu L, Roe BA, Wiley G, Hobbie SE, Reich PB. 2012. Elevated carbon dioxide alters the structure of soil microbial communities. *Appl Environ Microbiol* 78:2991–2995. <https://doi.org/10.1128/AEM.06924-11>.
- Dunbar J, Eichorst SA, Gallegos-Graves LV, Silva S, Xie G, Hengartner N, Evans RD, Hungate BA, Jackson RB, Megonigal JP. 2012. Common bacterial responses in six ecosystems exposed to 10 years of elevated atmospheric carbon dioxide. *Environ Microbiol* 14:1145–1158. <https://doi.org/10.1111/j.1462-2920.2011.02695.x>.
- Weber CF, Zak DR, Hungate BA, Jackson RB, Vilgalys R, Evans RD, Schadt CW, Megonigal JP, Kuske CR. 2011. Responses of soil cellulolytic fungal communities to elevated atmospheric CO₂ are complex and variable across five ecosystems. *Environ Microbiol* 13:2778–2793. <https://doi.org/10.1111/j.1462-2920.2011.02548.x>.
- Austin EE, Castro HF, Sides KE, Schadt CW, Classen AT. 2009. Assessment of 10 years of CO₂ fumigation on soil microbial communities and function in a sweetgum plantation. *Soil Biol Biochem* 41:514–520. <https://doi.org/10.1016/j.soilbio.2008.12.010>.
- Grüter D, Schmid B, Brandl H. 2006. Influence of plant diversity and elevated atmospheric carbon dioxide levels on belowground bacterial diversity. *BMC Microbiol* 6:68. <https://doi.org/10.1186/1471-2180-6-68>.
- Zak DR, Pregitzer KS, Curtis PS, Holmes WE. 2000. Atmospheric CO₂ and the composition and function of soil microbial communities. *Ecol Appl* 10:47–59. [https://doi.org/10.1890/1051-0761\(2000\)010\[0047:ACATCA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0047:ACATCA]2.0.CO;2).
- Castro HF, Classen AT, Austin EE, Norby RJ, Schadt CW. 2010. Soil microbial community responses to multiple experimental climate change drivers. *Appl Environ Microbiol* 76:999–1007. <https://doi.org/10.1128/AEM.02874-09>.
- Zak DR, Pregitzer KS, King JS, Holmes WE. 2000. Elevated atmospheric CO₂, fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytol* 147:201–222. <https://doi.org/10.1046/j.1469-8137.2000.00687.x>.
- van der Heijden MGA, Bardgett RD, van Straalen NM. 2008. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Lett* 11:296–310. <https://doi.org/10.1111/j.1461-0248.2007.01139.x>.
- Xiong J, He Z, Van Nostrand JD, Luo G, Tu S, Zhou J, Wang G. 2012. Assessing the microbial community and functional genes in a vertical soil profile with long-term arsenic contamination. *PLoS One* 7:e50507. <https://doi.org/10.1371/journal.pone.0050507>.
- He Z, Xiong J, Kent AD, Deng Y, Xue K, Wang G, Wu L, Van Nostrand JD, Zhou J. 2014. Distinct responses of soil microbial communities to elevated CO₂ and O₃ in a soybean agro-ecosystem. *ISME J* 8:714–726. <https://doi.org/10.1038/ismej.2013.177>.
- Griffiths RI, Whiteley AS, O'Donnell AG, Bailey MJ. 2003. Influence of depth and sampling time on bacterial community structure in an upland grassland soil. *FEMS Microbiol Ecol* 43:35–43. <https://doi.org/10.1111/j.1574-6941.2003.tb01043.x>.
- Leavitt S, Pendall E, Paul E, Brooks T, Kimball B, Pinter P, Johnson H, Matthias A, Wall G, LaMorte R. 2001. Stable-carbon isotopes and soil organic carbon in wheat under CO₂ enrichment. *New Phytol* 150:305–314. <https://doi.org/10.1046/j.1469-8137.2001.00113.x>.
- Norby RJ, Hanson PJ, O'Neill EG, Tschaplinski TJ, Weltzin JF, Hansen RA, Cheng W, Wullschlegel SD, Gunderson CA, Edwards NT, Johnson DW. 2002. Net primary productivity of a CO₂-enriched deciduous forest and the implications for carbon storage. *Ecol Appl* 12:1261–1266. [https://doi.org/10.1890/1051-0761\(2002\)012\[1261:NPPOAC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2002)012[1261:NPPOAC]2.0.CO;2).
- McCarthy HR, Oren R, Johnson KH, Gallet-Budynek A, Pritchard SG, Cook CW, LaDeau SL, Jackson RB, Finzi AC. 2010. Re-assessment of plant carbon dynamics at the Duke free-air CO₂ enrichment site: interactions of atmospheric [CO₂] with nitrogen and water availability over stand

- development. *New Phytol* 185:514–528. <https://doi.org/10.1111/j.1469-8137.2009.03078.x>.
35. Pritchard SG, Strand AE, McCormack M, Davis MA, Finzi AC, Jackson RB, Matamala R, Rogers HH, Oren R. 2008. Fine root dynamics in a loblolly pine forest are influenced by free-air-CO₂-enrichment: a six-year-minirhizotron study. *Glob Chang Biol* 14:588–602. <https://doi.org/10.1111/j.1365-2486.2007.01523.x>.
 36. Norby RJ, Zak DR. 2011. Ecological lessons from free-air CO₂ enrichment (FACE) experiments. *Annu Rev Ecol Syst* 42:181–203. <https://doi.org/10.1146/annurev-ecolsys-102209-144647>.
 37. Jastrow JD, Michael Miller R, Matamala R, Norby RJ, Boutton TW, Rice CW, Owensby CE. 2005. Elevated atmospheric carbon dioxide increases soil carbon. *Glob Chang Biol* 11:2057–2064. <https://doi.org/10.1111/j.1365-2486.2005.01077.x>.
 38. Iversen CM, Keller JK, Garten CT, Norby RJ. 2012. Soil carbon and nitrogen cycling and storage throughout the soil profile in a sweetgum plantation after 11 years of CO₂-enrichment. *Glob Chang Biol* 18:1684–1697. <https://doi.org/10.1111/j.1365-2486.2012.02643.x>.
 39. He Z, Deng Y, Van Nostrand JD, Tu Q, Xu M, Hemme CL, Li X, Wu L, Gentry TJ, Yin Y. 2012. GeoChip 3.0 as a high-throughput tool for analyzing microbial community composition, structure and functional activity. *ISME J* 4:1167–1179. <https://doi.org/10.1038/ismej.2010.46>.
 40. Lu Z, He Z, Parisi VA, Kang S, Deng Y, Van Nostrand JD, Masoner JR, Cozzarelli IM, Suflita JM, Zhou J. 2012. GeoChip-based analysis of microbial functional gene diversity in a landfill leachate-contaminated aquifer. *Environ Sci Technol* 46:5824–5833. <https://doi.org/10.1021/es300478j>.
 41. Zhang Y, Tian Z, Liu M, Shi ZJ, Hale L, Zhou J, Yang M. 2015. High concentrations of the antibiotic spiramycin in wastewater lead to high abundance of ammonia-oxidizing archaea in nitrifying populations. *Environ Sci Technol* 49:9124–9132. <https://doi.org/10.1021/acs.est.5b01293>.
 42. Cai M, Nie Y, Chi C-Q, Tang Y-Q, Li Y, Wang X-B, Liu Z-S, Yang Y, Zhou J, Wu X-L. 2015. Crude oil as a microbial seed bank with unexpected functional potentials. *Sci Rep* 5:16057. <https://doi.org/10.1038/srep16057>.
 43. Yu H, Chen C, Ma J, Liu W, Zhou J, Lee D-J, Ren N, Wang A. 2014. GeoChip-based analysis of the microbial community functional structures in simultaneous desulfurization and denitrification process. *J Environ Sci (China)* 26:1375–1382. <https://doi.org/10.1016/j.jes.2014.05.001>.
 44. Yu H, Chen C, Ma J, Xu X, Fan R, Wang A. 2014. Microbial community functional structure in response to micro-aerobic conditions in sulfate-reducing sulfur-producing bioreactor. *J Environ Sci (China)* 26:1099–1107. [https://doi.org/10.1016/S1001-0742\(13\)60589-6](https://doi.org/10.1016/S1001-0742(13)60589-6).
 45. Yu Z, He Z, Tao X, Zhou J, Yang Y, Zhao M, Zhang X, Zheng Z, Yuan T, Liu P. 2016. The shifts of sediment microbial community phylogenetic and functional structures during chromium(VI) reduction. *Ecotoxicology* 25:1759–1770. <https://doi.org/10.1007/s10646-016-1719-6>.
 46. Norby RJ, Warren JM, Iversen CM, Medlyn BE, McMurtrie RE. 2010. CO₂ enhancement of forest productivity constrained by limited nitrogen availability. *Proc Natl Acad Sci U S A* 107:19368–19373. <https://doi.org/10.1073/pnas.1006463107>.
 47. Norby RJ, Iversen CM, Childs J, Tharp ML. 2010. ORNL net primary productivity data. Carbon Dioxide Information Analysis Center, U.S. Department of Energy, Oak Ridge National Laboratory, Oak Ridge, TN. <http://cdiac.ornl.gov>.
 48. Morales SE, Holben WE. 2014. Simulated geologic carbon storage leak reduces bacterial richness and alters bacterial community composition in surface soil. *Soil Biol Biochem* 76:286–296. <https://doi.org/10.1016/j.soilbio.2014.05.018>.
 49. Sinsabaugh R, Saiya-Cork K, Long T, Osgood M, Neher D, Zak D, Norby R. 2003. Soil microbial activity in a *Liquidambar* plantation unresponsive to CO₂-driven increases in primary production. *Appl Soil Ecol* 24:263–271. [https://doi.org/10.1016/S0929-1393\(03\)00002-7](https://doi.org/10.1016/S0929-1393(03)00002-7).
 50. Zhou J, He Z, Van Nostrand JD, Wu L, Deng Y. 2010. Applying GeoChip analysis to disparate microbial communities. *Microbe Wash DC* 5:60–65.
 51. Pritchard S. 2011. Soil organisms and global climate change. *Plant Pathol* 60:82–99. <https://doi.org/10.1111/j.1365-3059.2010.02405.x>.
 52. Weber CF, Vilgalys R, Kuske CR. 2013. Changes in fungal community composition in response to elevated atmospheric CO₂ and nitrogen fertilization varies with soil horizon. *Front Microbiol* 4:78. <https://doi.org/10.3389/fmicb.2013.00078>.
 53. Chung H, Zak DR, Reich PB, Ellsworth DS. 2007. Plant species richness, elevated CO₂, and atmospheric nitrogen deposition alter soil microbial community composition and function. *Glob Chang Biol* 13:980–989. <https://doi.org/10.1111/j.1365-2486.2007.01313.x>.
 54. Liu Y, Zhang H, Xiong M, Li F, Li L, Wang G, Pan G. 2017. Abundance and composition response of wheat field soil bacterial and fungal communities to elevated CO₂ and increased air temperature. *Biol Fertil Soils* 53:3–8. <https://doi.org/10.1007/s00374-016-1159-8>.
 55. Okubo T, Liu D, Tsurumaru H, Ikeda S, Asakawa S, Tokida T, Tago K, Hayatsu M, Aoki N, Ishimaru K, Ujiie K, Usui Y, Nakamura H, Sakai H, Hayashi K, Hasegawa T, Minamisawa K. 2015. Elevated atmospheric CO₂ levels affect community structure of rice root-associated bacteria. *Front Microbiol* 6:136. <https://doi.org/10.3389/fmicb.2015.00136>.
 56. Inubushi K, Cheng W, Aonuma S, Hoque MM, Kobayashi K, Miura S, Kim HY, Okada M. 2003. Effects of free-air CO₂ enrichment (FACE) on CH₄ emission from a rice paddy field. *Glob Chang Biol* 9:1458–1464. <https://doi.org/10.1046/j.1365-2486.2003.00665.x>.
 57. Reich PB, Hobbie SE, Lee T, Ellsworth DS, West JB, Tilman D, Knops JM, Naem S, Trost J. 2006. Nitrogen limitation constrains sustainability of ecosystem response to CO₂. *Nature* 440:922–925. <https://doi.org/10.1038/nature04486>.
 58. Aerts R, Bakker C, De Caluwe H. 1992. Root turnover as determinant of the cycling of C, N, and P in a dry heathland ecosystem. *Biogeochemistry* 15:175–190. <https://doi.org/10.1007/BF00002935>.
 59. Guo DL, Mitchell RJ, Hendricks JJ. 2004. Fine root branch orders respond differentially to carbon source-sink manipulations in a longleaf pine forest. *Oecologia* 140:450–457. <https://doi.org/10.1007/s00442-004-1596-1>.
 60. Finzi AC, Moore DJ, DeLucia EH, Lichter J, Hofmockel KS, Jackson RB, Kim H-S, Matamala R, McCarthy HR, Oren R. 2006. Progressive nitrogen limitation of ecosystem processes under elevated CO₂ in a warm-temperate forest. *Ecology* 87:15–25. <https://doi.org/10.1890/04-1748>.
 61. Baggs E, Richter M, Hartwig U, Cadisch G. 2003. Nitrous oxide emissions from grass swards during the eighth year of elevated atmospheric pCO₂ (Swiss FACE). *Glob Chang Biol* 9:1214–1222. <https://doi.org/10.1046/j.1365-2486.2003.00654.x>.
 62. Robinson D, Conroy JP. 1998. A possible plant-mediated feedback between elevated CO₂, denitrification and the enhanced greenhouse effect. *Soil Biol Biochem* 31:43–53. [https://doi.org/10.1016/S0038-0717\(98\)00102-3](https://doi.org/10.1016/S0038-0717(98)00102-3).
 63. Knelman JE, Legg TM, O'Neill SP, Washenberger CL, González A, Cleveland CC, Nemergut DR. 2012. Bacterial community structure and function change in association with colonizer plants during early primary succession in a glacier forefield. *Soil Biol Biochem* 46:172–180. <https://doi.org/10.1016/j.soilbio.2011.12.001>.
 64. Bezemer T, Lawson CS, Hedlund K, Edwards AR, Brook AJ, Igual JM, Mortimer SR, van der Putten WH. 2006. Plant species and functional group effects on abiotic and microbial soil properties and plant-soil feedback responses in two grasslands. *J Ecol* 94:893–904. <https://doi.org/10.1111/j.1365-2745.2006.01158.x>.
 65. Allison V J YZ, Miller RM, Jastrow JD, Matamala R. 2007. Using landscape and depth gradients to decouple the impact of correlated environmental variables on soil microbial community composition. *Soil Biol Biochem* 39:505–516. <https://doi.org/10.1016/j.soilbio.2006.08.021>.
 66. Wieland G, Neumann R, Backhaus H. 2001. Variation of microbial communities in soil, rhizosphere, and rhizoplane in response to crop species, soil type, and crop development. *Appl Environ Microbiol* 67:5849–5854. <https://doi.org/10.1128/AEM.67.12.5849-5854.2001>.
 67. Norby RJ, Todd DE, Fuels J, Johnson DW. 2001. Allometric determination of tree growth in a CO₂-enriched sweetgum stand. *New Phytol* 150:477–487. <https://doi.org/10.1046/j.1469-8137.2001.00099.x>.
 68. van Miegroet H, Norby R, Tschaplinski T. 1994. Nitrogen fertilization strategies in a short-rotation sycamore plantation. *For Ecol Manage* 64:13–24. [https://doi.org/10.1016/0378-1127\(94\)90123-6](https://doi.org/10.1016/0378-1127(94)90123-6).
 69. Norby RJ, Iversen CM, Tharp ML. 2012. ORNL FACE nitrogen concentrations. Oak Ridge National Laboratories, Oak Ridge, TN.
 70. Nelson DW, Sommers LE. 1996. Total carbon, organic carbon, and organic matter, p 961–1010. In Sparks DL, Page AL, Helmke PA, Loeppert RH, Soltanpour PN, Tabatabai MA, Johnston CT, Sumner ME (ed), *Methods of soil analysis. Part 3. Chemical methods*. Soil Science Society of America, Inc., American Society of Agronomy, Inc., Madison, WI.
 71. Zhou J, Bruns MA, Tiedje JM. 1996. DNA recovery from soils of diverse composition. *Appl Environ Microbiol* 62:316–322.
 72. Wu L, Liu X, Schadt CW, Zhou J. 2006. Microarray-based analysis of subnanogram quantities of microbial community DNAs by using whole-community genome amplification. *Appl Environ Microbiol* 72:4931–4941. <https://doi.org/10.1128/AEM.02738-05>.
 73. He Z, Zhou J. 2008. Empirical evaluation of a new method for calculating signal-to-noise ratio for microarray data analysis. *Appl Environ Microbiol* 74:2957–2966. <https://doi.org/10.1128/AEM.02536-07>.