Bioremediation potential of hydrocarbon-utilizing fungi from select marine niches of India

Natasha Maria Barnes¹ · Vishwas B. Khodse¹ · Nikita P. Lotlikar¹ · Ram Murti Meena¹ · Samir R. Damare¹

Received: 12 April 2017 / Accepted: 8 December 2017 / Published online: 18 December 2017
© Springer-Verlag GmbH Germany, part of Springer Nature 2017

Abstract

Ten fungal isolates with an ability to degrade crude oil were isolated from select marine substrates, such as mangrove sediments, Arabian Sea sediments, and tarballs. Out of the ten isolates, six belonged to *Aspergillus*, two to *Fusarium* and one each to *Penicillium* and *Acremonium* as identified using ITS rDNA sequencing. The selected ten fungal isolates were found to degrade the long-chain *n*-alkanes as opposed to short-chain *n*-alkanes from the crude oil. Mangrove fungus #NIOSN-M126 (*Penicillium citrinum*) was found to be highly efficient in biodegradation of crude oil, reducing the total crude oil content by 77% and the individual *n*-alkane fraction by an average of 95.37%, indicating it to be a potential candidate for the development into a bioremediation agent.

Keywords Bioremediation · Degradation · Mangrove fungi · Marine pollution · Oil spills · Tarballs

Introduction

Oil spillages into the environment have become one of the biggest problems in the last century. The hydrocarbons spread horizontally on the ground/water surface and partition into groundwater, soil pore airspace and on the surfaces of soil particles (Pohl et al. 2002), causing contamination of large volumes of soil and groundwater. Oil spills, tank leakages, and wastewater disposal are the key routes for the entry of tons of hydrocarbons into the environment, and which occur at every stage of production, transportation, and handling of petroleum products. Crude oil is a complex mixture of hydrophobic components such as *n*-alkanes, aromatics, resins, and asphaltenes. Some fractions of crude oil are toxic to living organisms (Costello 1979). The leakage of crude oil into soil damages the biota residing in the soil, including microorganisms and plants, and becomes increasingly toxic beyond a concentration of 3% (Onuoha et al. 2003).

In recent years, various technologies have emerged to manage hydrocarbon contamination. Bioremediation comprises a set of technologies that aid in the removal of contaminants or render them less harmful using biological activity (Silva et al. 2015). This can be achieved with the help of unique microorganisms, such as bacteria, microalgae, and fungi, either isolated from a different environment or introduced into the contaminated sites or by enriching the organisms already present at the contaminated sites (Adebusoye et al. 2007). Bioremediation of hydrocarbons is preferred to chemical and conventional treatment for several reasons. In bioremediation, the selected and potential microbes utilize crude oil as carbon substrate and biodegrade the hydrocarbons into products that are non-toxic and comparatively safer to the environment and living beings, such as carbon dioxide and water (Toledo et al. 2006). The process is economical and can be directly applied to the contaminated site. Thus, microbial remediation is a promising, practical and economical method for complete mineralization of hydrocarbons into carbon dioxide and water (Wang et al. 2015). Hence, the present study aimed at harnessing the potential of fungi isolated from three marine niches of India, towards efficient hydrocarbon degradation and detoxification.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13205-017-1043-8>) contains supplementary material, which is available to authorized users.

✉ Samir R. Damare
samir@nio.org

¹ Biological Oceanography Division, CSIR-National Institute of Oceanography, Dona Paula, Panaji 403004, India

Materials and methods

Fungal isolates used in the study

In our previous studies, a total of 342 marine fungi were isolated. Out of these 342 fungal isolates, 167 were isolated from sediments collected from Divar mangroves (out-skirts of Panaji, Goa, on the shores of Mandovi estuary) and 165 from off-shore sediments collected from the Arabian Sea (68°E 21°N) when cultured on Czapek Dox agar, malt extract agar, corn meal agar, Sabouraud dextrose agar and potato dextrose agar (all media prepared in seawater). A total of ten marine fungi were isolated from tarball samples (collected from Betul beach, South Goa) cultured on Czapek Dox agar and malt extract agar. Sediment samples from mangrove vegetated area were collected during low tide, and approximately 0.5 g of soil was suspended in sterile seawater, and 100 µL aliquots were spread-plated on various growth media as mentioned above. Crude oil used in this study was obtained from off-shore drilling site operated by Oil and Natural Gas Corporation (ONGC) at Bombay High, India.

Screening

A total of 51 fungal isolates were screened for their ability to utilize hydrocarbon as sole carbon source. The inocula were prepared by growing the fungus in Czapek Dox broth at 30 °C under static conditions until a mycelial mat was formed. The mycelial mat was then broken down into fine fragments using sterile glass beads. This suspension was used as a 1% (v/v) inoculum for 20 mL Mineral Salt medium [0.83 g L⁻¹ potassium dihydrogen phosphate (KH₂PO₄); 10 g L⁻¹ sodium chloride (NaCl); 0.42 g L⁻¹ magnesium sulfate (MgSO₄·7H₂O); 0.29 g L⁻¹ potassium chloride (KCl); 1.25 g L⁻¹ disodium hydrogen phosphate (Na₂HPO₄); 0.42 g L⁻¹ sodium nitrate (NaNO₃)] containing 5% petrol as sole carbon source, without any other nutrient as additive. Petrol as C-source was used for initial screening and not crude oil as purified hydrocarbon products are more difficult to degrade compared to crude oil. Fungal isolates capable of growing in the medium and showing a visible increase in biomass were selected for the further studies to estimate the quantitative and qualitative degradation of crude oil.

Identification of fungal isolates

The pure fungal isolates were sub-cultured and allowed to grow for 7 days on Czapek Dox agar (CDA) medium (Fig. 1). Total genomic DNA of the isolates was extracted from the mycelia using manufacturer specified protocol

(ZR Fungal/Bacterial DNA Microprep Kit, Zymo Research, USA). The isolates were identified by ITS and beta-tubulin sequence analysis. For ITS sequence analysis, the ITS gene region was amplified using PCR with primer set: ITS1 (5'-TCCGTAGGTGAACCTGCCG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White et al. 1990). Amplification cycle involved an initial denaturation step at 95 °C for 5 min followed by 35 cycles with a denaturing step at 95 °C for 1 min, annealing temperature of 52 °C for 45 s; elongation step at 72 °C for 1 min and final elongation step at 72 °C for 10 min. The 600–750 bp amplicons obtained were stored at 4 °C and then purified using manufacturer specified protocol (Wizard SV Gel and PCR clean-up system, Promega, USA). For beta-tubulin sequence analysis, the partial beta-tubulin gene was amplified using primer set: Bt2a (5'-GGTAACCAAATCGGTGCTGCTTTC) and Bt2b (5'-ACCCTCAGTGTAGTGACCCTTGGC) (Glass and Donaldson 1995). Amplification cycle was the same as that used for ITS region, except the annealing temperature was set to 58 °C. The 550–600 bp amplicons obtained were stored at 4 °C and then purified using manufacturer specified protocol (Wizard SV Gel and PCR clean-up system, Promega, USA). The PCR purified products were then sequenced using Applied Biosystems (ABI) 3130 DNA stretch sequencer. DNA sequences obtained were analyzed using the BLAST tool to get the closest reference sequence (Altschul et al. 1990).

A multiple sequence alignment was carried out using the ITS sequences obtained in this study and the reference sequences obtained from NCBI-GenBank using with MEGA version 5.2 (Tamura et al. 2011). The phylogenetic trees were inferred using the Neighbour-Joining method (Saitou and Nei 1987), and bootstrap analyses were performed. The evolutionary distances were computed using the Maximum Composite Likelihood method (Tamura et al. 2004).

Quantitative estimation of crude oil degradation using gravimetry

The comparative abilities of ten fungal isolates (NIOSN-M113, NIOSN-M126, NIOSN-M142, NIOSN-M109, NIOSN-SK56S57, NIOSN-SK56S32, NIOSN-SK56S22, NIOSN-SK56C42, NIOSN-T5, and NIOSN-T4) to degrade crude oil were tested. Of the ten fungal isolates, four were obtained from mangrove environments (NIOSN-M113, NIOSN-M126, NIOSN-M142, and NIOSN-M127), four from marine environments (NIOSN-SK56S57, NIOSN-SK56S32, NIOSN-SK56S22 and NIOSN-SK56C42) and two from tarballs (NIOSN-T5 and NIOSN-T4). The fungal inoculum was prepared by sub-culturing into Czapek Dox broth as described above. For the biodegradation experiments, the fungal mycelia were cultivated under shaker conditions in 100-mL Erlenmeyer flasks containing 20 mL

Fig. 1 Growth of all the ten isolates used in the study on Czapek Dox agar. Inset

of mineral salts medium. Two milliliter of crude oil (corresponding to 2.67 g, which amounted to 13.35% w/v) was used as the sole carbon source. The culture flasks were incubated at 28 °C on a rotary shaker at 80 rpm for 23 days (till visible growth was observed). Uninoculated media served as a control. All the experiments were carried out in triplicates.

Following incubation, the residual crude oil was sequentially extracted three times from the culture medium with dichloromethane (1:1) for exhaustive separation. Sample with dichloromethane was placed in a separating funnel with continuous shaking, after which the contents were allowed to settle; two layers were formed: aqueous layer and organic dichloromethane layer containing the residual crude oil. The organic layer was decanted in pre-weighed vials and air dried. After dichloromethane evaporation, the residual oil was quantified gravimetrically (modified from Elshafie et al. 2007). The percentage degradation of the crude oil was then calculated as:

$$\frac{[\text{Weight of crude oil (initial)} - \text{weight of crude oil (after treatment)}]}{\text{Weight of crude oil (initial)}} \times 100.$$

Qualitative analysis of crude oil degradation using gas chromatography

After gravimetric quantification, the residual crude oil was fractionated into alkane fractions on a silica gel column. For this purpose, samples were dissolved in *n*-hexane and the soluble fraction loaded onto a silica column and alkane fraction eluted with *n*-hexane (modified from Mishra et al. 2001). The solvent fractions were then evaporated to partial dryness with a rotary evaporator followed by complete drying using nitrogen gas. Dried samples were re-dissolved in 500 µL *n*-hexane and spiked with internal standard *n*-methyl nona-decanoate and quantified via capillary gas chromatography using Shimadzu GC 2010 plus Chromatograph equipped with a flame-ionization detector (FID), fused silica capillary column of 30 m length, 0.32 mm internal diameter, and 0.25 µm film thickness. The detector and injector temperatures were maintained at 325 and 300 °C, respectively.

The column temperature was programmed to increase from 60 to 280 °C with a rate of 5 °C min⁻¹ and a final time of 15 min. High-purity nitrogen (99.9995%, oxygen-free) was used as a carrier gas at a flow rate of 3 mL min⁻¹. The method was validated using C₇–C₃₀ *n*-alkane standards obtained from Sigma Chemicals.

Toxicity assay using *Artemia salina* larvae

The bioremediation products were tested for their toxicity using *Artemia* larvae. The *Artemia* cysts were allowed to hatch in seawater 24 h before conducting the assay. Following the incubation period of 23 days, the culture broth spiked with crude oil was tested for toxic biodegradation by-products. For this, *A. salina* (10 numbers) were inoculated into mineral salts medium from the test as well as the control flasks and microscopic observations were made daily to record the percent survival of the animals.

Results

Screening of fungi for hydrocarbon degradation

Of the total 342 marine fungi isolated in our previous studies, 51 fast-growing fungal isolates were screened for their ability to utilize and degrade hydrocarbons. Among the 51 fungal isolates, 10 isolates exhibited a potential to use crude oil as the sole carbon source and selected for quantification of the degradation.

Identification of fungal isolates

The evolutionary relationships of the fungal isolates are presented in the NJ tree (Figs. 2, 3).

The trees as obtained from MEGA software are reproduced here. Figure 2 involves 34 ITS sequences in all, out of which 10 were obtained in this study, while Fig. 3 involves 30 β -tubulin sequences, out of which 10 are from this study. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) is shown next to the branches. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Maximum Composite Likelihood method (Tamura et al. 2004). Figure 4 shows the generic distribution of the selected 51 isolates where more than half are contributed by *Aspergillus* (12), *Cladosporium* (11) and *Penicillium* (11) (Table 1).

Quantitative estimation of crude oil degradation using gravimetric methods

The selected ten fungal isolates were able to grow and degrade crude oil. Percent degradation estimation following gravimetric analysis demonstrated that isolate *Penicillium citrinum* NIOSN-M126 obtained from Divar mangrove sediments showed maximum ability to utilize crude oil, giving the highest percent degradation of 77%, followed by *Aspergillus flavus* NIOSN-SK56S22 isolated from Arabian Sea sediments indicating a 62% degradation (Fig. 5).

Qualitative analysis of crude oil degradation using gas chromatography

The residual oil at the end of the incubation time was analyzed qualitatively and quantitatively using gas chromatography. The chromatograms showed a number of peaks, each representing the residual hydrocarbon in crude oil (Supplementary Figure). The *n*-alkanes detected in crude oil were in the range C₈–C₃₀. C₇ *n*-alkane was not detected in any of the samples. In general, the higher alkanes (C₃₀–C₁₄) showed over 80% degradation for all the isolates (Supplementary Table). *P. citrinum* NIOSN-M126 showed the highest efficiency of degradation of the long-chain *n*-alkanes (C₂₉–C₂₀), while *Aspergillus sydowii* NIOSN-SK56C42 was least efficient. Carbon chain C30 was completely degraded by isolate *A. flavus* NIOSN-SK56S22 giving 100% degradation (Supplementary Table). Among the short-chain *n*-alkanes, *Aspergillus* sp. NIOSN-SK56S32 and *Acremonium sclerotigenum* NIOSN-M109 showed an increase in the concentration of carbon number 12. This could be due to the degradation of higher chain alkanes to C₁₂ increasing the final concentration of Carbon 12 (Supplementary Table). Among the ten isolates screened, *P. citrinum* NIOSN-M126 proved to be most efficient in degradation of the higher chain mutagenic *n*-alkanes and a promising bioremediating agent.

Toxicity assay using *A. salina* larvae

Investigations for the production of toxic by-products from degradation of crude oil by the fungal isolates were carried out with *A. salina*, the brine shrimp, as a test organism. The chief advantage of using these organisms lies in the ease of handling, relatively small amounts of effluents required for tests and the ease by which data may be obtained. The survival percentages of test organism for the treated and untreated experimental flasks are given in Table 2. The results indicate that there were no toxic intermediates or bioremediation products produced by the fungi tested.

Fig. 2 The phylogenetic relationships of marine hydrocarbon-degrading fungi inferred from NJ analysis of the ITS sequences using MEGA

Discussion

Though there have been quite a few studies on hydrocarbon-degrading fungi, our understanding of the potential of fungi as bioremediation agents need to be improved. Fungi are the decomposers in the global cycle of nutrients. Fungi possess decomposing abilities to deal with an array of naturally occurring compounds that serve as potential carbon sources. Hydrocarbon pollutants have similar or analogous molecular structures which enable fungi to act on them as well.

Fungal genera, namely, *Amorphotheca*, *Neosartorya*, *Talaromyces*, and *Graphium* and yeast genera, namely, *Candida*, *Yarrowia*, and *Pichia* were isolated from petroleum-contaminated soil and proven to be potential organisms for hydrocarbon degradation (Chaillan et al. 2004). Singh (2006) also reported a group of terrestrial fungi, namely, *Aspergillus*, *Cephalosporium*, and *Penicillium* which were also found to be the potential degraders of crude oil hydrocarbons.

Table 1 Source of isolation and GenBank accession numbers obtained from NCBI for the fungal isolates capable of hydrocarbon degradation

S. no.	Isolate number	Source of isolation	Closest match in NCBI		GenBank accession number	
			ITS	β -Tubulin	ITS	β -Tubulin
1	NIOSN-SK56S32	Off-shore sediments (Arabian Sea)	<i>Aspergillus</i> sp.	<i>Aspergillus oryzae</i>	KX845389	KY852305
2	NIOSN-SK56S22	Off-shore sediments (Arabian Sea)	<i>Aspergillus flavus</i>	<i>Aspergillus flavus</i>	KX845388	KY852310
3	NIOSN-SK56S57	Off-shore sediments (Arabian Sea)	<i>Aspergillus sydowii</i>	<i>Aspergillus sydowii</i>	KX845391	KY852303
4	NIOSN-SK56C42	Off-shore sediments (Arabian Sea)	<i>Aspergillus sydowii</i>	<i>Aspergillus terreus</i>	KX845390	KY852309
5	NIOSN-T4	Tar ball	<i>Fusarium equiseti</i>	<i>Fusarium equiseti</i>	KX845382	KY852311
6	NIOSN-T5	Tar ball	<i>Fusarium circinatum</i>	<i>Penicillium chermesinum</i>	KX845383	KY852312
7	NIOSN-M109	Mangrove sediments	<i>Acremonium sclerotigenum</i>	<i>Shiraia</i> sp.	KX845384	KY852306
8	NIOSN-M126	Mangrove sediments	<i>Penicillium citrinum</i>	<i>Penicillium citrinum</i>	KX845386	KY852307
9	NIOSN-M142	Mangrove sediments	<i>Aspergillus polyporicola</i>	<i>Lecanicillium</i> sp.	KX845387	KY852308
10	NIOSN-M113	Mangrove sediments	<i>Aspergillus versicolor</i>	<i>Aspergillus steynii</i>	KX845385	KY852304

Fig. 5 The percent degradation of crude oil (2.67 g) by the ten selected fungal isolates after 23 days of incubation at 28 °C on a rotary shaker at 80 rpm as estimated by gravimetric quantification. The error bars represent standard deviation for set of triplicates for each isolate.

Deep-sea isolates.
 Tarball isolates.
 Mangrove isolates

ability to degrade kerosene. They reported *A. terreus* and *Rhizopus* sp. from the tarball samples and observed that *Rhizopus* sp. exhibited the highest growth diameter in 5% kerosene. Recently, Sanyal et al. (2016) isolated tarball-associated fungi belonging to the genera *Alternaria*, *Cladosporium*, *Curvularia*, *Issatchenkia*, and *Candida* from tarballs collected from Candolim beach in Goa, India.

In this study, we focused on the *n*-alkanes from C_7 to C_{30} . The higher molecular weight alkanes have greater toxicity compared to the lower molecular weight alkanes. This is because lower molecular weight alkanes do not persist for long periods in the contaminated site due to their ability to rapidly volatilize. Also, the vapour phase of short-chain alkanes is less toxic than the liquid phase (Bartha and Atlas 1977). Among the ten isolates screened, *P. citrinum*

NIOSN-M126 proved to be most efficient in degradation of the higher chain mutagenic *n*-alkanes and a promising bioremediating agent. As opposed to the current study, previous studies available have reported the preferential degradation of short-chain *n*-alkanes. *Aspergillus* sp. isolated from flare-pit burning petroleum waste (April et al. 1998), *Aspergillus fumigatus* present in diesel storage tanks (Bento et al. 2005) and *A. terreus* and *Fusarium solani* isolated from oil contaminated soils (Colombo et al. 1996) have shown preferential degradation of short-chain alkanes. The marine fungal isolates *Glomastix* sp., *Paecilomyces* sp., and *Verticillium* sp. have also shown preferential degradation of shorter chain *n*-alkanes (Fedorak et al. 1984). *Fusarium* sp., *Aspergillus* sp., and *Scopulariopsis* sp. isolated from shorelines of the Gulf of Mexico have also shown an increased degradation

Table 2 The percent survival of *Artemia salina* larvae after exposure to fungal remediated residual crude oil mixture

Percent survival ^a														Crude oil control	Seawater Control
Days of incubation	NIOSN-SK56S32	NIOSN-SK56S22	NIOSN-SK56S57	NIOSN-SK56C42	NIOSN-T4	NIOSN-T5	NIOSN-M109	NIOSN-M126	NIOSN-M142	NIOSN-M113					
Day 0	100	100	100	100	100	100	100	100	100	100					100
Day 1	100	90	100	100	100	100	93.33	100	100	93.33					100
Day 2	80	66.66	93.33	83.33	53.33	70	80	83.33	73.33	76.66					85
Day 3	70	36.66	63.33	60	43.33	53.33	56.66	26.66	66.66	56.66					75

^a The values denoted are average values of five replicates

of short-chain alkanes compared to longer chain alkanes (Simister et al. 2015).

Among the three different environments sampled for potential hydrocarbon-degrading fungi, the isolate *P. citrinum* NIOSN-M126 obtained from mangrove sediments has shown to be more efficient in total oil degradation when compared to those isolated from tarballs and deep-sea sediments. Mangroves are often situated in areas of high anthropogenic influence, being exposed to pollutants. Divar mangroves are located on the banks of Mandovi estuary, which is used for the barge movement, ferry-boats, and other tourist boats. No previous reports are available on hydrocarbon-degrading fungi from these regions. But in a previous study, Fernandes et al. (2014) have reported the presence of a bacterium, *Marinobacter* sp. from these mangrove sediments, which is known to utilize a variety of hydrocarbons as the sole source of carbon and energy.

Oil spills are the most destructive source of pollution impacting sandy beaches, affecting all trophic levels (Defeo et al. 2009). Persistence and breakdown of stranded oil depend on sand particle size, wave energy, temperature and other factors (Owens et al. 2008), including fungal degradation (Elshafie et al. 2007). Many isolated bacterial and fungal species have been reported to be capable of biodegrading petroleum hydrocarbons (Márquez-Rocha et al. 2005). Fungi, however, have been found to be better degraders of petroleum than traditional bioremediation techniques including bacteria (Thomas 2000). Also, fungi have an advantage over bacteria as fungal hyphae can penetrate the contaminated soil to reach the hydrocarbons that have spread beyond the top layer of the soil (Novotny et al. 1999).

Furthermore, fungi have a higher tolerance to the toxicity of hydrocarbons due to their physiology and adaptation to such variations in the environment and also the presence of mechanisms for the elimination of spilled oil from the environment. A few research studies carried out in the screening, isolation and characterization of fungi capable of degrading hydrocarbons and their derivatives recognized the enormous potential of the fungal isolates for use and application in the bioremediation of hydrocarbons (Prenafeta-Boldú et al. 2001; Clemente et al. 2001; Mesyami and Baheri 2003; Yamada et al. 2002; Santos et al. 2008). Also, the isolate *P. citrinum* NIOSN-M126 is a sporulating culture, and the application of spores to contaminated sites is more feasible when compared to bacterial cells. The isolate *P. citrinum* NIOSN-M126 can be further studied for the possible use in clean-ups of oil spills and tarballs that appear on beaches, and also its amalgamation as a component in the Oil-Zapper (TERI, India) which currently comprises a bacterial consortium only. A better understanding of the metabolism, enzyme mechanisms, and genetics of hydrocarbon degrading fungi is critical for the optimization of bioremediation

processes. It is not acceptable that in any clean-up process, the by-products generated as a result of the degradation turn out to be more toxic. Fathalla (2007) had shown that photo-oxidation of crude oil in seawater could lead to the destruction of existing toxic components leading to the generation of new toxic compounds. In this study, the breakdown products generated by the fungal action were not toxic as shown by the toxicity tests using *Artemia* larvae, thereby rendering their use for degrading crude oil as safe.

Conclusions

Environmental pollution caused due to oil spillage is one of the major problems of this century which needs to be tackled with all possible means. Bioremedial measures using fungi can emerge as a key to control this problem owing to their natural abilities to degrade crude oil. Fungi isolated from different marine niches reported here have shown ability to utilize hydrocarbon as sole C-source and in the process degrade the toxic long-chain *n*-alkanes to short-chain alkanes which are easier to handle and volatilize. Fungal degradation can be considered as a key component in the clean-up strategy for petroleum hydrocarbon remediation. They need to be evaluated as possible bioremediation agents.

Acknowledgements The authors thank the Director of CSIR-National Institute of Oceanography and Head, Biological Oceanography Division for their support and providing all the facilities to carry out this study. We are thankful to Dr. Belle Damodara Shenoy for his valuable suggestions. The authors gratefully acknowledge the funding support received from the project BSC0111 funded by Council of Scientific and Industrial Research (CSIR), India. We are also thankful to the anonymous reviewers whose suggestions and constructive comments helped us to improve the manuscript. This is NIO Contribution number #####.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

References

- Adebusoye SA, Ilori MO, Amund OO, Teniola OD, Olatope SO (2007) Microbial degradation of petroleum hydrocarbons in a polluted tropical stream. *World J Microbiol Biotechnol* 23:1149–1159
- Al-Nasrawi H (2012) Biodegradation of crude oil by fungi isolated from Gulf of Mexico. *J Bioremediat Biodegrad* 3:147. <https://doi.org/10.4172/2155-6199.1000147>
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- April TM, Abbott SP, Foght JM, Currah RS (1998) Degradation of hydrocarbons in crude oil by the ascomycete *Pseudallescheria boydii* (Microasaceae). *Can J Microbiol* 44:270–278
- Bartha R, Atlas RM (1977) The microbiology of aquatic oil spills. *Adv App Microbiol* 22:225–266
- Bento FM, Beech IB, Gaylarde CC, Englert GE, Muller IL (2005) Degradation and corrosive activities of fungi in a diesel–mild steel–aqueous system. *World J Microb Biotechnol* 21:135–142
- Bik HM, Halanynch KM, Sharma J, Thomas WK (2012) Dramatic shifts in benthic microbial eukaryote communities following the deep-water horizon oil spill. *PLoS One* 7:e38550
- Chaillan FA, Le Flèche EB, Phantavong YH, Grimont PAS, Oudot J (2004) Identification and biodegradation potential of tropical aerobic hydrocarbon-degrading microorganisms. *Res Microbiol* 155:587–595
- Clemente AR, Anazawa TA, Durrant LR (2001) Biodegradation of polycyclic aromatic hydrocarbons by soil fungi. *Braz J Microbiol* 32:255–261
- Colombo JC, Cabello M, Arambarri AM (1996) Biodegradation of aliphatic and aromatic hydrocarbons by natural soil microflora and pure cultures of imperfect and lignolytic fungi. *Environ Pollut* 94:355–362
- Costello J (1979) Morbidity and mortality study of shale oil workers in the United States. *Environ Health Perspect* 30:205–208
- Defeo O, McLachlan A, Schoeman DS, Schlacher TA, Dugan J, Jones A, Lastra M, Scapini F (2009) Threats to sandy beach ecosystems: a review. *Estuar Coast Shelf Sci* 81:1–12
- Elshafie A, AlKindi AY, Al-Busaidi S, Bakheit C, Albahry SN (2007) Biodegradation of crude oil and *n*-alkanes by fungi isolated from Oman. *Mar Pollut Bull* 54:1692–1696
- Fathalla EM (2007) Degradation of crude oil in the environment: toxicity arising through photochemical oxidation in the aqueous phase. PhD thesis Munster University, Germany
- Fedorak PM, Semple KM, Westlake DWS (1984) Oil-degrading capabilities of yeasts and fungi isolated from coastal marine environments. *Can J Microbiol* 30:565–571
- Fernandes SO, Kirchman DL, Michotey VD, Bonin PC, LokaBharathi PA (2014) Bacterial diversity in relatively pristine and anthropogenically-influenced mangrove ecosystems (Goa, India). *Braz J Microb* 45:1161–1171
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol* 61:1323–1330
- Lotfinasab S, Gunale VR, Rajurkar NS (2012) Assessment of petroleum hydrocarbon degradation from soil and tarball fungi. *Biosci Discov* 3:186–192
- Márquez-Rocha FJ, Olmos-Soto J, Rosano-Hernández MC, Muriel-García M (2005) Determination of the hydrocarbon-degrading metabolic capabilities of tropical bacterial isolates. *Int Biodeterior Biodegrad* 55:17–23
- Mesyami P, Baheri M (2003) Pre-screening of fungi and bulking agents for contaminated soil bioremediation. *Adv Environ Res* 7:881–887
- Mishra S, Jyot J, Kuhad RC, Lal B (2001) In situ bioremediation potential of an oily sludge-degrading bacterial consortium. *Curr Microbiol* 43:328–335
- Novotny C, Erbanova P, Sasek V, Kubatova A, Cajthaml T, Lang E et al (1999) Extracellular oxidative enzyme production and PAH removal in soil by exploratory mycelium of white rot fungi. *Biodegradation* 10:159–168
- Onuoha CI, Arinze AE, Ataga AE (2003) Evaluation of growth of some fungi in crude oil polluted environment. *Glob J Agric Sci* 2:1596–2903
- Owens EH, Taylor E, Humphrey B (2008) The persistence and character of stranded oil on coarse-sediment beaches. *Mar Pollut Bull* 56:14–26
- Plohl K, Leskovsek H, Bricelj M (2002) Biological degradation of motor oil in water. *Acta Chim Slov* 49:279–289

- Prenafeta-Boldú FX, Andrea K, Luykx DM, Heidrun A, Van Groenestijn JW (2001) Isolation and characterisation of fungi growing on volatile aromatic hydrocarbons as their sole carbon and energy source. *Mycol Res* 105:477–484
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Santos EO, Centeno da Rosa CF, dos Passos CT, Sanzo AVL, de Medeiros Burkert JF, Kalil SJ et al (2008) Pre-screening of filamentous fungi isolated from a contaminated site in Southern Brazil for bioaugmentation purposes. *Afr J Biotechnol* 7:1314–1317
- Sanyal O, Shinde VL, Meena RM, Damare S, Shenoy BD (2016) The ITS-based phylogeny of fungi associated with tarballs. *Mar Pollut Bull*. <https://doi.org/10.1016/j.marpolbul.2016.09.052>
- Silva DP, Cavalcanti DD, de Melo EV (2015) Bio-removal of diesel oil through a microbial consortium isolated from a polluted environment. *Int Biodeterior Biodegrad* 97:85–89
- Simister RL, Poutasse CM, Thurston AM, Reeve JL, Baker MC, White HK (2015) Degradation of oil by fungi isolated from Gulf of Mexico beaches. *Mar Pollut Bull* 100:327–333
- Singh H (2006) *Mycoremediation: fungal bioremediation*. Wiley-Interscience, New York, p 592
- Tamura K, Nei M, Kumar S (2004) Prospects for inferring very large phylogenies by using the neighbor-joining method. *Proc Natl Acad Sci USA* 101:11030–11035
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28:2731–2739
- Thomas SA (2000) Mushrooms: higher macrofungi to clean up the environment. *Battelle Environmental Issues*, Fall 2000
- Toledo FL, Calvo C, Rodelas B, Lopez JG (2006) Selection and identification of bacteria isolated from waste crude oil with polycyclic aromatic hydrocarbons removal capacities. *Syst Appl Microbiol* 29:244–252
- Wang X, Wang X, Liu M et al (2015) Adsorption–synergic biodegradation of diesel oil in synthetic seawater by acclimated strains immobilized on multifunctional materials. *Mar Pollut Bull* 92:195–200
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis M, Gelfand D, Sninsky J, White T (eds) *PCR protocols: a guide to methods and applications*. Academic Press, New York, pp 315–322
- Yamada K, Mukumoto H, Katsuyama Y, Tani Y (2002) Degradation of long-chain alkanes by a polyethylene-degrading fungus, *Penicillium simplicissimum* YK. *Enzyme Microb Technol* 30:828–831