Published in final edited form as:

Top Curr Chem. 2015 ; 356: 123–164. doi:10.1007/128_2013_499.

The Photosynthesis and Photo-Stability of Nucleic Acids in Prebiotic Extraterrestrial Environments

Scott A. Sandford^{1,*}, Partha P. Bera^{1,2}, Timothy J. Lee³, Christopher K. Materese¹, and Michel Nuevo^{1,4}

¹NASA Ames Research Center, Space Science and Astrobiology Division, MS 245-6, Moffett Field, CA 94035, USA

²Bay Area Environmental Research Institute, 596 1st St. W, Sonoma, CA 95476, USA

³NASA Ames Research Center, Space Science and Astrobiology Division, MS 245-1, Moffett Field, CA 94035, USA

⁴SETI Institute, 189 N. Bernardo Ave., Suite 100, Mountain View, CA 94043, USA

Abstract

Laboratory experiments have shown that the UV photo-irradiation of low-temperature ices of astrophysical interest leads to the formation of organic molecules, including molecules important for biology such as amino acids, quinones, and amphiphiles. When pyrimidine is introduced in these ices, the products of irradiation include the nucleobases uracil, cytosine, and thymine, the informational sub-units of DNA and RNA, as well as some of their isomers. The formation of these compounds, which has been studied both experimentally and theoretically, requires a succession of additions of OH, NH₂, and CH₃ groups to pyrimidine. Results show that H₂O ice plays key roles in the formation of the nucleobases, as an oxidant, as a matrix in which reactions can take place, and as a catalyst that assists proton abstraction from intermediate compounds. As H₂O is also the most abundant icy component in most cold astrophysical environments, it probably plays the same roles in space for the formation of biologically relevant compounds. Results also show that although the formation of uracil and cytosine from pyrimidine in ices is fairly straightforward, the formation of thymine is not. This is mostly due to the fact that methylation is a limiting step for its formation, particularly in H₂O-rich ices, where methylation must compete with oxidation. The relative inefficiency of the abiotic formation of thymine to that of uracil and cytosine, coupled with the fact that thymine has not been detected in meteorites are not inconsistent with the RNA world hypothesis. Indeed, a lack of abiotically produced thymine delivered to the early Earth may have forced the choice for an RNA world, in which only uracil and cytosine are needed, but not thymine.

Keywords

Nucleobases; Ice Irradiation; UV Irradiation; Astrochemistry; Extraterrestrial Abiotic Nucleobase Synthesis

* Corresponding author: Scott A. Sandford, Scott.A.Sandford@nasa.gov, Phone: (+1) 650 604 6849, Fax: (+1) 650 604 6779.

1. Introduction

The processes by which life on Earth originated are not well understood and have been the subject of considerable speculation. While numerous pathways for the emergence of life have been suggested, it was presumably preceded by some sort of prebiotic chemical evolution that set the stage. In this chapter we will concentrate on only one aspect of this potential prebiotic chemistry, namely the synthesis of nucleobases in extraterrestrial environments.

1.1. Prebiotic Synthesis of the Building Blocks of Life

It is generally assumed that before life emerged on the primitive Earth, there must have been a period of time during which its basic 'building blocks' were synthesized and confined in an environment where they could be protected and interact with each other. There is a priori no requirement for all of the building blocks to be synthesized in the same environment, so it cannot be excluded that the individual components in the mixture of materials that lead to the emergence of life could have been formed in different locations and at different times. Indeed, individual classes of components, such as amino acids, could have been made and provided from multiple prebiotic environments. Presumably newly forming life had no concern about the different origins of these compounds and instead used whatever was available in the surrounding environment.

1.1.1. Abiotic Chemistry and the Origin of Life—The chemical steps that progressively led to life are still unknown, but were probably driven by the inventory of molecules that were present on the primitive Earth and/or the astrophysical environment where they took place. All of these ingredients, together with the principles of thermodynamics, may have played an important role in the processes changing non-living molecules into living entities.

The origin of the building blocks of life has been extensively debated since the 1950s, at which time Miller produced small quantities of amino acids and other organic molecules of prebiotic interest in his famous experiment, from the spark-induced chemistry of a gas mixture that was assumed to simulate the primitive atmosphere of the Earth (Miller 1953). However, more recent studies suggest that this was probably not the most relevant model because the actual primitive atmosphere was not as reducing as the gas mixture used by Miller (Levine et al. 1982). The discovery in the 1960s and 1970s of a rich variety of organic molecules such as amino acids, carboxylic acids, hydrocarbons, sugar-related compounds, as well as puric and pyrimidic bases in carbonaceous meteorites like Murchison (e.g., Hayes 1967; Kvenvolden et al. 1970; Mullie 1987; Cronin and Chang 1993; Irvine 1998; Cooper et al. 2001) supported an extraterrestrial delivery of the building blocks of life rather than an indigenous production of these molecules on the primitive Earth. In particular, the amino acids and nucleobases found in the Murchison meteorite were shown to be of extraterrestrial origin, as they display enrichments in ^{13}C relative to standard terrestrial values (Pizzarello and Huang 2003; Martins et al. 2007, 2008). This scenario of an extraterrestrial delivery of the building blocks of life to the primitive Earth via meteorites, originating from comets and asteroids, is strengthened by the fact that meteoritic bombardment was significantly heavier

during the first millions of hundreds of years in the Earth's history (Oró 1961; Delsemme 1998; Prinn and Fegley 1987; Chyba 1990; Chyba and Sagan 1992).

A few chemical mechanisms were proposed for the formation of prebiotic molecules under abiotic conditions. For example, amino acids are usually thought to have formed via the Strecker synthesis in the meteorites' parent bodies, during phases of aqueous alteration, or in a primitive ocean. This synthesis involves an aldehyde that reacts with hydrogen cyanide (HCN) and ammonia (NH₃) to form an aminonitrile that can be hydrolyzed into the corresponding amino acid (Strecker 1850; Taillades et al. 1998). Other chemical pathways to the formation of amino acids involve the formation of N-carboylamino acids as intermediate species via the Bücherer-Bergs reaction from an aldehyde, HCN, NH₃, and CO₂, also leading to amino acids via subsequent hydrolysis (Bücherer and Fischbech 1934; Taillades et al. 1998). The polymerization and/or oligomerization of HCN in H₂O is another potential pathway to the formation of both amino acids and nucleobases, as HCN is abundant in comets (Schloerb et al. 1986; Ip et al. 1990; Jones et al. 2006; Biver et al. 2007; DiSanti et al. 2007; Lis et al. 2008; Jehin et al. 2009; Gibb et al. 2012) and can also be found in interstellar ices (Loren and Wooten 1978; Zuckerman and Dyck 1986; McMullin et al. 1994; Jørgensen et al. 2004; Thi et al. 2004; Decin et al. 2010). However, while this chemical route has been extensively studied both in the laboratory (Matthews and Moser 1967; Toupance et al. 1970; Matthews 1979; Ferris 1979; Ferris and Hagan 1984) and more recently from a theoretical point of view (Roy et al. 2007), it has also been strongly debated. A similar mechanism involving cyanoacetylene (HC₃N) reacting with either the cyanide anion (OCN⁻) or water in the presence of urea (NH₂CONH₂) may also lead to the formation of the nucleobases cytosine and uracil (Ferris et al. 1968), which are of great interest to this chapter. Finally, sugars, in particular glucose, ribose, and deoxiribose, as well as their derivatives, could be formed via the formose reaction, i.e., a polymerization of formaldehyde (H₂CO) (Butlerow 1961; Müller et al. 1990; Weber 1998; Jalbout et al. 2007, 2008).

The final step from a collection of non-living molecules to those capable of self-replication and evolution is probably the least understood of all. Indeed, even if the primitive Earth contained an abundance of all the molecules required for life, constituting the so-called 'primitive soup' (Oró 1961), it was also probably supplied with an abundance of compounds that did not have the right properties for self-replication and evolution that may have inhibited the emergence of life. The reactions that led to self-replication and evolution are believed to have occurred in a solvent, most probably H₂O, to allow molecules to meet and react under favorable thermodynamic conditions (Ben-Naim 1980). In addition, H₂O probably played an important role in the formation of prebiotic membranes, as shown by some experiments in which organic molecules produced from the UV irradiation of simple ices spontaneously lead to the formation of vesicles, a phenomenon also observed with organic extracts of the Murchison meteorite (Dworkin et al. 2001). Membranes provide considerable advantages for chemical evolution, as they can protect molecules inside the vesicles from external lethal radiation, and because enclosed molecules can evolve under specific chemical conditions (solvent, temperature, pH) that can be different from the outside medium. Reactions inside a membrane-enclosed vesicle can occur under conditions of non-thermodynamic equilibrium, which, when coupled to prebiotic reactions and exchange of entropy with the external medium, can lead to complex reactions including

polymerization of amino acids and sugars. Finally, exothermic reactions between organic compounds, in particular polymerizations, may also have occurred on the surface of minerals (Nissenbaum 1976; Siegel and Siegel 1981; Maurette 1998; Schulte et al. 2006; Vázquez-Mayagoitia et al. 2011), which can play the role of a third body capable of absorbing the excess reaction energy and preventing the dissociation of newly formed products. Clays, in particular montmorillonite, is often cited as one of the minerals likely to have played such a role because of its chemical and physical properties (Ertem 2004; Arribas et al. 2005).

1.1.2. Potential Environments for Prebiotic Synthesis—While numerous environments for prebiotic chemistry have been suggested, most fall into one of two main categories, namely, extraterrestrial environments and environments found on the early Earth. We briefly comment on both of these environments below, though the bulk of the discussion in this chapter will be devoted to prebiotic chemistry that occurred prior to incorporation of the material into the Earth. Within this category, we will comment on the abiotic synthesis of a number of classes of compounds of astrobiological interest, but will specifically concentrate on the prebiotic extraterrestrial synthesis of the nucleobases.

1.1.2.1. Extraterrestrial Environments: ISM Dense Clouds, Protosolar Nebula, Comets, and Asteroids: Prior to the middle of the last century it was generally assumed that the conditions found in space were hostile to molecules, that chemistry could not proceed efficiently in space, and that most of the material in space would exist as either atoms or amorphous dust grains. The advent of radio and infrared telescopes, detectors, and spectrometers ultimately showed this to be untrue. We now know that much of the material in circumstellar and interstellar space is in molecular form, and that a variety of chemical processes occur as these materials cycle through a range of astrophysical environments (Figure 1). In the paragraphs below we describe several environments in which chemistry is observed to occur in space.

Many stars, late in their life cycles, undergo phases in which they lose significant portions of their mass through either gradual (stellar winds) or explosive (nova and supernova) processes. Where the outflow conditions allow, these materials can form molecules and dust grains. Depending on the local C/O ratios in the ejectae, the main products can be minerals (O-rich ejecta) or carbonaceous molecules and grains (C-rich ejecta) (Tielens and Allamandola 1987; Dorschner and Henning 1995).

Much of the material ejected from stars is subsequently destroyed or modified in the diffuse interstellar medium (ISM) where it is subjected to shock waves from supernova, sputtering, photo-destruction, etc. However, there are several lines of evidence that some of these materials survive transit through the diffuse ISM. Indeed, astronomical observations clearly indicate that the diffuse ISM contains silicate grains and solid organic materials that contain both aromatic and aliphatic components (Tielens and Allamandola 1987; Sandford et al. 1991; d'Hendecourt et al. 1996; Pendleton and Allamandola 2002). Most individual molecules are unable to survive in the gas phase in the diffuse ISM, with the notable exception of polycyclic aromatic hydrocarbons (PAHs) and related species. These molecules are seen in the outflows of dying stars and are sufficiently robust to resist complete

photolytic destruction in the diffuse ISM. As a result, they are the dominant form of molecular carbon found in many astrophysical environments (Allamandola et al. 1989; Puget and Léger 1989; Roelfsema et al. 1996; Galiano et al. 2008).

Perhaps the most compelling and interesting proof that at least some material survives the trip from stellar synthesis through the diffuse ISM is the existence of presolar grains in meteorites. Presolar grains having a number of different compositions (aluminum oxides, silicates, SiC, graphite, etc.) have been identified and their circumstellar formation is demonstrated by the presence of non-solar isotopic ratios that indicate formation in a variety of nucleosynthetic environments.

However, from the perspective of prebiotic chemistry and astrobiology, the most interesting type of extraterrestrial environment is that of dense interstellar molecular clouds. Such clouds contain enough material to be optically thick so that they screen out much of the stellar radiation that destroys most molecules in the diffuse ISM. The large optical depths of these clouds allow their interiors to cool to temperatures as low as 10–15 K. At these temperatures, most gas phase species other than H, H₂, He, and Ne will condense out onto dust grains in the form of ice mantles (Figure 2). Little ‘normal’ chemistry of the sorts we are familiar with on Earth can take place under such conditions, but chemical reactions occur nonetheless.

Some chemistry occurs via gas-grain reactions in which individual gas-phase atoms like H, C, N, and O collide with grains, react with resident surface species on the grains and form new compounds (d’Hendecourt et al. 1985). None of the species made in this manner are particularly complex. In dense cloud environments where $H > H_2$, these reactions tend to create simply hydrides like CH₄, CH₃OH, NH₃, and H₂O, and this process is thought to be the principle source of interstellar H₂ via reactions between two H atoms on the surface of cold dust grains. When $H < H_2$, reactions involving heteroatoms become important and species like N₂ and O₂ predominate.

More complex species can be created when ionizing radiation is present (Figure 2). The large optical depth of dense clouds screens out most stellar radiation, but cosmic rays and the energetic photons they create when colliding with interstellar matter produce some ionization even in the densest clouds (Shen et al. 2004). Newly forming stars within dense interstellar clouds can also irradiate nearby materials. Ionizing radiation produces ions and radicals that are able to react even at the low temperatures common in these clouds. Ions created in the gas phase can react in an extensive chain of ion-molecule reactions that can lead to numerous gas-phase species (Millar et al. 1995; Charnley 1997). To date more than 175 specific molecules have been identified in the gas phase in such clouds (Müller et al. 2013), although most are relatively small (< 7–8 atoms). However, the majority of the material other than H and He in dense clouds is trapped in the dust grains and their icy mantles, and it is in these ices that ionizing radiation likely produces the most complex molecular species (Greenberg 1984; d’Hendecourt et al. 1985; Bernstein et al. 1995; Dworkin et al. 2004). Such a radiation process will be the subject of much of the discussion in this chapter, particularly in the context of the abiotic formation of the nucleobases.

Because of their biological significance, astronomers have sought to confirm the presence of small *N*-heterocycles, including pyrimidine, purine, and their derivatives, in the gas phase of the ISM. To date, none of these compounds have been detected (Simon and Simon 1973; Kuan et al. 2003, 2004; Charnley et al. 2005; Brünken et al. 2006). Nonetheless, PAHs are known to be ubiquitous in galactic and extragalactic interstellar/circumstellar environments, and polycyclic aromatic nitrogen heterocycles (PANHs) are expected to be present as well (Allamandola et al. 1989; Puget and Léger 1989; Roelfsema et al. 1996; Galliano et al. 2008). It is therefore reasonable to assume that *N*-heterocycles, including pyrimidine-based species, may be present in space where they can condense on the surfaces of cold, icy grains, such as those found in dense molecular clouds (Sandford et al. 2004; Bernstein et al. 2005). These materials are therefore expected to participate in the types of ice irradiation process mentioned above.

1.1.2.2. Delivery of Extraterrestrial Organics to the Early Earth: Of course, the creation of organics in extraterrestrial environments is moot if these materials are unable to survive incorporation into planetary environments. Organics incorporated into the early Earth during its initial molten accretion were presumably destroyed – arriving organics would do little beyond contributing part of the inventory of important elements like C, N, O, and H. The actual delivery and survival of astrobiologically important compounds would have had to wait for the surface of the Earth to become a more hospitable place. It is therefore likely that the majority of complex organics delivered intact to the surface of the early Earth were first incorporated into small bodies like comets and asteroids and only delivered to the Earth at a later stage (Chyba et al. 1990).

The efficiency of successful delivery of extraterrestrial organics to the surfaces of planets depends critically on the nature of the body that delivers them (Chyba and Sagan 1992). While comets and asteroids can contain large amounts of organics, as whole bodies they are not a very efficient way to deliver organics intact to planetary surfaces. The enormous energy releases that occur during hypervelocity impact of such large objects is likely to destroy or modify much of their organic molecules, not to mention mediating the destruction, alteration, and creation of compounds on the planet. However, over the course of their orbital evolution, bodies like asteroids can shed material in the form of smaller meteoroids or dust particles generated by impacts with other asteroidal bodies. Comets can also shed rocks and dust through collisions as well as lose dust during outgassing that occurs when cometary volatiles sublime during warming.

These smaller meteoroids and cosmic dust particles can deliver organics much more efficiently than their cometary and asteroidal parent bodies. When smaller bodies like meteorites and cosmic dust collide with the Earth, they are slowed to terminal velocity high in the atmosphere and can deliver organics to the surface of a planet with far less degradation (Baldwin and Sheaffer 1971, Fraundorf 1980). For example, while the exterior of a meteoroid suffers serious heating and ablation during atmospheric entry, the interiors of meteorites experience little or no changes in temperatures during the entire infall process because ablation removes their surface faster than heat can thermally conduct into the interior of the meteorite. For a typical meteorite that makes it to the surface of the Earth, atmospheric entry only affects the outer few millimeters of the surviving material.

Meteorites typically land on the surface of the Earth with their interior temperatures largely unchanged from what they were in space and can therefore preserve and deliver their initial organic contents (e.g., Hayes 1967; Kvenvolden et al. 1970; Mullie 1987; Cronin and Chang 1993; Irvine 1998; Cooper et al. 2001).

1.1.2.3. The Early, Prebiotic Earth: Since the focus of this chapter is on the possible extraterrestrial abiotic formation of nucleobases, we will not go into great detail regarding the possible synthesis of these compounds in terrestrial environments. Nonetheless, the origin of life would almost certainly have occurred in an environment in which compounds of both extraterrestrial and terrestrial origin could have been present. While the issue of terrestrial synthesis is not central to the subject of this chapter, the possible importance of extraterrestrial delivery of nucleobases (as well as many other molecules of biological importance, for that matter) should be assessed in context with terrestrial sources. There are a number of excellent papers that address prebiotic chemistry on the early Earth, including extensive discussions of the nucleobases, and the reader is invited to examine them for informative reviews (e.g., Deamer and Fleischaker 1994; Lazcano and Miller 1996; Orgel 2004; Miller and Cleaves 2006; and references therein).

We note that we also do not concern ourselves here with the formation of RNA and DNA in their macromolecular forms. While it is the ultimate formation of RNA and DNA that motivates much of the interest in the abiotic formation of nucleobases, it is likely that this advance in chemical complexity occurred in a planetary environment, which falls outside the purview of our discussion here. Indeed, it has even been suggested that the formation of RNA and DNA need not have required the preexistence of nucleobases (Powner et al. 2010).

1.2. Abiotic Extraterrestrial Chemistry and the Origin of Life

There are several lines of evidence that suggest abiotic extraterrestrial chemistry may have resulted in the delivery of numerous compounds of biological importance to the early Earth. Some of this evidence comes from astronomical observations of dense clouds and forming planetary systems, but the main lines of evidence come from laboratory simulation experiments and the detection of biologically important species in the organic materials found in primitive meteorites.

1.2.1. Evidence from Astronomical Observations and Laboratory Simulations

—Astronomical spectral observations of the ISM at infrared and longer wavelengths have demonstrated that the molecular composition of interstellar dense clouds is dominated by simple species. For example, absorption spectra taken along lines of sight towards embedded and background stars show that the ices in most dense clouds are dominated by H₂O along with varying amounts of other simple molecules like CH₃OH, CO, CO₂, CH₄, NH₃, HCN, etc. (d'Hendecourt et al. 1996; Whittet et al. 1996; Lacy et al. 1998; Ehrenfreund and Charnley 2000; Gibb et al. 2000, 2004; Dartois 2005). These ices are exposed to doses of ionizing radiation in the form of cosmic rays and high-energy photons that can create ions and radicals in the ices that can ultimately react to form more complex molecules (Figure 2). Evidence that interstellar ices are truly processed by ionizing radiation in this manner comes from the observation of a broad absorption feature near 4.62 μm (2165 cm^{-1}) in the infrared

spectra of the ices in many clouds (Lacy et al. 1984, 1998; d'Hendecourt et al. 1996; Whittet et al. 1996; Gibb et al. 2000; Thi et al. 2006). This feature is reliably reproduced in virtually all laboratory experiments involving the irradiation of ices containing sources of O, C, N, and H and is due to the $C\equiv N$ stretching mode in what is most likely OCN^- (Schutte and Greenberg 1997; Demyk et al. 1998; van Broekhuizen et al. 2004).

Numerous laboratory studies have demonstrated that irradiation of astrophysically relevant ices dominated by simple molecules like those listed above results in the production of numerous ions and radicals within the ices that can then combine as the ice warms to form a wide variety of new molecular species that are considerably more complex than those seen in the gas phase in dense clouds (Dworkin et al. 2004). Many of these species are of astrobiological interest and include amino acids (Bernstein et al. 2002a; Muñoz Caro et al. 2002; Nuevo et al. 2007, 2008), amphiphiles (Dworkin et al. 2001), as well as urea, glycerol, glyceric acid, hydantoin (Nuevo et al. 2010; de Marcellus et al. 2011), and, if aromatic species are present in the ices, quinones, ethers (Bernstein et al. 1999b, 2001, 2002b, 2003; Ashbourn et al. 2007), and nucleobases (Nuevo et al. 2009, 2012; Materese et al. 2013).

This chemistry appears to be very 'robust' in the sense that the types of organic materials that are produced are relatively insensitive to many of the experimental parameters of the irradiation. For example, the temperature of the ice during irradiation is relatively unimportant; all that is required is that the ice be cold enough that the starting materials remain condensed during irradiation. Since H_2O is the dominant species in many astrophysical ices, this means that ice irradiation processes yield similar products over the entire 10–150 K temperature range (Bernstein et al. 1995). This processing is similarly insensitive to the source of the ionization: identical ices will yield very similar products independent of whether they are irradiated by energetic protons or UV photons (Gerakines et al. 2000, 2001, 2004; Bernstein et al. 2003). Indeed, the basic suite of products is even relatively insensitive to the composition of the ices themselves provided they contain fragmentable sources of C, H, O, and N. For instance, ices containing widely different relative abundances of the same starting molecules will produce much the same set of products, although their relative and total abundances may vary. Similarly, laboratory experiments performed with ices in which the initial sources of C, H, O, and N are different will still often yield similar products. For example, irradiation of ices yields amino acids independent of whether the initial carrier of C in the ices is CH_3OH , CH_4 , or CO_2 (Bernstein et al. 2002a; Muñoz Caro et al. 2002; Nuevo et al. 2007, 2008), and whether the initial carrier of N is NH_3 or HCN (Elsila et al. 2007).

Of particular interest is the radiation processing of ices that contain aromatic molecules like polycyclic aromatic hydrocarbons (PAHs) and related aromatic species that contain heterocycles. In dense molecular clouds, they should be largely condensed out onto grains (Sandford et al. 2004; Bernstein et al. 2005). As previously stated, these molecules are relatively robust against photo-destruction in the gas phase in the ISM, but they can also participate in a rich chemistry when irradiated in mixed molecular ices. One of the chief outcomes of the irradiation of PAHs in ices is the addition of side groups to the aromatic rings of the PAHs. Since most interstellar ices are dominated by H_2O , this largely results in O and H addition reactions that make aromatic ketones, alcohols, and ethers (Bernstein et al.

1999b, 2001, 2002b, 2003; Ashbourn et al. 2007), and hydrogenated PAHs (H_n -PAHs; Sandford et al. 2013), but additional side groups like $-NH_2$, $-CH_3$, $-OCH_3$, $-C\equiv N$, etc. can be added if other molecules are present in the starting ice (Bernstein et al. 1999b, 2002b). As we will see, this process is likely to play an important role in the abiotic production of nucleobases in astrophysical environments.

1.2.2. Evidence from Meteorites—Meteorites that fall on the Earth are divided into a wide variety of classes having different compositions and evolutionary histories (Dodd 1981; Lauretta et al. 2006). Among these classes, carbonaceous chondrites are of the most interest for studies of extraterrestrial organics. These meteorites typically contain a few weight percent carbon (Pizzarello et al. 2006). Much of the carbon is in the form of insoluble organic matter (IOM), a material that consists of small aromatic moieties interlinked by a complex variety of bridging units (Cody et al. 2002; Alexander et al. 2007). Meteoritic IOM has a number of interesting properties, for example, it is the carrier of components having large D/H and $^{15}N/^{14}N$ isotopic anomalies (Kerridge et al. 1987; Alexander et al. 2007). However, other than being a major carrier of carbon in meteorites, it is of limited interest for astrobiology because it is chemically intractable.

Of more interest for astrobiology are the soluble carbonaceous components of these meteorites. A wide variety of soluble compounds are found in carbonaceous meteorites, including aliphatic and aromatic hydrocarbons, carboxylic and dicarboxylic acids, amines, and amides (Pizzarello et al. 2006). These individual classes of compounds are typically present at concentration falling in the range from a few ppm to a few hundred ppm.

A number of compounds are of clear astrobiological interest, including amino acids (Kvenvolden et al. 1970; Cronin and Pizzarello 1997; Sephton et al. 2002; Glavin et al. 2006; Martins et al. 2007; Burton et al. 2013), amphiphiles (Dworkin et al. 2001), and *N*-heterocycles (Hayatsu 1964; Folsome et al. 1971, 1973; Hayatsu et al. 1975; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981; Martins et al. 2008; Callahan et al. 2011), which are the building blocks of proteins, membranes, and nucleic acids, respectively. There is ample evidence that many of these materials are truly extraterrestrial, and not terrestrial contamination. For example, the extraterrestrial nature of the many amino acids detected in meteorites is supported by the presence of (i) amino acids not used by biological systems on Earth, (ii) amino acids that are largely (but not entirely) racemic, and (iii) amino acids having non-terrestrial isotopic ratios (Engel and Macko 1997; Cronin and Pizzarello 1997, 1999; Pizzarello et al. 2003; Pizzarello and Huang 2003; Martins et al. 2007; Glavin et al. 2012; Callahan et al. 2013).

Of particular interest to the discussion in this chapter are *N*-heterocycles. To date, a series of pyrimidines, quinolines, isoquinolines, benzoquinolines, and several of their methyl isomers have been identified in meteorites (Pizzarello et al. 2006, and references therein). Purine- and pyrimidine-based compounds have so far been detected in the carbonaceous chondrites Orgueil, Murchison, Murray, and Lonewolf Nunataks 94102 (Hayatsu 1964; Folsome et al. 1971, 1973; Hayatsu et al. 1975; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981; Callahan et al. 2011). The extraterrestrial origin of these compounds has been

confirmed in many cases by isotopic analysis (Martins et al. 2008), lending further credence to their extraterrestrial formation via at least one astrophysical, non-biological process.

Uracil is, to date, the only pyrimidine-based nucleobase that has been unequivocally identified in meteorites (Stoks and Schwartz 1979; Martins et al. 2008). The extraterrestrial nature of uracil in the Murchison meteorite is supported by $\delta^{13}\text{C}$ measurements by Martins et al. (2008), although it has been suggested that coeluting compounds could be confusing the issue (Callahan et al. 2011). The lack of detection of cytosine in meteorites might be explained by its conversion into uracil via hydrolysis when the organics are extracted from the meteorite, as cytosine was shown to be easily hydrolyzed in the laboratory (Shapiro 1999; Nelson et al. 2001). Thus, the presence of cytosine in meteorites cannot currently be ruled out.

So far, studies of the Allende, Orgueil, Murchison, and Murray have only managed to establish upper limits on the presence of thymine in these meteorites, but these upper limits are an order of magnitude *lower* than those for uracil (Stoks and Schwartz 1979). The non-detection of thymine cannot be explained by the same conversion process that may be limiting the detection of meteoritic cytosine. This suggests that there must be another reason for its lower abundance in meteorites relative to uracil. We will return to this point later in this chapter.

Several purines have been identified in meteorites, including xanthine, hypoxanthine, and the nucleobases guanine and adenine (Hayatsu 1964; Folsome et al. 1971, 1973; Hayatsu et al. 1975; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981; Callahan et al. 2010, 2011). The presence of pre-terrestrial pyrimidine and purine-based compounds in meteorites clearly indicates that some extraterrestrial abiotic processes are capable of making at least some of the nucleobases in space. In the sections that follow we will discuss some of the extraterrestrial environment in which these processes may occur.

2. The Synthesis/Stability of Nucleobases in Extraterrestrial Environments

The primary environments of interest for chemistry in the interstellar medium are dominated by materials in the gas phase and, in dense clouds and protostellar disks, in solids and ices. Chemistry in the liquid phase is largely restricted to planetary environments and, for brief periods in the early history of the Solar System, for some asteroids, and is therefore not considered in any detail here. In the sections that follow we will thus concentrate on gas- and solid-phase photochemistry.

2.1. Synthesis/Stability in the Gas Phase

A few mechanisms for the formation of both pyrimidine- and purine-based nucleobases have been proposed and tested experimentally and theoretically. Among them, the most popular is probably the formation of purine-based compounds from the polymerization or oligomerization of hydrogen cyanide (HCN) in the gas phase. HCN is a common component of astrophysical ices, in particular it is found in comets (Schloerb et al. 1986; Ip et al. 1990; Jones et al. 2006; Biver et al. 2007; DiSanti et al. 2007; Lis et al. 2008; Jehin et al. 2009; Gibb et al. 2012), but also in interstellar and circumstellar sources (Loren and Wooten 1978;

Zuckerman and Dyck 1986; McMullin et al. 1994; Jørgensen et al. 2004; Thi et al. 2004; Decin et al. 2010). In the 1960s, Matthews noticed that adenine, one of the purine-based biological nucleobases found in DNA and RNA, was a pentamer of HCN, i.e., that it could be obtained from the combination of five molecules of HCN. This mechanism was tested experimentally by his team and independently by Ferris' and Toupance's teams, but none of them could find a pathway that could lead to an efficient production yield of adenine (Matthews and Moser 1967; Toupance et al. 1970; Matthews 1979; Ferris 1979; Ferris and Hagan 1984). However, the probability for five molecules of HCN to encounter one another in low-density astrophysical gas phase environments is small, so such a process would not be very efficient in space. The formation of adenine from the pentamerization of HCN in the gas phase has also been studied theoretically via density functional theory (DFT) and singles and doubles coupled-cluster with perturbational triples [CCSD(T)] computations, which showed that such a process is energetically favored (Roy et al. 2007; Boulanger et al. 2013), and thus that adenine could be formed abiotically. Similarly, *ab initio* CCSD(T) quantum chemical computations indicate that pyrimidine could be formed in the gas phase via an ion-molecule mechanism through a structural isomer of vinyl cyanide ion (Bera et al. 2009).

Another mechanism, involving the reaction between cyanoacetylene (HC_3N) and the cyanate anion (OCN^-), could lead to the formation of the pyrimidine-based nucleobases uracil and cytosine (Raulin 1991). HC_3N is abundant in comets (Lis et al. 1997; Biver et al. 2011; Dello Russo et al. 2011), circumstellar disks (Jewell and Snyder 1982, 1984; Huang et al. 2005; Pardo et al. 2005), as well as in the atmosphere of Titan (Kunde et al. 1981; Coustenis et al. 1991; Bénilan et al. 1996; Khanna 2005), the largest satellite of Saturn, known to house a very complex chemistry that may be in some points similar to what existed on the primitive Earth several billions of years ago (Chang et al. 1979; Capone et al. 1981; Gupta et al. 1981; Raulin et al. 1998; Vuitton et al. 2012). OCN^- has been observed in the solid phase (ice) in several astrophysical sources (Lacy et al. 1984, 1998; d'Hendecourt et al. 1996; Whittet et al. 1996; Gibb et al. 2000; Thi et al. 2006), and is often considered to be the proof that photo-induced chemical reactions do occur at the surface of icy cold grains, since it can be formed from the irradiation of H_2O , CO , CO_2 , and NH_3 (Schutte and Greenberg 1997; Demyk et al. 1998). However, to date, OCN^- has not yet been observed in the gas phase. Alternatively, HC_3N could also react with H_2O to form an intermediate species, cyanoacetaldehyde ($\text{N}\equiv\text{CCH}_2\text{CHO}$), which, by reacting with urea (NH_2CONH_2), could lead to the formation of uracil and cytosine (Raulin 1991; Horn et al. 2008).

In dense environments, such as molecular clouds or Titan's atmosphere, *N*-heterocycles such as pyridine and pyrimidine may be formed in the gas phase via copolymerization of HCN and acetylene (C_2H_2), believed to be one of the precursors of PAHs in space (Frenklach and Feigelson 1989; Ricca et al. 2001). Acetylene and ethylene could also react with protonated hydrogen cyanide (HCNH^+), considered to be a major ion in Titan's ionosphere due to its high concentration, that could lead to synthesis of cyclic molecules involving nitrogen, and possible precursors of nucleobases (Bera et al. 2012). HCNH^+ is also known to exist in the gas phase in interstellar clouds, where it plays a role in the conversion of HCN to HNC (McElroy et al. 2013). Once formed, *N*-heterocycles and PAHs could condense on small, cold grains in the ISM (Sandford et al. 2004; Bernstein et al. 2005) and be photo-processed together with other ices present, such as H_2O , CH_3OH , CO , CO_2 , CH_4 , and NH_3 .

(d'Hendecourt et al. 1996; Whittet et al. 1996; Lacy et al. 1998; Ehrenfreund and Charnley 2000; Gibb et al. 2000, 2004; Dartois 2005), to subsequently form complex molecules of prebiotic and biological interest such as quinones (Bernstein et al. 1999b, 2002b) and nucleobases (Nuevo et al. 2009, 2012; Materese et al. 2013).

Once formed in astrophysical environments, nucleobases, like other gas-phase and ice components, will be subjected to strong ionizing radiation. While radiation can affect the chemical composition of a medium by breaking bonds, creating ions and radicals, and drive the formation of new, more complex products after recombination, it can also destroy these same compounds. Because of this, studying the photo-stabilities of compounds of interest is also important. Peeters et al. (2005) studied the photo-stability of pyridine, pyrimidine, and triazine in an argon matrix under astrophysically relevant conditions. Their results show that these molecules, if isolated in the gas phase and subjected to UV radiation, would have half-lives of 2–20 yr in the optically thin diffuse ISM, and up to 2 Myr in the dense ISM, which is mostly opaque to UV photons but subjected to cosmic-ray radiation and their secondary electrons as well as UV photons produced from the interaction between cosmic rays and grains (Peeters et al. 2005). Because of higher UV fluxes, these numbers drop to 4–30 min in the Solar System. All these values are smaller than those expected for small PAHs irradiated in astrophysical environments (Allamandola et al. 1989; Hudgins and Sandford 1998a,b; Ruiterkamp et al. 2005), probably due to the smaller bond energies and the polarization of the C–N bonds in heterocyclic compounds. The stability of nucleobases under UV irradiation, and the processes of radical and ion formation on them were also studied both experimentally (Satzger et al. 2006; Fondren et al. 2007) and theoretically (Bera and Schaefer 2006).

2.2. Synthesis of Pyrimidine-Based Nucleobases in Extraterrestrial Ices

Laboratory and theoretical work has been done on the ice-phase photochemistry of pyrimidine leading to the synthesis of all three pyrimidine-based nucleobases. In the sections that follow, we summarize the results of this work separately for uracil, cytosine, and thymine (Figure 3). We use the discussion of uracil to provide a general overview of the approaches that have been used for this kind of work. The subsequent discussions about cytosine and thymine are largely restricted to details unique to these individual molecules.

2.2.1. Synthesis of Uracil—Uracil (2,4-dihydropyrimidine) is a doubly oxidized pyrimidine derivative. It is one of the five biologically relevant nucleobases, and is found only in RNA. The 'RNA world' hypothesis proposes that RNA served as the original genetic material and as biological catalysts (Gilbert, 1986; Joyce 1989, 2002). Under this paradigm, understanding the abiotic formation of uracil is thus of great importance.

As mentioned earlier, the detection of pyrimidine- and purine-based nucleobases in meteorites suggests that such molecules may be present on the surfaces of cold grains in astrophysical environments and mixed into ices. If this is so, then biologically relevant molecules such as nucleobases might be formed via photo-processes of pyrimidine in ices, the same way other organic compounds such as amino acids are probably formed in similar environments (Bernstein et al. 2002a; Muñoz Caro et al. 2002; Nuevo et al. 2007, 2008).

The formation of uracil via the photo-irradiation of pyrimidine mixed with simple ices requires an oxygen source. Among the catalog of compounds present in interstellar ices, H₂O is by far the most abundant (Ehrenfreund and Charnley 2000; Gibb et al. 2000, 2004; Dartois 2005), and can be a strong oxidant. Consequently, H₂O:pyrimidine ices subjected to UV photon irradiation are simple, relevant models to simulate the oxidation of pyrimidine under astrophysical conditions in the laboratory.

2.2.1.1. Laboratory Synthesis of Uracil: In these laboratory experiments, H₂O:pyrimidine mixtures prepared in the gas phase are deposited in a controlled fashion onto a substrate cooled to temperatures as low as 15–20 K inside a cryogenic vacuum chamber evacuated to a pressure of a few 10⁻⁸ mbar. During deposition, the growing ice films are simultaneously irradiated with UV photons emitted by a microwave-powered H₂ discharge lamp. Such a lamp provides UV photons mainly at 121.6 nm (Lyman α) and a continuum centered around 160 nm, with a flux of about 2×10¹⁵ photons cm⁻² s⁻¹ (Warnek 1962) that simulates the UV radiation field observed in many astrophysical environments (Mathis et al. 1983; Prasad and Tarafdar 1983; Shen et al. 2004).

Following the simultaneous deposition and radiation process, each ice sample is warmed to 220 K to allow the original ice components to sublime away, at which temperature the resulting refractory residue material is extracted from the vacuum chamber for further analysis with high-performance liquid chromatography (HPLC) and gas chromatography coupled with mass spectrometry (GC-MS).

Two of the major compounds successfully identified in these residues include 4(3*H*)-pyrimidone (or 4-hydroxypyrimidine), a singly oxidized pyrimidine derivative and precursor of uracil, and uracil itself (Figure 4; Nuevo et al. 2009). It should be noted that 2-hydroxypyrimidine, another singly oxidized variant of pyrimidine and a uracil precursor, is only detected in very small quantities in this series of experiments, indicating that oxygen addition to pyrimidine occurs in a regioselective manner (Nuevo et al. 2009, 2012). These experiments also show that UV photons can break the pyrimidine ring, leading to the formation of small aliphatic molecules such as urea and the amino acid glycine in measurable quantities (Nuevo et al. 2012). As previously observed in other simple ice mixtures, for example ices in which amino acids are formed (Bernstein et al. 2002a; Muñoz Caro et al. 2002; Nuevo et al. 2007, 2008), the list of products formed is relatively insensitive to the ratios of the starting H₂O and pyrimidine, though their relative abundances vary.

Since data regarding the composition of the residues are obtained *ex post facto*, it is difficult to know at which point during the irradiation and/or subsequent warm-up products, such as 4(3*H*)-pyrimidone, uracil, and their isomers, are actually formed. However, it is known that the formation of uracil does not occur as a result of exposure to the aqueous phase while awaiting analysis, since its presence is confirmed in samples that were kept free of exposure to liquid water (Nuevo et al. 2009). While this does not answer the question of when uracil is produced, it shows that such compounds can form in the absence of liquid water.

There are limited experimental data related to the reaction mechanisms for the formation of oxidized pyrimidines and their derivatives. Such issues can be addressed with theoretical computations and are discussed in the next section. Nevertheless, experiments reveal that the formation of these compounds is relatively insensitive to temperature and wavelength, which suggests a radical and/or ion chemistry driven by the photo-decomposition of H₂O. Given this hypothesis, the unstable radicals and/or ions engage in a chemistry of opportunity that is not bound by thermodynamic favorability.

2.2.1.2. Theory of Uracil Synthesis in the Condensed Phase: Detailed quantum mechanical theoretical computations have been conducted to understand the process of photo-oxidation of pyrimidine in a pure H₂O ice matrix (Bera et al. 2010). B3LYP DFT, which contains Becke's three-parameter exchange functional (B3) (Becke 1993), and the correlation functional of Lee, Yang, and Parr (LYP) (Lee et al. 1988), was used in conjunction with Pople's 6-31G(d,p) split valence basis set (Hehre et al. 1972). Structures were optimized to obtain the lowest energy minima on the potential energy surfaces of the reactants, intermediates, and products, and were confirmed by a subsequent harmonic frequency calculation. Energy differences were then obtained using second order Møller-Plesset (MP2) perturbation theory for closed shell species and second order Z-averaged perturbation theory (ZAPT2) (Lee et al. 1993, 1994) for the open shell species, in conjunction with Dunning's correlation consistent valence triple zeta basis set (cc-pVTZ) (Dunning 1989). The Q-Chem 3.1 and 3.2 suites of *ab initio* and DFT quantum mechanical codes were used for all computations (Shao et al. 2006).

H₂O possesses a large photo-dissociation cross-section in the UV wavelength range (Engel et al. 1992). Hydroxyl (OH) radicals are therefore readily produced in the ices by the following process:

It is therefore reasonable to assume that the chemistry in these ices is dominated by reactions of OH radicals with pyrimidine and its derivatives. Other radicals and ions can also be generated from the pyrimidine itself via the following processes:

where [Py_{/H}][•] designates a radical formed from the cleavage of a C–H bond on the pyrimidine ring. OH radicals can react with either neutral pyrimidine molecules or pyrimidine cations to form cations and radicals of oxidized pyrimidines.

An important step in these mechanisms is the loss of either a proton or a hydrogen atom from the intermediates to form the final products. Reactions involving a pyrimidine cation

and an OH radical (first oxidation step), are presented in Figure 5a. Oxidation products of pyrimidine can be further oxidized by OH radicals, and produce doubly oxidized products:

The above reactions are ion-molecule or radical-molecule type in nature, and are typically barrierless and fast. Neutral-neutral reactions between H₂O and pyrimidine can also occur but, because of their high activation barriers, should be several orders of magnitude slower than reactions involving radicals or cations, particularly in cold ices.

Computations show that an ionic mechanism in which a pyrimidine cation reacts with an OH radical is energetically favorable and can lead to the formation of intermediate species followed by the subsequent loss of a proton to form the final products (Bera et al. 2010). Among all possible products, 4-hydroxypyrimidine and its tautomer 4(3*H*)-pyrimidone are the most favored singly oxidized products, followed by 2-hydroxypyrimidine and its tautomer 2(1*H*)-pyrimidone (Figure 5b). This result is important since any of these singly oxidized compounds can be further oxidized, leading to the formation of uracil. Additionally, computations show that a purely gas-phase proton loss from the ionic intermediate species is unfavorable, as illustrated in the energy diagram (Figure 5b). The presence of several H₂O molecules in the ice matrix, acting as a proton acceptor, allows this reaction to become favorable towards the formation of the products. In other words, a condensed-phase environment is critically important for these mechanisms to be viable. Oxidation of neutral pyrimidine via a reaction with an OH radical is also shown to be energetically favorable and lead to the expected products, although this mechanism is not as energetically favorable as that of an ionic mechanism according to the *ab initio* MP2/ZAPT2 calculations.

As illustrated in Figure 6a, Bera et al. (2010) showed that oxidation of 4(3*H*)-pyrimidone (and its tautomer 4-hydroxypyrimidine) via an ionic mechanism follows a similar path as that described for the first oxidation step via an ionic mechanism. That is, the initial hydroxylation step is highly exothermic and should be barrierless, and the subsequent proton loss is energetically favorable in the presence of the surrounding H₂O molecules (i.e., a condensed-phase environment is again required for the deprotonation step). From the energetic order of the formed intermediate species and products (Figure 6b), it is evident that the formation of *uracil* is favored over that of other doubly oxidized products. Additionally, the reaction between an OH radical and a neutral 4-hydroxypyrimidine molecule, which requires a neutral-radical intermediate species, is also a viable route for the formation of uracil and other doubly oxidized pyrimidine derivatives.

An alternate route for the formation of uracil is via the oxidation of 2-hydroxypyrimidine or its tautomer 2(1*H*)-pyrimidone. The reaction schemes for an ionic mechanism reaction are shown in Figure 7a, and the associated energies are shown in Figure 7b. It is interesting to

note that 2,4-dihydropyrimidine, the dienol tautomer of uracil, is found to be the most stable doubly oxidized product. Again, the presence of H₂O molecules to abstract a proton from the intermediate is essential for the oxidation reaction to proceed via an ionic mechanism. A similar result was found for a neutral-radical oxidation mechanism, which showed that the formation of uracil is still favored over that of other products.

Therefore, theoretical computations are in agreement with laboratory studies of the UV photo-irradiation of H₂O:pyrimidine ice mixtures at low temperature (Nuevo et al. 2009), in which 4(3*H*)-pyrimidone was found to be the most abundant singly oxidized pyrimidine. The detection of only small quantities of 2-hydroxypyrimidine in laboratory samples favor a route for the formation of uracil in which 4(3*H*)-pyrimidone is its main precursor, though computations show that the oxidation of 2-hydroxypyrimidine towards the formation of uracil is also energetically favored in an H₂O ice matrix (Bera et al. 2010). Similarly, uracil was found to be the most abundant doubly oxidized pyrimidine in laboratory samples (Nuevo et al. 2009), as well as the most favorable product in the *ab initio* quantum calculations, regardless of its singly oxidized precursors – 4-hydroxypyrimidine, 2-hydroxypyrimidine, or their tautomers. Finally, both experimental and theoretical results show a clear regioselectivity for the oxidation of pyrimidine, favoring position 4 of the ring over position 2 for the first oxidation, and the formation of uracil as the most stable doubly oxidized product.

Perhaps the single most important conclusion from these quantum mechanical computations is that the oxidation of pyrimidine is not expected to be energetically favorable in the gas phase, whereas it is feasible in an ice matrix. The presence of several H₂O molecules surrounding the reactants in the condensed phase is absolutely necessary to assist the proton abstractions leading to the formation of the final products for both oxidation steps of pyrimidine and its derivatives. This important result indicates that the formation of uracil, as well as other pyrimidine derivatives, via the oxidation of pyrimidine under astrophysical conditions is likely viable only in icy mantles, such as those observed at the surface of cold grains in the ISM, and in the presence of ionizing radiation (Bera et al. 2010).

2.2.2. Synthesis of Cytosine—Cytosine (4-amino-2-hydroxypyrimidine) is the second biologically relevant pyrimidine-based nucleobase, which is found in both DNA and RNA. Like uracil, this compound is a doubly substituted pyrimidine derivative, but has an amino (NH₂) group replacing the OH group on the 4 position (Figure 3). Therefore, the abiotic production of cytosine from the irradiation of pyrimidine requires the addition of both one OH group to position 2 of the pyrimidine ring and one NH₂ group to position 4. OH and NH₂ radicals are nucleophilic species that are readily formed from the photo-dissociation of H₂O and NH₃, respectively, both in laboratory ices and cold astrophysical environments such as diffuse clouds, protostellar disks, circumstellar sources, and comets (Ehrenfreund and Charnley 2000; Gibb et al. 2000, 2004; Dartois 2005). Because of the low NH₂–H and HO–H dissociation energies of 4.60 eV (Mordaunt et al. 1996) and 5.10 eV (Woon 2002), respectively, both OH and NH₂ radicals are expected to react with pyrimidine in a similar way under these experimental conditions.

2.2.2.1. Laboratory Synthesis of Cytosine: The irradiation of pyrimidine mixed with H₂O and NH₃ ices in the laboratory is thus expected to lead to the formation of cytosine, uracil, and a suite of other oxidized and/or aminated pyrimidine derivatives. Experiments in which H₂O:NH₃:pyrimidine ice mixtures with different relative proportions are irradiated following the same protocol as for the formation of uracil show that cytosine can be formed efficiently under similar conditions, and with a comparable yield to uracil (Nuevo et al. 2012). As was observed for the oxidation of pyrimidine, the addition of amino groups is regioselective and favors the amination of position 4 of the ring over the other positions to form 4-aminopyrimidine (one of cytosine's precursors), though small quantities of other isomers such as 2-aminopyrimidine are also formed.

However, even if 4-aminopyrimidine is a logical precursor of cytosine, its formation in H₂O + NH₃ is in competition with the formation of 4(3*H*)-pyrimidone via oxidation of pyrimidine (Nuevo et al. 2009, 2012; Section 2.2.1.1). The amination of 4(3*H*)-pyrimidone leads to the formation of isocytosine (2-amino-4-hydroxypyrimidine), an isomer of cytosine that has been detected in most of the residues produced from the UV photo-irradiation of H₂O:NH₃:pyrimidine ice mixtures (Nuevo et al. 2012).

Finally, it is interesting to note that although both OH and NH₂ groups have similar nucleophilic properties and are expected to react with pyrimidine in a similar manner, these experiments indicate that the presence of H₂O in the starting ice mixtures enhances the addition of NH₂ groups, in particular to increase the formation yield of products such as 2-aminopyrimidine, whose formation in an H₂O-poor ice matrix is not favored for regioselective reasons (Nuevo et al. 2012).

2.2.2.2. Theory of Cytosine Synthesis in the Condensed Phase: Investigating reaction mechanisms in mixed ices such as H₂O+NH₃ is more complicated than in separate pure H₂O or pure NH₃ ices. In order to perform theoretical computations aimed at understanding such processes, one needs to identify the most probable sequence of reactions in these ices. Since OH and NH₂ radicals are known to be readily formed from the photo-degradation of H₂O and NH₃, respectively, it appears logical that the two main types of reactions that will affect pyrimidine in these systems are oxidation (i.e., hydroxyl group addition) and amination (i.e., amino group addition).

Following what is known about the oxidation of pyrimidine from both the experimental and theoretical studies of uracil formation, it is reasonable to assume that functional group additions on pyrimidine (neutral, cation, or radical) take place one after the other rather than simultaneously. Because OH and NH₂ radicals have a reasonably similar probability to react with pyrimidine, the most important factor concerning which groups is most likely to be added first will largely depend on the relative abundance of their respective parent molecules (H₂O and NH₃) in the ices. For most astrophysical ices and the studies described here, where H₂O is generally the dominant ice component, oxidation will be favored over amination.

When oxidation takes place first, it favors the formation of 4(3*H*)-pyrimidone over that of 2-hydroxypyrimidine, for regioselective reasons (Nuevo et al. 2009; Bera et al. 2010; Section 2.2.1). The subsequent amino group addition, if it occurs at position 2 of the pyrimidine

ring, will lead to the formation of isocytosine (2-amino-4-hydroxypyrimidine), an isomer of the nucleobase cytosine. Of course, 4(3*H*)-pyrimidone and its isomers can also undergo further oxidation, towards the formation of uracil and its isomers, as shown for pyrimidine in a pure H₂O ice (Nuevo et al. 2009; Bera et al. 2010; Section 2.2.1).

In the case where amination takes place first, the most favored product may be 4-aminopyrimidine, similar to the oxidation step, which may then be either aminated to form doubly aminated pyrimidine derivatives, or oxidized to form cytosine and its isomers. Similar to what happens for the oxidation of 4(3*H*)-pyrimidone, the oxidation of 4-aminopyrimidine may favor the 2 position on the ring, resulting in the formation of cytosine. The formation of other isomers will likely also be exothermic, although they may be less abundant than cytosine if the formation of cytosine follows that of uracil.

The detailed *ab initio* quantum chemical calculations describing these reactions for three types of mechanisms, namely, ionic, neutral-radical, and radical-radical, are currently in progress (Rice et al. 2013).

2.2.3. Synthesis of Thymine—Thymine (2,4-dihydroxy-5-methylpyrimidine) is the third and last biologically relevant pyrimidine-based nucleobase. Thymine is found only in DNA and thus is not necessary for the emergence of an RNA world (Gilbert 1986; Joyce et al. 1989, 2002). The formation of this tri-substituted pyrimidine derivative requires a double oxidation in the same positions on the ring as uracil, and the addition of a methyl (CH₃) group to position 5. Therefore, the abiotic production of thymine from pyrimidine requires both a source of oxygen and a source of methyl groups.

Since H₂O is by far the most abundant component of interstellar ices, its presence in any starting ice mixtures to be studied is essential. On the other hand, possible sources of methyl groups in interstellar ices could include both methanol and methane, as both are widely observed in cold astrophysical environments, although methanol is usually observed with abundances that are an order of magnitude larger than those of methane (Ehrenfreund and Charnley 2000; Gibb et al. 2000, 2004; Dartois 2005). Thus, H₂O:CH₃OH:pyrimidine and H₂O:CH₄:pyrimidine mixtures appear to be the simplest relevant ice analogs to be studied both experimentally and theoretically for the formation of thymine through photo-irradiation. Experiments in which H₂O-free ice mixtures (CH₃OH:pyrimidine and CH₄:pyrimidine) are irradiated have also been performed in order to better evaluate the methylation efficiency of CH₃OH and CH₄, as well as the role of H₂O in these experiments.

2.2.3.1. Laboratory Synthesis of Thymine: Experimental protocol and analysis techniques employed in the study of the formation of thymine are similar to those previously employed for studying the formation of uracil and cytosine (Materese et al. 2013). One notable exception, however, is that laboratory experiments on thymine formation also included experiments that employed higher UV photon doses than needed for the formation of uracil and cytosine.

Experimental results indicate that ice mixtures containing CH₃OH as a methyl group source do not yield measurable quantities of pure methylpyrimidines, although they may still

produce small quantities of thymine. The lack of methylpyrimidines may be partially explained by the fact that the photo-dissociation of pure CH₃OH is known to lead to branching ratios from which only 1 in 7 photolytic reactions will result in the production of a CH₃ group (Öberg et al. 2009).

In contrast, methane is a much better source of CH₃ groups than methanol. Experiments on the irradiation of CH₄:pyrimidine ice mixtures do lead to the production of 4-methylpyrimidine, although no other methylpyrimidine isomers were detected (Materese et al. 2013). It is interesting to note that similar HPLC analysis of residues produced from experiments involving quinoline rather than pyrimidine yielded similar results for the non-detection and detection of methyl addition in CH₃OH- and CH₄-containing ices, respectively (Elsila *et al.* 2006). The addition of H₂O to CH₄:pyrimidine mixtures results in the production of both 4- and 5-methylpyrimidine, once again highlighting the role H₂O can play as a catalyst for the formation of other less abundant methylpyrimidine isomers such as 5-methylpyrimidine, as is also observed for the formation of aminopyrimidines from H₂O:NH₃:pyrimidine ice mixtures (Nuevo et al. 2012).

While the production of thymine has been observed in these experiments, its overall abundance is relatively low in comparison to uracil or cytosine in similar previous experiments. One likely cause for this difference is the fact that thymine formation requires three substitutions, while both uracil and cytosine only require two each. This means that more photons are required, and this puts further constraints on the makeup of the surrounding ice to ensure that the proper radicals and/or ions are available. Additionally, since the formation of thymine via UV processing of pyrimidine in ices requires more photons than would be required for either uracil or cytosine, there are more opportunities to derail its formation with an alternative substitution or through further modification of the final product or its precursors.

2.2.3.2. Theory of Thymine Synthesis in the Condensed Phase: Detailed quantum chemical methods have been used to study the formation of thymine in a system where pyrimidine is surrounded by H₂O molecules, OH radicals, and CH₃ radicals, using the same computational methods as described in Section 2.2.1.2 for uracil (Bera et al. 2013). As mentioned earlier, the formation of thymine from pyrimidine requires the addition of three groups, two OH groups (positions 2 and 4 on the ring) and one CH₃ group (position 5).

As in the case of cytosine, investigating reaction mechanisms of thymine formation is complicated because the process occurs in mixed ices in which the addition of different groups can be done in several different orders (Figure 8). The two main types of reactions in these systems are oxidation and methylation. Three different routes have been explored: (i) the double oxidation of pyrimidine followed by a methylation, (ii) the methylation of pyrimidine followed by a double oxidation, and (iii) the oxidation of pyrimidine, followed by a methylation, followed by a second oxidation. As in the case of uracil, Bera et al. (2013) studied each of these routes for three types of mechanisms, namely, ionic, neutral-radical, and radical-radical.

In the first case, reactions will lead to the formation of uracil via the same route as described for pyrimidine in pure H₂O ice (Nuevo et al. 2009; Bera et al. 2010; Section 2.1.1). The subsequent reaction is the methylation of uracil, which can lead to thymine as well as to its isomers. Computations show no clear preference for the product of methylation of uracil, indicating that thymine has the same, or a smaller, probability to be formed relative to its isomer 6-methyluracil. In the second case, pyrimidine is first methylated and then oxidized to form thymine and its isomers. Interestingly, 5-methylpyrimidine, which is the methylated pyrimidine precursor of thymine, is not the most energetically favored product among all possible methylpyrimidine isomers. The oxidation steps have much more favorable reaction energies compared to the methylation steps, which is the key reason why oxidation is overwhelmingly favored. Moreover, although the addition of methyl groups to pyrimidine is energetically favorable, in an H₂O-rich ice, methylation will compete unfavorably against oxidation for two main reasons: (i) because OH radicals outnumber CH₃ radicals in these ice mixtures, and (ii) because calculations show that the oxidation step is significantly more exothermic and thus favored. It was also shown experimentally that 5-methylpyrimidine deposited in an H₂O ice and UV irradiated can lose its CH₃ group to form pyrimidine, which can then be oxidized to form 4(3*H*)-pyrimidone (Materese et al. 2013; Section 2.2.3.1). All these results are in agreement with what was seen experimentally, namely, methylation is the limiting step in the formation of thymine.

The theoretical investigations of Bera et al. (2013) also revealed two very interesting points. Firstly, following the reactions of a hydroxyl radical with the pyrimidine cation and its derivatives, their results show that, as is the case for uracil, the formation of thymine is not energetically favorable in the gas phase, and only becomes feasible in the condensed phase, where the H₂O ice matrix plays two roles, both as an oxidant and as a solvent for proton abstraction in both oxidation and methylation steps. Secondly, computations confirm that the oxidation of pyrimidine is significantly favored over its methylation. By combining all these results, it appears that the most probable pathway for the formation of thymine from pyrimidine is via the formation of 4(3*H*)-pyrimidone as a first step, then the formation of uracil after a second oxidation, and finally the methylation of uracil (Bera et al. 2013). The fact that methyl addition is the limiting step in this process, and that the formation of thymine in this route is not particularly favored over those of its isomers, could explain the very low quantities of thymine detected in laboratory samples (Materese et al. 2013) compared to what is seen for uracil and cytosine (Nuevo et al. 2009, 2012).

2.2.3.3. The Interesting Difference of Thymine: Experiments showed that thymine is not efficiently produced from the UV photo-irradiation of pyrimidine in ices containing H₂O and a methyl source such as the interstellar icy components CH₃OH and CH₄. This result may be attributed to the fact that the formation of thymine requires three substitutions, and thus higher doses of photons compared with the formation of uracil and cytosine, and to the fact that in an H₂O-rich ice mixture, as is the case in the ISM, methylation becomes the limiting step to form thymine. From a theoretical point of view, quantum chemical computations do not show any clear thermodynamic preference for the formation of thymine over other competing products; indeed, the methylation of uracil actually slightly favors the formation of 6-methyluracil over that of thymine (5-methyluracil). In an environment where

methylation competes with oxidation, oxidation is expected to dominate by virtue of its significantly larger reaction energies compared to that of methylation. Even if methylation takes place before oxidation in the process leading towards the formation of thymine, experiments starting with 5-methylpyrimidine in H₂O ice clearly showed that the CH₃ groups can be easily cleaved off to form pyrimidine, which subsequently will be most probably oxidized to uracil.

The fact that thymine is not formed as easily as the two other pyrimidine-based nucleobases under astrophysical conditions, as demonstrated by both experimental and theoretical studies, and the fact that thymine has not been detected in carbonaceous meteorites, raise an interesting question about its role in the origins of life. If we assume that photo-processing of icy grains plays an important role in the formation of nucleobases in astrophysical environments (Figure 2), we can imagine a scenario in which the pyrimidine-based nucleobases are formed in space, preserved in small objects such as asteroids and comets during the formation of the Solar System, and subsequently delivered to the early Earth and other telluric planets (Oró 1961; Chyba and Sagan 1992). However, since (i) methylation is an important step in the formation of thymine, (ii) this reaction appears inefficient in CH₃OH-rich ices like those seen in many interstellar dense clouds, and (iii) the formation of thymine requires at least three substitutions as opposed to two for uracil and cytosine, it is possible that thymine was not formed in the ISM as efficiently as the other pyrimidine-based nucleobases.

Therefore, the abundance of thymine in the comets and asteroids which seeded the primitive Earth may have been significantly smaller than those of uracil and cytosine. It is interesting to consider these results and assumptions in light of the fact that terrestrial thymine is only found in DNA, but not in RNA. Indeed, several scenarios for the emergence of life involve a period of time during which RNA was the molecule dominating biological reactions as a catalyst and a storage for a primitive genetic information (Gilbert 1986; Joyce 1989, 2002). It should be mentioned, however, that the RNA world hypothesis is still debated and alternative mechanisms involving small organic molecules which interact with each other via catalyzed reaction cycles driven by a flow of available free energy have also been proposed (Shapiro 2006).

Nonetheless, in the hypothesis that such an RNA world did lead to the origin of life, a paucity of thymine among the building blocks that were delivered to Earth could potentially have forced the emergence of replicable systems involving uracil rather than thymine, such as RNA. In such a scenario, the emergence of DNA, and thus the use of thymine, as a more stable, robust molecule to store genetic information, may have appeared at a later stage in the complexity of life, in part because it was not effectively delivered to the early Earth.

2.3. Synthesis of Purine-Based Nucleobases in Extraterrestrial Ices

Although the formation of pyrimidine-based nucleobases under extraterrestrial abiotic conditions has been extensively studied, the same cannot be said about the formation of purine-based nucleobases and their derivatives. Purine is the backbone of the two other biological nucleobases that constitute the genetic material of DNA and RNA, namely, adenine and guanine. There are also a wide variety of biological and non-biological

molecules based on the backbone of purine, such as xanthine, hypoxanthine, or caffeine. A majority of these compounds have been detected in meteorites (Hayatsu 1964; Folsome et al. 1971, 1973; Hayatsu et al. 1975; van der Velden and Schwartz 1977; Stoks and Schwartz 1979, 1981; Martins et al. 2008; Callahan et al. 2011), and it appears that meteoritic purine-based compounds are always more abundant than pyrimidine-based compounds, probably because they are most resistant to the UV radiation the meteorites' parent bodies experience before they are recovered on Earth.

Therefore, the study of the formation and photo-stability of purine-based compounds is important from both an astrochemical and astrobiological point of view. Unfortunately, to date there have been no experimental or theoretical studies of the photolytic formation of such purine-based compounds in ices under astrophysically relevant conditions. However, numerous experimental results have been obtained for radiation-induced functional group addition to small polycyclic aromatic hydrocarbons (PAHs; Bernstein et al. 1999b, 2001b, 2002b, 2003; Ashbourn et al. 2007), and small polycyclic aromatic nitrogen heterocycles (PANHs; Elsila et al. 2006) including pyrimidine-based compounds (Nuevo et al. 2009, 2012; Materese et al. 2013), and these allow us to comment on what might be expected for purine.

Given the fact that purine-based species are more stable to UV radiation than are pyrimidine-based species, it is reasonable to predict that experiments in which purine is mixed in ices of astrophysical interest containing compounds like H₂O, NH₃, CH₃OH, and CH₄ will lead to the formation of mostly purine-based compounds, including adenine and guanine, as opposed to pyrimidine, whose ring can be broken by UV radiation to form smaller, aliphatic compounds (Nuevo et al. 2012). Indeed, the abiotic formation of adenine from the UV irradiation of purine in ices would only require the addition of a single NH₂ group, while that of guanine would require the addition of only one OH group and one NH₂ group (Figure 9). Since previous irradiation studies of small PAHs and pyrimidine have shown that the addition of OH and NH₂ groups to aromatic cyclic molecules is an efficient process, the formation of adenine and guanine from purine ought to be straightforward. Furthermore, since the needed functional group additions involve only OH and NH₂, and not CH₃, one might expect production yields comparable or higher to those for uracil and cytosine under similar conditions, rather than the lower yields seen for thymine.

Finally, since astrophysical ices do not only contain H₂O and NH₃, but also a variety of carbon sources such as CO, CO₂, CH₃OH, or CH₄, it will be interesting to (i) understand the effect of these compounds on the formation of adenine, guanine, and other purine-based nucleobases such as xanthine and hypoxanthine, (ii) verify whether the inclusion of carbonaceous species leads to the methylation of purine, known to be the limiting reaction for the formation of methylated pyrimidine derivatives when formed in a H₂O-rich ice matrix, and (iii) verify whether carbon sources have any inhibiting effects on the addition of OH and/or NH₂ groups.

2.4. Photo-Stability of the Nucleobases in Ices

Although there is no extensive body of research examining the photo-stability of pyrimidine based nucleobases in ices, we can make some general observations. Little work has been

done to quantify the effect, but it is clear from the synthesis work that products are not simply made and left alone, but are continuously susceptible to further photolytic reactions, i.e., the abundance of any given species in the ices is the result of the competition of 'creation' and 'destruction', where this latter term can include complete disruption of the molecule or alteration of the functional groups to form other pyrimidine-based molecules. It is not currently established what radiation exposures are needed to bring the competing creation-destruction processes into quasi-equilibrium, but true equilibrium is likely never attained since irradiation will steadily drive H from the ices (Sandford and Allamandola 1993).

In most experimental mixtures, much of the original pyrimidine survives and is detected in HPLC chromatograms of the resulting residues (Nuevo et al. 2012; Materese et al. 2013), with the exception of H₂O:pyrimidine ice mixtures in which pyrimidine is nearly fully consumed by either photo-destruction or the formation of new products (Nuevo et al. 2009). Having said that, these experiments show that the pyrimidine ring is susceptible to photolytic destruction and rearrangement in ices, and photo-products that could not have been produced from any other source have been detected. For example, in experiments examining the formation of uracil, cytosine, and thymine, acyclic compounds including urea, glycine, N-formylglycine, and alanine are also detected (Nuevo et al. 2009, 2012; Materese et al. 2013). Experiments starting with only H₂O, CH₃OH/CH₄, and pyrimidine, where the only source of available nitrogen is from the destruction of pyrimidine itself, also show the presence of aminopyrimidines (Materese et al. 2013). Hydantoin, an oxidized carbon and nitrogen heterocyclic molecule with a 5-member ring, is also detected in many of these experiments. It is therefore clear that photolytic destruction of the pyrimidine ring occurs in these ices.

In addition to the destruction of the ring itself, we know that side groups can be cleaved from pyrimidine rings. A good example to demonstrate this is a series of experiments starting with H₂O:5-methylpyrimidine ice mixtures that were designed to test the efficiency of the conversion of the latter into thymine in the presence of H₂O. In these experiments, oxidized pyrimidines, such as 4(3H)-pyrimidone, that lacked any methyl group, are found among the products. The formation of these oxidized pyrimidines lacking any methyl side group requires that the initial CH₃ group in 5-methylpyrimidine is cleaved from the ring (Materese et al. 2013). Though in this case 5-methylpyrimidine is one of the initial ice components, we can expect to observe the same effect in molecules formed during the UV photolysis, further demonstrating the interaction of creation and destruction processes.

We are currently unaware of any data regarding the photolytic stability of purines in these types of ices. On the basis of irradiation experiments of normal polycyclic aromatic hydrocarbons (PAHs) and pyrimidine in similar ices (Bernstein *et al.* 2001, 2002; Ashbourn *et al.* 2007), we would expect that substitutions or rearrangement of side groups should occur. The degree to which the purine ring system is susceptible to photolytic destruction is less clear. In the case of PAHs, ring destruction is greatly diminished with an increasing number of rings. The extent to which this trend can be extended from pyrimidine to purine is unknown because of the heterocyclic nature of the rings.

3. Conclusions

Numerous studies have demonstrated that the radiation processing of astrophysically relevant ices can lead to the production of a host of organic molecules of astrobiological interest, including amino acids, quinones, and amphiphiles. Recent experimental and theoretical work has demonstrated that the presence of pyrimidine in such ices is expected to lead to the formation of the three pyrimidine-based nucleobases uracil, cytosine, and thymine along with a number of their isomers.

Formation of the pyrimidine-based nucleobases requires various combinations of oxidation, amination, and methylation, all of which are observed experimentally. However, theoretical work makes it clear that the presence of condensed phase H₂O in the ice matrices plays a key role in formation of the nucleobases. Its most obvious role is as a source of oxygen for the oxidation reactions needed to make all of the pyrimidine-based nucleobases. However, it also plays a key role in the addition of hydroxyl, amino, *and* methyl groups by mediating the abstraction of either a proton or a hydrogen atom from intermediaries to form the final products. Since H₂O is the most abundant component observed in most astrophysical ices, this critical compound is likely to be available to play these roles in most cold astrophysical environments.

The relative abundance of uracil, cytosine, and thymine produced during ice photolysis depends on both the composition of the ices and on the photon dose. In general, larger photon doses will lead to greater degrees of ice processing, but do not necessarily lead to greater yields of any particular photo-product. This is because the abundance of any given photo-product is the result of competing creation and destruction reactions, where 'destruction' includes both the rupture of the pyrimidine ring and the replacement of a previously added functional group with a different group. Thymine represents a good example of this issue. While experiments show that photolysis of pyrimidine in CH₄-rich ices can lead to methylation of the pyrimidine, and that thymine can be made in H₂O:CH₄:pyrimidine ices, thymine is generally seen to be produced in far lower abundances than uracil, presumably because relevant astrophysical ices are dominated by H₂O, not CH₄, methylation is less energetically favorable, and the formation of thymine requires the addition of three functional groups instead of the two required for uracil or cytosine. Understanding the true range of relative production efficiencies of uracil, cytosine, and thymine will require the completion of studies of pyrimidine photolysis in more complex, realistic astrophysical ice analogs.

It is interesting to note that the lower production efficiency of thymine relative to uracil is consistent with what is seen in the organics in meteorites. A paucity of abiotically produced thymine delivered to, and available on the early Earth is also of interest because it is consistent with the idea that modern DNA-based life may have been preceded by an earlier 'RNA world' that did not involve any thymine.

While a growing body of work exists for the photoprocessing of pyrimidine in ices, little comparable work has been done for purine in ices. As a result, it is not known whether similar ice photolysis processes can produce the two purine-based nucleobases adenine and

guanine. However, based on the earlier work on pyrimidine, as well as ice photolysis experiments on PAHs and other aromatic heterocycles, it is anticipated that such processing should create both adenine and guanine. Since the only reactions needed to make these two nucleobases involve the addition of a few hydroxyl and amino groups, and no methyl groups, these two nucleobases are expected to be formed with efficiencies similar to those seen for uracil, rather than the lower efficiencies associated with the production of thymine.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors are grateful for support from the NASA Origins of Solar Systems, Exobiology, Astrophysics Research and Analysis, and Astrobiology Programs. Much of the work reported in this paper also benefited from support of postdoctoral researchers under the NASA Postdoctoral Program. PPB and TJL would like to acknowledge financial support from NASA to investigate the formation and evolution of carbon based material in the universe. This manuscript benefited from the helpful comments of an anonymous reviewer.

References

- Alexander CMO'D, Fogel M, Yabuta H, Cody GD. The origin and evolution of chondrites recorded in the elemental and isotopic compositions of their macromolecular organic matter. *Geochim Cosmochim Acta*. 2007; 71:4380–4403.
- Allamandola LJ, Tielens AGGM, Barker JR. Interstellar polycyclic aromatic hydrocarbons - The infrared emission bands, the excitation/emission mechanism, and the astrophysical implications. *Astrophys J Suppl Ser*. 1989; 71:733–775. [PubMed: 11542189]
- Arribas M, de Vicente A, Arias A, Lázaro E. Effect of metallic cations on the efficiency of DNA amplification. Implications for nucleic acid replication during early stages of life. *Intl J Astrobiol*. 2005; 4:115–123.
- Ashbourn SFM, Elsila JE, Dworkin JP, Bernstein MP, Sandford SA, Allamandola LJ. Ultraviolet photolysis of anthracene in H₂O interstellar ice analogs: Potential connection to meteoritic organics. *Meteorit Planet Sci*. 2007; 42:2035–2041.
- Baldwin B, Sheaffer Y. Ablation and breakup of large meteoroids during atmospheric entry. *Journal of Geophysical Research - Space Physics*. 1971; 76(#19):4653–4668.
- Becke AD. Density functional thermochemistry. III. The role of exact exchange. *J Chem Phys*. 1993; 98:5. id. 5648.
- Ben-Naim, A. Hydrophobic interactions. Plenum Press; New York: 1980.
- Bénilan Y, Andrieux D, Khelifi M, Bruston P, Raulin F, Guillemin JC, Cossart-Magos C. Temperature dependence of HC₃N, C₆H₂, and C₄N₂ mid-UV absorption coefficients. Application to the interpretation of Titan's atmospheric spectra. *Astrophys Space Sci*. 1996; 236:85–95.
- Bera PP, Head-Gordon M, Lee TJ. Initiating molecular growth in the interstellar medium via complexes of observed ions and molecules. *Astron & Astrophys*. 2011; 535:12. id. A74.
- Bera PP, Lee TJ, Schaefer HF III. Are Isomers of vinyl cyanide missing links of interstellar pyrimidine formation? *J Chem Phys*. 2009; 131:7. id. 074303.
- Bera PP, Nuevo M, Mataerese CK, Sandford SA, Lee TJ. The formation of thymine under astrophysical conditions constrains its role in the origin of life: theoretical study. 2013 To be submitted to *Astrobiology*.
- Bera PP, Nuevo M, Milam SN, Sandford SA, Lee TJ. Mechanism for the abiotic synthesis of uracil via UV-induced oxidation of pyrimidine in pure H₂O ices under astrophysical conditions. *J Chem Phys*. 2010; 133:7. id. 104303.

- Bera, PP., Schaefer, HF, III. Trends and Perspectives in Modern Computational Science. Vol. 6. Brill Academic Publishers; Boston: 2006. Lesions in DNA Subunits: The Nucleic Acid Bases; p. 254-264.
- Bernstein MP, Dworkin JP, Sandford SA, Allamandola LJ. Ultraviolet irradiation of naphthalene in H₂O ice: Implications for meteorites and biogenesis. *Meteorit Planet Sci.* 2001; 36:351–358.
- Bernstein MP, Dworkin JP, Sandford SA, Cooper GW, Allamandola LJ. Racemic amino acids from the ultraviolet photolysis of interstellar ice analogues. *Nature.* 2002a; 416:401–403. [PubMed: 11919623]
- Bernstein MP, Elsila JE, Dworkin JP, Sandford SA, Allamandola LJ, Zare RN. Side group addition to the PAH coronene by UV photolysis in cosmic ice analogs. *Astrophys J.* 2002b; 576:1115–1120.
- Bernstein MP, Moore MH, Elsila JE, Sandford SA, Allamandola LJ, Zare RN. Side group addition to the PAH coronene by proton irradiation in cosmic ice analogs. *Astrophys J.* 2003; 582:L25–L29.
- Bernstein MP, Sandford SA, Allamandola LJ. Life's far-flung raw materials [interstellar organic molecules]. *Scientific American.* 1999a; 281:26–33. [PubMed: 10920765]
- Bernstein MP, Sandford SA, Allamandola LJ. The mid-infrared absorption spectra of neutral polycyclic aromatic hydrocarbons in conditions relevant to dense interstellar clouds. *Astrophys J Suppl Ser.* 2005; 161:53–64.
- Bernstein MP, Sandford SA, Allamandola LJ, Chang S, Scharberg MA. Organic compounds produced by photolysis of realistic interstellar and cometary ice analogs containing methanol. *Astrophys J.* 1995; 454:327–344.
- Bernstein MP, Sandford SA, Allamandola LJ, Gillette JS, Clemett SJ, Zare RN. UV irradiation of polycyclic aromatic hydrocarbons in ices: Production of alcohols, quinones, and ethers. *Science.* 1999b; 283:1135–1138. [PubMed: 10024233]
- Biver N, Bockelée-Morvan D, Boissier J, Crovisier J, Colom P, Lecacheux A, Moreno R, Paubert G, Lis DC, Sumner M, Frisk U, Hjalmarson Å, Olberg M, Winnberg A, Florén HG, Sandqvist A, Kwok S. Radio observations of Comet 9P/Tempel 1 before and after Deep Impact. *Icarus.* 2007; 191:494–512.
- Biver N, Bockelée-Morvan D, Colom P, Crovisier J, Paubert G, Weiss A, Wiesemeyer H. Molecular investigations of comets C/2002 X5 (Kudo-Fujikawa), C/2002 V1 (NEAT), and C/2006 P1 (McNaught) at small heliocentric distances. *Astron Astrophys.* 2011; 528:19. id. A142.
- Boulanger E, Anoop A, Nachtigallova D, Thiel W, Barbatti M. Photochemical steps in the prebiotic synthesis of purine precursors from HCN. *Angew Chem Int Ed.* 2013; 52:8000–8003.
- van Broekhuizen FA, Keane JV, Schutte WA. A quantitative analysis of OCN[−] formation in interstellar ice analogs. *Astron Astrophys.* 2004; 415:425–436.
- Brünken S, McCarthy MC, Thaddeus P, Godfrey PD, Brown RD. Improved line frequencies for the nucleic acid base uracil for a radioastronomical search. *Astron Astrophys.* 2006; 459:317–320.
- Bücherer HT, Fischbeck HTJ. *Prakt Chem, Advanced Synthesis & Catalysis.* 1934; 140:69–89.
- Burton AS, Elsila JE, Hein JE, Glavin DP, Dworkin JP. Extraterrestrial amino acids identified in metal-rich CH and CB carbonaceous chondrites from Antarctica. *Meteorit Planet Sci.* 2013; 48:390–402.
- Butlerow A. Formation synthétique d'une substance sucrée. *C R Acad Sci.* 1961; 53:145–147. (In French).
- Callahan MP, Burton AS, Elsila JE, Baker EM, Smith KE, Glavin DP, Dworkin JP. A search for amino acids and nucleobases in the Martian meteorite Roberts Massif 04262 using liquid chromatography-mass spectrometry. *Meteorit Planet Sci.* 2013; 48:786–795.
- Callahan MP, Smith KE, Cleaves HJ II, Ruzicka J, Stern JC, Glavin DP, House CH, Dworkin JP. Carbonaceous meteorites contain a wide range of extraterrestrial nucleobases. *Proc Natl Acad Sci.* 2011; 108:13995–13998. [PubMed: 21836052]
- Callahan, MP., Stern, JC., Glavin, DP., Whelley, KE., Martin, MG., Dworkin, JP. Distribution of nucleobases in CM and CR carbonaceous chondrites; *Astrobiology Science Conference 2010: Evolution and Life: Surviving Catastrophes and Extremes on Earth and Beyond*; April 26–30, 2010; League City, Texas. 2010. p. 5160LPI Contribution No. 1538
- Capone LA, Prasad SS, Huntress WT, Whitten RC, Dubach J, Santhanam K. Formation of organic molecules on Titan. *Nature.* 1981; 293:45–46.

- Chang S, Scattergood T, Aronowitz S, Flores J. Organic chemistry on Titan. *Rev Geophys Space Phys.* 1979; 17:1923–1933.
- Charnley, S. On the nature of interstellar organic chemistry. In: Cosmovici, CB, Bowyer, S., Werthimer, D., editors. *Astronomical and Biochemical Origins and the Search for Life in the Universe*, Proc 5th International Conf on Bioastronomy. Kluwer Academic; Dordrecht: 1997. p. 89-96. IAU Coll #161
- Charnley SB, Kuan YJ, Huang HC, Botta O, Butner HM, Cox N, Despois D, Ehrenfreund P, Kisiel Z, Lee YY, Markwick AJ, Peeters Z, Rodgers SD. Astronomical searches for nitrogen heterocycles. *Adv Space Res.* 2005; 36:137–145.
- Chyba CF. Impact delivery and erosion of planetary oceans in the early inner Solar System. *Nature.* 1990; 343:129–133.
- Chyba CF, Thomas PJ, Brookshaw L, Sagan C. Cometary delivery of organic molecules to the early earth. *Science.* 1990; 249:366–373. [PubMed: 11538074]
- Chyba C, Sagan C. Endogenous production, exogenous delivery and impact-shock synthesis of organic molecules: An inventory for the origins of life. *Nature.* 1992; 355:125–132. [PubMed: 11538392]
- Cody GD, Alexander CMO'D, Tera F. Solid-state (^1H and ^{13}C) nuclear magnetic resonance spectroscopy of insoluble organic residue in the Murchison meteorite: a self-consistent quantitative analysis. *Geochimica et Cosmochimica Acta.* 2002; 66:1851–1865.
- Cooper G, Kimmich N, Belisle W, Sarinana J, Brabham K, Garrel L. Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature.* 2001; 414:879–883. [PubMed: 11780054]
- Coustenis A, Bézard B, Gautier D, Marten A, Samuelson R. Titan's atmosphere from Voyager infrared observations. III – Vertical contributions of hydrocarbons and nitriles near Titan's north pole. *Icarus.* 1991; 89:152–167.
- Cronin JR, Chang S. Organic matter in meteorites: molecular and isotopic analyses of the Murchison meteorite. *NATO Advanced Study Institute and International School of Space Chemistry: The chemistry of life's origins.* 1993:209–258.
- Cronin JR, Pizzarello S. Enantiomeric excesses in meteoritic amino acids. *Science.* 1997; 275:951–955. [PubMed: 9020072]
- Cronin JR, Pizzarello S. Amino acid enantiomer excesses in meteorites: origin and significance. *Adv Space Res.* 1999; 23:293–299.
- Dartois E. The ice survey opportunity of ISO. *Space Sci Rev.* 2005; 119:293–310.
- Deamer, DW., Fleischaker, GR. *Origins of life: the central concepts.* Jones & Bartlett Pub; 1994.
- Decin L, De Beck E, Brünken S, Müller HSP, Menten KM, Kim H, Willacy K, de Koter A, Wyrowski F. Circumstellar molecular composition of the oxygen-rich AGB star IK Tauri. II. In-depth non-LTE chemical abundance analysis. *Astron Astrophys.* 2010; 516:23. id. A69.
- Delsemme, AH. Cosmic origin of the biosphere. In: Brack, A., editor. *The molecular origin of life.* Cambridge University Press; 1998.
- Dello Russo N, Vervack RJ Jr, Lisse CM, Weaver HA, Kawakita H, Kobayashi H, Cochran AL, Harris WM, McKay AJ, Biver N, Bockelée-Morvan D, Crovisier J. The volatile composition and activity of Comet 103 P/Hartley 2 during the EPOXI closest approach. *Astrophys J Lett.* 2011; 734:6. id. L8.
- Demyk K, Dartois E, d'Hendecourt L, Jourdain de Muizon M, Heras AM, Breitfellner M. Laboratory identification of the 4.62 μm solid state absorption band in the ISO-SWS spectrum of RAFGL 7009S. *Astron Astrophys.* 1998; 339:553–560.
- Disanti MA, Villanueva GL, Bonev BP, Magee-Sauer K, Lyke JE, Mumma MJ. Temporal evolution of parent volatiles and dust in Comet 9P/Tempel 1 resulting from the Deep Impact experiment. *Icarus.* 2007; 191:481–493.
- Dodd, RT. *Meteorites: A Petrologic-Chemical Synthesis.* Cambridge Univ. Press; Cambridge: 1981.
- Dorschner J, Henning Th. Dust Metamorphosis in the Galaxy. *Astron Astrophys Rev.* 1995; 6:271–333.
- Dunning TH. Gaussian-Basis Sets for Use in Correlated Molecular Calculations. 1 The Atoms Boron through Neon and Hydrogen. *J Chem Phys.* 1989; 90:1007–1024.

- Dworkin JP, Deamer DW, Sandford SA, Allamandola LJ. Self-assembling amphiphilic molecules: Synthesis in simulated interstellar/precometary ices. *Proc Natl Acad Sci*. 2001; 98:815–819. [PubMed: 11158552]
- Dworkin JP, Gillette JS, Bernstein MP, Sandford SA, Allamandola LJ, Elsila JE, McGlothlin DR, Zare RN. An Evolutionary Connection Between Interstellar Ices and IDPs? Clues from Mass Spectroscopy Measurements of Laboratory Simulations. *Adv Spa Res*. 2004; 33:67–71.
- Ehrenfreund P, Charnley SB. Organic molecules in the interstellar medium, comets, and meteorites: A voyage from dark clouds to the early Earth. *Ann Rev Astron Astrophys*. 2000; 38:427–483.
- Elsila JE, Dworkin JP, Bernstein MP, Martin MP, Sandford SA. Mechanisms of amino acid formation in interstellar ice analogs. *Astrophys J*. 2007; 660:911–918.
- Elsila JE, Hammond MR, Bernstein MP, Sandford SA, Zare RN. UV photolysis of quinoline in interstellar ice analogs. *Meteorit Planet Sci*. 2006; 41:785–796.
- Engel V, Staemmler V, van der Wal RL, Crim FF, Sension RJ, Hudson B, Anderson P, Hennig S, Weide K, Shinke R. Photodissociation of water in the first absorption band: A prototype for dissociation in a repulsive potential energy surface. *J Phys Chem*. 1992; 96:3201–3213.
- Ertem G. Montmorillonite, oligonucleotides, RNA and origin of life. *Orig Life Evol Biosph*. 2004; 34:549–570. [PubMed: 15570708]
- Ferris JP. HCN did not condense to give heteropolypeptides on the primitive earth. *Science*. 1979; 16:1135–1137.
- Ferris JP, Hagan WJ. HCN and chemical evolution: The possible role of cyano compounds in prebiotic synthesis. *Tetrahedron*. 1984; 40:1093–1120. [PubMed: 11541961]
- Ferris JP, Sanchez RA, Orgel LE. Studies in prebiotic synthesis – III. Synthesis of pyrimidine from cyanoacetylene and cyanate. *J Mol Biol*. 1968; 33:693–704. [PubMed: 5700419]
- Folsome CE, Lawless J, Romiez M, Ponnampuruma C. Heterocyclic compounds indigenous to the Murchison meteorite. *Nature*. 1971; 232:108–109. [PubMed: 16062864]
- Fondren LD, McLain J, Jackson DM, Adams NG, Babcock LM. Studies of reactions of a series of ions with nitrogen containing heterocyclic molecules using a selected ion flow tube. *Int J Mass Spectrom*. 2007; 265:60–67.
- Fraundorf P. The distribution of temperature maxima for micrometeorites decelerated in the Earth's atmosphere without heating. *Geophys Res Lett*. 1980; 10:765–768.
- Frenklach M, Feigelson ED. Formation of polycyclic aromatic hydrocarbons in circumstellar envelopes. *Astrophys J*. 1989; 341:372–384.
- Galliano F, Madden SC, Tielens AGGM, Peeters E, Jones AP. Variations of the mid-IR aromatic features inside and among galaxies. *Astrophys J*. 2008; 679:310–345.
- Gerakines PA, Moore MH, Hudson RL. Carbonic acid production in H₂O:CO₂ ices. UV photolysis vs proton bombardment. *Astron Astrophys*. 2000; 357:793–800.
- Gerakines PA, Moore MH, Hudson RL. Energetic processing of laboratory ice analogs: UV photolysis versus ion bombardment. *J Geophys Res*. 2001; 106:33381–33386.
- Gerakines PA, Moore MH, Hudson RL. Ultraviolet photolysis and proton irradiation of astrophysical ice analogs containing hydrogen cyanide. *Icarus*. 2004; 170:202–213.
- Gibb EL, Whittet DCB, Boogert ACA, Tielens AGGM. Interstellar ice: the infrared space observatory legacy. *Astrophys J Suppl Ser*. 2004; 151:35–73.
- Gibb EL, Whittet DCB, Schutte WA, Boogert ACA, Chiar JE, Ehrenfreund P, Gerakines PA, Keane JV, Tielens AGGM, van Dishoeck EF, Kerkhof O. An inventory of interstellar ices toward the embedded protostar W33A. *Astrophys J*. 2000; 536:347–356.
- Glavin DP, Dworkin JP, Aubrey A, Botta O, Doty JH, Martins Z, Bada JL. Amino acid analyses of Antarctic CM2 meteorites using liquid chromatography-time of flight-mass spectrometry. *Meteorit Planet Sci*. 2006; 41:889–902.
- Glavin DP, Elsila JE, Burton AS, Callahan MP, Dworkin JP, Hilt RW, Herd CDK. Unusual nonterrestrial L-proteinogenic amino acid excesses in the Tagish Lake meteorite. *Meteorit Planet Sci*. 2012; 47:1347–1364.
- Greenberg MJ. The structure and evolution of interstellar grains. *Scient Am*. 1984; 250:96–107.

- Gupta SK, Ochiai E, Ponnampereuma C. Organic synthesis in the atmosphere of Titan. *Nature*. 1981; 293:725–727.
- Hayatsu R. Orgueil meteorite: Organic nitrogen contents. *Science*. 1964; 146:1291–1293. [PubMed: 17810143]
- Hayatsu R, Anders E, Studier MH, Moore LP. Purines and triazines in the Murchison meteorite. *Geochim Cosmochim Acta*. 1975; 39:471–488.
- Hayes JM. Organic constituents of meteorites – A review. *Geochim Cosmochim Acta*. 1967; 31:1395–1440.
- Hehre WJ, Ditchfield R, Pople JA. Self-consistent molecular orbital methods. XII Further extensions of Gaussian-type basis sets for use in molecular orbital studies of organic molecules. *J Chem Phys*. 1972; 56:2257–2261.
- d'Hendecourt LB, Allamandola LJ, Greenberg JM. Time dependent chemistry in dense molecular clouds. I – Grain surface reactions, gas/grain interactions and infrared spectroscopy. *Astron Astrophys*. 1985; 152:130–150.
- d'Hendecourt L, Jourdain de Muizon M, Dartois E, Breittellner M, Ehrenfreund P, Bénit J, Boulanger F, Puget JL, Habing HJ. ISO-SWS observations of solid state features towards RAFGL 7009S. *Astron Astrophys*. 1996; 315:L365–L368.
- Horn A, Møllendal H, Guillemin JC. A quantum chemical study of the generation of a potential prebiotic compound, cyanoacetaldehyde, and related sulfur containing species. *J Phys Chem A*. 2008; 112:11009–11016. [PubMed: 18831542]
- Huang HC, Kuan YJ, Charnley SB, Hirano N, Takakuwa S, Bourke TL. Organic molecules in the hot corinos and circumstellar disks of IRAS 16293-2422. *Adv Space Res*. 2005; 36:146–155.
- Hudgins DM, Sandford SA. Infrared spectroscopy of matrix-isolated polycyclic aromatic hydrocarbons 1. PAHs containing 2 to 4 rings. *J Phys Chem A*. 1998a; 102:329–343.
- Hudgins DM, Sandford SA. Infrared spectroscopy of matrix-isolated polycyclic aromatic hydrocarbons 2. PAHs containing 5 or more rings. *J Phys Chem A*. 1998b; 102:344–352.
- Ip WH, Balsiger H, Geiss J, Goldstein BE, Kettmann G, Lazarus AJ, Meier A, Rosenbauer H, Schwenn R, Shelley E. Giotto IMS measurements of the production rate of hydrogen cyanide in the coma of Comet Halley. *Ann Geophys*. 1990; 8:319–325.
- Irvine WM. Extraterrestrial Organic Matter: A review. *Orig Life Evol Biosph*. 1998; 28:365–383. [PubMed: 9742722]
- Jalbout AF. Prebiotic synthesis of simple sugars by an interstellar formose reaction. *Orig Life Evol Biosph*. 2008; 38:489–497. [PubMed: 18998238]
- Jalbout AF, Abrell L, Adamowicz L, Polt R, Apponi AJ, Ziurys LM. Sugar synthesis from a gas-phase formose reaction. *Astrobiology*. 2007; 7:433–442. [PubMed: 17630839]
- Jehin E, Bockelée-Morvan D, Dello Russo N, Manfroid J, Hutsemékers D, Kawakita H, Kobayashi H, Schulz R, Smette A, Stüwe J, Weiler M, Arpigny C, Biver N, Cochran A, Crovisier J, Magain P, Rauer H, Sana H, Vervack RJ, Weaver H, Zucconi JM. A multi-wavelength simultaneous study of the composition of the Halley family comet 8P/Tuttle. *Earth Moon Planets*. 2009; 105:343–349.
- Jewell PR, Snyder LE. New circumstellar cyanoacetylene sources. *Astrophys J*. 1982; 255:L69–L73.
- Jewell PR, Snyder LE. Observations and analysis of circumstellar cyanoacetylene. *Astrophys J*. 1984; 278:176–185.
- Jones PA, Sarkissian JM, Burton MG, Voronkov MA, Filipovi MD. Radio observations of comet 9P/Tempel 1 with the Australia Telescope facilities during the Deep Impact encounter. *Month Not R Astron Soc*. 2006; 369:1995–2000.
- Jørgensen JK, Hogerheijde MR, van Dishoeck EF, Blake GA, Schöier FL. The structure of the NGC 1333-IRAS2 protostellar system on 500 AU scales. An infalling envelope, a circumstellar disk, multiple outflows, and chemistry. *Astron Astrophys*. 2004; 413:993–1007.
- Kerridge JF, Chang S, Shipp R. Isotopic characterization of kerogen-like material in the Murchison carbonaceous chondrite. *Geochim Cosmochim Acta*. 1987; 51:2527–2540. [PubMed: 11542082]
- Khanna RK. Condensed species in Titan's stratosphere: Confirmation of crystalline cyanoacetylene (HC₃N) and evidence for crystalline acetylene (C₂H₂) on Titan. *Icarus*. 2005; 178:165–170.

- Kuan YJ, Yan CH, Charnley SB, Kisiel Z, Ehrenfreund P, Huang HC. A search for interstellar pyrimidine. *Month Not R Astron Soc.* 2003; 345:650–656.
- Kuan YJ, Charnley SB, Huang HC, Kisiel Z, Ehrenfreund P, Tseng WL, Yan CH. Searches for interstellar molecules of potential prebiotic importance. *Adv Space Res.* 2004; 33:31–39.
- Kunde VG, Aikin AC, Hanel RA, Jennings DE, Maguire WC, Samuelson RE. C_4H_2 , HC_3N and C_2N_2 in Titan's atmosphere. *Nature.* 1981; 292:686–688.
- Kvenvolden K, Lawless J, Pering K, Peterson E, Flores J, Ponnamperna C. Evidence for extraterrestrial amino-acids and hydrocarbons in the Murchison meteorite. *Nature.* 1970; 228:923–926. [PubMed: 5482102]
- Lacy JH, Baas F, Allamandola LJ, van de Bult CEP, Persson SE, McGregor PJ, Lonsdale CJ, Geballe TR. 4.6 micron absorption features due to solid phase CO and cyano group molecules toward compact infrared sources. *Astrophys J.* 1984; 276:533–543.
- Lacy JH, Faraji H, Sandford SA, Allamandola LJ. Unraveling the 10 micron “silicate” feature of protostars: The detection of frozen interstellar ammonia. *Astrophys J Lett.* 1998; 501:L105–L109.
- Lauretta, DS., McSween, HY., Jr, editors. *Meteorites and the Early Solar System II.* Univ. Ariz. Press; Tucson: 2006.
- Lazcano A, Miller SL. The origin and early evolution of life: Prebiotic chemistry, the pre-RNA world, and time. *Cell.* 1996; 85:793–798. [PubMed: 8681375]
- Lee, TJ., Scuseria, GE. Achieving chemical accuracy with coupled-cluster calculations. In: Langhoff, SR., editor. *Quantum Chemical Electronic Structure Calculations with Chemical Accuracy.* Kluwer Academic Publishers; Dordrecht: 1995. p. 47-108.
- Lee CT, Yang WT, Parr RG. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys Rev B.* 1988; 37:785–789.
- Lee TJ, Jayatilaka D. An open-shell restricted Hartree-Fock perturbation theory based on symmetric spin orbitals. *Chem Phys Lett.* 1993; 201:1–10.
- Lee TJ, Rendell AP, Dyall KG, Jayatilaka D. Open-Shell Restricted Hartree-Fock Perturbation Theory: Some Considerations and Comparisons. *J Chem Phys.* 1994; 100:10. id. 7400.
- Levine JS, Augustsson TR, Natarajan M. The prebiological paleoatmosphere: Stability and composition. *Orig Life.* 1982; 12:245–259. [PubMed: 7162799]
- Lis DC, Bockelée-Morvan D, Boissier J, Crovisier J, Biver N, Charnley SB. Hydrogen isocyanide in Comet 73P/Schwassmann-Wachmann (Fragment B). *Astrophys J.* 2008; 675:931–936.
- Lis DC, Mehringer DM, Benford D, Gardner M, Phillips TG, Bockelée-Morvan D, Biver N, Colom P, Crovisier J, Despois D, Rauer H. New molecular species in Comet C/1995 O1 (Hale-Bopp) observed with the Caltech Submillimeter Observatory. *Earth Moon Planets.* 1997; 78:13–20.
- Loren RB, Wootten HA. Star formation in the bright-rimmed molecular cloud IC 1848 A. *Astrophys J.* 1978; 225:L81–L84.
- de Marcellus P, Bertrand M, Nuevo M, Westall F, Le Sergeant d'Hendecourt L. Prebiotic significance of extraterrestrial ice photochemistry: Detection of hydantoin in organic residues. *Astrobiology.* 2011; 11:847–854. [PubMed: 22059641]
- Martins Z, Alexander CMO'D, Orzechowska GE, Fogel ML, Ehrenfreund P. Indigenous amino acids in primitive CR meteorites. *Meteor Planet Sci.* 2007; 42:2125–2136.
- Martins Z, Botta O, Fogel ML, Sephton MA, Glavin DP, Watson JS, Dworkin JP, Schwartz AW, Ehrenfreund P. Extraterrestrial nucleobases in the Murchison meteorite. *Earth Planet Sci Lett.* 2008; 270:130–136.
- Materese CK, Nuevo M, Bera PP, Lee TJ, Sandford SA. Thymine and other Prebiotic Molecules Produced from the Ultraviolet Photo-Irradiation of Pyrimidine in Simple Astrophysical Ice Analogs. 2013 Submitted to *Astrobiology.*
- Mathis JS, Mezger PG, Panagia N. Interstellar radiation field and dust temperatures in the diffuse interstellar matter and in giant molecular clouds. *Astron Astrophys.* 1983; 128:212–229.
- Matthews CN. HCN did not condense to give heteropolypeptides on the primitive Earth (Reply to Ferris 1979). *Science.* 1979; 16:1136–1137.

- Matthews CN, Moser RE. Peptide synthesis from hydrogen cyanide and water. *Nature*. 1967; 215:1230–1234. [PubMed: 6052719]
- Maurette M. Carbonaceous micrometeorites and the origin of life. *Orig Life Evol Biosph*. 1998; 4/6:385–412.
- McElroy D, Walsh C, Markwick AJ, Cordiner MA, Smith K, Millar TJ. The UMIST database for astrochemistry 2012. *Astron Astrophys*. 2013; 550:A36. 13 Pages.
- McMullin JP, Mundy LG, Blake GA. The circumstellar environment of IRAS 05338-0624. *Astrophys J*. 1994; 437:305–316. [PubMed: 11539169]
- Millar TJ, Roueff E, Charnley SB, Rodgers SD. The chemistry of complex molecules in interstellar clouds. *Int J Mass Spectrom & Ion Proc*. 1995; 149/150:389–402.
- Miller SL. A production of amino acids under possible primitive Earth conditions. *Science*. 1953; 117:528–529. [PubMed: 13056598]
- Miller, SL., Cleaves, HJ. Prebiotic chemistry on the primitive Earth. In: Rigoutsos, I., Stephanopoulos, G., editors. *Systems Biology, Genomics*. Vol. I. Oxford University Press; Oxford: 2006. p. 4-56.
- Mordaunt DH, Dixon RN, Ashfold MNR. Photodissociation dynamics of $\tilde{A}$ state ammonia molecules. II The isotopic dependence for partially and fully deuterated isotopomers. *J Chem Phys*. 1996; 104:6472–6481.
- Müller D, Pitsch S, Kittaka A, Wagner E, Wintner CE, Eschenmoser A. Chemie von α -Aminonitrilen. Aldomerisierung von Glycolaldehyd-Phosphat zu racemischen Hexose-2,4,6-Triphosphaten und (in gegenwart von Formaldehyd) racemischen Pentose-2,4-Diphosphaten: *rac*-Allose-2,4,6-Triphosphat und *rac*-Ribose-2,4,-Diphosphat sind die Reaktionshauptprodukte. *Helv Chim Acta*. 1990; 73:1410–1468. (In German).
- Müller, HSP., Endres, CP., Stutzki, J., Schlemmer, S. Molecules in space. Physikalisches Institut, Universität zu Köln; 2013. <http://www.astro.uni-koeln.de/cdms/molecules> Accessed 10 June 2013
- Mullie F, Reisse J. Organic matter in carbonaceous chondrites. *Topics in Current Chemistry*. 1987; 139:85–117.
- Muñoz Caro GM, Meierhenrich UJ, Schutte WA, Barbier B, Arcones Segovia A, Rosenbauer H, Thiemann WHP, Brack A, Greenberg JM. Amino acids from ultraviolet irradiation of interstellar ice analogues. *Nature*. 2002; 416:403–406. [PubMed: 11919624]
- Nissenbaum A. Scavenging of soluble organic matter from the prebiotic oceans. *Orig Life*. 1976; 7:413–416. [PubMed: 1023140]
- Nuevo M, Auger G, Blanot D, d'Hendecourt L. A detailed study of the amino acids produced from the vacuum UV irradiation of interstellar ice analogs. *Orig Life Evol Biosph*. 2008; 38:37–56. [PubMed: 18175206]
- Nuevo M, Bredehöft JH, Meierhenrich UJ, d'Hendecourt L, Thiemann WHP. Urea, glycolic acid, and glycerol in an organic residue produced by ultraviolet irradiation of interstellar/pre-cometary ice analogs. *Astrobiology*. 2010; 10:245–256. [PubMed: 20402585]
- Nuevo M, Chen YJ, Yih TS, Ip WH, Fung HS, Cheng CY, Tsai HR, Wu CYR. Amino acids formed from the UV/EUV irradiation of inorganic ices of astrophysical interest. *Adv Space Res*. 2007; 40:1628–1633.
- Nuevo M, Milam SN, Sandford SA. Nucleobases and prebiotic molecules in organic residues produced from the ultraviolet photo-irradiation of pyrimidine in NH_3 and $\text{H}_2\text{O}+\text{NH}_3$ ices. *Astrobiology*. 2012; 12:295–314. [PubMed: 22519971]
- Nuevo M, Milam SN, Sandford SA, Elsila JE, Dworkin JP. Formation of uracil from the ultraviolet photo-irradiation of pyrimidine in pure H_2O ices. *Astrobiology*. 2009; 9:683–695. [PubMed: 19778279]
- Öberg KI, Garrod RT, van Dishoeck EF, Linnartz H. Formation rates of complex organics in UV irradiated CH_3OH -rich ices. I Experiments. *Astron Astrophys*. 2009; 504:891–913.
- Orgel LE. Prebiotic chemistry and the origin of the RNA world. *Critical Reviews in Biochemistry and Molecular Biology*. 2004; 39:99–123. [PubMed: 15217990]
- Oró J. Comets and the formation of biochemical compounds on the primitive Earth. *Nature*. 1961; 190:389–390.
- Oró J. Mechanisms of synthesis of adenine from hydrogen cyanide under possible primitive Earth conditions. *Nature*. 1961; 191:1193–1194. [PubMed: 13731264]

- Pardo JR, Cernicharo J, Goicoechea JR. Observational evidence of the formation of cyanopolyynes in CRL 618 through the polymerization of HCN. *Astrophys J.* 2005; 628:275–282.
- Peeters Z, Botta O, Charnley SB, Kiesel Z, Kuan YJ, Ehrenfreund P. Formation and photostability of N-heterocycles in space. I The effect of nitrogen on the photostability of small aromatic molecules. *Astron Astrophys.* 2005; 433:583–590.
- Pendleton YJ, Allamandola LJ. The organic refractory material in the diffuse interstellar medium: Mid-infrared spectroscopic constraints. *Astrophys J Suppl Ser.* 2002; 138:75–98.
- Pizzarello, S., Cooper, GW., Flynn, GJ. The nature and distribution of the organic material in carbonaceous chondrites and interplanetary dust particles. In: Lauretta, DS., McSween, HY., Jr, editors. *Meteorites and the Early Solar System II*. University of Arizona Press; Tucson: 2006. p. 625–651.
- Pizzarello S, Huang Y. Carbon isotopic analyses of individual Murchison amino acids. *Geochim Cosmochim Acta Suppl.* 2003; 67:380.
- Pizzarello S, Zolensky M, Turk KA. Nonracemic isovaline in the Murchison meteorite: chiral distribution and mineral association. *Geochim Cosmochim Acta.* 2003; 67:1589–1595.
- Prasad SS, Tarafdar SP. UV radiation field inside dense clouds—its possible existence and chemical implications. *Astrophys J.* 1983; 267:603–609.
- Prinn RG, Fegley B Jr. Bolide impacts, acid rain, and biospheric traumas at the Cretaceous-Tertiary boundary. *Earth Planet Sci Lett.* 1987; 83:1–15.
- Puget JL, Léger A. A new component of the interstellar matter – Small grains and large aromatic molecules. *Ann Rev Astron Astrophys.* 1989; 27:161–198.
- Raulin, F. *Lecture Notes in Physics*. Vol. 390. Springer; Berlin, Heidelberg: 1991. *Bioastronomy: The search for extraterrestrial life—The exploration broadens*; p. 141–148.
- Raulin F, Coll P, Coscia D, Gazeau MC, Sternberg R, Bruston P, Israël G, Gautier D. An exobiological view of Titan and the Cassini-Huygens mission. *Adv Space Res.* 1998; 22:353–362.
- Ricca A, Bauschlicher CW, Bakes ELO. A computational study of the mechanisms for the incorporation of a nitrogen atom into polycyclic aromatic hydrocarbons in the Titan haze. *Icarus.* 2001; 154:516–521.
- Rice J, Bera PP, Lee TJ. Ab initio quantum chemical study of formation of cytosine under astrophysical conditions. 2013 To be submitted to *J Chem Phys*.
- Roelfsema PR, Cox P, Tielens AGGM, Allamandola LJ, Baluteau JP, Barlow MJ, Beintema D, Boxhoorn DR, Cassinelli JP, Caux E, Churchwell E, Clegg PE, de Graauw T, Heras AM, Huygen R, van der Hucht KA, Hudgins DM, Kessler MF, Lim T, Sandford SA. SWS observations of IR emission features towards compact HII regions. *Astron Astrophys.* 1996; 315:L289–L292.
- Roy D, Najafian K, von Ragué Schleyer P. Chemical evolution: The mechanism of the formation of adenine under prebiotic conditions. *Proc Natl Acad Sci.* 2007; 104:17272–17277. [PubMed: 17951429]
- Ruiterkamp R, Peeters Z, Moore MH, Hudson RL, Ehrenfreund P. A quantitative study of proton irradiation and UV photolysis of benzene in interstellar environments. *Astron Astrophys.* 2005; 440:391–402.
- Sandford SA, Allamandola LJ, Tielens AGGM, Sellgren K, Tapia M, Pendleton Y. The Interstellar C-H stretching band near 3.4 μm : Constraints on the composition of organic material in the diffuse interstellar medium. *Astrophys J.* 1991; 371:607–620. [PubMed: 11538103]
- Sandford SA, Bernstein MP, Allamandola LJ. The mid-infrared laboratory spectra of naphthalene (C_{10}H_8) in solid H_2O . *Astrophys J.* 2004; 607:346–360.
- Sandford SA, Bernstein MP, Materese CK. The infrared spectra of polycyclic aromatic hydrocarbons with excess peripheral H atoms (H_n -PAHs) and their relation to the 3.4 and 6.9 μm PAH emission features. *Astrophys J.* 2013; (Suppl. 205):30. id. 8.
- Satzger H, Townsend D, Zgierski MZ, Patchkovskii S, Ullrich S, Stolow A. Primary processes underlying the photostability of isolated DNA bases: Adenine. *Proc Natl Acad Sci.* 2006; 103:10196–10201. [PubMed: 16803961]
- Schloerb FP, Kinzel WM, Swade DA, Irvine WM. HCN production from comet Halley. *Astrophys J.* 1986; 310:L55–L60.

- Schulte M, Blake D, Hoehler T, McCollom T. Serpentinization and its implications for life on the early Earth and Mars. *Astrobiology*. 2006; 6:364–376. [PubMed: 16689652]
- Schutte WA, Greenberg JM. Further evidence for the OCN[−] assignment to the XCN band in astrophysical ice analogs. *Astron Astrophys*. 1997; 317:L43–L46.
- Sephton MA, Wright IP, Gilmour I, de Leeuw JW, Grady MM, Pillinger CT. High molecular weight organic matter in Martian meteorites. *Planet Space Sci*. 2002; 50:711–716.
- Shao Y, Molnar LF, Jung Y, Kussmann J, Ochsenfeld C, Brown ST, Gilbert ATB, Slipchenko LV, Levchenko SV, O'Neill DP, DiStasio RA, Lochan RC, Wang T, Beran GJO, Besley NA, Herbert JM, Lin CY, Van Voorhis T, Chien SH, Sodt A, Steele RP, Rassolov VA, Maslen PE, Korambath PP, Adamson RD, Austin B, Baker J, Byrd EFC, Dachsel H, Doerksen RJ, Dreuw A, Dunietz BD, Dutoi AD, Furlani TR, Gwaltney SR, Heyden A, Hirata S, Hsu CP, Kedziora G, Khalliulin RZ, Klunzinger P, Lee AM, Lee MS, Liang W, Lotan I, Nair N, Peters B, Proynov EI, Pieniazek PA, Rhee YM, Ritchie J, Rosta E, Sherrill CD, Simmonett AC, Subotnik JE, Woodcock HL, Zhang W, Bell AT, Chakraborty AK, Chipman DM, Keil FJ, Warshel A, Hehre WJ, Schaefer HF, Kong J, Krylov AI, Gill PMW, Head-Gordon M. Advances in methods and algorithms in a modern quantum chemistry program package. *Phys Chem Chem Phys*. 2006; 8:3172–3191. [PubMed: 16902710]
- Shapiro R. Small molecule interactions were central to the origin of life. *Q Rev Biol*. 2006; 81:105–126. [PubMed: 16776061]
- Shen CJ, Greenberg JM, Schutte WA, van Dishoeck EF. Cosmic ray induced explosive chemical desorption in dense clouds. *Astron Astrophys*. 2004; 415:203–215.
- Shimoyama A, Hagishita S, Harada K. Search for nucleic-acid bases in carbonaceous chondrites from Antarctica. *Geochem J*. 1990; 24:343–348.
- Siegel BZ, Siegel SM. Enzyme-mimicking properties of silicates and other minerals. *Adv Space Res*. 1981; 1:27–36. [PubMed: 11541715]
- Simon MN, Simon M. Search for interstellar acrylonitrile, pyrimidine, and pyridine. *Astrophys J*. 1973; 184:757–762.
- Stoks PG, Schwartz AW. Uracil in carbonaceous meteorites. *Nature*. 1979; 282:709–710.
- Stoks PG, Schwartz AW. Nitrogen-heterocyclic compounds in meteorites – Significance and mechanisms of formation. *Geochim Cosmochim Acta*. 1981; 45:563–569.
- Strecker A. Über die künstliche Bildung der Milchsäure und einen neuen, dem Glycocol homologen Körper. *Justus Liebigs Ann Chem*. 1850; 75:27–45. (In German).
- Taillades J, Beuzelin I, Garrel L, Tabacik V, Bied C, Commeyras A. *N*-Carbamoyl- α -amino acids rather than free α -amino acids formation in the primitive hydrosphere: A novel proposal for the emergence of prebiotic peptides. *Orig Life Evol Biosph*. 1998; 28:61–77. [PubMed: 11536856]
- Thi WF, van Dishoeck EF, Dartois E, Pontoppidan KM, Schutte WA, Ehrenfreund P, d'Hendecourt L, Fraser HJ. VLT-ISAAC 3–5 μ m spectroscopy of embedded young low-mass stars. III Intermediate-mass sources in Vela. *Astron Astrophys*. 2006; 449:251–265.
- Thi WF, van Zadelhoff GJ, van Dishoeck EF. Organic molecules in protoplanetary disks around T Tauri and Herbig Ae stars. *Astron Astrophys*. 2004; 425:955–972.
- Tielens, AGGM., Allamandola, LJ. Composition, structure, and chemistry of interstellar dust. In: Hollenbach, D., Thronson, H., editors. *Interstellar Processes*. D. Reidel; Dordrecht: 1987. p. 397–469.
- Toupance G, Sebban G, Buvet R. Etape initiale de la polymérisation de l'acide cyanhydrique et synthèses prébiologiques. *J Chim Phys*. 1970; 67:1870–1874. (In French).
- Vázquez-Mayagoitia Á, Horton SR, Sumpter BG, Sponer J, Sponer JE, Fuentes-Cabrera M. On the stabilization of ribose by silicate minerals. *Astrobiology*. 2011; 11:115–121. [PubMed: 21391822]
- van der Velden W, Schwartz AW. Search for purines and pyrimidines in the Murchison meteorite. *Geochim Cosmochim Acta*. 1977; 41:961–968.
- Vuitton V, Yelle RV, Lavvas P, Klippenstein SJ. Rapid association reactions at low pressure: Impact on the formation of hydrocarbons on Titan. *Astrophys J*. 2012; 744:7. id. 11.
- Warnek P. A microwave-powered hydrogen lamp for vacuum ultraviolet photochemical research. *Applied Optics*. 1962; 1:721–726.

- Weber AL. Prebiotic amino acid thioester synthesis: Thiol-dependent amino acid synthesis from formose substrates (formaldehyde and glycolaldehyde) and ammonia. *Orig Life Evol Biosph.* 1998; 28:259–270. [PubMed: 9611766]
- Whittet DCB, Schutte WA, Tielens AGGM, Boogert ACA, de Graauw T, Ehrenfreund P, Gerakines PA, Helmich FP, Prusti T, van Dishoeck EF. An ISO SWS view of interstellar ices: First results. *Astron Astrophys.* 1996; 315:L357–L360.
- Woon DE. Pathways to glycine and other amino acids in ultraviolet irradiated astrophysical ices determined via quantum chemical modeling. *Astrophys J.* 2002; 571:177–180.
- Zuckerman B, Dyck HM. Dust grains and gas in the circumstellar envelopes around luminous red giant stars. *Astrophys J.* 1986; 311:345–359.

Figure 1.

The life cycle of matter in galaxies. New atoms and molecules formed in stars are ejected into the diffuse interstellar medium (ISM). Some of these materials can be subsequently gathered into dense molecular clouds, the formation sites of new stars and planets. During the formation of new stellar systems, these materials can be modified, and new materials made, as they are incorporated into small objects such as asteroids, comets, and planetesimals. Organic matter contained in asteroids and comets can then be delivered to telluric planets such as the Earth via meteorites and interplanetary dust particles (IDPs). Those organic compounds may constitute an inventory of prebiotic molecules that

subsequently play a role in the emergence of life if favorable chemical and physical conditions are available (from Dworkin et al. 2001).

Figure 2.

(Top) Spitzer telescope image of the Eta Carinae molecular cloud. (Bottom) Schematic of a cold interstellar dust grain. Volatiles such as H₂O, CO, CO₂, CH₃OH, and larger molecules such as polycyclic aromatic hydrocarbons (PAHs) can condense on the surface of cold dust grains in interstellar clouds. Such ice mantles can be processed by ionizing radiation, including UV photons, leading to the formation of more complex, refractory molecules (adapted from Bernstein et al. 1999a).

Figure 3.

The molecular structures of pyrimidine, purine, and the five biological nucleobases whose backbones are based on these compounds: uracil (2,4-dihydroxypyrimidine), cytosine (4-amino-2-hydroxypyrimidine), and thymine (2,4-dihydroxy-5-methylpyrimidine) for pyrimidine, and adenine (6-aminopurine) and guanine (2-amino-6-hydroxypurine) for purine.

Figure 4.

Total-ion chromatogram (top trace) of a residue produced from the UV photo-irradiation of an H_2O :pyrimidine = 20:1 ice mixture at low temperature followed by a subsequent warmup to room temperature. The middle and bottom traces show the single-ion chromatograms of the same residue corresponding to the masses for the singly and doubly oxidized pyrimidine derivatives, respectively, and show the identification of 4(3*H*)-pyrimidone and uracil in this sample (from Nuevo et al. 2009).

Figure 5.

(a) Mechanistic routes of pyrimidine cation reaction with hydroxyl radicals. The first step is a nucleophilic attack of a hydroxyl radical to pyrimidine cation. The second step is the loss of a proton assisted by H_2O . The third step is a tautomerization. (b) Energy diagram of the first oxidation step (energies are in kcal mol^{-1}). On the left hand side are the energies of intermediates + H_2O , and on the right hand side are the energies of singly oxidized products + H_3O^+ (see Bera et al. 2010).

Figure 6.

(a) Mechanistic routes of hydroxyl radical attacking the 4(3H)-pyrimidone cation. The first step is a nucleophilic attack of a hydroxyl radical to pyrimidine cation. The second step is the loss of a proton. (b) Energy diagram of the second oxidation step starting from 4(3H)-pyrimidone (energies are in kcal mol⁻¹). On the left hand side are the energies of intermediates + H₂O, and on the right hand side are the energies of singly oxidized products + H₃O⁺ (see Bera et al. 2010).

Figure 7.

(a) Mechanistic routes for the oxidation of starting from 2(1H)-pyrimidone towards the formation of uracil and its isomers. The first step is a nucleophilic attack of a hydroxyl radical to the 2(1H)-pyrimidone cation. The second step is the loss of a proton. (b) Energy diagram of 2(1H)-pyrimidone cation and hydroxyl radical (energies are in kcal mol⁻¹). On the left hand side are the energies of intermediates + H₂O, and on the right hand side are the energies of doubly oxidized products + H₃O⁺ (see Bera et al. 2010).

Figure 8.

Pathways leading to the formation of the three nucleobases uracil, cytosine, and thymine from the UV photo-irradiation of pyrimidine in H_2O , NH_3 , CH_3OH , and/or CH_4 ices, via the successive addition of OH , NH_2 , and CH_3 groups.

Figure 9.

Schematic showing how the UV photo-irradiation of purine in simple ices consisting of H_2O and/or NH_3 can be expected to lead to the formation of the biological nucleobases adenine and guanine, as well as several other derived compounds of astrobiological interest such as xanthine (2,6-dihydroxypurine) and hypoxanthine (6-hydroxypurine).