Published in final edited form as:

Dev Psychopathol. 2017 October ; 29(4): 1319–1331. doi:10.1017/S0954579417000293.

The Interaction Between Parenting and Children's Cortisol Reactivity at Age Three Predicts Increases in Children's Internalizing and Externalizing Symptoms at Age Six

Chelsey S. Barrios, B.A.¹, Sara J. Bufferd, Ph.D.², Daniel N. Klein, Ph.D.³, and Lea R. Dougherty, Ph.D.¹

¹Department of Psychology, University of Maryland College Park

²Department of Psychology, California State University San Marcos

³Department of Psychology, Stony Brook University

Abstract

Little is known about the role of stress reactivity in the emergence of psychopathology across early childhood. In this longitudinal study, we tested the hypothesis that child cortisol reactivity at age three moderates associations between early parenting and children's internalizing and externalizing symptoms from age three to age six. 160 children were assessed at age three and 135 children were reassessed at age six. At age three, we exposed children to stress-inducing laboratory tasks, during which we obtained four salivary cortisol samples, and parental hostility was assessed using an observational parent-child interaction task. At ages three and six, child psychiatric symptoms were assessed using a clinical interview with parents. Results indicated that the combination of high child cortisol reactivity and high observed parental hostility at age three was associated with greater concurrent externalizing symptoms at age three and predicted increases in internalizing and externalizing symptoms from age three to age six. Findings highlight that increased stress reactivity, within the context of hostile parenting, plays a role in the emergence of psychopathology from preschool to school entry.

Keywords

Cortisol Reactivity; Parenting; Preschool; Psychopathology

Recent research indicates that psychiatric symptoms and disorders in preschool-aged children constitute a significant public health problem (for reviews see Dougherty et al., 2015a and Egger & Angold, 2006). Preschool psychopathology occurs at similar rates to those observed in older children, is associated with significant impairment, and demonstrates stability over time (Bufferd, Dougherty, Carlson, & Klein, 2011; Bufferd, Dougherty, Carlson, Rose, & Klein, 2012; Lavigne et al., 1998, 2001; Luby, Gaffrey, Tillman, April, & Belden, 2014; Mesman, Bongers & Koot, 2001; Wichstrøm et al., 2012). Preschool mental

Corresponding Author: Lea Dougherty, Ph.D., Department of Psychology, University of Maryland, College Park, MD 20742, USA; Tel: +1-301-405-5464; Fax: +1-301-314-9566; ldougher@umd.edu.

Conflict of interest: The authors report no conflicts of interest.

health problems have also been found to predict psychopathology and impairment across early childhood (Bufferd et al., 2012; Dougherty et al., 2013a; Lavigne et al., 1998; Luby, Si, Belden, Tandon & Spitznagel, 2009), middle childhood (Dougherty et al., 2015b; Lahey et al., 2004) and into adolescence (Chronis-Tuscano et al., 2010; Luby et al., 2014). However, there is a paucity of prospective, longitudinal studies investigating the developmental pathways and mechanisms underlying psychopathology across early childhood to school entry.

The significance of life stress in the development of psychopathology across the lifespan has been well documented (Latimer et al., 2012; Monroe & Reid, 2009). Specifically, stress associated with the early caregiving environment has been found to be a potent predictor of psychiatric disorders across childhood (Humphreys & Zeanah, 2015) and into adulthood (Carr, Martins, Stingel, Lemgruber, & Jurueña, 2013). The hypothalamic pituitary adrenal (HPA) axis, one of the body's primary stress response systems, has been hypothesized to play a role in the etiology of numerous forms of psychopathology (Gunnar & Vazquez, 2006). The HPA axis is activated in the face of perceived threat or stress, resulting in the synthesis and release of the glucocorticoid cortisol, the primary stress hormone in humans. A dysregulated cortisol response to a psychosocial stressor has been linked concurrently to psychopathology in preschoolers, children, adolescents, and adults (Gunnar and Vazquez, 2006; Murri et al., 2014; Pariante, 2003). Nevertheless, evidence has been mixed as both cortisol hyper-reactivity and cortisol hypo-reactivity have been linked to youths' internalizing (hyper-reactivity: Hankin, Badanes, Abela, & Watamura, 2010; Kryski, Smith, Sheikh, Singh, & Hayden, 2013b; Luby et al., 2003; Rao, Hammen, Ortiz, Chen & Poland, 2008; hypo-reactivity: Hankin et al., 2010; Harkness, Stewart, & Wynne-Edwards, 2011; Hastings et al., 2011) and externalizing (hyper-reactivity: Obradovi, Bush, Stamperdahl, Adler, & Boyce, 2010; hyporeactivity: van Leeuwen et al., 2011) problems; however, it is unclear what role methodological differences across studies contribute to these divergent findings. Surprisingly, little research has examined whether cortisol reactivity prospectively predicts psychopathology across the lifespan, and there is a particular dearth of studies in children. In a study of adults with depression, increased cortisol reactivity to a relatively minimal psychosocial stressor predicted increases in depressive symptoms six months later (Morris, Rao, & Garber, 2012). In adolescent populations, increased cortisol reactivity predicted increased emotional and/or behavioral problems across six-month (Granger, Weisz, McCracken, Ikeda, & Douglas, 1996) and one-year (Susman, Dorn, Germain, Nottelmann, & Chrousos, 1997) follow-ups.

A growing body of research has focused on vulnerability-stress models, in which dysregulated stress reactivity serves as a vulnerability marker rendering individuals more susceptible to negative outcomes when exposed to adverse environments (Belsky, 2005; Boyce & Ellis, 2005; Monroe & Simons, 1991). It has been proposed that children demonstrating abnormalities in stress reactivity are at increased risk for maladaptive outcomes when they are exposed to adverse environments, including harsh parenting and high levels of family stress. For example, in a sample of adolescents, the combination of increased cortisol reactivity and greater cumulative childhood family aggression was associated with greater concurrent post-traumatic stress symptoms and antisocial behavior (Saxbe, Margolin, Spies Shapiro, & Baucom, 2012). Early childhood is a particularly critical

developmental period in which to study these effects as children rely heavily on their primary caregiver for most basic needs, and thus the parent-child relationship is a highly influential aspect of young children's environment. A few cross-sectional studies in preschoolers have demonstrated that the combination of increased cortisol reactivity and higher levels of family stress was associated with greater emotional symptoms (von Klitzing et al., 2012), poorer prosocial behavior (Obradovi et al., 2010), and greater concurrent externalizing symptomatology for boys but not girls (Hastings et al., 2011). In contrast, a recent study reported that the combination of preschoolers' lower cortisol reactivity and greater life stress in the past year was associated with greater externalizing symptoms and poorer psychosocial functioning (Kushner, Barrios, Smith, & Dougherty, 2016). Thus, similar to the cortisol reactivity-psychopathology associations, these cross-sectional findings show that both higher and lower cortisol reactivity in preschoolers may incur greater vulnerability to the early environment.

Very little research has examined these models using longitudinal designs. In a sample of children ages 9 to 15 years old, Badanes, Watamura, and Hankin (2011) reported that children who demonstrated lower cortisol reactivity and high levels of life stress had increased depressive symptoms one-year later after controlling for prior symptoms. In contrast, although cross-sectional findings were observed in two samples of five-year old children (Obradovi et al., 2010; von Klitzing et al., 2012), the interactive effect in these studies did not predict symptoms or functioning across the kindergarten school year. However, the cortisol reactivity assessments used in the early childhood studies collected cortisol samples at only two time points (i.e., pre- and post-stressor) (Obradovi et al., 2010; von Klitzing et al., 2012), which may limit the extent to which the assessments captured individual differences in children's cortisol responses to stress. In addition, the few longitudinal studies have only assessed children's symptoms over relatively short follow-up periods (e.g., one year or less), which may reduce the ability to observe changes in symptoms over time. Furthermore, previous longitudinal studies have not examined outcome measures of children's internalizing and externalizing symptoms in a single study, despite the fact that these prospective associations may vary across these different dimensions. Internalizing and externalizing psychopathology, though moderately correlated, can be differentiated even in early childhood (Olino, Dougherty, Bufferd, Carlson, & Klein, 2014; Sterba, Egger, & Angold, 2007); nevertheless, models of preschool psychopathology have also supported a single common factor shared by both internalizing and externalizing symptoms (Olino et al., 2014). Thus, it is unknown whether cortisol reactivity represents a vulnerability marker that is specific to one dimension or represents a general shared risk marker for both internalizing and externalizing problems across early childhood.

Using a multi-method assessment, the current study aimed to build upon this research to test the hypothesis that child cortisol reactivity at age three moderates associations between early negative parenting and children's internalizing and externalizing symptoms from age three to age six. Drawn from a larger longitudinal study ($N=559$), 160 three-year old children were randomly recruited to participate in the early childhood cortisol reactivity assessment (Dougherty, Klein, Rose, & Laptook, 2011b). Of the 160 children, 135 (84.4%) were followed-up three years later at age six. At the age three assessment, children were exposed to stress-inducing laboratory tasks, during which we obtained four salivary cortisol samples,

and parental hostility was assessed using an observational parent-child interaction task. We chose to focus on parental hostility because this parenting dimension shows consistent and strong associations with children's internalizing (McLeod, Weisz, & Wood, 2007) and externalizing (McKee, Colletti, Rakow, Jones, & Forehand, 2008) psychopathology and predicts mental and physical health problems into adulthood (Repetti, Taylor, & Seeman, 2002). In addition, given that research suggests that other family stressors (e.g., socio-economic status) impact the child via parenting (Bradley & Corwyn, 2002), our observational measure of parenting may capture broader environmental family stressors. At the age three and six assessments, child internalizing and externalizing symptoms were assessed using a structured clinical interview with parents. We hypothesized that children at age three who had high levels of cortisol reactivity and were exposed to high levels of parental hostility would demonstrate the highest levels of internalizing and externalizing symptoms concurrently at age three, and the greatest increases in internalizing and externalizing symptoms from age three to six years.

Method

Participants

The sample included 160 children and their primary caregivers recruited from the community as part of a larger longitudinal study ($N=559$) on early temperament and risk for psychopathology (Dougherty et al., 2011a; Olino, Klein, Dyson, Rose, & Durbin, 2010). Children and families were assessed at ages three and six years. At the age three assessment, a random sample of 160 children (80 females) participated in a laboratory-based cortisol reactivity assessment. The mean age of our subsample was 3.58 years ($SD = .24$) at the age three assessment and 6.33 years ($SD = .34$) at the age six assessment. Participants identified themselves as White (95.6%), African-American/Black (1.9%), Asian (1.9%), and other race (.6%). The majority of participants came from two-parent households (98.1%) and was middle-class, as measured by the Four Factor Index of Social Status ($M = 46.1$, $SD = 10.3$; Hollingshead, 1975). Children were of average cognitive ability, as indexed by the Peabody Picture Vocabulary Test ($M=105.0$, $SD=14.1$; Dunn & Dunn, 1997). Of the 160 participants, 135 (84.4%) parents completed the parent-reported clinical interview at the age six assessment. Only one significant difference emerged comparing the subsample presented in this report to the larger sample: children in the subsample had fewer internalizing symptoms at age three ($M=8.18$, $SD=6.30$) compared to the larger sample ($M=10.25$, $SD=8.38$), $t(367.66)=3.12$, $p=.002$.

Procedure

At age three, all children attended an initial laboratory session approximately 2.5 hours in duration, during which they participated with a female experimenter in 12 standardized tasks selected from the Laboratory Temperament Assessment Battery (Lab-TAB; Goldsmith, Reilly, Lemery, Longley, & Prescott, 1995), and cortisol samples were obtained. The tasks were designed to elicit a range of emotions and behaviors from the child and are described in detail elsewhere (see Dougherty et al., 2011a; Olino et al., 2010). Approximately two weeks later, parents and children returned for a second laboratory session and completed an observational parent-child interaction task to assess parental observed hostility. In between

laboratory sessions, parents completed a diagnostic clinical interview on the telephone to assess children's current internalizing and externalizing symptoms at age three. At the age six assessment, parents completed the clinical interview in person.

Child cortisol sampling procedure—Salivary cortisol was collected four times during the laboratory assessment. Saliva for cortisol assay was obtained by having the children dip 2-inch-long cotton dental rolls into small cups containing approximately .025 g of cherry Kool-Aid®. Children were then instructed to chew the cotton rolls until they were saturated with saliva. Previous work shows that the use of Kool-Aid® does not compromise the quality of the assays when used sparingly, as it does not significantly alter the pH of the saliva (Talge, Donzella, Kryzer, Gierens, & Gunnar, 2005). The collection of each sample took approximately 1–2 minutes. After each sample was collected, the saliva was expunged from the cotton roll into a micro tube and stored at –20°C until assayed.

The total duration of the 12 Lab-TAB tasks was approximately two hours. The length of time for each Lab-TAB task was standardized across all participants, and each episode was followed by a brief play break (2 minutes) to allow the child to return to a baseline affective state. The timing of the salivary cortisol samples was determined based on findings that salivary cortisol levels reflect the level of stress experienced in the prior 20–40 min (Dickerson & Kemeny, 2004) and on previous studies using similar stress-inducing paradigms, which have been shown to be sensitive to individual differences in cortisol reactivity in preschool-aged children (Luby et al., 2003; Talge, Donzella, & Gunnar, 2008). Based on these considerations, the first sample was taken after the informed consent process, approximately 20 minutes after arrival (baseline; Time 1). The second sample was collected 30 minutes following the Stranger Approach task of the Lab-TAB, the most stressful episode in the battery, during which the child was separated from his/her parent and a stranger entered the room (baseline + 60 min, Time 2). The third salivary cortisol sample was taken 30 minutes after Transparent Box, a frustration-inducing task in which the child was unable to unlock a box with a desirable toy inside (baseline + 90 min; Time 3). The fourth and final sample was collected 20 minutes after completion of all Lab-TAB tasks (baseline + 130 min; Time 4). To control for non-stress related elevations of cortisol, laboratory assessments were conducted at either 10 am (69% of the assessments) or 2 pm. Families were instructed prior to coming to the laboratory that the child should not eat within one hour before the scheduled lab visit, and that children should avoid caffeine for at least two hours and dairy products for at least 15 minutes prior to arrival. No child was taking corticosteroids at the time of the assessment.

Samples were assayed using a time-resolved fluorescence immunoassay with fluorometric end point detection (DELFI). All samples were assayed in duplicate. Samples yielding values above 44 nanomoles per liter (nmol/L) were excluded, which applied to four laboratory samples from four different individuals. The inter- and intra- assay coefficients of variation were between 7.1% and 9.0% and 4.0% and 6.7%, respectively.

Means and standard deviations of children's cortisol values for each of the four samplings are included in Table 1. Children's cortisol responses to the laboratory stressors in this sample have been described extensively in previous papers (see Dougherty, Klein, Congdon,

Canli, & Hayden, 2010; Dougherty et al., 2011b; Kryski et al., 2013a). As previously detailed, using multilevel modeling in HLM 6 (Scientific Software International Inc., IL), we observed a mean decrease (unstandardized coefficient = $-.085$, $SE = .016$, $t = -5.184$, $p < .001$) in children's cortisol from arrival to the laboratory to children's cortisol levels following the separation stressor (sample 2). Following the separation stressor, children's mean cortisol levels began to rise for sample 3 (30 min after a frustrating task) and sample 4 (20 min after the last Lab-TAB task), as indicated by a significant positive (i.e., concave up) quadratic effect (unstandardized coefficient = $.030$, $SE = .004$, $t = 8.174$, $p < .001$). The random error terms associated with the intercept, slope, and curvature were significant ($p < .001$), demonstrating variability among children's cortisol responses. While counterintuitive, it is important to note that this pattern is frequently observed in laboratory studies of cortisol reactivity in young children (Gotlib, Joorman, Minor & Cooney, 2006; Luby et al., 2003; for a review see Gunnar, Talge, & Herrera, 2009a), and may reflect stress-related increases related to anticipating the laboratory visit (Gunnar & Talge, 2007), such that the baseline sample is elevated and then declines. Conversely, it has been posited that this pattern of mean cortisol responses may reflect developmental differences in how children's HPA axis responds to stress, and that hyporesponsivity may be the normative response to stress in early to middle childhood (Gunnar, Wewerka, Fenn, Long & Griggs, 2009b).

To quantify a composite measure of cortisol reactivity, we calculated the area under the curve (AUC) with respect to increase (AUC_i), derived from the trapezoid formula from the four individual cortisol samples (Pruessner, Kirschbaum, Meinlschmid, & Hellhammer, 2003). The AUC_i provides a measure of the total change in cortisol levels across the four time points, and has been used widely in the literature as an index of HPA axis response (e.g. Booji, Bouma, de Jonge, Ormel, & Oldehinkel, 2013; Brennan et al., 2008; Dougherty et al., 2011b; Dougherty, Tolep, Smith, & Rose, 2013b; Kushner et al., 2016). Children's mean AUC_i was 0.36 ($SD = 12.90$), and scores ranged from -88.04 to 78.66 , demonstrating significant variability in children's AUC_i scores. 62.8% ($n = 98$) of children demonstrated a positive AUC_i (increase) and 37.2% ($n = 58$) demonstrated a negative AUC_i (decrease). AUC_i was positively skewed and was log10 transformed. The standardized z-score of the log10 transformed AUC_i variable was used in all analyses.

Observed parental hostility—At age three, 149 of the 160 children participated in a laboratory-based parent-child interaction task with one parent (96% mothers). The observational assessment was based on a modified version of the Teaching Tasks Battery and included six standardized tasks (book reading, block building, naming objects with wheels, matching shapes, completing a maze using an etch-a-sketch, and gift presentation) designed to elicit parent and child behaviors (Egeland et al., 1995). Parental hostility, which captures a parent's expression of anger, frustration, and criticism toward the child, was rated on a 5-point scale for each task, and ratings were averaged across tasks ($M = 1.17$, $SD = 0.27$, range = 1–3). A “1” indicated very low hostility (no signs of anger, annoyance, frustration, or rejection), a “3” indicated “moderate hostility” (signs of hostility or rejection occurred on several occasions during the session and at least one was clear or overt), and a “5” indicated “very high hostility” (frequent and consistent expressions of hostility or rejection towards

the child). Coders were unaware of the data on child psychopathology and cortisol reactivity. The internal consistency ($\alpha = .76$) and interrater reliability (intraclass correlation coefficient [ICC] = .83, $n = 55$) of the parental hostility scale were acceptable.

Child psychopathology—At the age three and age six assessments, parents were interviewed using the Preschool Age Psychiatric Assessment (PAPA; Egger, Ascher, & Angold, 1999). The PAPA is a parent-reported structured diagnostic interview designed to assess a range of psychiatric disorders from the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (American Psychiatric Association, 2000) in children ages two to six years old. Interviews were conducted by telephone by advanced graduate students in clinical psychology at the age three assessment and in person by a master's level clinician at the age six assessment. Diagnostic interviews with parents regarding their children have yielded similar results when administered by telephone and face-to-face (Lyneham & Rapee, 2005). For information on the interview's psychometric properties, see Egger et al. (2006). Rates of psychiatric disorders in the current subsample are reported in Table 1 and are consistent with the rates in the full sample (see Bufferd et al., 2011; 2012).

As described elsewhere (Dougherty et al., 2013a, 2015b), dimensional symptom scales for depression (major depressive disorder, dysthymic disorder, or depression-not otherwise specified [NOS]), anxiety (specific phobia, separation anxiety, social phobia, generalized anxiety disorder, agoraphobia, panic disorder, selective mutism), attention-deficit/hyperactivity disorder (ADHD), and oppositional defiant disorder (ODD) were created by summing items in each diagnostic category. We then created total internalizing and externalizing symptom scales by summing the depression and anxiety symptom scales, and the ADHD and ODD symptom scales, respectively. To examine inter-rater reliability, 21 audiotaped interviews from the age three assessment and 35 audiotaped interviews from the age six assessment were randomly selected and rated by a second interviewer. Inter-rater reliability and internal consistency were good for the age three internalizing ($ICC = .98$; $\alpha = .84$) and externalizing ($ICC = .81$; $\alpha = .89$) symptom scales and for the age six internalizing ($ICC = .73$; $\alpha = .86$) and externalizing ($ICC = .99$; $\alpha = .89$) symptom scales.

Due to concerns about administration time at the age three assessment, in the first 53.2% of the full sample ($n = 246$) the interviewer used the Early Childhood Inventory-4 (ECI-4; Gadow & Sprafkin, 2000) ADHD and ODD scales to serve as a screen to help determine whether to complete the ADHD and ODD sections of the PAPA. The ECI-4 is a parent rating scale used to screen DSM-IV emotional and behavioral disorders in three- to six-year-olds (Gadow & Sprafkin, 2000). Sprafkin and colleagues reported that the correct classification rates for ADHD and ODD with respect to chart diagnoses were 60% and 74%, respectively (Sprafkin, Volpe, Gadow, Nolan, & Kelly, 2002). Coefficient alphas for the ECI-4 were .79 (ADHD-Inattention), .82 (ADHD-Hyperactivity/Impulsivity), and .85 (ODD) in the sample. All ECI-4 ODD and ADHD items were reviewed by the interviewer. When parent-reports on the ECI-4 indicated a low likelihood of ODD or ADHD symptoms (i.e., most items were endorsed as “never” or “sometimes”), interviewers probed the broad domains of oppositionality, inattention, hyperactivity, and impulsivity to confirm the absence of symptoms before skipping out. When parent-reports on the ECI-4 indicated a potential likelihood of ODD or ADHD symptoms (i.e., items endorsed as “often” or “very often”), the

corresponding PAPA sections were administered in their entirety. In the remaining 46.8% of the sample ($n = 216$) and in the entire sample at age six, the PAPA ADHD and ODD sections were administered to all parents. At age three, ADHD and ODD dimensional scores were estimated for children for whom these sections were skipped using the ECI-4 ADHD and ODD items and maximum likelihood estimation procedures for missing values (Acock, 2005). This is less biased than pairwise and list-wise deletion procedures, even with large amounts of missing data (Navarro, 2003). Given the differences in the assessment of externalizing symptoms at age three and age six, we cannot make direct comparisons of the scores.

Data Analysis Plan

We examined whether child cortisol reactivity at age three interacted with observed parental hostility at age three to predict internalizing and externalizing symptoms across early childhood. All analyses used standardized (z-scored) variables as independent variables and unstandardized variables as dependent variables. In each model, independent variables included child cortisol reactivity assessed at age three, parental hostility assessed at age three, and the cortisol reactivity X parental hostility interaction term. Dependent variables included internalizing symptoms at age three and age six, and externalizing symptoms at age three and age six. Separate models were run for each of the four dependent variables. For models with age six symptom scales as the dependent variable, we controlled for levels of children's internalizing and externalizing symptoms scales at age three. Dependent variables therefore represent residuals; that is, the effects of the predictors on the dependent variables reflect change in that variable from one time point to the next. Child age at baseline, gender, and time of visit were also included as covariates if they were associated with the dependent variable.

Significant interactions were probed using simple slopes tests according to Aiken and West (1991). To better examine the pattern of moderation, Hayes and Matthes' guidelines (Hayes & Matthes, 2009) were used to test the regions of significance according to the Johnson-Neyman technique (Johnson & Fay, 1950). This approach uses the asymptotic variances, covariances, and other regression parameters to determine the upper and lower boundaries of the moderator (cortisol reactivity) at which the relation between the independent and dependent variable is significant ($p < .05$).

Results

Means and standard deviations of the study variables are reported in Table 1, and correlations among study variables are presented in Table 2. Child age at baseline was positively associated with children's externalizing symptoms at age six; thus, child age at baseline was included as a covariate in analyses with age six externalizing symptoms as the dependent variable.¹ Child gender and time of visit were not significantly associated with children's symptoms at ages three or six and were not included as covariates.² Child cortisol reactivity at age three was significantly positively associated with externalizing symptoms at

¹Child age at the age six assessment was not significantly associated with concurrent symptoms and was not included as a covariate.

²Given the diurnal rhythm of cortisol across the day, we re-ran all analyses controlling for time of day, and all results were similar.

age three and age six. Observed parental hostility at age three was significantly positively associated with externalizing symptoms at age three and age six. Internalizing and externalizing symptoms demonstrated moderate stability from age three to age six, and internalizing and externalizing symptoms were significantly positively correlated from age three to age six.

Early Childhood Cortisol Reactivity, Parental Hostility, and Internalizing Symptoms at Ages Three and Six Years

Results of the multiple linear regression models for child internalizing symptoms are presented in Table 3. The interaction between cortisol reactivity and parental hostility at age three was associated with age three internalizing symptoms at a trend-level ($p = .06$), and the form of the interaction was consistent with the significant interaction described next. At age six, the interaction between child cortisol reactivity and parental hostility at age three predicted internalizing symptoms at age six after controlling for internalizing and externalizing symptoms at age three. As seen in Figure 1, for children with high levels of cortisol reactivity at age three, parental hostility at age three predicted increases in internalizing symptoms at age six ($b = 2.79$, $SE = 1.17$, $pr = .21$, $p = .02$), whereas for children with low levels of cortisol reactivity at age three, there was no significant association between parental hostility at age three and internalizing symptoms at age six ($b = -3.64$, $SE = 2.20$, $pr = -.15$, $p = .10$). Regions of significance testing indicated that for children with cortisol reactivity at levels greater than .80 (standardized z-score), parental hostility at age three predicted increases in internalizing symptoms at age six. Reversing the moderator, regions of significance tests also demonstrated that this moderated effect was specific to levels of parental hostility greater than 1.06 (standardized z-score).

Early Childhood Cortisol Reactivity, Parental Hostility, and Externalizing Symptoms at Ages Three and Six Years

Results of the multiple linear regression models for child externalizing symptoms are presented in Table 4. The interaction between child cortisol reactivity and parental hostility at age three was significantly associated with concurrent externalizing symptoms at age three. For children with high levels of cortisol reactivity at age three, parental hostility at age three was significantly positively associated with externalizing symptoms at age three ($b = 5.13$, $SE = .92$, $pr = .43$, $p < .001$). In contrast, for children with low levels of cortisol reactivity at age three, there was no significant concurrent association between parental hostility at age three and externalizing symptoms at age three ($b = -2.49$, $SE = 1.80$, $pr = -.12$, $p = .17$). Regions of significance tests indicated that for children with cortisol reactivity at levels greater than .15 (standardized z-score), parental hostility at age three was positively associated with externalizing symptoms at age three. Reversing the moderator, regions of significance tests also demonstrated that this moderated effect was specific to levels of parental hostility greater than .28 (standardized z-score).

Similarly, the interaction between child cortisol reactivity and parental hostility at age three significantly predicted increases in externalizing symptoms at age six after controlling for internalizing and externalizing symptoms at age three. As seen in Figure 2, for children with high levels of cortisol reactivity at age three, parental hostility at age three significantly

predicted increases in externalizing symptoms at age six ($b = 3.21$, $SE = .60$, $pr = .44$, $p < .001$). In contrast, for children with low levels of cortisol reactivity at age three, parental hostility at age three did not significantly predict externalizing symptoms at age six ($b = -1.41$, $SE = 1.11$, $pr = -.11$, $p = .21$). Regions of significance tests indicated that for children with age three cortisol reactivity at levels greater than .22 (standardized z-score), parental hostility at age three significantly predicted increases in externalizing symptoms at age six. Reversing the moderator, regions of significance tests also demonstrated that this moderated effect was specific to levels of parental hostility greater than .61 (standardized z-score).³

In order to test whether our findings could be attributed to differences in children's baseline cortisol levels, all models reported in Table 3 and Table 4 were rerun replacing the AUCi statistic with children's baseline (sample 1) cortisol levels. No main or moderated effects involving children's baseline cortisol levels were observed. The interaction between children's baseline cortisol levels and parental hostility was not significantly associated with children's internalizing (age 3: $b = -.43$, $SE = .89$, $pr = -.04$, $p = .63$; age 6: $b = -1.87$, $SE = .128$, $pr = -.13$, $p = .14$) or externalizing (age 3: $b = -.27$, $SE = 1.19$, $pr = -.02$, $p = .82$; age 6: $b = -.70$, $SE = .66$, $pr = -.09$, $p = .29$) symptoms at age three or age six. In addition, in Appendix A we present supplementary data that includes parental supportive presence at age three as an additional independent variable. The interactions between parental hostility and children's cortisol reactivity remained significant in all models, whereas none of the interactions between parental support and children's cortisol reactivity were significant, indicating that our findings are specific to negative aspects of the parenting environment.

Lastly, we explored whether child gender moderated associations between early child cortisol reactivity and parental hostility and children's psychiatric symptoms. We reran each of the four models described above and added child gender, the two-way interaction terms, child gender X child cortisol reactivity and child gender X parental hostility, and the three-way interaction term, child gender X child cortisol reactivity X parental hostility, as independent variables. No gender differences were observed for any of the four dependent variables.

Discussion

The current study tested whether child cortisol reactivity moderated the effects of parental hostility on children's internalizing and externalizing symptoms both concurrently at age three and prospectively at age six after accounting for prior symptoms. We found that preschoolers who exhibited high cortisol reactivity and experienced high levels of parental hostility at age three evidenced greater concurrent externalizing symptoms at age three and increases in both internalizing and externalizing symptoms from age three to age six. Our findings suggest that increased cortisol reactivity in early childhood may render young children more susceptible to negative parenting and increase risk for developing emotional and behavioral problems by school entry. Identifying young children who are at greatest risk

³All results were similar for the subsample that received the full ADHD and ODD sections at age three. When restricting analyses to only those subjects who had complete PAPA ADHD and ODD sections at age three, the interactive effects of age three cortisol reactivity and age three parental hostility continued to be associated with externalizing symptoms at age three ($b = 2.29$, $SE = .86$, $pr = .39$, $p = .01$) and externalizing symptoms at age six ($b = 2.48$, $SE = 1.09$, $pr = .34$, $p = .03$).

for early mental health problems is critical given that early psychopathology adversely affects young children's functioning and predicts later psychopathology and impairment across development (Bufferd et al., 2012; Dougherty et al., 2013a; 2015b; Mesman et al., 2001).

Our findings are consistent with cross-sectional studies in young children indicating that the combination of dysregulated cortisol reactivity and exposure to negative family contexts is associated with concurrent child internalizing symptoms, externalizing symptoms, and psychosocial difficulties (Hastings et al., 2011; Kushner et al., 2016; Obradovi et al., 2010; von Klitzing et al., 2012). However, in contrast to our findings, the only other study that observed a significant longitudinal effect found that the combination of low, rather than high, cortisol reactivity and greater family stress predicted greater internalizing symptoms one year later in youth ages 9–15 (Badanes et al., 2011). Taken together, these findings suggest that dysregulated cortisol reactivity may serve as a vulnerability marker across childhood. However, the association of risk with high versus low stress reactivity may vary across development, or it may vary as a function of prior exposure to earlier chronic stressors. It is important to note that to date, none of the longitudinal studies have considered whether abnormalities in children's cortisol reactivity were present prior to the environmental stressor. It will be important for future research to establish cortisol reactivity as a pre-existing vulnerability marker and to test the temporal or causal associations. Nevertheless, our findings support that these early childhood markers of risk predict psychiatric symptoms three years later, which is the longest follow-up in the literature to date. Moreover, it is especially noteworthy that these effects were evident over and above prior symptoms given the stability of symptoms across the assessments.

We observed significant interactive effects in longitudinal models for both internalizing and externalizing symptoms, even after controlling for prior internalizing and externalizing symptoms. In concurrent models, we found a significant interactive effect on age three externalizing symptoms only; however, there was a trend level interactive effect on children's concurrent internalizing symptoms at age three. These findings suggest that cortisol reactivity to environmental stressors may represent a common or shared risk factor for both internalizing and externalizing symptoms across early childhood. Our findings also suggest that externalizing symptoms may be more sensitive than internalizing symptoms to variations in cortisol reactivity and parental hostility, in that the regions of significance tests demonstrated that it takes more extreme levels of both to produce significant effects for internalizing symptoms. It will be important to determine whether variations in cortisol reactivity and parental hostility continue to serve as a common vulnerability marker for both internalizing and externalizing problems across development or whether it becomes differentiated and specific to one form of psychopathology over time.

Our findings may have implications for understanding the developmental trajectory of psychopathology. Importantly, cortisol responses to stress appear to change across development with particularly marked changes occurring with pubertal onset and hormone fluctuations (Gunnar et al., 2009a; Gunnar et al., 2009b). Similarly, trajectories of internalizing and externalizing symptoms also change across development, with internalizing symptoms showing a marked rise in adolescence (McLaughlin & King, 2015)

and child externalizing symptoms possibly manifesting as mood dysregulation or conduct problems in adolescence (Okado & Bierman, 2014). Thus, it is possible that the associations observed in the current study may also change across development. However, to date, studies examining the interactive effect of cortisol reactivity and stress on psychopathology have been mixed. In preschool-age, school-age, and adolescent youth, the combination of environmental stress and both heightened (Hastings et al., 2011; Obradović et al., 2010; Saxbe et al., 2012; Steeger, Cook, & Connell, 2016; von Klitzing et al., 2012) and blunted (Badanes et al., 2011; Kushner et al., 2016) cortisol reactivity have been linked to both greater internalizing and externalizing symptoms. These studies have been largely limited to cross-sectional designs and have typically not examined both children's internalizing and externalizing psychopathology in the same study, making it difficult to determine whether these associations vary over time. Nevertheless, a recent study reported that in emerging adults exposed to childhood maltreatment, increased cortisol reactivity was linked to greater internalizing problems, whereas blunted cortisol reactivity was linked to greater externalizing problems (Hagan, Roubinov, Mistler, & Luecken, 2014), suggesting that distinct associations may emerge later in development. Taken together, these findings suggest that these associations are dynamic, and it will be important for future research to use long-term follow-up designs into adulthood to delineate the developmental progression of these associations.

Our findings suggest that child cortisol reactivity may sensitize young children to negative features of the parent-child relationship. Research suggests that cortisol reactivity is related to emotion regulation abilities (de Veld, Riksen-Walraven, de Weerth, 2012; Lam, Dickerson, Zoccola, & Zaldivar, 2009), as well as other physiological indices of stress sensitivity, including respiratory sinus arrhythmia (Doussard-Roosevelt, Montgomery, & Porges, 2003), neural responses to psychosocial stress (Dedovic, Duchesne, Andrews, Engert, & Pruessner, 2009), and genes involved in the susceptibility to life stress (Dougherty et al., 2010). Therefore, children with dysregulated cortisol reactivity may have multiple underlying biological and behavioral markers of stress sensitivity that render them less able to cope with parents' expression of anger, frustration, and criticism. Increased stress sensitivity may also contribute to children's behavioral responses within the parent-child relationship (e.g., greater distress, avoidance, or noncompliance) that evoke or exacerbate negative parenting. It will be important for future research to examine child-level risks across multiple levels of analysis and within the reciprocal context of the parent-child dyad to determine whether they represent common or unique risk markers and how they impact the parent-child relationship and child outcomes.

Although our study focused on negative aspects of the parenting environment, we ran supplementary analyses (see Appendix A) that show that parental support did not moderate children's cortisol reactivity. Nevertheless, it is possible that if we examined the effects of enriched environments (e.g., teacher/mentor supportive presence, positive sibling relationships) in children with identified vulnerability markers, our results may have differed (e.g., Essex, Armstrong, Burk, Goldsmith & Boyce, 2011; Morgan, Shaw, & Olino, 2012). Recent theoretical models including biological sensitivity to context (Boyce & Ellis, 2005) and differential susceptibility (Belsky & Pluess, 2009) suggest that children with vulnerability markers may respond in a "for better or for worse" manner to the environment,

experiencing poor outcomes when exposed to negative environments but optimal outcomes when exposed to positive environments. This work has important clinical implications as intervention studies suggest that children with increased cortisol reactivity may benefit most from treatment (van de Wiel, van Goozen, Matthys, Snoek, & van Engeland, 2004) and that children's HPA axis functioning may mediate the effects of a family intervention on change in children's behavior problems (O'Neal et al., 2010). Taken together, this work highlights that children's cortisol reactivity may be involved in multiple pathways to child adjustment and holds promise in identifying children who may be most responsive to parent interventions or possibly as a mechanisms of behavioral change.

The current study had several notable strengths. This study included a three-year follow-up assessment, which is the longest follow-up to date examining the interactive effects of cortisol reactivity and negative family influences on child, adolescent and adult outcomes. In addition, the sample was randomly recruited from the community, and we used a multi-method approach, including an observational assessment of early parenting, multiple post-stressor cortisol samplings to capture individual differences in stress responses, and a developmentally-sensitive, parent-reported clinical interview to assess child internalizing and externalizing symptoms from preschool to school entry.

The study also had limitations. First, primary caregivers, typically mothers, were the sole informants regarding children's psychopathology. Future work should incorporate a multi-informant approach, including co-parent and teacher reports. Second, since the combination of child cortisol reactivity and parental hostility was assessed at age three only, we cannot test whether their change across early childhood or whether the continued presence of high cortisol reactivity and parental hostility at age six influence children's symptoms. Third, children in our subsample had significantly fewer internalizing symptoms at age three as compared to the larger sample; this may have decreased our power to detect significant effects, as we found only a trend level interaction effect for children's internalizing symptoms at age three. Fourth, although we did not observe any gender differences in our sample, it is possible that our sample size was too small to detect significant differences. Further work with larger samples is needed to examine whether gender moderates the complex associations between children's stress reactivity, parenting, and child psychopathology (e.g., Hastings et al., 2011).

Fifth, we did not have sufficient power to test whether the interaction between children's cortisol reactivity and parental hostility predict psychiatric diagnoses. It will be important for future studies to examine both psychiatric symptoms and diagnoses in larger samples. Sixth, we used a screener for ADHD and ODD to reduce administration time for a portion of the sample at the age three assessment and employed maximum likelihood estimation procedures for missing values. However, results were comparable for the subsample of participants who did complete the full PAPA at ages three and six. Nonetheless, it is possible that the use of the screener resulted in some additional error variance. Seventh, although the range of the mean parental hostility scores may be somewhat restricted to the lower end, the level of hostility is expected. The nature of the observational task, which consists of the parent and child working on teaching tasks together, would not necessarily evoke extremely

high levels of parental hostility. Nevertheless, any hostility in this context appears to be meaningful.

Another possible limitation of this study is the way in which cortisol reactivity was indexed. As in other studies on young children (e.g., Luby et al., 2003, Gunnar et al., 2009a), the laboratory paradigm did not evoke increases in cortisol responses in all children, a circumstance that creates difficulty in interpreting the results in terms of cortisol reactivity (Gunnar et al., 2009a); nevertheless, variability in children's cortisol responses was observed, and this afforded the examination of how individual differences in children's cortisol responses, alone or in conjunction with early parenting, predict children's psychiatric symptoms across early childhood. In addition, the use of the AUC_i statistic as a measure of cortisol reactivity may obscure important nuances in the cortisol trajectories that may be important for understanding the emergence of psychopathology (e.g., individual differences in the initial engagement of the stress response or how prolonged the cortisol response is following the offset of the stressor). Lastly, the sample was largely comprised of White, two-parent, middle class families. Thus, we cannot generalize our results to more ethnically diverse samples, clinical samples, or those with extreme forms of parental hostility (i.e., maltreatment). Future research should extend this work to more diverse samples.

In summary, we found that children with high levels of cortisol reactivity to a laboratory stressor and who were exposed to high levels of parental hostility at age three demonstrated greater concurrent externalizing symptoms at age three and increases in internalizing and externalizing symptoms from age three to age six. Our findings suggest that preschoolers with increased cortisol reactivity represent a particularly vulnerable group of children when exposed to adverse parenting contexts, and might be targeted for early intervention to prevent the development of psychopathology. These interventions may be particularly effective during early childhood, a period of development characterized by a high degree of neuroplasticity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to all of the families who took part in the study and to the Stony Brook Temperament Study research team.

Financial support: This research was supported by R01MH069942 (Klein), F31MH075484 (Dougherty), F31MH084444 (Bufferd) and General Clinical Research Center grant M01RR10710 (Stony Brook University, National Center for Research Resources). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health or the National Institutes of Health.

References

Acock AC. Working with missing values. *Journal of Marriage and Family*. 2005; 67:1012–1028. DOI: 10.1111/j.1741-3737.2005.00191.x

- Aiken, L.S., West, S.G. Multiple regression: Testing and interpreting interactions. Newbury Park, CA: Sage; 1991.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Washington, DC: American Psychiatric Press; 2000.
- Badanes LS, Watamura SE, Hankin BL. Hypocortisolism as a potential marker of allostatic load in children: Associations with family risk and internalizing disorders. *Development and Psychopathology*. 2011; 23:881–896. DOI: 10.1017/S095457941100037X [PubMed: 21756439]
- Belsky, J. Differential susceptibility to rearing influences: An evolutionary hypothesis and some evidence. In: Ellis, B., Bjorklund, D., editors. *Origins of the social mind: Evolutionary psychology and child development*. New York: Guilford Press; 2005. p. 139–163.
- Belsky J, Pluess M. Beyond diathesis stress: Differential susceptibility to environmental influences. *Psychological Bulletin*. 2009; 135:885–908. DOI: 10.1037/a0017376 [PubMed: 19883141]
- Booij SH, Bouma EC, de Jonge P, Ormel J, Oldehinkel AJ. Chronicity of depressive problems and the cortisol response to psychosocial stress in adolescents: The TRAILS study. *Psychoneuroendocrinology*. 2013; 38:659–666. DOI: 10.1016/j.psyneuen.2012.08.004 [PubMed: 22963816]
- Boyce WT, Ellis BJ. Biological sensitivity to context: I. An evolutionary developmental theory of the origins and functions of stress reactivity. *Development and Psychopathology*. 2005; 17:271–301. DOI: 10.1017/S0954579405050145 [PubMed: 16761546]
- Bradley RH, Corwyn RF. Socioeconomic status and child development. *Annual Review of Psychology*. 2002; 53:371–399. DOI: 10.1146/annurev.psych.53.100901.135233
- Brennan PA, Pargas R, Walker E, Green P, Newport DJ, Stowe Z. Maternal depression and infant cortisol: Influences of timing, comorbidity and treatment. *Journal of Child Psychology and Psychiatry*. 2008; 49:1099–1107. DOI: 10.1111/j.1469-7610.2008.01914.x [PubMed: 18492036]
- Bufferd SJ, Dougherty LR, Carlson GA, Klein DN. Parent-reported mental health in preschoolers: Findings using a diagnostic interview. *Comprehensive Psychiatry*. 2011; 52:359–369. DOI: 10.1016/j.comppsy.2010.08.006 [PubMed: 21683173]
- Bufferd SJ, Dougherty LR, Carlson GA, Rose S, Klein DN. Psychiatric disorders in preschoolers: Continuity from ages 3 to 6. *American Journal of Psychiatry*. 2012; 169:1157–1164. DOI: 10.1176/appi.ajp.2012.12020268 [PubMed: 23128922]
- Carr CP, Martins CM, Stingel AM, Lemgruber VB, Juruena MF. The role of early life stress in adult psychiatric disorders: A systematic review according to childhood trauma subtypes. *The Journal of Nervous and Mental Disease*. 2013; 201:1007–1020. DOI: 10.1097/NMD.0000000000000049 [PubMed: 24284634]
- Chronis-Tuscano A, Molina BG, Pelham WE, Applegate B, Dahlke A, Overmyer M, Lahey B. Very early predictors of adolescent depression and suicide attempts in children attention-deficit/hyperactivity disorder. *Archives of General Psychiatry*. 2010; 67:1044–1051. DOI: 10.1001/archgenpsychiatry.2010.127 [PubMed: 20921120]
- de Veld DM, Riksen-Walraven JM, de Weerth C. The relation between emotion regulation strategies and physiological stress responses in middle childhood. *Psychoneuroendocrinology*. 2012; 37:1309–1319. DOI: 10.1016/j.psyneuen.2012.01.004 [PubMed: 22309825]
- Dedovic K, Duchesne A, Andrews J, Engert V, Pruessner JC. The brain and the stress axis: the neural correlates of cortisol regulation in response to stress. *Neuroimage*. 2009; 47:864–871. DOI: 10.1016/j.neuroimage.2009.05.074 [PubMed: 19500680]
- Dickerson SS, Kemeny ME. Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin*. 2004; 130:355–391. DOI: 10.1037/0033-2909.130.3.355 [PubMed: 15122924]
- Dougherty LR, Bufferd SJ, Carlson GA, Dyson M, Olinio TM, Durbin EC, Klein DN. Preschoolers' observed temperament and psychiatric disorders assessed with a parent diagnostic interview. *Journal of Clinical Child & Adolescent Psychology*. 2011a; 40:295–306. DOI: 10.1080/15374416.2011.546046 [PubMed: 21391025]
- Dougherty LR, Klein DN, Congdon E, Canli T, Hayden EP. Interaction between 5-HTTLPR and BDNF Val66Met polymorphisms on HPA axis reactivity in preschoolers. *Biological Psychology*. 2010; 83:93–100. DOI: 10.1016/j.biopsycho.2009.10.009 [PubMed: 19914329]

- Dougherty LR, Klein DN, Rose S, Laptook RS. Hypothalamic-pituitary-adrenal axis reactivity in the preschool-age offspring of depressed parents: Moderation by early parenting. *Psychological Science*. 2011b; 22:650–658. DOI: 10.1177/0956797611404084 [PubMed: 21460339]
- Dougherty LR, Leppert KA, Merwin SM, Smith VC, Bufferd SJ, Kushner MR. Advances and directions in preschool mental health research. *Child Development Perspectives*. 2015a; 9:14–19. DOI: 10.1111/cdep.12099
- Dougherty LR, Smith VC, Bufferd SJ, Kessel E, Carlson GA, Klein DN. Preschool irritability predicts child psychopathology, functional impairment, and service use at age nine. *Journal of Child Psychology and Psychiatry*. 2015b; 56:999–1007. DOI: 10.1111/jcpp.12403 [PubMed: 26259142]
- Dougherty LR, Smith VC, Bufferd SJ, Stringaris A, Leibenluft E, Carlson GA, Klein DN. Preschool irritability: Longitudinal associations with psychiatric disorders at age 6 and parental psychopathology. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2013a; 52:1304–1313. DOI: 10.1016/j.jaac.2013.09.007 [PubMed: 24290463]
- Dougherty LR, Tolep MR, Smith VC, Rose S. Early exposure to parental depression and parenting: Associations with young offspring's stress physiology and oppositional behavior. *Journal of Abnormal Child Psychology*. 2013b; 41:1299–1310. DOI: 10.1007/s10802-013-9763-7 [PubMed: 23722864]
- Doussard-Roosevelt JA, Montgomery LA, Porges SW. Short-term stability of physiological measures in kindergarten children: Respiratory sinus arrhythmia, heart period, and cortisol. *Developmental Psychobiology*. 2003; 43:230–242. DOI: 10.1002/dev.10136 [PubMed: 14558045]
- Dunn, LM., Dunn, LM. Peabody Picture Vocabulary Test. 3. Circle Pines, Minnesota: American Guidance Service; 1997.
- Egeland, B., Weinfield, N., Hiester, M., Lawrence, C., Pierce, S., Chippendale, K. Teaching tasks administration and scoring manual. University of Minnesota; 1995.
- Egger HL, Angold A. Common emotional and behavioral disorders in preschool children: Presentation, nosology, and epidemiology. *Journal of Child Psychology and Psychiatry*. 2006; 47:313–337. DOI: 10.1111/j.1469-7610.2006.01618.x [PubMed: 16492262]
- Egger, HL., Ascher, BH., Angold, A. The Preschool Age Psychiatric Assessment: Version 1.1. Durham, NC: Center for Developmental Epidemiology, Department of Psychiatry and Behavioral Sciences, Duke University Medical Center; 1999.
- Egger HL, Erkanli A, Keeler G, Potts E, Walter BK, Angold A. Test-retest reliability of the Preschool Age Psychiatric Assessment (PAPA). *Journal of the American Academy of Child & Adolescent Psychiatry*. 2006; 45:538–549. DOI: 10.1097/01.chi.0000205705.71194.b8 [PubMed: 16601400]
- Essex MJ, Armstrong JM, Burk LR, Goldsmith HH, Boyce WT. Biological sensitivity to context moderates the effects of the early teacher–child relationship on the development of mental health by adolescence. *Development and Psychopathology*. 2011; 23:149–161. DOI: 10.1017/S0954579410000702 [PubMed: 21262045]
- Gadow, KD., Sprafkin, J. Early childhood inventory-4 screening manual. Stony Brook, NY: Checkmate Plus; 2000.
- Goldsmith HH, Reilly J, Lemery KS, Longley S, Prescott A. The Laboratory Temperament Assessment Battery: Preschool version. 1995 Unpublished manuscript.
- Gotlib, IH., Joormann, J., Minor, KL., Cooney, RE. Cognitive and biological functioning in children at risk for depression. In: Canli, T., Canli, T., editors. *Biology of Personality and Individual Differences*. New York, NY, US: Guilford Press; 2006. p. 353-382.
- Granger DA, Weisz JR, McCracken JT, Ikeda SC, Douglas P. Reciprocal influences among adrenocortical activation, psychosocial processes, and the behavioral adjustment of clinic-referred children. *Child Development*. 1996; 76:3250–3262. DOI: 10.2307/1131777
- Gunnar, MR., Talge, NM. Neuroendocrine measures in developmental research. In: Schmidt, LA., Segalowitz, SJ., editors. *Developmental Psychophysiology: Theory, Systems, and Methods*. Cambridge, England: Cambridge University Press; 2007. p. 343-366.
- Gunnar MR, Talge NM, Herrera A. Stressor paradigms in developmental studies: What does and does not work to produce mean increases in salivary cortisol. *Psychoneuroendocrinology*. 2009a; 34:953–967. DOI: 10.1016/j.psyneuen.2009.02.010 [PubMed: 19321267]

- Gunnar, MR., Vazquez, DM. Stress neurobiology and developmental psychopathology. In: Cicchetti, D., Cohen, DJ., editors. *Developmental psychopathology, Vol 2: Developmental neuroscience*. 2. Hoboken, NJ: John Wiley; 2006. p. 533-577.
- Gunnar MR, Wewerka S, Frenn K, Long JD, Griggs C. Developmental changes in hypothalamus–pituitary–adrenal activity over the transition to adolescence: normative changes and associations with puberty. *Development and Psychopathology*. 2009b; 21:69–85. DOI: 10.1017/S0954579409000054 [PubMed: 19144223]
- Hagan MJ, Roubinov DS, Mistler AK, Luecken LJ. Mental health outcomes in emerging adults exposed to childhood maltreatment: The moderating role of stress reactivity. *Child Maltreatment*. 2014; 19:156–167. DOI: 10.1177/1077559514539753 [PubMed: 24920249]
- Hankin BL, Badanes LS, Abela JR, Watamura SE. Hypothalamic–pituitary–adrenal axis dysregulation in dysphoric children and adolescents: Cortisol reactivity to psychosocial stress from preschool through middle adolescence. *Biological psychiatry*. 2010; 68:484–490. DOI: 10.1016/j.biopsych.2010.04.004 [PubMed: 20497900]
- Harkness KL, Stewart JG, Wynne-Edwards KE. Cortisol reactivity to social stress in adolescents: Role of depression severity and child maltreatment. *Psychoneuroendocrinology*. 2011; 36:173–181. DOI: 10.1016/j.psyneuen.2010.07.006 [PubMed: 20688438]
- Hastings PD, Ruttle PL, Serbin LA, Mills RS, Stack DM, Schwartzman AE. Adrenocortical responses to strangers in preschoolers: Relations with parenting, temperament, and psychopathology. *Developmental Psychobiology*. 2011; 53:694–710. DOI: 10.1002/dev.20545 [PubMed: 21432849]
- Hayes AF, Matthes J. Computational procedures for probing interactions in OLS and logistic regression: SPSS and SAS implementations. *Behavioral Research Methods*. 2009; 41:924–936. DOI: 10.3758/BRM.41.3.924
- Humphreys KL, Zeanah CH. Deviations from the expectable environment in early childhood and emerging psychopathology. *Neuropsychopharmacology*. 2015; 40:154–170. DOI: 10.1038/npp.2014.165 [PubMed: 24998622]
- Johnson PO, Fay LC. The Johnson–Neyman technique, its theory and application. *Psychometrika*. 1950; 15:349–367. DOI: 10.1038/npp.2014.165 [PubMed: 14797902]
- Kryski KR, Dougherty LR, Dyson MW, Olino TM, Laptook RS, Klein DN, Hayden EP. Effortful control and parenting: Associations with HPA axis reactivity in early childhood. *Developmental Science*. 2013a; 16:531–541. DOI: 10.1111/desc.12050 [PubMed: 23786471]
- Kryski KR, Smith HJ, Sheikh HI, Singh SM, Hayden EP. HPA axis reactivity in early childhood: Associations with symptoms and moderation by sex. *Psychoneuroendocrinology*. 2013b; 38(10): 2327–2336. [PubMed: 23764193]
- Kryski KR, Smith HJ, Sheikh HI, Singh SM, Hayden EP. Assessing stress reactivity indexed via salivary cortisol in preschool-aged children. *Psychoneuroendocrinology*. 2011; 36:1127–1136. DOI: 10.1016/j.psyneuen.2011.02.003 [PubMed: 21388745]
- Kushner M, Barrios C, Smith VC, Dougherty LR. Physiological and behavioral vulnerability markers increase risk to early life stress in preschool-aged children. *Journal of Abnormal Child Psychology*. 2016; 44:859–870. DOI: 10.1007/s10802-015-0087-7 [PubMed: 26424217]
- Lahey BB, Pelham WE, Loney J, Kipp H, Ehrhardt A, Lee SL, ... Massetti G. Three-year predictive validity of DSM-IV attention deficit hyperactivity disorder in children diagnosed at 4–6 years of age. *American Journal of Psychiatry*. 2004; 161:2014–2020. DOI: 10.1176/appi.ajp.161.11.2014 [PubMed: 15514401]
- Lam S, Dickerson SS, Zoccola PM, Zaldivar F. Emotion regulation and cortisol reactivity to a social-evaluative speech task. *Psychoneuroendocrinology*. 2009; 34:1355–1362. DOI: 10.1016/j.psyneuen.2009.04.006 [PubMed: 19464808]
- Latimer K, Wilson P, Kemp J, Thompson L, Sim F, Gillberg C, ... Minnis H. Disruptive behaviour disorders: A systematic review of environmental antenatal and early years risk factors. *Child: Care, Health and Development*. 2012; 38:611–628. DOI: 10.1111/j.1365-2214.2012.01366.x
- Lavigne JV, Arend R, Rosenbaum D, Binns HJ, Christoffel KK, Gibbons RD. Psychiatric disorders with onset in the preschool years: I. Stability of diagnoses. *Journal Of The American Academy Of Child & Adolescent Psychiatry*. 1998; 37:1246–1254. DOI: 10.1097/00004583-199812000-00007 [PubMed: 9847496]

- Lavigne JV, Cicchetti C, Gibbons RD, Binns HJ, Larsen L, DeVito C. Oppositional defiant disorder with onset in preschool years: Longitudinal stability and pathways to other disorders. *Journal Of The American Academy Of Child & Adolescent Psychiatry*. 2001; 40:1393–1400. DOI: 10.1097/00004583-200112000-00009 [PubMed: 11765284]
- Luby JL, Gaffrey MS, Tillman R, April LM, Belden AC. Trajectories of preschool disorders to full DSM depression at school age and early adolescence: Continuity of preschool depression. *American Journal of Psychiatry*. 2014; 171:768–776. DOI: 10.1176/appi.ajp.2014.13091198 [PubMed: 24700355]
- Luby JL, Heffelfinger A, Mrakotsky C, Brown K, Hessler M, Spitznagel E. Alterations in stress cortisol reactivity in depressed preschoolers relative to psychiatric and no-disorder comparison groups. *Archives of General Psychiatry*. 2003; 60:1248–1255. DOI: 10.1001/archpsyc.60.12.1248 [PubMed: 14662557]
- Luby JL, Si X, Belden AC, Tandon M, Spitznagel E. Preschool depression: Homotypic continuity and course over 24 months. *Archives of General Psychiatry*. 2009; 66:897–905. DOI: 10.1001/archgenpsychiatry.2009.97 [PubMed: 19652129]
- Lyneham HJ, Rapee RM. Agreement between telephone and in-person delivery of a structured interview for anxiety disorders in children. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2005; 44:274–282. DOI: 10.1097/00004583-200503000-00012 [PubMed: 15725972]
- McKee L, Colletti C, Rakow A, Jones DJ, Forehand R. Parenting and child externalizing behaviors: Are the associations specific or diffuse? *Aggression and Violent Behavior*. 2008; 13:201–215. DOI: 10.1016/j.avb.2008.03.005 [PubMed: 19122818]
- McLaughlin KA, King K. Developmental trajectories of anxiety and depression in early adolescence. *Journal of Abnormal Child Psychology*. 2015; 43:311–323. DOI: 10.1007/s10802-014-9898-1 [PubMed: 24996791]
- McLeod BD, Weisz JR, Wood JJ. Examining the association between parenting and childhood depression: A meta-analysis. *Clinical Psychology Review*. 2007; 27:986–1003. DOI: 10.1016/j.cpr.2007.03.001 [PubMed: 17449154]
- Mesman J, Bongers IL, Koot HM. Preschool developmental pathways to preadolescent internalizing and externalizing problems. *Journal of Child Psychology and Psychiatry*. 2001; 42:679–689. DOI: 10.1111/1469-7610.00763 [PubMed: 11464972]
- Monroe SM, Reid MW. Life stress and major depression. *Current Directions in Psychological Science*. 2009; 18:68–72. DOI: 10.1111/j.1467-8721.2009.01611.x
- Monroe SM, Simons AD. Diathesis–stress theories in the context of life stress research: Implications for the depressive disorders. *Psychological Bulletin*. 1991; 110:406–425. DOI: 10.1037/0033-2909.110.3.406 [PubMed: 1758917]
- Morgan JK, Shaw DS, Olino TM. Differential susceptibility effects: The interaction of negative emotionality and sibling relationship quality on childhood internalizing problems and social skills. *Journal of Abnormal Child Psychology*. 2012; 40:885–899. DOI: 10.1007/s10802-012-9618-7 [PubMed: 22366882]
- Morris CM, Rao U, Garber J. Cortisol responses to psychosocial stress predict depression trajectories: Social-evaluative threat and prior depressive episodes as moderators. *Journal of Affective Disorders*. 2012; 143:223–230. DOI: 10.1016/j.jad.2012.05.059 [PubMed: 22858210]
- Murri M, Pariante C, Mondelli V, Masotti M, Atti AR, Mellacqua A, ... Amore M. HPA axis and aging in depression: Systematic review and meta-analysis. *Psychoneuroendocrinology*. 2014; 41:46–62. DOI: 10.1016/j.psyneuen.2013.12.004 [PubMed: 24495607]
- Navarro, Pastor JB. Methods for the analysis of explanatory linear regression models with missing data not at random. *Quality and Quantity*. 2003; 37:363–376. DOI: 10.1023/A:1027323122628
- Obrodovi J, Bush NR, Stamperdahl J, Adler NE, Boyce WT. Biological sensitivity to context: The interactive effects of stress reactivity and family adversity on socio-emotional behavior and school readiness. *Child Development*. 2010; 81:270–289. DOI: 10.1111/j.1467-8624.2009.01394.x [PubMed: 20331667]

- Okado Y, Bierman KL. Differential risk for late adolescent conduct problems and mood dysregulation among children with early externalizing behavior problems. *Journal of Abnormal Child Psychology*. 2015; 43:735–747. [PubMed: 25183553]
- Olino TM, Dougherty LR, Bufferd SJ, Carlson GA, Klein DN. Testing models of psychopathology in preschool-aged children using a structured interview-based assessment. *Journal of Abnormal Child Psychology*. 2014; 42:1201–1211. DOI: 10.1007/s10802-014-9865-x [PubMed: 24652485]
- Olino TM, Klein DN, Dyson MW, Rose SA, Durbin EC. Temperamental emotionality in preschool-aged children and depressive disorders in parents: Associations in a large community sample. *Journal of Abnormal Psychology*. 2010; 119:468–478. DOI: 10.1037/a0020112 [PubMed: 20677836]
- O’Neal CR, Brotman LM, Huang KY, Gouley KK, Kamboukos D, Calzada EJ, Pine DS. Understanding relations among early family environment, cortisol response, and child aggression via a prevention experiment. *Child Development*. 2010; 81:290–305. DOI: 10.1111/j.1467-8624.2009.01395.x [PubMed: 20331668]
- Pariante CM. Depression, stress, and the adrenal axis. *Journal of Neuroendocrinology*. 2003; 15(8): 811–812. [PubMed: 12834443]
- Pruessner JC, Kirschbaum C, Meinlschmid G, Hellhammer DH. Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*. 2003; 28:916–931. DOI: 10.1016/S0306-4530(02)00108-7 [PubMed: 12892658]
- Rao U, Hammen C, Ortiz LR, Chen LA, Poland RE. Effects of early and recent adverse experiences on adrenal response to psychosocial stress in depressed adolescents. *Biological Psychiatry*. 2008; 64(6):521–526. DOI: 10.1016/j.biopsych.2008.05.012 [PubMed: 18597740]
- Repetti RL, Taylor SE, Seeman TE. Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*. 2002; 128:330–366. DOI: 10.1037/0033-2909.128.2.330 [PubMed: 11931522]
- Saxbe DE, Margolin G, Spies Shapiro LA, Baucom BR. Does dampened physiological reactivity protect youth in aggressive family environments? *Child Development*. 2012; 83:821–830. DOI: 10.1111/j.1467-8624.2012.01752.x [PubMed: 22548351]
- Sprafkin J, Volpe RJ, Gadow KD, Nolan EE, Kelly K. A DSM-IV referenced screening instrument for preschool children: The Early Childhood Inventory-4. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2002; 41:604–612. DOI: 10.1097/00004583-200205000-00018 [PubMed: 12014793]
- Steege CM, Cook EC, Connell CM. The interactive effects of stressful family life events and cortisol reactivity on adolescent externalizing and internalizing behaviors. *Child Psychiatry & Human Development*. 2016; Advance online publication. doi: 10.1007/s10578-016-0635-6
- Sterba S, Egger HL, Angold A. Diagnostic specificity and nonspecificity in the dimensions of preschool psychopathology. *Journal of Child Psychology and Psychiatry*. 2007; 48:1005–1013. DOI: 10.1111/j.1469-7610.2007.01770.x [PubMed: 17915001]
- Susman EJ, Dorn LD, Inoff-Germain G, Nottelmann ED, Chrousos GP. Cortisol reactivity, distress behavior, and behavioral psychological problems in young adolescents: A longitudinal perspective. *Journal of Research on Adolescence*. 1997; 7:81–105. DOI: 10.1207/s15327795jra0701_5
- Talge NM, Donzella B, Gunnar MR. Fearful temperament and stress reactivity among preschool-aged children. *Infant and Child Development*. 2008; 17:427–445. DOI: 10.1002/icd.585 [PubMed: 19122850]
- Talge NM, Donzella B, Kryzer EM, Gierens A, Gunnar MR. It’s not that bad: Error introduced by oral stimulants in salivary cortisol research. *Developmental Psychobiology*. 2005; 47:369–376. DOI: 10.1002/dev.20097 [PubMed: 16284967]
- van De Wiel NH, van Goozen SM, Matthys W, Snoek H, van Engeland H. Cortisol and treatment effect in children with disruptive behavior disorders: A preliminary study. *Journal of The American Academy of Child & Adolescent Psychiatry*. 2004; 43:1011–1018. DOI: 10.1097/01.chi.0000126976.56955.43 [PubMed: 15266196]

- van Leeuwen AP, Creemers HE, Greaves-Lord K, Verhulst FC, Ormel J, Huizink AC. Hypothalamic–pituitary–adrenal axis reactivity to social stress and adolescent cannabis use: the TRAILS study. *Addiction*. 2011; 106:1484–1492. DOI: 10.1111/j.1360-0443.2011.03448.x [PubMed: 21631618]
- von Klitzing KV, Perren S, Klein AM, Stadelmann S, White LO, Groeben M, et al. The interaction of social risk factors and HPA axis dysregulation in predicting emotional symptoms of five- and six-year-old children. *Journal of Psychiatric Research*. 2012; 46:290–297. DOI: 10.1016/j.jpsychires.2011.12.004 [PubMed: 22226637]
- Wichstrøm L, Berg-Nielsen TS, Angold A, Egger HL, Solheim E, Sveen TH. Prevalence of psychiatric disorders in preschoolers. *Journal of Child Psychology and Psychiatry*. 2012; 53:695–705. DOI: 10.1111/j.1469-7610.2011.02514.x [PubMed: 22211517]

Figure 1. The interaction between child cortisol reactivity and observed parental hostility at age three predicts increases in children's internalizing symptoms at age six (error bars included).

Figure 2.

The interaction between child cortisol reactivity and observed parental hostility at age three predicts increases in children's externalizing symptoms at age six (error bars included).

Table 1

Characteristics of the Study Sample

	Age 3		Age 6	
<i>Demographic Characteristics</i>				
Child mean age: years (SD)	3.58	(.24)	6.33	(.34)
Child sex: female <i>n</i> (%)	80	(50)		
Child race: <i>n</i> (%)				
White	153	(95.6)		
Black/African-American	3	(1.9)		
Asian	3	(1.9)		
Other	1	(.6)		
Child Hispanic ethnicity: <i>n</i> (%)	16	(10)		
Biological parents' marital status: <i>n</i> (%)				
Married	150	(93.8)		
Divorced, separated, or widowed	2	(1.3)		
Never married	7	(4.4)		
Parents' education: graduated college <i>n</i> (%)				
Mother	83	(51.9)		
Father	76	(47.5)		
<i>Salivary cortisol indicators (nmol/L)</i>				
Mean cortisol level at sample 1 (SD)	4.55	(7.11)		
Mean cortisol level at sample 2 (SD)	3.95	(5.97)		
Mean cortisol level at sample 3 (SD)	3.91	(4.23)		
Mean cortisol level at sample 4 (SD)	5.76	(5.53)		
Mean <i>AUCi</i> (SD)	.36	(12.90)		
<i>AUCi</i> positive <i>n</i> (%)	98	(62.8%)		
<i>AUCi</i> negative <i>n</i> (%)	58	(37.2%)		
<i>Mean child symptom scales (SD); Range^a</i>				
Internalizing symptoms	8.18 (6.30)	0–41	14.41 (10.55)	0–51
Externalizing symptoms	14.62 (8.96)	0–65.93	5.60 (6.57)	0–50
<i>Child psychiatric diagnoses <i>n</i> (%)</i>				
At least 1 DSM diagnosis	38	(23.8)	37	(23.1)
At least 1 Emotional diagnosis	27	(16.9)	22	(13.8)
At least 1 Behavioral diagnosis	14	(8.8)	19	(11.9)
Any anxiety disorder diagnosis	25	(15.6)	19	(11.9)
Any depressive disorder diagnosis	3	(1.9)	5	(3.1)
ADHD diagnosis	4	(2.5)	9	(5.6)
ODD diagnosis	13	(8.1)	14	(8.8)

Note:

^aThe total possible scores for the PAPA internalizing symptoms scale range from 0 to 254, and the total possible scores for the PAPA externalizing symptoms scale range from 0 to 50.

Table 2**Correlations Among All Study Variables**

	1	2	3	4	5	6	7	8	9
1. Age 3 child cortisol reactivity	-								
2. Age 3 observed parental hostility	.12	-							
3. Age 3 internalizing symptoms	.08	-.05	-						
4. Age 3 externalizing symptoms	.22**	.34**	.27**	-					
5. Age 6 internalizing symptoms	.12	.11	.50**	.22*	-				
6. Age 6 externalizing symptoms	.23**	.44**	.32**	.64**	.44**	-			
7. Child age at baseline	.09	-.02	.02	.09	.11	.18*	-		
8. Child gender	-.07	-.05	.01	.12	.05	-.05	.01	-	
N	156	149	153	153	135	135	160	160	160

Note: Child age reported in months at baseline (age three assessment); Child Gender: male = 1 ($n = 80$) and female = 2 ($n = 80$);

* $p < .05$,

** $p < .01$.

Table 3

Early Childhood Cortisol Reactivity, Parental Hostility, and Internalizing Symptoms at Ages Three and Six Years

	Age 3 Internalizing Symptoms		Age 6 Internalizing Symptoms	
	b (SE)	B	b (SE)	B
Age 3 internalizing symptoms	--	--	6.21 (1.02)	.48***
Age 3 externalizing symptoms	--	--	-.39 (1.06)	-.03
Age 3 child cortisol reactivity	.39 (1.09)	.03	-.01 (1.54)	.00
Age 3 parental hostility	-1.31 (.84)	-.16	-.42 (1.29)	-.03
Age 3 parental hostility $\times$ cortisol reactivity	1.48 (.79)	.20 [†]	3.22 (1.20)	.28**

Note:

[†]
 $p < .10$;

*
 $p < .05$;

**
 $p < .01$;

 $p < .001$.

Table 4

Early Childhood Cortisol Reactivity, Parental Hostility, and Externalizing Symptoms at Ages Three and Six Years

	Age 3 Externalizing Symptoms		Age 6 Externalizing Symptoms	
	b (SE)	B	b (SE)	B
Age 3 externalizing symptoms	--	--	2.95 (.54)	.40***
Age 3 internalizing symptoms	--	--	1.29 (.52)	.16*
Age 3 child cortisol reactivity	1.49 (1.34)	.09	.03 (.78)	.00
Age 3 parental hostility	1.32 (1.04)	.12	.90 (.65)	.11
Age 3 parental hostility $\times$ cortisol reactivity	3.81 (.98)	.37***	2.31 (.61)	.32***

Note:

* $p < .05$;

** $p < .01$;

*** $p < .001$.