Video Article

Single-molecule Manipulation of G-quadruplexes by Magnetic Tweezers

Huijuan You^{*1}, Shimin Le^{*2,3}, Hu Chen^{2,4}, Linyan Qin¹, Jie Yan^{2,4,5}

¹School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology

²Mechanobiology Institute, National University of Singapore

³Department of Physics, National University of Singapore

⁴Research Institute for Biomimetics and Soft Matter, Fujian Provincial Key Lab for Soft Functional Materials Research, Department of Physics, Xiamen University

⁵Center for Bioimaging Sciences, National University of Singapore

*These authors contributed equally

Correspondence to: Huijuan You at youhuijuan@hust.edu.cn, Jie Yan at phyjy@nus.edu.sg

URL: <https://www.jove.com/video/56328>

DOI: [doi:10.3791/56328](https://doi.org/10.3791/56328)

Keywords: Biochemistry, Issue 127, Single-molecule manipulation, magnetic tweezers, G-quadruplexes, folding-unfolding, helicase

Date Published: 9/19/2017

Citation: You, H., Le, S., Chen, H., Qin, L., Yan, J. Single-molecule Manipulation of G-quadruplexes by Magnetic Tweezers. *J. Vis. Exp.* (127), e56328, doi:10.3791/56328 (2017).

Abstract

Non-canonical nucleic acid secondary structure G-quadruplexes (G4) are involved in diverse cellular processes, such as DNA replication, transcription, RNA processing, and telomere elongation. During these processes, various proteins bind and resolve G4 structures to perform their function. As the function of G4 often depends on the stability of its folded structure, it is important to investigate how G4 binding proteins regulate the stability of G4. This work presents a method to manipulate single G4 molecules using magnetic tweezers, which enables studies of the regulation of G4 binding proteins on a single G4 molecule in real time. In general, this method is suitable for a wide scope of applications in studies for proteins/ligands interactions and regulations on various DNA or RNA secondary structures.

Video Link

The video component of this article can be found at <https://www.jove.com/video/56328/>

Introduction

Four-stranded DNA or RNA G4 structures play critical roles in many important biological processes¹. Many proteins are involved in G4 binding and regulation, including telomere binding proteins (telomerase, POT1, RPA, TEBPs, TRF2)^{1,2}, transcription factors (nucleolin, PARP1)³, RNA processing proteins (hnRNP A1, hnRNP A2)⁴, helicases (BLM, FANCD1, RPA, WRN, Dna2, Pif1)⁵, and DNA replication related proteins (Rif1, REV1, PrimPol)⁶. Protein binding can stabilize or destabilize G4 structures; thus regulating the subsequent biological functions. The stability of G4 was measured by thermal melting using ultraviolet (UV) or circular dichroism (CD) methods⁷. However, such conditions are not physiological relevant and are difficult to apply to studying the effects of binding proteins⁷.

The rapid development in single-molecule manipulation technologies has enabled studies of folding and unfolding of a biomolecule, such as a DNA or a protein, at a single-molecule level with nanometer resolution in real time⁸. Atomic force microscopy (AFM), optical tweezers, and magnetic tweezers are the most commonly used single-molecule manipulation methods. Compared to AFM and optical tweezers⁹, magnetic tweezers allow stable measurements of folding-unfolding dynamics of a single molecule over days by using an anti-drift technique^{10,11}.

Here, a single-molecule manipulation platform using magnetic tweezers to study the regulation of G4 stability by binding proteins is reported^{12,13}. This work outlines the basic approaches, including sample and flow channel preparation, the setup of magnetic tweezers, and the force calibration. The force control and the anti-drift protocols as described in step 3 allow for long time measurements under various force controls, such as constant force (force clamp) and constant loading rate (force-ramp), and force-jump measurement. The force calibration protocol described in step 4 enables force calibration of < 1 μm short tethers over a wide force range up to 100 pN, with a relative error within 10%. An example of regulation of the stability of the RNA Helicase associated with AU-rich element (RHAU) helicase (alias DHX36, G4R1) that plays essential roles in resolving RNA G4 is used to demonstrate the applications of this platform¹³.

Protocol

1. Preparation of G4 DNA for Single-molecule Stretching

1. Prepare 5'-thiol labeled and 5'-biotin labeled dsDNA handles by PCR using DDNA polymerase on a lambda phage DNA template using 5'-thiol and 5'-biotin primers¹⁴ (**Figure 1**). Both dsDNA handles have high GC content (> 60%) to prevent DNA melting when DNA is held at high forces or during DNA overstretching transition¹⁵.

- Purify PCR products using a commercial purification kit and digest with BstXI restriction enzyme according to the manufacturer's protocol.
- Ligate G4 forming ssDNA, and flank ssDNA and dsDNA handles using T4 DNA ligase according to the manufacturer's protocol. Purify the ligated product by gel extraction using a commercial purification kit according to the manufacturer's protocol.

2. Preparation of Flow Channel

- Coverslip cleaning and surface functionalization
 - Place bottom coverslips (#1.5, 22 mm × 32 mm) and top coverslips (#1.5, 20 mm × 20 mm) into cover glass staining jars (each jar can hold 7 pieces of coverslips, the volume is ~ 20 mL). Rinse the coverslips in the jars with distilled water 2-5 times.
 - Add ~ 20 mL of 5-40% detergent solution into each jar, and then place in an ultrasonic cleaning bath for 30 min. Rinse with distilled water for > 10 times to remove the detergent.
 - Dry the coverslips in the jars in oven (~ 150 °C; **CAUTION**, hot), or by N₂ gas. Store the dried top coverslips in a dry cabinet.
 - Use plasma (O₂ gas) to clean the coverslips in the jar for 10 min. During the 10 min, prepare 20 mL of 1% (3-Aminopropyl)triethoxysilane (APTES) solution in methanol (**CAUTION**, toxic/flammable). Use a chemical fume hood to dissolve APTES in methanol.
NOTE: Avoid humidity when storing APTES solution. Old APTES often causes problem in surface functionalizing of coverslips.
 - Immediately after the plasma cleaning, add all of the 1% APTES solution into the jars and incubate for 1 h. Pour the waste into the waste bottle specific for 1% APTES methanol. Perform the methanol-related steps in the fume hood for flammable chemicals.
 - Rinse the jars once with methanol and > 10 times with distilled water, and then dry by oven (~150 °C; **CAUTION**, hot). Store the APTES-coated bottom coverslips in a dry cabinet if not in use for up to two weeks.
NOTE: After each sub-step from 2.1.1 to 2.1.4, the cleaning process can be paused before the next step.
- Assemble the flow channel
 - Prepare two "spacers", *i.e.*, parafilm or double-side tape (~ 4 mm × 20 mm) for each channel. Place the two pieces of spacers on a bottom coverslip along the long-edge. Place a top coverslip on the spacer, forming a flow cell in between (~ 10 mm × 20 mm area, **Figure 2A, B**).
 - If parafilm is used as a spacer, place the flow channel on a heater (60-120 °C; **CAUTION**, hot) for 5-10 s while gently pressing the sides of the top coverslip to stick the two coverslips together by parafilm. The resulting flow channel has a height of ~ 100 µm, and thereby the volume of the channel is ~ 20 µL.
 - Seal the long edge of the channel with silicone glue to avoid leakage. Use silicone glue to make a small sink-like structure at each open edge of the flow channel, which serves as an entry and an exit of solution.
NOTE: The entry and exit can also be made by other ways, *e.g.*, by adhering small plastic rings using wax.
 - Store channels in a dry cabinet for up to 4 weeks.
- Tether DNA onto the bottom surface of the flow channel
 - Dilute the amino-coated polystyrene beads (diameter: 3 µm) by 200X in distilled 1X phosphate buffered saline (PBS) buffer. Vortex the bead solution and then flow into channels. Incubate the bead solution in channels for ~ 30 min. Remove any unstuck beads by washing with 200 µL of 1X PBS buffer.
 - Adjust (increase/decrease) the incubation time to achieve a surface density of 1-5 beads per 50 mm × 50 mm area. Store the channels deposited with the reference beads for up to 3 days.
 - Dilute sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (Sulfo-SMCC) powder into 1X PBS solution (~ 0.5 mg/mL). Vortex the solution and then flow into the channel.
NOTE: Prepare fresh Sulfo-SMCC solution before use and add it immediately to the channel to avoid hydrolyzation of the SMCC.
 - Incubate the SMCC solution in the channel for 30 min. Remove the SMCC solution by washing with a large amount (1 mL, ~ 50X the channel volume) of 1X PBS solution.
NOTE: Wash the excess SMCC carefully. This step is very essential for the binding of DNA on the coverslip.
 - DNA tethering**
 - Dilute the thiol-biotin labeled DNA into 1X PBS, with a resulting DNA concentration of ~ 0.3 nM. Gently pipette to mix the solution, flow the DNA solution into the SMCC-coated channel, and incubate for 30 min at room temperature (~ 23 °C).
 - Gently wash away free DNA with 200 µL of blocking solution that contains 1X PBS with 10 mg/mL bovine serum albumin (BSA) and 0.01% 2-Mercaptoethanol.
 - Block the channel surface by incubating the channel in blocking solution (10 mg/mL BSA, 0.01% 2-Mercaptoethanol) for > 2 h; after this step, the channel is ready for experiments. The prepared channel can be kept at 4 °C for ~ 1 day.
NOTE: The blocking step is important for reducing the non-specific binding of DNA and magnetic beads to the coverslips.

3. Magnetic Tweezers Setup and Identification of Single dsDNA Tether

- Magnetic tweezers setup**
 - Start the magnetic tweezers control program. Here, the magnetic tweezers were controlled by an in-house-written LabVIEW program.
 - Align the magnet centers before mounting the channel. Use a 4X objective lens to adjust the x- and y-axis of the magnets in the optical axis of the microscope.
 - Use a computer-controlled motorized manipulator to move the magnets along the z-direction (**Figure 2A**) and set the distance as $d = 0$ (d : distance between the magnets and coverslip) when the magnets attach to a coverslip on the microscope.
 - Program the movement of the magnets through the manipulator to achieve control of the force, including constant force (*i.e.*, constant d) and time varying force $F(t)$ (*i.e.*, time varying $d(t)$).

5. Use bright a light-emitting diode (LED) light source for back-scattered illumination of the bead through the objective. Collect the bead images at a sampling rate of 100 Hz by a charge-coupled device (CCD) camera.
2. **Sample setup and identify single dsDNA tether**
 1. **Tether formation**
 1. Gently flow in 200X diluted M-280, paramagnetic beads in assay solution (100 mM KCl, 2 mM MgCl₂, 10 mM Tris-pH 7.4) into a channel. Incubate for 10 min to allow the beads to bind to biotin-labeled DNA molecules immobilized on the SMCC-coated surface through the thiol-labeled end. Gently wash away un-tethered beads using 200 µL of standard reaction solution.
NOTE: The M-280 beads show lower non-specific binding to the surface compared to other commercial magnetic beads such as M-270.
 2. Mount the channel onto the microscope stage. Search for the beads on the bottom surface using the 100X oil immersion objective.
 3. Select a reference bead on the surface and a moving tethered bead. Build the initial image libraries of both the reference bead and tethered bead at different defocus planes.
 4. **Calibration of beads image**
 1. Before experiments, use an objective piezo actuator to obtain images of both reference and tethered beads at different defocus planes spaced by 20-50 nm, which are stored as two separate bead image libraries and are respectively indexed with the defocus distance (**Figure 2D**).
 5. **x, y, z position determination**
 1. During experiments, determine the position of the bead in the x-y plane by the bead centroid. Determine the height change of the tethered bead by comparing the current bead image with those stored in the library. Use the auto-correlation function analysis of the power spectrum of the Fourier transform of the bead images¹⁶.
 6. **Anti-drift feedback using reference bead**
 1. During experiments, use the piezo to "lock" the distance between the objective and a specific reference bead image stored in the library through a low frequency feedback control, so that the stuck bead image has the best correlation with a specific image stored in the library (**Figure 2D**).
 7. Determine whether the tether is a single dsDNA molecule by applying ~ 65 pN force Determine a single dsDNA molecule if the tether undergoes the characteristic DNA overstretching transition^{17,18,19}. Repeat the process until a single dsDNA tether is found. (Please refer to the next section for force calibration and DNA overstretching transition).

4. Magnetic Tweezers Force Calibration

1. Directly determine the force (up to 100 pN) for long DNA (48,502 bp λ-DNA) molecules using bead fluctuation through:

$$F \approx \frac{k_B T}{\delta_y^2} (l + r_0)$$

, where k_B is the Boltzmann constant, T is the temperature in Kelvin scale, δ_y^2 is the variance of the bead fluctuation in the direction perpendicular to the magnetic field and the force direction. In this direction, the motion of the bead can be described as a pendulum fluctuation with a length of $l + r_0$, where l is the end-to-end distance along the force direction (extension) of the DNA, and r_0 is the radius of the bead¹⁰ (**Figure 3A**).

2. Determine the calibration curve for force F as a function of magnets to bead distance d , and $F(d)$ for different beads using the long λ-DNA. Usually, the distance between the magnet and the coverslip is used as the DNA tether length can be ignored. Fit the F - d curve by the double-exponential decay function:

$$F = C \left(\alpha_1 \exp\left(\frac{d}{\gamma_1}\right) + \alpha_2 \exp\left(\frac{d}{\gamma_2}\right) \right)$$

, where the fitting parameters α_1 , α_2 , γ_1 , γ_2 are determined by the magnetic tweezers and the parameter C is determined by the heterogeneous property of the magnetic beads. By shifting the C , the F - d curve obtained from different beads can be overlapped¹⁰ (**Figure 3B**).

3. **Calibration of parameter C for individual magnetic beads for short DNA experiments.**

1. Determining the force based on the fluctuation requires recording the bead position at a frequency higher than the corner frequency:

$$f_c = \frac{F}{2\pi(l + r_0)\gamma}$$

, where γ is the drag coefficient of the bead. For short DNA at high force, it requires recording frequency faster than 100 Hz, therefore, the force cannot be directly measured based on fluctuation with the camera. However, the parameter C can be determined by measuring the force at a low force range (< 15 pN) using fluctuation and fitting the data with a calibrated F - d curve to obtain the parameter C ²⁰.

2. Alternatively, for experiments with dsDNA, determine the parameter C using DNA overstretching transition at a force of ~ 65 pN (**Figure 3C**)^{21,22,23} by recording the d position at which dsDNA showed extension jump (0.6X length increase of contour length). After determining the parameter C , calculate the force for the tethered beads.
4. Control the loading rate by programming the movement of the magnets through the inverse function of $F(d)$. The magnet should approach the sample double exponentially.

NOTE: The relative error of force determined by such an extrapolation method is ~ 10%, and is mainly caused by the bead radius heterogeneity²⁰.

5. Single-molecule Manipulation of G4 in the Presence and Absence of Binding Proteins

1. Force-ramp experiments

1. After identifying the dsDNA tether, perform a force-increase scan at a loading rate of 0.2 pN/s followed by a force-decrease scan at -0.2 pN/s. After each stretching cycle, hold the DNA molecule at 1 pN for 30 s to allow the ssDNA to refold to G4.

2. Force-jump experiments

1. Cycle the force between 54 pN for 30 s under which a folded G4 could be unfolded, and 1 pN for 60s, under which a folded G4-15T could refold.

3. Flowing protein solution

1. Flow the protein solution when at ~ 20 pN force to avoid attachment of the beads to the coverslip. The DEAH-box RHAU helicase, which showed high specificity to the G4 structure, was used²⁴. The recombinant *Drosophila melanogaster* RHAU (DmRHAU) helicase was expressed in *Escherichia coli* and purified as described previously²⁵. The G4 unwinding activity of DmRHAU was assayed on a tetramolecular G4 DNA substrate¹³.

4. Analyze the unfolding force using an in-house written Matlab program¹⁴.

Representative Results

The experiment setup for stretching a single G4 molecule is shown in **Figure 4**. A single-stranded G4 forming sequence spanned between two dsDNA handles was tethered between a coverslip and a paramagnetic bead. To find a single dsDNA tethered bead, an overstretching assay was performed by increasing the force at constant loading rates. Three types of measurements were often used for studying the folding and unfolding of biomolecules: (i) constant force measurement, (ii) force-ramp measurement, and (iii) force-jump measurement. Due to the extremely slow unfolding rates of this G4 structure, the equilibrium folding-unfolding transition could only occur for the time over days, so force-ramp and force-jump measurements were used for characterizing the unfolding kinetics and stability of G4 structures.

Force-ramp Measurements for G4 Unfolding in the Presence or Absence of RHAU Protein:

Figure 4B shows typical force-extension curves obtained by a force-increase scan at a loading rate of 0.2 pN/s followed by a force-decrease scan at -0.2 pN/s. After each stretching cycle, the DNA molecule was held at 1 pN for 30 s to allow ssDNA to refold to G4. The extension jump in the force-increase scan indicated an unfolding transition of G4. When using a non-G4 forming sequence, the sudden extension jump cannot be observed. The unfolding force distribution could be obtained by recording the forces at which unfolding occurred. The unfolding force distribution of G4-15T obtained at 0.2 pN/s showed a single peak at ~ 52 pN. However, in the presence of 10 nM RHAU, G4-15T remained folded during the force-increase scan for up to 60 pN, indicating G4 stabilization by RHAU binding. In the presence of 10 nM RHAU and 1 mM ATP, the unfolding force distribution was shifted to a lower force, indicative of an ATP-dependent destabilization of G4 by RHAU.

Force-jump Measurements of G4 Unfolding in the Presence or Absence of RHAU Protein:

Figure 5 shows a representative trace of the bead height during four force-jump cycles (**Figure 5**, left panel); the force applied to the molecule was cycled between 54 pN for 30 s under which a folded G4 could be unfolded, and 1 pN for 60 s under which a folded G4-15T could refold. The average lifetime of folded G4-15T at 54 pN was ~ 6.4 s, estimated by fitting the lifetime histogram with an exponential decay function¹³. After flowing in a solution of 10 nM RHAU helicase without ATP, the extension of tethered DNA remained at a folded level throughout the 30 s holding time at 54 pN, indicating that the RHAU strongly stabilizes the G4-15T structure against mechanical unfolding in the absence of ATP. After flowing in 10 nM RHAU and 1 mM ATP, the extension of the same molecule right after jumping from 1 pN to 54 pN was at the level of unfolded ssDNA, indicating that the RHAU destabilizes the G4 structure in the presence of ATP.

Figure 1: Preparation of G4 DNA for single-molecule stretching experiments.

The dsDNA handles were prepared by PCR using 5'-thiol and 5'-biotin primers. PCR products were purified and digested with the BstXI restriction enzyme and were purified by gel extraction. G4 forming ssDNA, two flank ssDNA and dsDNA handle were ligated using T4 DNA ligase. The ligated product was purified by gel extraction. [Please click here to view a larger version of this figure.](#)

Figure 2: Sketches of basic magnetic tweezers apparatus.

(A) Sketch of a flow channel with a pair of magnets placed above and a microscopy objective placed below. (B) Sketch of an assembled flow channel where the blue colored rectangles represent the top and bottom coverslips, and gray elliptical lines represent the entry and exit of the flow channel. (C) Sketch of the force generation and calibration of the magnetic tweezers setup. (D) Examples of images of both the reference bead and moving bead at different de-focus positions stored in the bead image libraries. The red sign denotes the reference bead image obtained at a particular de-focus plane used for the focal plane locking. [Please click here to view a larger version of this figure.](#)

Figure 3: Force calibration of magnetic tweezers.

(A) Calibration of force using the fluctuations of bead along a rotation-free y-direction. X-axis is along the same direction as the magnetic field. Force is along the z-axis and is controlled by adjusting the distance, d , between the permanent magnet pair and the coverslip. (B) Calibration of using a long 48,502 bp -DNA. Heterogeneous magnetic bead property can be described by a single-parameter C . This figure has been modified from¹⁰. (C) Calibration of parameter C of a magnetic bead attached on a 2 kb-DNA using B to S overstretching transition. [Please click here to view a larger version of this figure.](#)

Figure 4: Single-molecule manipulation of a G4.

(A) Sketch of a stretching single G4 structure and measuring the unfolding events. (B) Typical example of force-increase (cyan) and force-decrease curve of a tethered G4. This figure has been modified from¹². (C) Force-extension curve of G4 in the absence (grey) or presence (pink) of RHAU protein. (D) Unfolding force distribution of G4 in the absence (grey) ($n = 140$) or presence (red) of RHAU and ATP ($n = 117$). This figure has been modified from¹³. [Please click here to view a larger version of this figure.](#)

Figure 5: Force-jump measurements of G4 unfolding.

(A) Representative time trace of the bead height change during force jump cycles between 1 pN and 54 pN measured without DmRHAU, with 10 nM DmRHAU but without ATP, and with both 10 nM DmRHAU and 1 mM ATP, as indicated in the figure panel. (B) Representative time trace of extension change of a G4 molecule (grey data) after jumping to 54 pN in several force-jump cycles taken from data in (A). The extensions corresponding to folded (F) and unfolded (U) G4 are also indicated; G4 unfolding events are indicated by arrows. This figure has been modified from¹³. [Please click here to view a larger version of this figure.](#)

Discussion

As described above, a platform for studying the mechanical stability of G4 DNA and the interactions of protein to G4 using single-molecule magnetic tweezers is reported. Accompanying the platform, highly efficient protocols of finding G4 DNA tether, and measurement of the folding-unfolding dynamics and stability of the G4 structure with nanometers special resolution are developed. The focal plane locking enables highly stable anti-drift control, which is important for detecting a small structure transition such as G4 (step size ~ 7 nm) and the interactions with proteins in buffer exchange experiments. This platform has recently been employed to studies of the folding and unfolding dynamics of telomeric G4¹⁴ and c-Myc promoter G4¹², as well as studies of G4 binding protein RHAU helicase¹³.

A typical experiment error is the non-specific binding of DNA and beads on the coverslip or multiple DNAs bind to magnetic beads. To reduce the non-specific binding, first, the storage of chemicals such as APTES and SMCC need to be routinely checked. Second, it is important to select a proper type of magnetic bead and block the non-specific binding. Using 10 mg/mL BSA buffer or small DNA oligo, such as poly T, to block the hydrophobic surface of the beads can significantly reduce the non-specific binding. Finally, a single DNA tether can be easily distinguished from multiple DNA tethers by the characteristic DNA overstretching transition^{17,18,19}.

Currently, the position determination of the platform is based on the bead imaging analysis, which has a limited spatial resolution of ~ 2 nm. In addition, it has a limited sampling rate of ~ 100 Hz, mainly due to limited acquisition rate of typical CCD cameras and correlation analysis of the images. These limitations can be overcome by using total internal reflection microscopy (TIRF) illumination. It produces a thin layer of evanescent light that exponentially decays with the height from the surface, which can be used to determine the extension change of short molecules at a high signal-to-noise ratio, while avoiding bead image acquisition and analysis²⁶. Another limitation is that it is challenging to directly visualize the tethered molecule or ligand bound to the tethered molecule using fluorescence imaging in the vertical stretching design, which can be overcome by stretching molecules in the focal plane using transverse magnetic tweezers²⁷. Finally, the current design of the reaction channel does not allow rapid solution exchange due to the flow perturbation to the molecule. Furthermore, a large drag force generated at high speed flow may even break the tethers. This limitation can be overcome by manipulating tethers inside the microwells placed on the bottom surface of the channel²⁸.

G4 stabilizing compounds are currently considered as potential therapeutic targets for diseases, including cancer, virus related diseases, and neurodegeneration diseases^{29,30,31}. The protocol can be applied to a wide range of G4 binding proteins² and small ligands³². In addition to applications in G4 studies, this platform as well as the protocols, in principle, can be used for studies of other nucleic acids secondary structures, including DNA and RNA hairpins, triplex, and i-motif³³.

Disclosures

The authors have nothing to disclose.

Acknowledgements

The authors thank Meng Pan for proofreading the manuscript. This work is supported by Singapore Ministry of Education Academic Research Fund Tier 3 (MOE2012-T3-1-001) to J.Y.; the National Research Foundation through the Mechanobiology Institute Singapore to J.Y.; the National Research Foundation, Prime Minister's Office, Singapore, under its NRF Investigatorship Programme (NRF Investigatorship Award No. NRF-NRFI2016-03 to J.Y.; the Fundamental Research Fund for the Central Universities (2017KFYXJJ153) to H. Y.

References

1. Rhodes, D., & Lipps, H. J. G-quadruplexes and their regulatory roles in biology. *Nucleic Acids Res.* **43** (18), 8627-8637 (2015).
2. Brazda, V., Haronikova, L., Liao, J. C., & Fojta, M. DNA and RNA quadruplex-binding proteins. *Int J Mol Sci.* **15** (10), 17493-17517 (2014).
3. Gonzalez, V., & Hurley, L. H. The C-terminus of nucleolin promotes the formation of the c-MYC G-quadruplex and inhibits c-MYC promoter activity. *Biochemistry.* **49** (45), 9706-9714 (2010).
4. Wang, F. *et al.* Telomere- and telomerase-interacting protein that unfolds telomere G-quadruplex and promotes telomere extension in mammalian cells. *Proc Natl Acad Sci U S A.* **109** (50), 20413-20418 (2012).
5. Mendoza, O., Bourdoncle, A., Boule, J. B., Brosh, R. M., Jr., & Mergny, J. L. G-quadruplexes and helicases. *Nucleic Acids Res.* **44** (5), 1989-2006 (2016).
6. Schiavone, D. *et al.* PrimPol Is Required for Replicative Tolerance of G Quadruplexes in Vertebrate Cells. *Mol Cell.* **61** (1), 161-169 (2016).
7. Lane, A. N., Chaires, J. B., Gray, R. D., & Trent, J. O. Stability and kinetics of G-quadruplex structures. *Nucleic Acids Res.* **36** (17), 5482-5515 (2008).
8. Woodside, M. T., & Block, S. M. Reconstructing folding energy landscapes by single-molecule force spectroscopy. *Annu Rev Biophys.* **43**, 19-39 (2014).
9. Neuman, K. C., & Nagy, A. Single-molecule force spectroscopy: optical tweezers, magnetic tweezers and atomic force microscopy. *Nat Methods.* **5** (6), 491-505 (2008).
10. Chen, H. *et al.* Improved high-force magnetic tweezers for stretching and refolding of proteins and short DNA. *Biophys J.* **100** (2), 517-523 (2011).
11. Chen, H. *et al.* Dynamics of equilibrium folding and unfolding transitions of titin immunoglobulin domain under constant forces. *J Am Chem Soc.* **137** (10), 3540-3546 (2015).
12. You, H., Wu, J., Shao, F., & Yan, J. Stability and kinetics of c-MYC promoter G-quadruplexes studied by single-molecule manipulation. *J Am Chem Soc.* **137** (7), 2424-2427 (2015).

13. You, H., Lattmann, S., Rhodes, D., & Yan, J. RHAU helicase stabilizes G4 in its nucleotide-free state and destabilizes G4 upon ATP hydrolysis. *Nucleic Acids Res.* **45** (1), 206-214 (2017).
14. You, H. *et al.* Dynamics and stability of polymorphic human telomeric G-quadruplex under tension. *Nucleic Acids Res.* **42** (13), 8789-8795 (2014).
15. Fu, H., Chen, H., Marko, J. F., & Yan, J. Two distinct overstretched DNA states. *Nucleic Acids Res.* **38** (16), 5594-5600 (2010).
16. Gosse, C., & Croquette, V. Magnetic tweezers: micromanipulation and force measurement at the molecular level. *Biophys. J.* **82** (6), 3314-3329 (2002).
17. Fu, H. *et al.* Transition dynamics and selection of the distinct S-DNA and strand unpeeling modes of double helix overstretching. *Nucleic Acids Res.* **39** (8), 3473-3481 (2011).
18. Zhang, X., Chen, H., Fu, H., Doyle, P. S., & Yan, J. Two distinct overstretched DNA structures revealed by single-molecule thermodynamics measurements. *Proc Natl Acad Sci U S A.* **109** (21), 8103-8108 (2012).
19. Zhang, X. *et al.* Revealing the competition between peeled ssDNA, melting bubbles, and S-DNA during DNA overstretching by single-molecule calorimetry. *Proc Natl Acad Sci U S A.* **110** (10), 3865-3870 (2013).
20. Chen, H. *et al.* Improved High-Force Magnetic Tweezers for Stretching and Refolding of Proteins and Short DNA. *Biophys. J.* **100** (2), 517-523 (2011).
21. Fu, H. X. *et al.* Transition dynamics and selection of the distinct S-DNA and strand unpeeling modes of double helix overstretching. *Nucleic Acids Res.* **39** (8), 3473-3481 (2011).
22. Zhang, X., Chen, H., Fu, H., Doyle, P. S., & Yan, J. Two distinct overstretched DNA structures revealed by single-molecule thermodynamics measurements. *Proc. Natl. Acad. Sci. U.S.A.* **109** (21), 8103-8108 (2012).
23. Zhang, X. *et al.* Revealing the competition between peeled ssDNA, melting bubbles, and S-DNA during DNA overstretching by single-molecule calorimetry. *Proc. Natl. Acad. Sci. U.S.A.* **110** (10), 3865-3870 (2013).
24. Vaughn, J. P. *et al.* The DEXH protein product of the DHX36 gene is the major source of tetramolecular quadruplex G4-DNA resolving activity in HeLa cell lysates. *J Biol Chem.* **280** (46), 38117-38120 (2005).
25. Giri, B. *et al.* G4 resolvase 1 tightly binds and unwinds unimolecular G4-DNA. *Nucleic Acids Res.* **39** (16), 7161-7178 (2011).
26. De Vlaminc, I., & Dekker, C. Recent advances in magnetic tweezers. *Annu Rev Biophys.* **41** 453-472 (2012).
27. Yan, J., Skoko, D., & Marko, J. F. Near-field-magnetic-tweezer manipulation of single DNA molecules. *Phys Rev E Stat Nonlin Soft Matter Phys.* **70** (1 Pt 1), 011905 (2004).
28. Le, S. *et al.* Disturbance-free rapid solution exchange for magnetic tweezers single-molecule studies. *Nucleic Acids Res.* **43** (17), e113 (2015).
29. Neidle, S. Quadruplex Nucleic Acids as Novel Therapeutic Targets. *J Med Chem.* **59** (13), 5987-6011 (2016).
30. Simone, R., Fratta, P., Neidle, S., Parkinson, G. N., & Isaacs, A. M. G-quadruplexes: Emerging roles in neurodegenerative diseases and the non-coding transcriptome. *FEBS Lett.* **589** (14), 1653-1668 (2015).
31. Balasubramanian, S., Hurley, L. H., & Neidle, S. Targeting G-quadruplexes in gene promoters: a novel anticancer strategy? *Nat Rev Drug Discov.* **10** (4), 261-275 (2011).
32. Amato, J. *et al.* Toward the Development of Specific G-Quadruplex Binders: Synthesis, Biophysical, and Biological Studies of New Hydrazone Derivatives. *J Med Chem.* **59** (12), 5706-5720 (2016).
33. Wells, R. D. Non-B DNA conformations, mutagenesis and disease. *Trends Biochem Sci.* **32** (6), 271-278 (2007).