almost unknown;—that even under the age of twenty, the cases do not compose above $5\frac{1}{2}$ per cent.;—and that what is usually believed the most critical period for women, namely, between forty and fifty, when the catamenia cease, is either less critical than has been thought, or at all events equally so for the male sex; for the proportion attacked between these years is in both sexes very nearly the same as the proportion attacked during the ten previous years.

We must here leave our author for the present. The consideration of the subsequent parts of the work, which treat of the several Forms of Insanity, its various Terminations, its Prognosis, its Treatment, and its relations to Medical Evidence, must be reserved till our next number.

ART. II.—*Pathological and Practical Researches on the Diseases of the Stomach, the Intestinal Canal, the Liver, and other Viscera of the Abdomen.* By JOHN ABERCROMBIE, M.D. &c. Edinburgh, 1828. Pp. 396.

THE present volume consists of five parts, in which the author treats successively of the pathology of the stomach, the intestinal canal, the liver, the spleen, and the pancreas. The inquiry is introduced by some general observations on the nature of the structures concerned in the diseases of the two first parts, and on the morbid conditions to which each is respectively liable. The author then enters on the investigation of the pathology of the stomach, of which he treats in three sections, under the heads of inflammatory affections, organic diseases, and dyspepsia. The survey of these subjects corresponds so closely with that which the learned author already traced in this Journal,* that to enter into any detailed account would only give rise to much repetition. Some idea of the subjects discussed, and of the order in which they are examined, may be communicated by the following outline.

The author, after defining *gastritis* as inflammation of the gastric mucous membrane, proceeds to illustrate its progress and effects by various cases, in which it produced ulceration in definite patches, or diffusely spread thickening and induration, hemorrhage by erosion of vessels, and perforation with fatal peritonitis. One of the most interesting varieties of this occurrence is presented in a case by Dr Combe of Leith, in which the ulcer had formed an opening into the transverse arch of the

* Contributions to the Pathology of the Stomach, the Pancreas, and the Spleen, Medical and Surgical Journal, Vol. xxi. p. 1.

colon, in consequence of which, the prominent symptom during life was occasional vomiting of feculent matter.

Under the head of inflammation of the gastric mucous membrane, the chief novelty consists in some observations on the peculiar albuminous inflammation, denominated by M. Bretonneau, and other foreign authors, *Diphtherite*, or *inflammation of the mucous corion*, causing an exudation like a pellicle or covering.* It is certainly unfortunate that this author, notwithstanding all his endeavours to distinguish with accuracy, nevertheless confounds the albuminous with various other inflammatory affections of the mucous corion. So far, however, as we understand his distinctions, and more especially the facts disclosed by his cases, we fear we cannot agree with Dr Abercrombie in thinking that the disease is different from croup. It is manifest, we imagine, that several of the cases of M. Bretonneau are examples of genuine croup; and when their situation in the pharynx and œsophagus prevents them from being referable to this head, it is further manifest that they are allied to it in producing a very similar inflammation of the œsophageal mucous membrane.† It must be further observed that the aphthous disease described by Dr Hamilton, and by the present author, is evidently distinct from the true albuminous inflammation of croup; and that M. Bretonneau has not at all times been able to observe this distinction, we agree with Dr Abercrombie in thinking. But while we willingly make this admission on the one hand, we must further say that we are not prepared to agree with the latter author, when he seems in speaking of this subject, from mere oversight, we presume, to represent croup as an “idiopathic inflammation of the membrane of the larynx.” That it is idiopathic and inflammatory is evident from a variety of circumstances. But that it neither originates in, nor is confined to the larynx, is proved by the evidence of numerous dissections. We believe also, that it is unnecessary for us to remind Dr Abercrombie, that inflammation of the larynx, properly so called, is a very different disease from the tracheo-bronchial inflammation, which in infants gives rise to the peculiar symptoms known by the name of croup.

The third section contains some valuable observations on the pathology of that multiform and anomalous state of the stomach and bowels termed *dyspepsia*. After enumerating the circumstances necessary to the healthy state of the process, Dr Abercrombie shows that the different forms of indigestion may depend either on impaired action of the muscular coat, deficiency

* Des Inflammations du Tissu Muqueux et en particulier de la Diphthérie, ou inflammation pelliculaire. Par P. Bretonneau, Medecin en chef de l'Hôpital de Tours. A Paris, 1826. P. 41.

† Ibidem, p. 144, et ensuite.

of the consentaneous intestinal action, changes in the fluids of the stomach, or morbid irritability of the gastric mucous membrane, separately or conjointly. Applying the inferences which these views suggest to those which may be derived from the circumstance of the necessity of confining the quantity of food within those limits within which digestion may be easily effected, from the comparative digestibility of different articles, and from the propriety of never interrupting the process by the addition of fresh alimentary matter, Dr Abercrombie delivers some clear and simple observations on the management of this disorder. For these, as well as for sundry judicious remarks on *gastrodynia*, on chronic vomiting, on *pyrosis* or water-brash, on *hæmatemesis*, and on sympathetic affections of the heart, we refer the reader to the work itself, in which he will find much useful matter to reward perusal.

In an appendix to this part, the first subject which is shortly examined, is “derangement of the function of the stomach, by tumours attached to its external surface, without disease of the coats.” The second section, on diseases of the œsophagus, consists of two subdivisions, inflammation of that canal, and the pathology of *dysphagia*. Of the former, the author gives one example, in which part of the larynx, the whole pharynx, and the inner surface of the œsophagus, as far as the cardia, were covered by a loose soft adventitious membrane. On the subject of *dysphagia*, Dr Abercrombie gives a short general view of the causes of that disease, chiefly from the classification of Dr Monro, to whose work he refers. On this arrangement we take the present opportunity of remarking, that since dysphagia is only a symptom or effect of several diseases of the œsophagus, or contiguous parts by no means similar, every classification of this kind is liable to the objection of being much more nosological than pathological, and the individual parts are liable to be accidentally assembled by very casual connection. To the practitioner, doubtless, this mode of classification is exceedingly useful, since it enables him to perceive at one glance all the possible changes in organization, structure or position, which may cause *dysphagia*, and on this ground we imagine it is adopted by the present author. The observations and cases of Bleuland on this subject* are not mentioned by Dr Abercrombie.

In the third section of the appendix, which is entitled Diseases of the Duodenum, Dr Abercrombie, after regretting the want of facts, quotes a case of tubercular disease and ulceration from Dr Irvine of Philadelphia, a case of ulcer laying open the

* Jani Bleuland, Observationes Anatomico-Medicæ de Œsophagi structurâ sanâ et Morbosâ. 4to. Lugduni Bat. 1785.

hepatic artery, and causing fatal hemorrhage from Broussais, and one case of perforation from the Museum of the College of Surgeons, and another from M. Robert, (Nouvelle Bibliotheque Medicale, June 1828.) The subject is concluded with notices of two cases of ventral hernia, in consequence of accidental rupture, or giving way of the abdominal parietes. The effects caused by both were relieved by the use of pressure.

The second general division of his subject Dr Abercrombie investigates under three heads; *ileus*; the inflammatory affections of the external textures, including peritonitis and enteritis; and diseases of the mucous membrane. The chronic and organic diseases are combined with one or other of these heads according to their natural connections.

Ileus, which the author regards, with Cullen and other systematists, as a different degree or more advanced stage of the affection denominated colic, he represents as occurring under three modifications;—1st, either as simple ileus without previous disease; 2^d, as ileus with previous disease of that character that it operates by deranging the muscular power without inducing mechanical obstruction; and 3^{dly}, ileus with mechanical obstruction.

To illustrate the first variety of the disease the author relates two cases in which the usual symptoms of ileus took place, and proceeded to the fatal termination, yet without presenting, on inspection, any change in the intestinal canal beyond unusual distension, and such slight injection of its capillaries as was totally inadequate to render it a fatal disorder. By the appearances of other two cases he then traces the transition of the disease from the state of distension to that of lividity, and eventually of gangrene, yet without inflammatory exudation,—demonstrating that the mere mechanical injury occasioned by distension is adequate, without the intervention of a distinct or active inflammatory attack, to produce mortification of the intestinal tissues. By the phenomena of a fifth case, however, in which the intestine was not only distended, but highly inflamed, and very lacerable, and its peritonæal surface covered by inflammatory exudation, he shows that this inflammatory process may be established in consequence of the inordinate distension, and may eventually proceed to mortification of the intestinal tissues. From two cases in the appendix to this head, it appears that the primary state of simple ileus consists in that extensive distension of the intestine by air, which is known nosologically by the name of *tympanites*, and which appears not only to operate on the intestinal tissues like a mechanical power of great violence, but by neutralizing or annihilating the power of the muscular layer, to arrest the peristaltic motion, and to prevent the bowel from being capable of unloading itself, and thus place it in a

situation highly favourable to the developement of gangrenous inflammation.

Ileus of the second variety is in general the consequence of previous inflammation or some one of its numerous products. In organs of so complicated figure and disposition, and so liable to change of situation, as the intestines, the effects of the inflammatory process are liable to sundry singular modifications. Thus, albuminous exudation taking place from the peritonæal surface of a fold of ileum, may unite it to a contiguous fold so firmly, that every motion of the one portion causes motion of the other, however distant, and often with manifest injury of the intermediate portion of the tube. In other instances a portion of ileum may be connected to the muscular peritonæum, to the colon, or any other viscus, the situation of which is fixed, and may thus be prevented from undergoing the changes of situation necessary to the healthy performance of its functions. Upon the same principle the old adhesions induced by hernial protrusions are a fertile source of intestinal obstruction, or rather of suspension or annihilation of the peristaltic motion.

Each of these various sources of disease are well illustrated in the second section by the author of this treatise. The necroscopic inspections which he adduces under this head, are extremely valuable in demonstrating the efforts of these membranous bands,—the frequent consequence of previous inflammation. In one instance, § iv. p. 115, the same effect had been produced by an unnatural twist of the sigmoid flexure upon the rectum, in consequence of which a valvular obstruction to the transition of the contents of the former bowel into the latter had taken place, and caused very great distension, proceeding to gangrene, in the sigmoid flexure and other parts of the colon.

The third section comprehends the following varieties:—old disease of the intestine connected with hernia, and artificial anus; internal hernia; intus-susception; ileus from gall-stone; contraction of the calibre of the intestine; and stricture of the bowel. The case and inspection related under the first head, furnishes an example of the usual appearances found in cases of artificial anus after hernia. Under the second head are given three examples of strangulation of portions of intestine through preternatural spaces formed by pseudo-membranous adhesions. Under that of intus-susception, are recorded five cases in which portions more or less extensive of the intestinal tube had been invaginated within other parts, with the effect of inducing either inflammation, or gangrene, or both. Of ileus from gall-stone, the author gives only one instance, in which a calculus of unusual dimensions, four inches in its large circumference, and three and a-half in the small one, impacted near the end of the upper

half of the small intestine, had produced great distension of the bowel, followed by inflammation and albuminous effusion. Of contraction of the calibre of the intestine, the first instance is one in which the contraction was produced by thickening of the intestinal tunics, and fungous excrescences or granulations from its villous membrane. The second is one in which it was produced in the upper end of the rectum by a firm, flat band passing in front of the bowel from a mass of diseased glands situate behind. In the third it was the effect of inflammatory induration and thickening; and in the fourth it was the joint effect of these changes and of adhesions on the peritonæal surface, mutually connecting several folds of ileum. Of stricture one example only in the arch of the colon is given.

This general view of the cases and necroscopic reports of Dr Abercrombie we give thus shortly, because we regard it as out of place here to enter into any of their details, and because we consider it undoubted, that whoever wishes to understand thoroughly their practical tendency, will not fail to peruse them in the words of the author. Our present concern is more especially with the fourth section, which contains the inferences and deductions, pathological and practical, which the author thinks himself justified in deriving from these facts.

The subject of ileus has at all times been the source of much perplexity both to the practitioner and the pathologist. The hypothesis of spasm, so liberally used by Hoffmann, Cullen, and others of the same school, may be easily shown to be not only gratuitous, but absolutely contradicted by facts. For not only have all the phenomena of ileus been observed to take place without the smallest indication of spasm, either during life or after death; but in many instances ileus has appeared to be connected rather with absolute loss of power than with spasmodic augmentation. Of this fact several of the cases detailed by the present author are remarkable examples; and others are recorded in the writings of physicians and pathological collectors. Again, though the phenomena of ileus are in very many instances connected with distinct traces of the inflammatory process, not only as effect but as cause, in other circumstances it is impossible to establish any connection of this nature; and ileus has been observed to proceed to a fatal termination without being attended either with inflammation or gangrene. The idea, therefore, which is either suggested or maintained by Speranza and other authors, that ileus depends on inflammation, is, like others, of partial application only. The only method of establishing a genuine theory of the disease, and one of universal application, therefore, is by comparing all the known possible cases and varieties in which it has been known to occur,

and finding some principle, not of partial, but of general resemblance.

The views of the present author on this subject, though already partially known to the profession through the medium of this Journal, * are perhaps more fully and clearly developed in the present volume. After remarking, that in the earliest period in which the anatomist has an opportunity of inspecting the necroscopic phenomena of ileus, it appears to consist in simple distension, without visible change in the structure of the intestinal canal; that a little later a vivid tinge of redness is visible; and that, at a period still more advanced, the parts assume a leaden or livid hue, and eventually present the soft lacerable condition of gangrene;—he directs the attention of the reader to the fact—that these changes occur without any appearances of peritonæal inflammation. From these circumstances, he infers, that though the disease may be, and is actually observed to be, occasionally connected or complicated with peritonæal inflammation, the appearances now alluded to are seated, not in the peritonæal, but in the muscular coat.

Having established this point, the next is to ascertain the precise condition or morbid change of the muscular coat. At advanced periods of the disease, Dr Abercrombie observes, it is manifest that all muscular power is gone; “for we find the part which has been distended fallen flat when it is emptied, and presenting a broad surface like an empty bag, without any tendency to contraction. This case, indeed, presents the disease in a state of perfect gangrene; but it appears that the same loss of muscular power may take place at a much earlier period, and in connection with a much lower state of the disease.” The justice of this inference he exemplifies by reference to the 35th case, in which, though the patient lived for eighteen hours, with a free external opening directly communicating with the distended intestine, which contained only air and fluid excrement, yet no discharge took place,—a circumstance which, he conceives, indicates that the intestine, for a considerable space above the opening, must have lost its muscular action. Combining these considerations with those derived from the phenomena of mechanical obstruction, as exemplified in cases 33d and 51st, in the last of which, after all the most powerful means had failed, the intestine was made to contract and empty itself by the repeated application of galvanism, he draws the following conclusion:—“It is probable, therefore, that there occurs in the state of ileus, a certain loss of the muscular power in a portion of the canal, in consequence

* *Researches on the Pathology of the Intestinal Canal.* Part i. *Med. and Surgical Journal*, Vol. xvi. p. 2—7, and 11—23.

of which it does not act in concert with the other parts, but becomes distended by the impulse from the parts above, which in the healthy state would have excited it to contraction."

The opinion now stated, which is here rather more distinctly and specifically expressed than it formerly was in the pages of this Journal, Dr Abercrombie proceeds to confirm by various facts and arguments, derived from a view of the comparative states of musculo-membranous organs in the collapsed and distended conditions. These arguments are very much of the same nature with those already adduced by Dr Abercrombie in his investigation of this subject in the essay already quoted,* and are, upon the whole, so little modified that we do not feel justified in reproducing them, either in the form of enumeration or abridgement. It is sufficient to observe, that by more pointed reference to the results of inspection, and by deducing conclusions from a greater variety of examples of the disease, the author renders his hypothesis of ileus both more consistent and upon the whole more intelligible.

The chief point, it may be remembered, in which the hypothesis of Dr Abercrombie is at variance with that commonly received, is that the empty, collapsed, or contracted part is not the morbid but the *sound* portion of the intestine; whereas the distended portion which, according to the hypothesis of spasm, ought not to be diseased, is in a state of inordinate and morbid distension. Without pretending to espouse either side, or to scrutinize rigidly the arguments of the author, we may remark that the most certain proofs of the truth of this hypothesis may be reasonably expected in two circumstances.

First, If the distended part is invariably the real seat of disease, and the contracted part the normal state of the intestine, the latter ought never to be found either red, livid, or in a state of gangrenous softness; and these appearances, when they occur, ought always to be confined to the distended portion of the intestine. This test, so far as we can discover, all the cases adduced by the present author will undergo without objection; and since in each of them, therefore, the pathologist must admit that the contracted portion does not become the seat of inflammation, it results that the morbid state of this part is at least not established. Whether it ever becomes inflamed, livid or gangrenous, must depend upon the evidence of further and more numerous examples of the disease.

Secondly, If the contracted portion is not in its natural state, but is so in consequence of spasmodic action of the muscular fibres, then this contraction ought either to disappear, or at least to be remarkably diminished after death. Spasm is ad-

mitted by all observers to be an action of muscle during life only, and either disappears when life ceases, or leaves only its effects in the form of congestion of vessels, laceration of fibres, extravasation, or some similar change. In the case of the intestinal canal, it might be supposed, though however rigidly, forcibly, and permanently the muscular fibres may be contracted during life, relaxation would take place at death or soon after it. In none of the examples inspected by the present author does this relaxation appear to have taken place; in all, the portion below the distended part was much contracted and generally empty; and, so far as the evidence of these inspections goes, it must be inferred that the unusual contraction was the result, not of spasm, but of the natural action of the muscular fibres, when not antagonized by the successive propulsion of the contents of the upper portion of the intestine.

The subsequent speculations regarding the manner in which the intestine becomes inordinately distended, and in which the peristaltic motion, when once weakened in any given portion of intestine, allows this distortion to be effected, are so similar to the observations at p. 4 and 5 of the essay already referred to, that it is unnecessary to do more than simply notice them. The general conclusions also which Dr Abercrombie digests in thirteen separate propositions, though somewhat modified, and enumerated on the whole with greater precision, are substantially the same with those given at p. 22 and 23 of the same volume.*

Upon the whole, the opinion which Dr Abercrombie entertains of the nature of ileus, is nearly the following.—Though considerable variety in the necroscopic appearances is observed, and though there may be much diversity in the primary state of the parts concerned in the disease, in most instances the bowel is more or less distended beyond its natural capacity; and in several this distension is extreme, and may alone be the cause of the fatal termination. The disease in cases of this description, Dr Abercrombie ascribes to simple loss of muscular power, which, however, may pass at a later period into the inflammatory stage. In another class of cases, in which traces of inflammation, with much less distension, are recognized, he regards inflammation as part of the primary disease; but this inflammation he represents as different from that of *enteritis*, chiefly in the absence of febrile symptoms, and in the disease passing speedily into gangrene, unaccompanied with the flocculent or albuminous deposition which invariably attends the former disorder. Assuming the principle that gangrene is a disease chiefly of the muscular coat, he infers that, as in this malady gangrene is found uncombined with any other disease, it

* Med. and Surgical Journal, Vol. xvi. p. 22.

is reasonable to conjecture that the muscular coat is the seat of inflammation. This inflammation of the muscular coat, Dr Abercrombie regards as quite distinct from enteritis, and as constituting ileus of the most formidable kind, and proceeding rapidly to the fatal termination.

Of the theory thus stated, we do not feel entirely competent to examine the merits within the short space which we can set apart for this purpose, nor perhaps is it requisite that we should. When it is once made known, which is more especially the duty of these pages, its accuracy will not be long in being subjected to scrutiny, with various degrees of strictness. It is due, nevertheless, to the author to say, that it is plausible, and appears to accord well with the facts—better it may be said than any yet adduced. Of some of the intermediate links of the chain of reasoning, the connection may be questionable; but as these, so far as we see, are not necessary, we are, on the whole, disposed to admit the general truth of the conclusions. It may be doubted, for example, whether gangrene of the intestinal canal is confined chiefly to the muscular tissue; but we do not, on this account, deny that the muscular coat is seriously, and perhaps irreparably injured in cases of ileus. The great influence, extreme severity, and the pernicious effects of this injury we can easily perceive, from knowing the limited distensibility of muscular fibre, and how liable it is, during great distension, either to give way, or to be so much injured in its texture, as to become the seat of a bad or gangrenous inflammation. When the experiments of the iatro-mathematical physiologists on this substance, and the numerous repetitions of them by modern authors are remembered, it is easy to see that inordinate distension of the intestinal canal may not only impair or even annihilate the power of the muscular coat, as maintained by Dr Abercrombie, but injure its organization so severely and irreparably, as to induce gangrene with unequalled rapidity. In short, the more we reflect on the effects of this violence, on a tissue so lacerable as the muscular fibre, the more extraordinary does it appear that it is not more frequently succeeded by inflammatory and gangrenous disorganization, than that this process takes place only in the circumstances now mentioned by Dr Abercrombie.

In admitting the competency of the hypothesis now advanced, two circumstances merit inquiry. The first is the cause of the inordinate distension; the second is in what manner, or by what circumstances this distension differs from that admitted by nosologists in intestinal tympany.

On the first point the speculations of Dr Abercrombie at p. 4 and 5 of the 16th volume of this Journal, and at p. 136 and 137 of the present volume, contain all that he thinks the

nature of the subject justifies. Of the explanation afforded in these passages the leading feature is the supposition of unequal or unharmonious action of the intestinal canal. One portion he conceives may, from various accidental causes, for instance accumulation of solid contents or of elastic fluid, so far lose its inherent muscular power, as to be unable to contract and propel these contents into the next portion. As the portions of the canal above this, however, are supposed healthy, and consequently capable of propelling their contents, the portion of which the power is impaired is undergoing increasing distension, with the same or a less adequate power. As this distension proceeds, the bowel is every moment becoming less capable of overcoming the resistance; and at a certain point it may be conceived to be so distended, that, as already stated, it is utterly incapable of exercising any contraction whatever. It is at this period that the distension may be conceived adequate to injure the organization of the muscular fibres, and from this time that the commencement of the disorganizing or gangrenous process of inflammation may be dated.

In this supposition the chief difficulty is to conceive the first or incipient trace of impaired action. By what is it induced, and how does it take place in any one part more than another, that is to say, without organic injury or mechanical obstruction? We know well, that in the latter case particularly, there is an evident cause of impaired or impeded action, and one of which we can readily appreciate the influence. But by what means, and in what manner one portion of the canal should suddenly lose its contractile power without previous injury of its muscular tissue, is certainly not very obvious. Must we admit, as the only cause, the mere presence of air or of alimentary matter itself, as adequate, especially during the fermentation of indigestion to produce this effect? Still it may be replied the muscular fibres contracting on these ought to propel them to the next portion of intestine. Unless, indeed, the previous loss of power be admitted, or it be said that the distension caused by the presence of air is sufficient to neutralize the muscular action, no resource is left but this, which is in truth *reasoning in a circle*. The idea that the inability to contract is occasioned by mere mechanical distension from the presence of air, receives some support from the phenomena of lead-colic, in which the action of the muscular fibres is directly impaired by the poisonous operation of the mineral.

On the second point, viz. by what circumstances the distension maintained by the present author in ileus, differs from that of intestinal tympany, we have no information whatever, unless that it actually occurs as a symptom of ileus, and a part of fatal diarrhoea. (See p. 308.) Is it in the circumstance, that in

the former one part is excessively distended, and another is contracted,—while in the latter the distension is uniform,—that the difference consists? In the absence of positive facts we abstain from conjecture, and merely state the difference as one on which it would be desirable to obtain some satisfactory information. The reader, however, may consult with advantage the second section of the appendix to this part, for some good observations on the acute and chronic forms of tympany.

In entering on the consideration of the treatment of ileus, Dr Abercrombie most properly directs the attention of the practitioner to the question of the presence or the absence of hernia. It is indeed a point which cannot be too frequently or too forcibly impressed, that a much larger proportion of cases of ileus than is commonly imagined really depend on hernial protrusion and strangulation, and that in every case of abdominal pain with vomiting and obstruction of the intestines, the first duty of the practitioner is to determine, by accurate examination, whether the symptoms arise from hernia or not. In conducting this inquiry, Dr Abercrombie reminds the practitioner of two circumstances;—*1st*, that hernia may exist without the patient being aware of it, or making any complaint that might lead to the supposition of its existence: *2d*, that the hernial protrusion may be so small as to include a minute portion only of one side of the intestine, (the anti-mesenteric,) and yet be the cause of fatal ileus. Of course, when the existence of external hernia is recognized, it is easy to know what is to be done. When no external protrusion can be detected, and when one of two suppositions must rush on the mind, either that of internal strangulation, or of the state which, according to the author of the present work, constitutes simple ileus, the choice and application of practical measures becomes a task of much greater difficulty.

It is doubtless a mortifying proof of the frequent inadequacy of pathological inquiry, that, under the circumstances now mentioned, the treatment must, notwithstanding, be conducted upon empirical principles. Without attempting to ascertain from the results of his pathological investigations what is the form of treatment calculated to restore the action of the intestinal muscular fibres, and to counteract the injurious effects of over-distension, Dr Abercrombie confines himself chiefly to the statement of the remedial means which he has found most beneficial. These consist in blood-letting, the tobacco-injection, the application of cold, either in the form of cloths immersed in vinegar and water applied to the belly, or injections of cold water, opiates, and the stimulant power of aloetic wine in full doses, or of the tincture less liberally.

Of the merits of blood-letting, the tobacco-injection, and the local application of cold, he speaks still more confidently

and forcibly than in the essay to which we have already so often referred; though without attempting to trace any connection between the state intended to be removed, and the operation of the remedy, unless in regard to blood-letting. Upon this point we believe all practical observers are agreed; and as a large proportion of cases of ileus are attended with inflammatory symptoms, prudence will always suggest the free use of blood-letting. With the hypothesis of Dr Abercrombie, however, there is this difficulty, that as in cases of simple ileus the disease appears to originate in *loss* of power, or at least in irregular and unharmonious action of the muscular coat, the detraction of blood may not appear a very rational method of repairing its energy, or re-establishing the balance which is thus imagined to be lost. In regard also to the influence deduced from the benefit resulting during or immediately after venesection, there is this difficulty, that it is not unfrequently impossible to determine till after death, whether of any given case of abdominal pain, with vomiting and alvine obstruction, the pathological cause be inflammation or the state of distension maintained by the author. When loss of blood is beneficial, it is not irrational to infer, that inflammation is the chief cause of the symptoms.

On the efficacy of opiates, in which Dr Abercrombie is disposed to place much confidence, different opinions will be entertained. So far as the operation of opium and of all its preparations on the muscular fibre is known, it ought to be the last remedy employed in such a state of the intestinal canal as that maintained by Dr Abercrombie, and an agent from which it is impossible, rationally speaking, to expect beneficial effects. On this subject, nevertheless, one or two good experiments is worth a thousand conjectures, unless these also are derived from experience or observation. On this point, therefore, we observe, that though the fact of the efficacy of an opiate after a full bleeding, as stated by the ingenious author, is undoubted, and though in some cases the exhibition of an opiate alone enables purgatives medicine, which previously seemed unavailing, to produce their specific effects;—yet there are cases in which opium, instead of operating favourably, seems to be directly injurious, by actually destroying the little contractile power that the intestinal muscular tissue retains. We have in cases of ileus seen opiates administered till they produced stupefaction, with no other effect than that of abating, or rendering the patient less capable of feeling, the abdominal pain. Opium, in short, notwithstanding the strong testimony of the present author in its favour, must be regarded as a remedy of ambiguous effect in ileus; and whenever it is thought to be indicated, its place may in all cases be advantageously occupied by the tobacco-injection,

which, when properly managed, is both a safer remedy, and one in which more confidence may be reposed.

In the second part of this division, Dr Abercrombie proceeds to the consideration of the inflammatory disorders of the external tissues of the intestinal tube, including under this head *peritonitis* and *enteritis*. Of these subjects the investigation is at once comprehensive and accurate. Without attempting to follow the steps of the author through each of his sections, it is sufficient to say, that he considers these diseases under four different heads;—*peritonitis*, *enteritis*, *erysipelatous peritonitis*, and chronic *peritonitis*.

The history and illustrations of the phenomena and pathological characters of the two first varieties of disease correspond so much in materials and general arrangement with those of the Second Essay of the author, formerly published in this Journal, that it is quite unnecessary to give any minute account of them here. In general results also they are so similar to those of other authors on this subject, that it is impossible to hope to interest the reader by an enumeration of symptoms and morbid appearances with which he is already quite familiar. It is proper, nevertheless, to mention, that Dr Abercrombie has made, in his arrangement of this subject, various improvements, suggested doubtless by further observation and reflection. Thus, though his general description of the characteristic phenomena of *peritonitis* and *enteritis*, distinguished according to his idea of these disorders, is much the same, in his divisions and practical illustrations of the subject, he has, in conformity with opinions expressed in a note at the end of the number, discarded the distinction of *peritonitis with diarrhæa* and *peritonitis with a natural state of the bowels*. In that note, Dr Abercrombie was led justly to express the doubt, whether the peritonæal inflammation were the primary disease, which there was reason to suspect originated in the mucous membrane, and afterwards proceed to affect the peritonæum. We confess that the distinction there made always appeared to us erroneous; and nothing but the known character of the author for accurate and sound observation, withheld us from saying so. We have, therefore, some pleasure in perceiving that its deficiency, though thus early recognized by the author himself, is at length fully admitted.

In the present volume, the author considers the two diseases, *peritonitis* and *enteritis*, under the following modifications:—1. simple *peritonitis*; 2. *peritonitis* confined nearly to the descending colon and rectum; 3. local *peritonitis* of very small extent; 4. *peritonitis* terminating by extensive suppuration; 5. *peritonitis* passing into *enteritis*; and, 6. *enteritis*.

To this mode of division, which is certainly sufficiently mi-

nute, the principal objection is, that it is not altogether perspicuous, and constructed according to the established rules of rational classification. Some of the heads are evidently different degrees of the same morbid process; and if circumstances of this description be employed as one basis of division, the mode of termination, it may be said, cannot with propriety be admitted as another. *Enteritis*, our readers are aware, consists, according to Dr Abercrombie, in simultaneous inflammation of the peritonæal and muscular coats; and therefore, though not with perfect accuracy, is placed under the same head. These objections, which are mentioned not because we have either space or inclination to subject the method to rigid scrutiny, probably belong to the subject itself, which, from the complicated character of the textures composing the intestinal canal, renders any attempt to classify with precision difficult in the extreme. How forcibly soever, therefore, the ingenious author himself may have felt this inconvenience, it is manifest that he regards it as completely counterbalanced, in the circumstance that it enables him to give a view of the subject of practical utility.

In the treatment of these forms of intestinal disease, Dr Abercrombie, besides the ordinary rules for subduing inflammation by large and repeated blood-letting, is anxious to impress the advantage to be derived from small bleedings, at short intervals, when the effect of the first begins to subside. "In this manner," he continues, "we prolong, as it were, the impression which is made by the first bleeding, and a two-fold advantage arises from the practice; namely, that the disease is checked at an early period, and that the quantity of blood lost is in the end much smaller than probably would be required under other circumstances. If we allow the patient to lie after the first bleeding 10 or 12 hours, or even a shorter period, the effect of it is entirely lost, and a repetition of it to the extent of 20 ounces may be required for producing that effect upon the disease, which, by the former method, might be produced by 5; and besides, the disease has in the interval been gaining ground, its duration is protracted, and the result consequently rendered more uncertain. The inflammation of a vital organ should not be lost sight of above an hour or two at a time, until the force of it be decidedly broken; and unless this takes place within 24 hours, the termination must be considered as doubtful."

These observations we give in the words of the author, because we regard it impossible to impress too strongly upon the mind, especially of the young practitioner, the important lesson which they convey. Often have we had occasion to see the most extensive mischief produced by neglecting the performance of venesection, at that period of the disease at which alone it can be either beneficial or sanative. Though we have no wish to improve upon the small secondary blood-lettings of

the author, many circumstances induce us to regard the quantity as of infinitely less moment than the season of the venesection. Whether a patient with acute peritonæal or intestinal inflammation lose, after the first large evacuation, 5, 10, 15, or 20 ounces of blood, is a matter of very little moment, provided he lose it at a *short* interval after the first. The error of most practitioners, after performing a large blood-letting, is to wait too long before a second is attempted; and during this space the disease too often recovers as much as it had lost, and renders very nearly the same measures necessary as at first. It is chiefly while the system is still under the immediate operation of the first blood-letting, that a second or third even is useful in promoting that state which is favourable to unloading the capillary vessels of an inflamed organ; and then chiefly should the practitioner, by the energy and promptitude of his measures, endeavour to extinguish completely the morbid action. On this topic, however, this is not the place to enlarge; and we have only to recommend this part of the work of Dr Abercrombie to the serious attention of the reader. The class of cases, in particular, in which, after evacuations have subdued the active inflammation, the use of wine and other stimulants becomes necessary, possesses much practical interest.

The principal novelty in this part of the work is the section on *erysipelalous peritonitis*. The characters of this form of peritonæal disorder are not very distinctly defined; and all that we can perceive the author gives as a reason for the denomination is, that it is connected with erysipelas and other diseases of an erysipelalous character. He warns the reader, nevertheless, that he attaches to the name no other importance, than that it affords a convenient title to the section in the general arrangement of the subject.

The morbid effects of this form of disorder furnish the most distinctive characters of its nature. It terminates chiefly in effusion of fluid, with little or none of the albuminous exudation by which the common form of the disease is distinguished. The effused fluid may be bloody serum, either alone or mixed with a proportion of opaque purulent fluid which subsides; or whey-coloured fluid with albuminous flakes; or occasionally distinct purulent fluid. The appearance of the affected membrane varies considerably from the usual sound state to one of slight vascularity, or of a uniform dark red. It has generally a thickened softened appearance like a part which has been boiled; and in some instances the omentum is the chief seat of disease. These circumstances, combined with the history of the disease and the kind of subjects in which chiefly it occurs, show that it is allied somewhat to the peritonæal inflammation

of puerperal females, in some of the varieties of which the fluid effused is abundant, thin, serous, and puriform.

Besides two cases, both in females, adduced to illustrate the nature of this disorder, Dr Abercrombie gives a sketch of an epidemic form in which it appeared in the Merchants' Maiden Hospital in this city in March 1824. Among the girls educated in this institution, the disease commenced with slight fever, vomiting, and erysipelatous affection of the throat, accompanied in some cases with swelling of the glands of the neck. It spread with such rapidity, that from 15 to 20 girls were affected at once, but disappeared with little treatment in the course of 3 or 4 days. The disease went on in this manner for about a week, when on the 13th March, a girl aged 10, after being affected with similar mild symptoms, was suddenly attacked by vomiting, diarrhoea, abdominal pain, and mortal sinking, which terminated fatally on the evening of the 15th. The necroscopic appearances were marks of extensive peritonæal inflammation, with albuminous flakes deposited in various parts of the intestines, a considerable quantity of milky puriform fluid in the cavity, and a dark-coloured appearance of the vermiform appendix, which was large and turgid.

The epidemic went on still without any urgent case till the 4th of April, when a girl of 12, who after 3 or 4 days of illness, was considered convalescent, was suddenly attacked with abdominal pain, and frequent desire to go to stool. These symptoms went on, notwithstanding the use of remedies, till the 7th, on the evening of which mortal sinking appeared, and soon terminated fatally. Inspection showed marks of inflammation of the peritonæal membrane, with albuminous flakes, and much puriform fluid in the cavity.

From these and similar facts the author infers, not without reason, that this variety of peritonæal inflammation has some alliance with erysipelatous inflammation of the skin, and several of the forms of the disease incident to puerperal females. Dr Abercrombie observes, that at the time at which the disease now mentioned prevailed in the Merchants' Maiden Hospital, he saw in private many cases of the erysipelatous inflammation of the throat in persons of all ages; and in several of these the formation of aphthous crusts and morbid mucus indicated some connection with the diphtheral inflammation of M. Bretonneau. Its resemblance to the subacute and latent forms of puerperal peritonitis is too obvious to require particular remark. The description of the third species of this disease given by Dr Cusack in the present number, will enable the reader to trace this resemblance with greater facility.

Though examples of *chronic peritonitis* are mentioned by

various authors, from Morgagni and Lieutaud to John Hunter and Baillie, the first good account of the pathognomonic symptoms and anatomical characters of this disease as a distinct affection is given by Pemberton; * and the observations of this author have since been well confirmed by the dissections of Dr Black of Newry†, and M. Scoutteten.‡ The account given in the fifth section of the second part of the present volume is still more comprehensive, and illustrates well the various modifications which this disease assumes, and the effects which it produces.

Commencing either with abdominal pain and uneasiness, which continue or increase, and may undergo occasional abatement and aggravation,—or merely with a sense of fulness, which proceeds sometimes with, and not unfrequently without dull pain,—chronic inflammation of the peritonæum advances with a slow but steady pace to injure seriously the membrane in all its divisions. Lymph appears to be effused early, and separating into its coagulating and its serous portions, at once unites all the turns of the intestines with each other, and with the muscular peritonæum, and gives rise to sero-purulent or purulent cysts, either in spaces and cavities formed by the new membranous productions, or in the general cavity, or in both at once. According as the early or late occurrence of the fatal termination allows the physician to inspect early or late after the commencement of the disease, the adhesion resulting from this exudation is soft and lacerable, or hard, firm, and not easily detached. In some instances the peritonæum is occupied extensively with tubercular deposition and accretion; and in some it is thickened and cartilaginous. The mesenteric glands are very generally enlarged.

This process gives rise to many symptoms, some constant, some variable. The most constant are wasting, quick pulse, peculiar dryness of the skin and tongue, a sense of stiffness, and weight of the belly, often abdominal pains, and tympanitic tension and swelling of the belly. The most variable are the degree of pain, constipation or looseness, and affection of the urinary organs.

The tendency of this disease is very generally to the fatal termination; and unless remedial measures be employed early and decidedly, little or nothing can be done, after the disease is established, to avert the fatal issue. In the experience of Dr Abercrombie, who has seen it occur most frequently in young

* On the Abdominal Viscera.

† Clinical and Pathological Reports.

‡ Sur l'Anatomie Pathologique, &c. Archives Generales.

persons from 10 to 15, it has seldom assumed so acute a character as to admit of general blood-letting; and he, therefore, trusts chiefly to free and repeated topical bleeding, blistering, confinement, rest, antiphlogistic regimen, and the mildest diet. Though he admits that when, under such treatment, the symptoms ascribed to chronic peritonitis terminate favourably, it is not always possible to decide with confidence on the exact nature of the malady; he, nevertheless, thinks there is reason to infer its existence when, in a young person, deranged health is connected with abdominal tenderness. Upon such evidence he thinks he has seen the disease terminate favourably in families, some members of which had formerly been cut off by this malady. In one instance, we may add, in which we had every reason to ascribe a sense of painful distension of the belly, with abdominal tenderness, and diffuse tense swelling to this disorder, in a boy of 15, after a full bleeding from the arm, followed at the interval of a few days by leeches, with the use of the antiphlogistic regimen generally, the symptoms gradually receded.

The third part, on “the inflammatory affections of the mucous membrane of the intestinal canal,” is introduced by some good preliminary observations on the morbid appearances of the intestinal mucous surface, in which the author traces the various shades and effects of the inflammatory process, from diffuse red or scarlet injection, red definite patches, or black or brown and softened spots, to ulcers; enlargement of the follicles, follicular ulceration, and tubercular growths. The whole subject is then divided into the three following heads; 1st, active inflammation of the intestinal mucous membrane; 2d, chronic diseases of the intestinal mucous membrane; and 3d, ulcers of the mucous membrane without prominent symptoms. In a subsequent section, the author considers the phenomena and characters of the disease as it occurs in infants. The consideration of these subjects, though more ample and detailed, and illustrated by a greater variety of cases than in the essay originally published by Dr Abercrombie on this subject,* follows, nevertheless, the same tract so closely, and further coincides in results so much with the researches of Andral, Latham, Louis, and Billard, that it is impossible, without much repetition, at the same time, to do the author justice, and avoid misapplying the time of the reader. After noticing briefly, therefore, the leading facts of the inquiry, we must be satisfied with a general reference to the work itself.

The acute form of inflammation of the intestinal mucous

* Medical and Surgical Journal, Vol. xvi. p. 321.

membrane, though generally commencing in the colon, may pass to the ileum, and produce there the same effects which it does in the former situation. Though dangerous, chiefly by its extent, it may also, by causing ulceration or gangrene within small spaces, prove certainly fatal, or become so by perforation and secondary peritonitis. In some instances it is shown to be contagious, or at least epidemic; and of this an example is given (case 85,) in which five individuals of the same family were affected in quick succession, and with fatal issue in two. Lastly, the symptoms vary according to the part of the canal affected. Thus, the affection of the rectum and the lower part of the colon constitutes the dysentery of systematic authors; that of the whole colon and lower part of the ileum appears to constitute the tropical dysentery and *colonitis* of practical authors; and a uniform affection of the small intestine especially, appears to Dr Abercrombie to constitute the fatal modification observed in Indian cholera.

The chronic affection of the intestinal mucous membrane consists either in irregular-shaped dark-red fungous patches, raised above the level of the surrounding membrane; or in small definite ulcers, with the intermediate membrane sound,—evidently the result of follicular inflammation; or irregular ulcerated patches, alternating with fungous elevations; or in induration and thickening, occasionally with distension.

The variety of intestinal ulcers, referred to the next section, belongs, as the author himself admits, to the same pathological head; but, differing in the effects produced, deserve, therefore, to be distinguished in a practical view. They terminate in one of two modes, either by hemorrhage or by peritonæal inflammation, generally in consequence of perforation. As we formerly had occasion to direct the attention of our readers to this subject in speaking of the work of M. Louis, to whom also Dr Abercrombie refers, it is superfluous to dwell longer on the subject.

On the subject of the treatment of the acute and chronic forms of intestinal mucous inflammation, to which the fourth section is devoted, Dr Abercrombie lays down very judicious precepts and directions. These, however, the reader will peruse with most interest in the work itself.

The fifth section, on the disease as it occurs in infants, is much the same with that of the essay already before the profession in this Journal. The whole subject is concluded with some observations on that doctrine, as yet little more than hypothetical, maintained by several of the French pathologists, that disease of the gastro-enteric mucous membrane, or of the duodenal and Peyerian glands, is the cause of the varieties of fever, and most of the forms of dyspeptic disorder. The evi-

dence adduced on this point Dr Abercrombie rejects, as derived partly from pseudo-morbid appearances, partly from changes which, though real, are not uniform, but mere accidental complications, and which are equally likely to be the effects as the cause of fever. Upon much the same grounds, he denies the competency of the disease said by Velpeau to have been observed so long by Bretonneau in the mucous glands (*dothineria*) and which this author seems to wish to identify with the malignant fever of Sydenham, the hospital fever of Pringle, and the typhus of Cullen, to account for the phenomena of that disease. The cases adduced by these authors he admits as facts, but justly regards it as entirely hypothetical to employ these as means to explain the phenomena of fevers.

Upon these points, the opinions of Dr Abercrombie are in general well founded. It is proper, however, to observe, that swelling and redness of the Peyerian glands is a much more frequent affection in the fevers of France, Germany, and other continental countries, than in those of this country. It is at the same time to be remarked, that in maintaining the dothineric affection to be the cause of fever, an error not only as to the pathology, but as to the nosological rank of the disease, is committed, in confounding the typhus of Cullen with the ordinary remittent. The observations of Roederer and Wagler show that this disease is very often attended with an affection of the mucous surfaces; and little doubt can be entertained that the mucous glands are disordered, as causes not of the disease, but of the symptoms. The affection of these glands in diarrhoea and dysentery was originally maintained and in some degree demonstrated by Peyer himself; * and if this takes place idiopathically, there is nothing to prevent the symptomatic affection. Such disorder, however, is not a cause but a complication or effect of fever. That a similar affection takes place in continued fever even in this country has been demonstrated by the inspections of Dr Bright.

In the appendix to this part are considered disease of the mesenteric glands as the effect of disease of the intestinal tissues; tympany, abdominal and intestinal, both of which are justly regarded as symptoms or effects of other disorders; arterial hemorrhage from the rectum; a peculiar species of atrophy dependent on the intestinal canal; and consecutive inflammation, suppuration, and perforation of the *cæcum*, in consequence of the presence of foreign matter in that bowel. These it is unnecessary to consider minutely. The first, second, and fifth sections might have been referred, we conceive, not only with safety but advantage to the heads of disease, either of the mucous

* Exercitatio I. in Parergis Anatomicis. Geneva, 1681.

or the peritoneal tissues, to one or both of which they properly belong. The variety of hemorrhage, treated in the third, appears to consist in aneurismal dilatation of the arteries of the rectum, which, from its functions and occasional extreme distension, is liable to this form of disorder. The case related under section fourth we confess our inability to explain.

The pathology of the liver is treated rather fully and at some extent in two sections, and an appendix subdivided into five sections. This division of the work, however, is entirely new, and therefore merits some particular attention.

In his preliminary remarks on this subject we are happy to find Dr Abercrombie expressing doubts regarding the truth of that fashionable doctrine which has for many years past led practitioners to ascribe to disease of the liver a considerable number of disorders, which are either seated in other organs, or which at all events are not in the least connected with that bulky and convenient gland.

In the first place, a large proportion of complaints, sometimes rather anomalous, but evidently dependent on disorder of the bowels and consecutive derangement in the functions of the whole alimentary canal, it has become fashionable, for a long series of years, but especially since the writings and opinions of Mr Abernethy have been more spoken of than studied, to ascribe to disorder of the liver. An individual whose bowels are slow or costive, is denominated in the nomenclature of this school *bilious*; and nothing but a course of antibilious medicines, but especially calomel and the blue pill, can be expected to cure him. Instead, however, of the liver being disordered in this description of cases, it is perfectly healthy; and in the few instances in which its function seems impaired, this is not the cause, but as sound anatomy and physiology can prove, the effect of the intestinal disorder. It is in many cases of this class that Dr Abercrombie justly observes that the complaints disappear under mild treatment, especially regulation of the bowels and attention to diet, and that those who have received no benefit from long and pernicious courses of mercurial medicines, recover under the use of laxatives.

Secondly, in certain instances of chronic disease in other organs, in which the symptoms are obscure, or from the nature of the affected parts or the stage of the disease, ill-defined, it is the fashion with many practitioners who aim at a prompt and compendious method of diagnosis, to ascribe all the symptoms of deranged health to disease of the liver. We do not wish to be censorious; but regard to the true interest of the art of observing, if not of healing disease, compels us to declare, that we have more than once seen tuberculated lungs in this manner set down to the account of diseased liver; and we

have seen the same unfortunate organ charged with a train of bad health, which dissection proved to be severe ulceration of the mucous membrane of the ileum and colon. We admit that diagnosis is the most difficult part of the task of the physician. But as the difficulties may be diminished, if not overcome, by a patient exercise of the talent of observation, it would be becoming in those practical persons, with whom chiefly these errors take place, to be a little more diffident regarding the exact organ affected, and regarding the exact nature of the affection. In short, we must speak out the truth, however painful to utter or to hear, that we know not a more pernicious species of quackery than that which, under the semblance of great practical experience and tact, and superior penetration, declares decidedly that any given set of symptoms which cannot be well explained, depends on disease of the liver, and must be treated accordingly.

Dr Abercrombie divides the affections of the liver in general into two great heads, those connected with acute disease, and those connected with chronic.

Under the former head we find inflammation, enlargement with dark colour of texture, abscess, inflammatory softening, black softening, white or encephaloid softening, deposition of gelatinous matter, and inordinate distension of the biliary vessels. The first three forms of disease are evidently different stages of the same morbid process. It is sufficient to say, that the varieties of abscess are well described, and that the simple softening is justly represented as an effect of inflammation. The black softening appears also to be an effect of capillary congestion; for from the testimony of Jackson, but especially of Chisholm, who describes many cases of it, it is very frequent in the acute hepatic disorder of the West Indies.* The white or encephaloid softening does not appear to be well named, for instead of resembling *fungus hæmatodes*, which the name would lead the reader to suppose, it consists in the peritonæal coat being occupied by innumerable close-set semitransparent tubercles, small as a split-pea, and in the substance being soft and pulpy like brain. The two last heads consist of short notices, the first from Portal, the second from Boissier.

Under the head of chronic affections, we find chronic inflammation, simple enlargement without change of texture, the *tubera circumscripta* of Dr Farre; pale degeneration without remarkable change of texture; pale degeneration with hardening; dark induration; tubercular disease of the surface (of the peritonæum?), tubercles and *tubera* in the substance; hydatids and watery cysts beneath the peritonæal coat.

* Manual, &c.

These various topics, the author with great candour admits, are enumerated rather in the manner of outline than thoroughly investigated. It must, nevertheless, be acknowledged, that the sketch here given is highly useful, especially in the illustrations which the learned author generally furnishes of the morbid change and its effects on the system. The pale or brick-coloured induration of the liver, in particular, is well illustrated under three heads; 1. the dull white and cartilaginous induration of natural size; 2. the ashen or brick-coloured cartilaginous induration with enlargement; and, 3. the pale coloured induration with diminished size. On the relation between these morbid states of the liver, we find no information in the present sketch. From several cases in which it has occurred to us to inspect examples of them, we have reason to believe that they are either different degrees of the same morbid change, or that they are, at most, different effects of the same morbid action. Almost invariably they may be traced to chronic inflammation of the hepatic texture, however excited, and terminating in a sort of contraction or obliteration of its minute vascular structure. It is quite impossible to inject the organ so minutely as in the healthy state.

In speaking, at the conclusion of this subject, of the treatment of these disorders, Dr Abercrombie justly impresses the necessity of general and local depletion in the acute forms; and inculcates the mildest or even entirely passive measures, and the necessity of dietetic means in the chronic forms of the disorder. In the latter also, he properly condemns the use of mercury, so indiscriminately administered by many practitioners wherever diseased liver is either proved or suspected. Local depletion or counter-irritation, where there is dull pain or weight in the region of the organ, affords a much safer and more effectual means of cure.

In the appendix are discussed the following subjects:—rupture of the liver by external violence, illustrated by one case; diseases of the gall-bladder, consisting of fatal inflammation from a gall-stone impacted in the common duct, and perforation or rupture of the gall-bladder; changes in the quality and quantity of the bile; and a good sketch of the pathological causes of jaundice.

The article on the pathology of the spleen, to which we now come, consists chiefly of the essay already published by the author in this Journal,* expanded, corrected, and in some respects changed in arrangement. In the present volume, Dr Abercrombie considers successively inflammation of the spleen;

* Contributions to the Pathology of the Stomach, the Pancreas, and the Spleen, Part iv. Diseases of the Spleen, Vol. xxii. p. 1, &c.

suppuration, and abscess; softening or black degeneration; simple enlargement; tubercles; colourless induration approaching to scirrhus; hydatids and hemorrhage, either from disease or by violence. On the subject of simple enlargement, Dr Abercrombie remarks the rapidity with which this takes place and disappears under proper management. The cases mentioned by Pemberton,[†] Baillie,[‡] and Bree,[§] in which this organ appears to have attained an unusual size, varying in weight from 3 to 11, or even 20 or 30 pounds, appear to have been of a more chronic character.

The last subject treated in the present volume is the pathology of the pancreas. This, like the last, consists of the essay already published, amplified, somewhat modified, and, upon the whole, improved. These changes, however, we find it impossible to specify; but we may be allowed to say in general, that they tend to render the sketch of the morbid states of the organ both more complete and more accurate.

The foregoing sketch will communicate some idea of what the reader may expect in the present volume. We have, in the course of it, been more studious to show what the research of the author has effected, than to express any opinion of the manner in which it is accomplished. Of the materials only one opinion we believe can be entertained,—that they demonstrate indefatigable industry in the collection, and much acuteness in the observation of the phenomena and effects of disease in the tissues and organs of the human body. We do not, however, conceive that it is necessary for us, to add to the merited reputation of the author by eulogy, which, in his case, is superfluous, and which, in other circumstances, may too often be interpreted as the language of mere civility. When we say that the value of the writings of Dr Abercrombie consists in recording facts which may at all times be referred to for the purpose of illustrating the nature and effects of morbid action, we know that we express the opinion of the most competent judges on topics of this nature.

To the language of indiscriminate panegyric, however, we are averse; and we must be allowed to say that one defect of the present work is its arrangement. This we have already had occasion to notice in the course of our observations; and we have at the same time admitted, that probably deficient or imperfect classification is inseparable from the nature of the subject, which is doubtless complicated. We can easily imagine that the author himself saw some of the difficulties of this de-

† On the Abdominal Viscera.

‡ The Morbid Anatomy, &c. Chapter xi.

§ Medico-Chirurgical Transactions, Vol. iii.

scription, but felt it also impossible to obviate them. On some occasions we think it is possible to trace them to the desire to be useful to the practitioner, and, in describing the progress of morbid action, not to lose sight altogether of the observation of its effects and symptoms, and the nosological importance which physicians assign to the latter. This particularly may be recognized in the appendixes attached to each division of the work, the number of which certainly gives it an appearance rather complicated.

Upon this subject, however, we shall not dwell. It may be safely said that it is rare indeed to meet with a work upon a subject so difficult and extensive, yet with so few blemishes. To the medical reader Dr Abercrombie has rendered an important service, by putting him in possession not only of the results of his own observation, but of the illustrative and confirmatory facts derived from the collateral inquiries of others. To pathological science he has also performed a meritorious duty, in tivating a field hitherto waste, in extending the bounds of knowledge, and in correcting several erroneous notions. These services give his work a high rank among the useful writings of the present age.

ART. III.—*Essai sur la Méthode Endermique, &c.—Essay on the Endermic Method of administering Remedies.* Read before the Royal Academy of Sciences on the 25th September 1826. By ANT. LEMBERT, Hospital-Assistant. Paris 1828. Pp. 125, 8vo.

BY the Endermic method of administering remedies is understood the application of them to the surface of the true skin, previously stripped of its cuticle by the operation of a blister. Of this plan of treatment, which appears to have been first suggested by M. Lember, and which has of late attracted considerable notice in Paris, we have already described an example in an analysis of a paper by M. de Martin on the treatment of intermittents by the external use of sulphate of kina.* M. Lember having now published his own researches on this and several other remedies, the whole subject appears to us to possess interest as well as novelty, and to be well deserving of general attention, as pointing out a new and powerful channel through which the physician may successfully combat disease and study the action of medicines.

The author commences with a short view of the difficulties

* See the Number of this Journal for April last, xxix. 448 ; where, however, the discovery of the endermic method of cure has been incorrectly ascribed to M. Lesueur.