Published in final edited form as:

Microcirculation. 2018 January ; 25(1): . doi:10.1111/micc.12426.

Coronary microvascular Kv1 channels as regulatory sensors of intracellular pyridine nucleotide redox potential

Marc M. Dwenger¹, Vahagn Ohanyan², Manuel F. Navedo³, and Matthew A. Nystoriak¹

¹Diabetes and Obesity Center, Institute of Molecular Cardiology, Department of Medicine, University of Louisville, Louisville, KY 40202

²Department of Integrative Medical Sciences, Northeast Ohio Medical University, Rootstown, OH, 44272

³Department of Pharmacology, University of California, Davis, Davis, CA, 95616

Abstract

Smooth muscle voltage-gated potassium (Kv) channels are important regulators of microvascular tone and tissue perfusion. Recent studies indicate that Kv1 channels represent a key component of the physiological coupling between coronary blood flow and myocardial oxygen demand. While the mechanisms by which metabolic changes in the heart are transduced to alter coronary Kv1 channel gating and promote vasodilation are unclear, a growing body of evidence underscores a pivotal role of Kv1 channels in sensing the cellular redox status. Here, we discuss current knowledge of mechanisms of Kv channel redox regulation with respect to pyridine nucleotide modulation of Kv1 function via ancillary Kv β proteins as well as direct modulation of channel activity via reactive oxygen and nitrogen species. We identify areas of additional research to address the integration of regulatory processes under altered physiological and pathophysiological conditions that may reveal insights into novel treatment strategies for conditions in which the matching of coronary blood supply and myocardial oxygen demand is compromised.

Keywords

ion channels; vascular smooth muscle; endothelium; vasodilation; NADH

1. Introduction

Excitable cells utilize ionic gradients across biological membranes to enable numerous life processes. In this respect, the intracellular environment is rich in potassium ions, and transmembrane voltage-dependent potassium (Kv) channels are widely expressed to control the electrical properties of both eukaryotic and prokaryotic cells. Kv channels, due to their broad phyletic distribution and genetic diversity, are thought to be the oldest of voltage-gated

*To whom correspondence should be addressed: Matthew A. Nystoriak, Diabetes and Obesity Center, Delia Baxter Bldg., 580 S. Preston St., Rm 421C, University of Louisville, Louisville KY 40202 (matthew.nystoriak@louisville.edu, Tel. 502-852-6024, Fax. 502-853-3663).

DR. MANUEL F NAVEDO (Orcid ID : 0000-0001-6864-6594)

DR. MATTHEW A NYSTORIAK (Orcid ID : 0000-0001-8872-158X)

ion channels. Mammalian excitable cells express a number of Kv channel subtypes that work in concert to regulate membrane potential [40]. Through their ability to sense changes in membrane voltage, these channels rapidly respond to depolarization by altering the conformation of their gating apparatus to allow potassium efflux, reducing intracellular positive charge and thus changing the membrane potential to a more hyperpolarized state. In this manner, the activity of Kv channels contributes to the regulation of neuronal action potential firing, muscle contraction, and hormonal secretion. In excitable cells of the cardiovascular system, the coordinated K⁺ efflux by numerous Kv channel subtypes promotes physiological processes such as cardiac action potential repolarization and vascular smooth muscle (VSM) relaxation.

Smooth muscle cells of the vasculature have been reported to express channel subtypes belonging to the Kv1, Kv2, Kv3, Kv4, Kv6, Kv7, Kv9 and Kv11 families (see reference [25], in this issue). In addition to their regulation by membrane voltage, Kv channels are sensitive to functional modulation by a wide variety of intracellular signaling pathways. In small arteries and arterioles of the microcirculation, these pathways tune local and regional blood flow in response to endogenous vasoconstrictor and vasodilator stimuli. In the coronary microcirculation, recent evidence has revealed the importance of redox-mediated Kv1 functional upregulation for physiological enhancement of blood flow (i.e., ‘functional’ or “metabolic” hyperemia), that occurs during periods of increased cardiac oxygen consumption (e.g., during increased heart rate and ventricular contractility) [19,46,47]. While the underlying cellular and molecular mechanisms connecting the level of oxygen demand of cardiomyocytes in an active heart to the redox regulation of Kv1 function in coronary arterial myocytes are not clear, this process likely involves the complex coordination of multiple contributing pathways. Indeed, a thorough understanding of these mechanisms could aid the rational development of novel strategies to improve the coupling between oxygen supply and demand in the heart, which is impaired in a number of conditions such as hypertension, diabetes mellitus, heart failure, and coronary artery disease [6,12,13,17,64,72].

Here, we discuss the known role for Kv1 in linking cellular metabolism to regulation of coronary blood flow. A brief discussion of redox-dependent mechanisms of Kv1 channel regulation with respect to the control of coronary blood flow is followed by a review of Kv1 structural features and the potential role for modulation of channel activity via sensing of changes in cellular pyridine nucleotide status by the regulatory auxiliary Kv β subunits. This is followed by a condensed review of direct influence of reactive oxygen species and reactive nitrogen species on Kv1 channel function. We emphasize cellular redox changes following altered myocardial workload and how these may be sensed by Kv1 channels in the coronary vasculature to promote VSM hyperpolarization and vasodilation to match blood flow with metabolic demand in the heart.

2. Physiological role of Kv1 channels in coronary functional hyperemia

Unlike other organs, the heart extracts most of the oxygen delivered to it by the arterial blood supply at rest [3]. With little reserve for further oxygen extraction, enhancement of pump activity (e.g., during exercise) and greater demand for oxygen by the myocardium

must be coupled with dilation of the coronary arteries and arterioles to instantaneously increase blood flow. This hyperemic response is crucial for maintenance of proper pump function via prevention of myocardial ischemia during periods of increased workloads. While extensive research efforts have aimed to reveal the signaling pathways responsible for metabolic hyperemia in the heart, the mechanisms underlying this process remain poorly understood. In the search for a molecular link between vascular function and myocardial metabolism, decades of pharmacological studies employing receptor and ion channel blockade have yielded controversial conclusions [3,4,15,36,37,55,63,73,74]. The vasoactive metabolite adenosine, which is released from active cardiomyocytes, was long postulated as a primary mediator of coronary artery dilation in response to increase cardiac workload [3]. Yet, studies have shown that administration of adenosine receptor blockers fails to disrupt the normal relationship between myocardial oxygen consumption, coronary vasodilation and blood flow during exercise [4,15,74]. Intriguing results of a study by Tune et al. demonstrated that simultaneous blockade of adenosine receptors, K_{ATP} channels, and nitric oxide synthesis, failed to lower coronary blood flow in exercising dogs [73], suggesting a critical role for an alternative mediator of hyperemia.

A growing body of evidence now supports the concept that coronary hyperemia requires activation of smooth muscle Kv1 channels via a redox-dependent mechanism. For example, a study by Dick et al. reported that the Kv channel blocker 4-aminopyridine decreases coronary blood flow at rest and reduces normalized debt repayment ratio during reactive hyperemia [14]. Moreover, mice in which Kv1.5 is genetically ablated have significantly blunted hyperemic responses to increases in cardiac work via administration of norepinephrine, yet the normal relationship between cardiac work and myocardial blood flow is restored when Kv1.5 is selectively reconstituted in smooth muscle on an otherwise Kv1.5-null background [46], strongly suggesting that loss of hyperemia in the Kv1.5-null animals occurs due to the absence of these channels from the coronary arterial myocytes. These results were corroborated in a separate investigation in which *in vivo* administration of correolide, a selective Kv1 channel blocker, decreased blood flow in response to dobutamine challenge and inhibited blood flow repayment during reactive hyperemia [19]. Interestingly, mice lacking Kv1.3 subunits also exhibit decreased myocardial blood flow in response to increases in cardiac work, [47], suggesting a potential role of heteromeric Kv1 channels in mediating metabolism-dependent vasodilation. *In vitro* work has also indicated that vasodilation of coronary arteries to H_2O_2 , a purported metabolic vasodilator of coronary arteries, is sensitive to the thiol reductant dithioerithritol and 4-aminopyridine, but not the BK_{Ca} channel inhibitor iberiotoxin [56,57]. These results suggest that Kv channels in VSM may respond to redox-modulating myocardial derived metabolites to induce coronary vasodilation.

While Kv1 channel activity has been implicated as an important regulator of human vascular tone, suppression of Kv1 function may contribute to vascular pathologies. Consistent with this, a recent study found that inhibition of H_2O_2 -induced dilation by DPO-1, a selective Kv1.5 channel blocker, was attenuated in arterioles from patients with coronary artery disease, suggesting a reduced capacity for Kv1-dependent vasodilator function [43]. Thus, an improved understanding of mechanisms linking cellular metabolism and vascular Kv1 channel function is essential, as this process likely becomes compromised during pathology.

In the following sections, we discuss redox regulation of Kv1 channels with respect to intracellular pyridine nucleotides and reactive oxygen and nitrogen species and how channel function may be altered to influence coronary vascular tone and myocardial blood flow via these mediators.

3. Mechanisms of Kv1 regulation by cellular pyridine nucleotide redox

Structurally, channels of the Kv1 family share a similar pore complex consisting of four individual α subunits. Each subunit consists of six transmembrane segments (S1–S6) with S1–S4 forming the voltage sensor complex while S5 and S6 form the pore region, together with the S4–S5 linker comprising the gating apparatus [33]. Voltage-sensitivity is conferred by a group of highly conserved positive residues within each S4 segment [1], which render the channel responsive to depolarization. Members of one Kv subfamily can co-assemble with each other to form heterotetrameric pore complexes, an attribute thought to contribute to greater functional diversity of Kv membrane potential regulation [9]. Pore-forming Kv subunits are known to assemble with a variety of intracellular ancillary subunits such as the Kv β s [52]. Like the α pore subunits, the Kv β proteins can also assemble as heterotetramers [54,61], and extend the structure of the channel by approximately 30 Å into the cytosolic compartment [33]. Each β protein binds an α subunit at a T1 docking domain (Fig. 1), which interacts with the voltage sensor and may constrain its conformation [11,34], thus providing a plausible means of regulating the voltage sensitivity of the channel.

A primary mode of regulation of Kv1 in the coronary arteries and arterioles could conceivably arise via specific α/β interactions. The Kv β proteins were found in an early study to share significant identity with proteins belonging to the aldo-keto reductase (AKR) superfamily [39]. The AKR enzymes catalyze the reduction of carbonyl substrates to primary and secondary alcohols in a manner requiring hydride transfer from a nicotinamide adenine dinucleotide (i.e., NAD(P)H) cofactor. Structurally, the Kv β s share key features with other AKR proteins, including an α_8/β_8 barrel and key amino acid residues that are required for catalysis and NAD(P)(H) binding. Consistent with other AKRs, the model three-dimensional structure of the Kv β 2 homotetramer revealed high affinity binding of NADP⁺ within a deep cleft of the active site [21,22]. The reasons for the pairing between a reductase and a voltage-gated potassium channel persisting in a number of excitable cell types are not known. However, interactions between the active site of Kv β with the α subunit voltage sensor support the notion that there may be functional coupling between Kv channels and β -subunits such that either (1) the catalytic activity of Kv β could be controlled by voltage-dependent channel gating via conformational changes in the active site, or perhaps more likely (2) that Kv activity could be differentially affected by catalysis or cofactor oxidation at the active site of Kv β via conformational changes in the voltage sensor affecting the Kv β protein.

Numerous reports support the concept that Kv β proteins strongly influence Kv1 channel function (see reviews [52] and [27]). Studies in heterologous expression systems demonstrate that co-expression of Kv β with Kv α subunits confers rapid inactivation to otherwise non-inactivating K⁺ currents and shifts the voltage-dependence of channel activation towards more negative membrane potentials [52]. Several lines of evidence now

suggest the potential importance of Kv β as a functional sensor of cellular metabolic status such that biochemical modification upon binding pyridine nucleotide cofactors modifies channel gating [27,50,77]. Purified Kv β proteins have been demonstrated to exhibit catalytic function with a wide range of aldehyde and ketone substrates, albeit their in vitro catalytic efficiency is remarkably low relative to other known AKRs [28,69]. Nonetheless, all Kv β proteins bind to both oxidized (i.e. NAD(P)⁺) and reduced (NAD(P)H) pyridine nucleotide cofactors with affinities in the low micromolar range, which is well below normal intracellular levels [5,18]. Moreover, the activation and inactivation properties of Kv1 α / β channels are sensitive to the redox state of NAD(P)(H) that interacts with the Kv β subunits. For example, Tipparaju et al. demonstrated that although non-inactivating currents mediated by Kv1.5 expressed alone in COS-7 cells are insensitive to pyridine nucleotides applied via the patch pipette solution, application of NAD(P)H increased total inactivation and shifted the voltage-dependence of inactivation and activation towards more negative membrane potentials relative to when NAD(P)⁺ was applied in cells co-expressing both Kv1.5 and Kv β 1.3 [71]. A reduction in Kv β 1-mediated channel inactivation upon application of NADP⁺ was also observed for Kv1.1 channels [49]. Similar results were shown for Kv1.5 channels co-expressed with Kv β 3 subunits [70], which similar to Kv β 1, also contain an N-terminal inactivation domain [52]. Redox modulation of channel inactivation by pyridine nucleotides appears to require both the C-terminal region of the Kv1 α protein and electrostatic interactions between the N-terminal region and AKR enzymatic core of the β protein [50,70]. In contrast to effects of pyridine nucleotides on cells expressing Kv1/Kv β 1 and Kv1/Kv β 3 couples, channel inactivation is unchanged by coexpression of Kv1/Kv β 2, which likely reflects the lack of an N-terminal inactivation domain on the Kv β 2 subunit (Fig. 1) [52]. Nonetheless, a robust negative shift in the voltage-dependence of activation of Kv1 channels interacting with Kv β 2 is observed in the presence of reduced pyridine nucleotides [70]. Despite the availability of data demonstrating differential regulation of Kv1 activity by oxidized versus reduced redox couples, the in vivo relevance of these phenomena has not been demonstrated.

The functional role for Kv β proteins in modulating vascular Kv1 channel activity in vivo and tone regulation are currently lacking, yet several studies have reported the expression profiles of Kv1/Kv β proteins in various vascular beds. Acute inhibition of Kv1 channel activity is considered a primary contributor to hypoxic pulmonary vasoconstriction (HPV) [2] and Kv β subunits have been proposed as determinants of redox sensing capacity of Kv1 channels for oxygen tension in the lung parenchyma [10]. Primary cultures of rat arterial myocytes derived from branches and left and right main pulmonary arteries express transcripts for several Kv1 α members as well as Kv β 1.1, Kv β 2, and Kv β 3 subunits [83], yet the specific contribution of these subunits to the HPV response remains unclear. In small mesenteric arterial smooth muscle of rats, mRNA encoding Kv β 1, Kv β 2 and Kv β 3 is present [79]. Whereas, in rabbit portal vein, mRNAs for Kv β 1.1, β 1.2, and β 2, but not β 3.1, are detectable [68]. A recent study examining Kv1 expression in human adipose arterioles revealed mRNA expression of three splice variants of Kv β 1 (i.e., β 1.1, β 1.2 and β 1.3) in these vessels [43]. Note that while many early studies investigating the molecular identity of delayed rectifier 4-aminopyridine-sensitive K⁺ currents in VSM have mostly examined gross mRNA and protein levels in crude ex vivo artery preparations, further characterization of

regional variations in subunit stoichiometry of heteromeric channel structures in the vasculature at the molecular level have remained largely unexplored. Considering the marked differences in channel regulation imparted by the diverse repertoire of channel subunits, distinct bed-specific expression patterns of $Kv\alpha$ and β subunits may indeed reflect differences in the demand for redox regulation for control of organ perfusion and should be revisited with newly available advanced technologies and experimental approaches. Furthermore, altered channel composition following more sustained changes in metabolic demand could represent an important microvascular adaptation in vivo that may become compromised to varying degrees in disease states, and adversely affect vascular resistance and tissue oxygenation.

To begin to examine the aforementioned issues, we recently examined the molecular identity of $Kv\beta$ proteins that are expressed in the murine coronary circulation [45]. In first and second order left anterior descending coronary arteries, we found mRNA transcripts for $Kv\beta 1$, $Kv\beta 2$ and $Kv\beta 3$. At the protein level, $Kv\beta 1$ and $Kv\beta 2$ were detected by Western blot, whereas in situ proximity ligation analyses suggested that both of these $Kv\beta$ isoforms interact with $Kv1.5$ subunits in native coronary Kv channels [45]. Additionally, our results suggest that $Kv\beta 1$ and $Kv\beta 2$ may assemble as heterotetramers in a subpopulation of $Kv1$ channels in coronary arterial myocytes. These findings may have important implications for arterial function in regards to redox regulation of smooth muscle contractility and blood flow. For example, as mentioned above, $Kv\beta 2$ is structurally and functionally unique among the $Kv\beta$ proteins in that it lacks the N-terminal inactivating domain found in $Kv\beta 1$ and $Kv\beta 3$, which occludes the channel pore to confer rapid inactivation to otherwise non-inactivating Kv currents. Thus, the functional influences of these ancillary subunits on $Kv1$ activity in response to changes in $NAD(P)H:NAD(P)^+$ could oppose each other. Experiments performed in COS cells suggest that $Kv\beta 2$ subunits can indeed mask the inactivation function of $Kv\beta 1$ [80], yet the net effect of $Kv\beta$ complexes consisting of more than one $Kv\beta$ isoform on the gating properties of native vascular $Kv1$ channels remains unknown. Nonetheless, given that smooth muscle contractility is strongly influenced by relatively small changes in membrane potential [41,42], $Kv\beta$ may couple vascular function to cellular metabolism by modulating the $Kv1$ window current (Fig. 2). Considering that a subpopulation of $Kv\beta$ subunits induces time-dependent inactivation of $Kv1$ channels, only a fraction of channels in association with these subunits may be available at steady state membrane potentials in coronary arterial myocytes. With the remainder of channels lacking the dominant inactivation function, the $Kv\beta$ -induced negative shift in voltage-dependence of activation may represent a key contributor to greater macroscopic current upon changes in cellular metabolic status. While the functional relevance of this balance is currently unclear, divergent functional regulation of $Kv1$ by distinct auxiliary subunits in association with the channel pore complex may also contribute to fine tuning of $Kv1$ function with respect to coupling of local blood flow with metabolic demands of the local tissue environment.

Importantly, a number of factors could participate in transient or sustained modulation of intracellular pyridine nucleotide redox ratio in VSM upon increases in heart rate and ventricular contractility. These include systemic circulating factors (e.g., bFGF, IL-1 β , TNF α , lactate) as well as diffusible factors released into the interstitium by the active myocardium (e.g., H_2O_2 , adenosine) and a set of myocyte or endothelial-derived growth

factors.[29,35,48,62,76] Additionally, while the heart does not itself undergo sustained hypoxia under physiological conditions, myocardial oxygen consumption is markedly increased during moderate to intense exercise and augmented demand for oxygen by cardiomyocytes could siphon oxygen from the vascular wall, thereby producing a local modest oxygen-deprived microenvironment to promote accumulation of reduced pyridine nucleotides in VSM [78]. However, it remains possible that discrepancies may exist in the regulation of pyridine nucleotide redox between vascular beds. Thus, a detailed assessment of the relationship between vascular smooth muscle pyridine nucleotide redox and cardiac oxygen demand would be valuable to future studies addressing the role of Kv β :NAD(P)(H) interactions in the control of blood flow to the heart. Moreover, future studies employing knock-in strategies to target key residues in the Kv β subunits (e.g., Tyr⁹⁰ of Kv β 2) [38,69] are needed to evaluate the specific in vivo roles for oxido-reductase catalytic function versus cofactor binding, in Kv1-mediated vasodilation.

4. Direct influence of reactive oxygen and nitrogen species on vascular Kv1 function

Elevated levels of reactive oxygen species (ROS) likely contribute to the control of coronary Kv channel function. Experiments examining human, rodent and large animal coronary arterioles have demonstrated that hydrogen peroxide (i.e., H₂O₂), a product of superoxide dismutation, functions as a cardiomyocyte- and endothelium-derived physiological vasoactive agent that can freely diffuse across the plasma membrane and hyperpolarize VSM membrane potential via direct and indirect modulation of K⁺ channel activity [31,56,57,60,84]. Direct functional regulation of Kv channels by oxidative agents may occur via modification of specific cysteine, methionine, histidine and tyrosine residues within the channel complex [59]. In particular, early studies using expression of cloned mammalian Kv subunits in *Xenopus* oocytes have suggested that oxidation of N-terminal cysteine residues within both the Kv α and β subunits inhibits channel inactivation, possibly due to rapid and reversible disulphide bridge formation between the inactivation gate and a neighboring site within the channel complex [54,58]. Application of H₂O₂ was also shown to augment peak Kv current, shift the voltage-dependence of channel activation toward more negative membrane potentials, and accelerate channel activation in CHO cells expressing Kv1.5, in the absence of Kv β co-expression [7]. A recent study in mesenteric arterial myocytes suggests that H₂O₂-mediated Kv potentiation may also occur via incorporation of glutathione in the channel protein at cysteine residues [51], which has also been suggested to promote activation of Cav1.2 voltage-dependent calcium channels [67]. However, in the presence of heightened oxidative stress, H₂O₂ fails to further activate Kv channels and may even lead to channel inhibition, thus supporting a bidirectional functional influence of oxidative modification, depending on the redox state of the cell.

Superoxide production by the mitochondria can also lead to the formation of peroxynitrite (ONOO⁻) via reaction with nitric oxide [26]. ONOO⁻ is known to be highly reactive with tyrosine residues and can alter the function of cellular proteins, including voltage-gated ion channels [32]. In addition to numerous pathways by which nitric oxide and reactive nitrogen species could influence Kv-mediated vasodilation [16,44,66,81], channel activity could be

directly altered by reaction of ONOO^- with channel proteins. In coronary arteries of rats, increased production of ONOO^- may impair Kv-mediated vasodilation via nitration of tyrosine residues in Kv1.2 subunits [30]. In addition to nitration of tyrosine residues, S-nitrosothiols, like S-nitroso-N-actylpenicillamine (SNAP), have been shown to modify cysteines in the L-type Ca^{2+} channel through multiple mechanisms including transnitrosation, mixed disulfide bonds, and disulfide bond formation between cysteines [24]. A biotin-switch assay revealed the presence of S-nitrosylated cysteine residues in SNAP-treated ltk^- cells stably expressing human Kv1.5. Through molecular modeling, two cysteines in the S2 segment (i.e., C331 and C346) were identified as a potential site of S-nitrosylation, as S-nitrosothiol groups at these cysteines could be stabilized by hydrogen bridge bonds with I262, located in S1, and R342, located in S2. Interestingly, this modification correlated with a reduction in Kv1.5 current density. Although direct modification identified thus far at cysteine and tyrosine residues generally appears to be associated with channel inhibition, it is unclear whether direct modification of Kv channels by NO-derived factors impacts the regulation of coronary vasomotor tone and myocardial blood flow. Nonetheless, nitration of Kv channels may impair vasodilatory function in the presence of pathological conditions in which superoxide production is elevated, such as hyperglycemia and diabetes. Future work is needed to address the precise functional consequences of specific tyrosine and cysteine modifications within the Kv channel complex and the differential effects conferred by α/β subunit compositions in the context of physiological and pathological regulation of coronary vasodilation.

6. Summary

Kv1 channels in the microcirculation are capable of sensing cellular redox state by a variety of mechanisms which ultimately act in concert to control VSM membrane potential to adapt blood flow to constantly changing metabolic conditions (Fig 3). In addition to their regulation by pyridine nucleotide redox and reactive oxygen/nitrogen species discussed here, Kv1 channel subunits can be functionally modulated by a number of other modifications including phosphorylation, palmitoylation, and glycosylation [20,23,75]. Future work using in silico modeling could help shape a better understanding of how this complex network of pathways works in an integrated fashion to influence vascular function in a variety of physiological and pathological states. Advanced gene editing technology now offers an exceptional tool to determine precise sites of redox sensing within the channel structure in the setting of native channels of the vasculature. Importantly, a number of both gain-of-function and loss-of-function mutations have been identified in Kv1 channel subunits [8,53,65,82]. In addition to the impact that these mutations likely have on cardiac and neuronal repolarization, it is plausible that altered coupling between myocardial metabolism and coronary blood flow may also contribute to cardiovascular disease in patients harboring these mutations. Thus, future advances in our understanding of redox regulation of coronary Kv channels during disease states could ultimately lead to improved strategies to enhance coupling between metabolic demand and myocardial blood flow for therapeutic benefit.

Acknowledgments

We thank Dr. Aruni Bhatnagar for critically reading the manuscript. This work was supported in part by grants from the National Institutes of Health (GM103492, HL098200, and HL121059), American Heart Association (16SDG27260070), and the University of Louisville School of Medicine.

References

1. Aggarwal SK, MacKinnon R. Contribution of the S4 segment to gating charge in the Shaker K⁺ channel. *Neuron*. 1996; 16:1169–1177. [PubMed: 8663993]
2. Archer SL, London B, Hampl V, Wu X, Nsair A, Puttagunta L, Hashimoto K, Waite RE, Michelakis ED. Impairment of hypoxic pulmonary vasoconstriction in mice lacking the voltage-gated potassium channel Kv1.5. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology*. 2001; 15:1801–1803. [PubMed: 11481235]
3. Bacchus AN, Ely SW, Knabb RM, Rubio R, Berne RM. Adenosine and coronary blood flow in conscious dogs during normal physiological stimuli. *The American journal of physiology*. 1982; 243:H628–633. [PubMed: 7124971]
4. Bache RJ, Dai XZ, Schwartz JS, Homans DC. Role of adenosine in coronary vasodilation during exercise. *Circulation research*. 1988; 62:846–853. [PubMed: 3349577]
5. Bessho M, Tajima T, Hori S, Satoh T, Fukuda K, Kyotani S, Ohnishi Y, Nakamura Y. NAD and NADH values in rapidly sampled dog heart tissues by two different extraction methods. *Anal Biochem*. 1989; 182:304–308. [PubMed: 2610348]
6. Brush JE Jr, Cannon RO 3rd, Schenke WH, Bonow RO, Leon MB, Maron BJ, Epstein SE. Angina due to coronary microvascular disease in hypertensive patients without left ventricular hypertrophy. *The New England journal of medicine*. 1988; 319:1302–1307. [PubMed: 3185633]
7. Caouette D, Dongmo C, Berube J, Fournier D, Daleau P. Hydrogen peroxide modulates the Kv1.5 channel expressed in a mammalian cell line. *Naunyn Schmiedeberg's Arch Pharmacol*. 2003; 368:479–486. [PubMed: 14614593]
8. Christophersen IE, Olesen MS, Liang B, Andersen MN, Larsen AP, Nielsen JB, Haunso S, Olesen SP, Tveit A, Svendsen JH, Schmitt N. Genetic variation in KCNA5: impact on the atrial-specific potassium current I_{Kur} in patients with lone atrial fibrillation. *Eur Heart J*. 2013; 34:1517–1525. [PubMed: 23264583]
9. Coetzee WA, Amarillo Y, Chiu J, Chow A, Lau D, McCormack T, Moreno H, Nadal MS, Ozaita A, Pountney D, Saganich M, Vega-Saenz de Miera E, Rudy B. Molecular diversity of K⁺ channels. *Ann N Y Acad Sci*. 1999; 868:233–285. [PubMed: 10414301]
10. Coppock EA, Tamkun MM. Differential expression of K(V) channel alpha- and beta-subunits in the bovine pulmonary arterial circulation. *Am J Physiol Lung Cell Mol Physiol*. 2001; 281:L1350–1360. [PubMed: 11704530]
11. Darman RB, Ivy AA, Ketty V, Blaustein RO. Constraints on voltage sensor movement in the shaker K⁺ channel. *The Journal of general physiology*. 2006; 128:687–699. [PubMed: 17101817]
12. Dayanikli F, Grambow D, Muzik O, Mosca L, Rubenfire M, Schwaiger M. Early detection of abnormal coronary flow reserve in asymptomatic men at high risk for coronary artery disease using positron emission tomography. *Circulation*. 1994; 90:808–817. [PubMed: 8044952]
13. Di Carli MF, Charytan D, McMahon GT, Ganz P, Dorbala S, Schelbert HR. Coronary circulatory function in patients with the metabolic syndrome. *J Nucl Med*. 2011; 52:1369–1377. [PubMed: 21849399]
14. Dick GM, Bratz IN, Borbouse L, Payne GA, Dincer UD, Knudson JD, Rogers PA, Tune JD. Voltage-dependent K⁺ channels regulate the duration of reactive hyperemia in the canine coronary circulation. *American journal of physiology Heart and circulatory physiology*. 2008; 294:H2371–2381. [PubMed: 18375717]
15. Duncker DJ, Stubenitsky R, Verdouw PD. Role of adenosine in the regulation of coronary blood flow in swine at rest and during treadmill exercise. *The American journal of physiology*. 1998; 275:H1663–1672. [PubMed: 9815074]

16. Favalaro JL, Kemp-Harper BK. Redox variants of NO (NO{middle dot} and HNO) elicit vasorelaxation of resistance arteries via distinct mechanisms. *Am J Physiol Heart Circ Physiol*. 2009; 296:H1274–1280. [PubMed: 19252101]
17. Fukuda D, Yoshiyama M, Shimada K, Yamashita H, Ehara S, Nakamura Y, Kamimori K, Tanaka A, Kawarabayashi T, Yoshikawa J. Relation between aortic stiffness and coronary flow reserve in patients with coronary artery disease. *Heart*. 2006; 92:759–762. [PubMed: 16216858]
18. Glock GE, McLean P. Levels of oxidized and reduced diphosphopyridine nucleotide and triphosphopyridine nucleotide in animal tissues. *Biochem J*. 1955; 61:388–390. [PubMed: 13269372]
19. Goodwill AG, Noblet JN, Sassoon D, Fu L, Kassab GS, Schepers L, Herring BP, Rottgen TS, Tune JD, Dick GM. Critical contribution of KV1 channels to the regulation of coronary blood flow. *Basic Res Cardiol*. 2016; 111:56. [PubMed: 27496159]
20. Gubitosi-Klug RA, Mancuso DJ, Gross RW. The human Kv1.1 channel is palmitoylated, modulating voltage sensing: Identification of a palmitoylation consensus sequence. *Proc Natl Acad Sci U S A*. 2005; 102:5964–5968. [PubMed: 15837928]
21. Gulbis JM, Mann S, MacKinnon R. Structure of a voltage-dependent K⁺ channel beta subunit. *Cell*. 1999; 97:943–952. [PubMed: 10399921]
22. Gulbis JM, Zhou M, Mann S, MacKinnon R. Structure of the cytoplasmic beta subunit-T1 assembly of voltage-dependent K⁺ channels. *Science*. 2000; 289:123–127. [PubMed: 10884227]
23. Hogan-Cann A, Li W, Guo J, Yang T, Zhang S. Proteolytic cleavage in the S1–S2 linker of the Kv1.5 channel does not affect channel function. *Biochim Biophys Acta*. 2016; 1858:1082–1090. [PubMed: 26874203]
24. Hu H, Chiamvimonvat N, Yamagishi T, Marban E. Direct inhibition of expressed cardiac L-type Ca²⁺ channels by S-nitrosothiol nitric oxide donors. *Circ Res*. 1997; 81:742–752. [PubMed: 9351448]
25. Jackson WF. KV channels and the regulation of vascular smooth muscle tone. *Microcirculation*. 2017
26. Kelm M. Nitric oxide metabolism and breakdown. *Biochim Biophys Acta*. 1999; 1411:273–289. [PubMed: 10320663]
27. Kilfoil PJ, Tipparaju SM, Barski OA, Bhatnagar A. Regulation of ion channels by pyridine nucleotides. *Circulation research*. 2013; 112:721–741. [PubMed: 23410881]
28. Kratzer R, Kavanagh KL, Wilson DK, Nidetzky B. Studies of the enzymic mechanism of *Candida tenuis* xylose reductase (AKR 2B5): X-ray structure and catalytic reaction profile for the H113A mutant. *Biochemistry*. 2004; 43:4944–4954. [PubMed: 15109252]
29. Lassegue B, Griendling KK. NADPH oxidases: functions and pathologies in the vasculature. *Arterioscler Thromb Vasc Biol*. 2010; 30:653–661. [PubMed: 19910640]
30. Li H, Gutterman DD, Rusch NJ, Bubolz A, Liu Y. Nitration and functional loss of voltage-gated K⁺ channels in rat coronary microvessels exposed to high glucose. *Diabetes*. 2004; 53:2436–2442. [PubMed: 15331556]
31. Liu Y, Bubolz AH, Mendoza S, Zhang DX, Gutterman DD. H₂O₂ is the transferrable factor mediating flow-induced dilation in human coronary arterioles. *Circulation research*. 2011; 108:566–573. [PubMed: 21233456]
32. Liu Y, Terata K, Chai Q, Li H, Kleinman LH, Gutterman DD. Peroxynitrite inhibits Ca²⁺-activated K⁺ channel activity in smooth muscle of human coronary arterioles. *Circulation research*. 2002; 91:1070–1076. [PubMed: 12456494]
33. Long SB, Campbell EB, Mackinnon R. Crystal structure of a mammalian voltage-dependent Shaker family K⁺ channel. *Science*. 2005; 309:897–903. [PubMed: 16002581]
34. Long SB, Campbell EB, Mackinnon R. Voltage sensor of Kv1.2: structural basis of electromechanical coupling. *Science*. 2005; 309:903–908. [PubMed: 16002579]
35. Lotscher HR, Winterhalter KH, Carafoli E, Richter C. Hydroperoxides can modulate the redox state of pyridine nucleotides and the calcium balance in rat liver mitochondria. *Proc Natl Acad Sci U S A*. 1979; 76:4340–4344. [PubMed: 41241]

36. Lynch FM, Austin C, Heagerty AM, Izzard AS. Adenosine- and hypoxia-induced dilation of human coronary resistance arteries: evidence against the involvement of K(ATP) channels. *British journal of pharmacology*. 2006; 147:455–458. [PubMed: 16341231]
37. Lynch FM, Austin C, Heagerty AM, Izzard AS. Adenosine and hypoxic dilation of rat coronary small arteries: roles of the ATP-sensitive potassium channel, endothelium, and nitric oxide. *American journal of physiology Heart and circulatory physiology*. 2006; 290:H1145–1150. [PubMed: 16243919]
38. McCormack K, Connor JX, Zhou L, Ho LL, Ganetzky B, Chiu SY, Messing A. Genetic analysis of the mammalian K⁺ channel beta subunit Kvbeta 2 (Kcnab2). *The Journal of biological chemistry*. 2002; 277:13219–13228. [PubMed: 11825900]
39. McCormack T, McCormack K. Shaker K⁺ channel beta subunits belong to an NAD(P)H-dependent oxidoreductase superfamily. *Cell*. 1994; 79:1133–1135. [PubMed: 8001150]
40. Moran Y, Barzilai MG, Liebeskind BJ, Zakon HH. Evolution of voltage-gated ion channels at the emergence of Metazoa. *J Exp Biol*. 2015; 218:515–525. [PubMed: 25696815]
41. Nelson MT, Patlak JB, Worley JF, Standen NB. Calcium channels, potassium channels, and voltage dependence of arterial smooth muscle tone. *The American journal of physiology*. 1990; 259:C3–18. [PubMed: 2164782]
42. Nelson MT, Quayle JM. Physiological roles and properties of potassium channels in arterial smooth muscle. *The American journal of physiology*. 1995; 268:C799–822. [PubMed: 7733230]
43. Nishijima Y, Cao S, Chabowski DS, Korishettar A, Ge A, Zheng X, Sparapani R, Gutterman DD, Zhang DX. Contribution of KV1.5 Channel to Hydrogen Peroxide-Induced Human Arteriolar Dilation and Its Modulation by Coronary Artery Disease. *Circulation research*. 2017; 120:658–669. [PubMed: 27872049]
44. Nunez L, Vaquero M, Gomez R, Caballero R, Mateos-Caceres P, Macaya C, Iriepa I, Galvez E, Lopez-Farre A, Tamargo J, Delpon E. Nitric oxide blocks hKv1.5 channels by S-nitrosylation and by a cyclic GMP-dependent mechanism. *Cardiovasc Res*. 2006; 72:80–89. [PubMed: 16876149]
45. Nystoriak MA, Zhang D, Jagatheesan G, Bhatnagar A. Heteromeric complexes of aldo-keto reductase auxiliary KVbeta subunits (AKR6A) regulate sarcolemmal localization of KV1.5 in coronary arterial myocytes. *Chem Biol Interact*. 2017
46. Ohanyan V, Yin L, Bardakjian R, Kolz C, Enrick M, Hakobyan T, Kmetz J, Bratz I, Luli J, Nagane M, Khan N, Hou H, Kuppusamy P, Graham J, Fu FK, Janota D, Oyewumi MO, Logan S, Lindner JR, Chilian WM. Requisite Role of Kv1.5 Channels in Coronary Metabolic Dilation. *Circ Res*. 2015; 117:612–621. [PubMed: 26224794]
47. Ohanyan V, Yin L, Bardakjian R, Kolz C, Enrick M, Hakobyan T, Luli J, Graham K, Khayata M, Logan S, Kmetz J, Chilian WM. Kv1.3 channels facilitate the connection between metabolism and blood flow in the heart. *Microcirculation*. 2017; 24
48. Oka S, Hsu CP, Sadoshima J. Regulation of cell survival and death by pyridine nucleotides. *Circ Res*. 2012; 111:611–627. [PubMed: 22904041]
49. Pan Y, Weng J, Cao Y, Bhosle RC, Zhou M. Functional coupling between the Kv1.1 channel and aldo-ketoreductase Kvbeta1. *The Journal of biological chemistry*. 2008; 283:8634–8642. [PubMed: 18222921]
50. Pan Y, Weng J, Levin EJ, Zhou M. Oxidation of NADPH on Kvbeta1 inhibits ball-and-chain type inactivation by restraining the chain. *Proceedings of the National Academy of Sciences of the United States of America*. 2011; 108:5885–5890. [PubMed: 21436029]
51. Park SW, Noh HJ, Sung DJ, Kim JG, Kim JM, Ryu SY, Kang K, Kim B, Bae YM, Cho H. Hydrogen peroxide induces vasorelaxation by enhancing 4-aminopyridine-sensitive Kv currents through S-glutathionylation. *Pflugers Arch*. 2015; 467:285–297. [PubMed: 24756196]
52. Pongs O, Schwarz JR. Ancillary subunits associated with voltage-dependent K⁺ channels. *Physiol Rev*. 2010; 90:755–796. [PubMed: 20393197]
53. Portero V, Le Scouarnec S, Es-Salah-Lamoureux Z, Burel S, Gourraud JB, Bonnaud S, Lindenbaum P, Simonet F, Violleau J, Baron E, Moreau E, Scott C, Chatel S, Loussouarn G, O'Hara T, Mabo P, Dina C, Le Marec H, Schott JJ, Probst V, Baro I, Marionneau C, Charpentier F, Redon R. Dysfunction of the Voltage-Gated K⁺ Channel beta2 Subunit in a Familial Case of Brugada Syndrome. *J Am Heart Assoc*. 2016; 5

54. Rettig J, Heinemann SH, Wunder F, Lorra C, Parcej DN, Dolly JO, Pongs O. Inactivation properties of voltage-gated K⁺ channels altered by presence of beta-subunit. *Nature*. 1994; 369:289–294. [PubMed: 8183366]
55. Richmond KN, Tune JD, Gorman MW, Feigl EO. Role of K⁺ATP channels in local metabolic coronary vasodilation. *The American journal of physiology*. 1999; 277:H2115–2123. [PubMed: 10600828]
56. Rogers PA, Chilian WM, Bratz IN, Bryan RM Jr, Dick GM. H₂O₂ activates redox- and 4-aminopyridine-sensitive K_v channels in coronary vascular smooth muscle. *Am J Physiol Heart Circ Physiol*. 2007; 292:H1404–1411. [PubMed: 17071731]
57. Rogers PA, Dick GM, Knudson JD, Focardi M, Bratz IN, Swafford AN Jr, Saitoh S, Tune JD, Chilian WM. H₂O₂-induced redox-sensitive coronary vasodilation is mediated by 4-aminopyridine-sensitive K⁺ channels. *Am J Physiol Heart Circ Physiol*. 2006; 291:H2473–2482. [PubMed: 16751285]
58. Ruppersberg JP, Stocker M, Pongs O, Heinemann SH, Frank R, Koenen M. Regulation of fast inactivation of cloned mammalian IK(A) channels by cysteine oxidation. *Nature*. 1991; 352:711–714. [PubMed: 1908562]
59. Sahoo N, Hoshi T, Heinemann SH. Oxidative modulation of voltage-gated potassium channels. *Antioxid Redox Signal*. 2014; 21:933–952. [PubMed: 24040918]
60. Saitoh S, Kiyooka T, Rocic P, Rogers PA, Zhang C, Swafford A, Dick GM, Viswanathan C, Park Y, Chilian WM. Redox-dependent coronary metabolic dilation. *American journal of physiology Heart and circulatory physiology*. 2007; 293:H3720–3725. [PubMed: 17965288]
61. Scott VE, Rettig J, Parcej DN, Keen JN, Findlay JB, Pongs O, Dolly JO. Primary structure of a beta subunit of alpha-dendrotoxin-sensitive K⁺ channels from bovine brain. *Proceedings of the National Academy of Sciences of the United States of America*. 1994; 91:1637–1641. [PubMed: 8127858]
62. Sistare FD, Haynes RC Jr. The interaction between the cytosolic pyridine nucleotide redox potential and gluconeogenesis from lactate/pyruvate in isolated rat hepatocytes. Implications for investigations of hormone action. *J Biol Chem*. 1985; 260:12748–12753. [PubMed: 4044607]
63. Stepp DW, Kroll K, Feigl EO. K⁺ATP channels and adenosine are not necessary for coronary autoregulation. *The American journal of physiology*. 1997; 273:H1299–1308. [PubMed: 9321819]
64. Strauer BE, Motz W, Vogt M, Schwartzkopff B. Impaired coronary flow reserve in NIDDM: a possible role for diabetic cardiopathy in humans. *Diabetes*. 1997; 46(Suppl 2):S119–124. [PubMed: 9285513]
65. Syrbe S, Hedrich UBS, Riesch E, Djemie T, Muller S, Moller RS, Maher B, Hernandez-Hernandez L, Synofzik M, Caglayan HS, Arslan M, Serratos JM, Nothnagel M, May P, Krause R, Loffler H, Detert K, Dorn T, Vogt H, Kramer G, Schols L, Mullis PE, Linnankivi T, Lehesjoki AE, Sterbova K, Craiu DC, Hoffman-Zacharska D, Korff CM, Weber YG, Steinlin M, Gallati S, Bertsche A, Bernhard MK, Merkenschlager A, Kiess W, Euro ERESc, Gonzalez M, Zuchner S, Palotie A, Suls A, De Jonghe P, Helbig I, Biskup S, Wolff M, Maljevic S, Schule R, Sisodiya SM, Weckhuysen S, Lerche H, Lemke JR. De novo loss- or gain-of-function mutations in KCNA2 cause epileptic encephalopathy. *Nat Genet*. 2015; 47:393–399. [PubMed: 25751627]
66. Tanaka Y, Tang G, Takizawa K, Otsuka K, Eghbali M, Song M, Nishimaru K, Shigenobu K, Koike K, Stefani E, Toro L. K_v channels contribute to nitric oxide- and atrial natriuretic peptide-induced relaxation of a rat conduit artery. *J Pharmacol Exp Ther*. 2006; 317:341–354. [PubMed: 16394199]
67. Tang H, Viola HM, Filipovska A, Hool LC. Ca(v)1.2 calcium channel is glutathionylated during oxidative stress in guinea pig and ischemic human heart. *Free Radic Biol Med*. 2011; 51:1501–1511. [PubMed: 21810465]
68. Thorneloe KS, Chen TT, Kerr PM, Grier EF, Horowitz B, Cole WC, Walsh MP. Molecular composition of 4-aminopyridine-sensitive voltage-gated K(+) channels of vascular smooth muscle. *Circulation research*. 2001; 89:1030–1037. [PubMed: 11717160]
69. Tipparaju SM, Barski OA, Srivastava S, Bhatnagar A. Catalytic mechanism and substrate specificity of the beta-subunit of the voltage-gated potassium channel. *Biochemistry*. 2008; 47:8840–8854. [PubMed: 18672894]

70. Tipparaju SM, Li XP, Kilfoil PJ, Xue B, Uversky VN, Bhatnagar A, Barski OA. Interactions between the C-terminus of Kv1.5 and Kvbeta regulate pyridine nucleotide-dependent changes in channel gating. *Pflugers Arch.* 2012; 463:799–818. [PubMed: 22426702]
71. Tipparaju SM, Saxena N, Liu SQ, Kumar R, Bhatnagar A. Differential regulation of voltage-gated K⁺ channels by oxidized and reduced pyridine nucleotide coenzymes. *American journal of physiology Cell physiology.* 2005; 288:C366–376. [PubMed: 15469953]
72. Tsagalou EP, Anastasiou-Nana M, Agapitos E, Gika A, Drakos SG, Terrovitis JV, Ntalianis A, Nanas JN. Depressed coronary flow reserve is associated with decreased myocardial capillary density in patients with heart failure due to idiopathic dilated cardiomyopathy. *Journal of the American College of Cardiology.* 2008; 52:1391–1398. [PubMed: 18940529]
73. Tune JD, Richmond KN, Gorman MW, Feigl EO. K(ATP)(+) channels, nitric oxide, and adenosine are not required for local metabolic coronary vasodilation. *American journal of physiology Heart and circulatory physiology.* 2001; 280:H868–875. [PubMed: 11158988]
74. Tune JD, Richmond KN, Gorman MW, Olsson RA, Feigl EO. Adenosine is not responsible for local metabolic control of coronary blood flow in dogs during exercise. *American journal of physiology Heart and circulatory physiology.* 2000; 278:H74–84. [PubMed: 10644586]
75. Watanabe I, Zhu J, Sutachan JJ, Gottschalk A, Recio-Pinto E, Thornhill WB. The glycosylation state of Kv1.2 potassium channels affects trafficking, gating, and simulated action potentials. *Brain Res.* 2007; 1144:1–18. [PubMed: 17324383]
76. Weiner MW, Lardy HA. Reduction of pyridine nucleotides induced by adenosine diphosphate in kidney mitochondria. The influence of sodium, magnesium, and inhibitors of oxidative phosphorylation. *J Biol Chem.* 1973; 248:7682–7687. [PubMed: 4147982]
77. Weng J, Cao Y, Moss N, Zhou M. Modulation of voltage-dependent Shaker family potassium channels by an aldo-keto reductase. *The Journal of biological chemistry.* 2006; 281:15194–15200. [PubMed: 16569641]
78. Wolin MS, Ahmad M, Gupte SA. Oxidant and redox signaling in vascular oxygen sensing mechanisms: basic concepts, current controversies, and potential importance of cytosolic NADPH. *Am J Physiol Lung Cell Mol Physiol.* 2005; 289:L159–173. [PubMed: 16002998]
79. Xu C, Lu Y, Tang G, Wang R. Expression of voltage-dependent K(+) channel genes in mesenteric artery smooth muscle cells. *The American journal of physiology.* 1999; 277:G1055–1063. [PubMed: 10564112]
80. Xu J, Li M. Kvbeta2 inhibits the Kvbeta1-mediated inactivation of K⁺ channels in transfected mammalian cells. *The Journal of biological chemistry.* 1997; 272:11728–11735. [PubMed: 9115226]
81. Yang Q, Chen SR, Li DP, Pan HL. Kv1.1/1.2 channels are downstream effectors of nitric oxide on synaptic GABA release to preautonomic neurons in the paraventricular nucleus. *Neuroscience.* 2007; 149:315–327. [PubMed: 17869444]
82. Yang Y, Li J, Lin X, Yang Y, Hong K, Wang L, Liu J, Li L, Yan D, Liang D, Xiao J, Jin H, Wu J, Zhang Y, Chen YH. Novel KCNA5 loss-of-function mutations responsible for atrial fibrillation. *J Hum Genet.* 2009; 54:277–283. [PubMed: 19343045]
83. Yuan XJ, Wang J, Juhaszova M, Golovina VA, Rubin LJ. Molecular basis and function of voltage-gated K⁺ channels in pulmonary arterial smooth muscle cells. *The American journal of physiology.* 1998; 274:L621–635. [PubMed: 9575881]
84. Zhang DX, Borbouse L, Gebremedhin D, Mendoza SA, Zinkevich NS, Li R, Gutterman DD. H₂O₂-induced dilation in human coronary arterioles: role of protein kinase G dimerization and large-conductance Ca²⁺-activated K⁺ channel activation. *Circulation research.* 2012; 110:471–480. [PubMed: 22158710]

Figure 1. K_vβ-dependent regulation of K_v1 activity by pyridine nucleotide cofactors
 (A) Regulation of K_v1 function by Kvβ1 and Kvβ3 subunits, which possess NH₂-terminal ball-and-chain-like inactivation. Inactivation is enhanced by bound pyridine nucleotides in their reduced form (i.e., NAD(P)H). (B) Regulation of K_v1 function by Kvβ2 subunits. Kvβ2 proteins lack the NH₂-terminal inactivation domain. Voltage-dependence of activation is shifted towards more negative membrane potentials in the presence of reduced pyridine nucleotide cofactors. Transition between states can occur by exchange of pyridine nucleotides or by catalytic activity to oxidize bound pyridine nucleotide cofactors.

Figure 2. Hypothetical model of coronary blood flow regulation by pyridine nucleotide-mediated activation of Kv1 channels

Flow diagram on the left shows role of smooth muscle Kv channels in opposing membrane potential depolarization and vasoconstriction. Expanded window shows proposed model of Kv1 activation during periods of elevated cardiac workload. In this scheme, increases in myocardial oxygen demand leads to changes in pyridine nucleotide redox status in coronary arterial myocytes. Accumulation of reduced pyridine nucleotides could shift the threshold and voltage-dependence of activation of available Kv1 channels further into the narrow window of physiological steady-state membrane potential in arterial myocytes of pressurized arteries and arterioles (−60 to −35 mV), thereby significantly increasing Kv1 activity leading to membrane hyperpolarization, vasodilation, and enhanced blood flow.

Figure 3. Coronary smooth muscle Kv1 channel modulation by cellular redox

Shown are the major discussed pathways which could contribute to functional alteration of Kv1 activity upon changes in myocardial oxygen demand.