Case report

Can an exostosis be free floating without any continuity to the underlying bone: A diagnostic dilemma

Manish Kanwat^{a,*}, Vivek Jangira^b, Bhavna Sharma^c, Dushyant Chauhan^d^a Department of Orthopedics, Lady Hardinge Medical College, India^b Department of Orthopedics, PGIMER, Dr. Ram Manohar Lohia Hospital, India^c Department of Pathology, PGIMER, Dr. Ram Manohar Lohia Hospital, India^d Department of Orthopedics, Lady Hardinge Medical College, India

ARTICLE INFO

Article history:

Received 24 July 2017

Received in revised form 2 September 2017

Accepted 9 September 2017

Available online 11 September 2017

Keywords:

Osteochondroma

Femur

Extraskeletal

ABSTRACT

Osteochondroma or exostosis is the most common primary bone tumor containing both bone and cartilage. Soft tissue osteochondromas have been described at various locations, however, to the best of our knowledge, there is only a single case report of a soft tissue osteochondroma around the femoral neck. We, hereby report second such case.

Case Report: A forty year old female presented with swelling on medial aspect of left thigh since four years. Radiographs showed a large mass at the inferior surface of the left femoral neck. CT scan revealed a well defined lobulated mass lesion involving muscles of adductor compartment of thigh. MRI showed lobulated periarticular, intramuscular, non enhancing lesion isointense to the bone, but without any continuity to the adjacent femur. Histopathology of the excised lesion turned out to be osteochondroma with origin from soft tissue. This case highlights an important differential diagnosis to be considered whenever an ossified mass is located in soft tissue.

© 2017

1. Introduction

Osteochondroma or exostosis is the most common primary bone tumor containing both bone and cartilage. It is usually an extra-articular lesion arising from the metaphysis of long bones most commonly at the knee and forearm [1]. Soft tissue osteochondromas have been described at various locations [2–6], however, to the best of our knowledge, there is only a single case report of a soft tissue osteochondroma around the femoral neck [7]. We, hereby report second such case.

2. Case report

A forty year old female presented with progressively increasing swelling on medial aspect of left thigh alongwith pain in left groin since four years. The swelling was associated with restricted motion of the left hip. Radiographs of the left hip showed a large mass at the inferior surface of the left femoral neck [Fig. 1]. CT scan

revealed a well defined lobulated mass lesion with extensive flocculent calcification measuring $8 \times 8 \times 7.5$ cm involving muscles of adductor compartment of thigh extending superiorly from the level of neck of femur upto 4.5 cm below the lesser trochanter inferiorly with <1 cm space between the mass and lesser trochanter [Fig. 2]. A diagnosis of benign pathology (? calcinosis/? heterotopic ossification) was rendered and MRI was advised. MRI showed lobulated periarticular, intramuscular, non enhancing lesion isointense to the bone but, to surprise without any continuity to the adjacent femur. Periostitis was suspected as there was minimal periosteal thickening in femoral neck. Possibility of heterotopic ossification was suggested [Fig. 3]. A surgical excision was performed using the Ludloff's approach. The patient was placed in supine position with left hip flexed, abducted and slightly externally rotated. A longitudinal incision was made on medial aspect of thigh beginning about 2.5 cm distal to pubic tubercle. A plane was made between adductor longus and gracilis and deep dissection was continued in between adductor brevis and adductor magnus till lesser trochanter was reached. Special care was taken to protect the posterior division of obturator nerve. The tumor was palpated and complete excision was done. Complete surprise was the histopathology of the excised lesion, which

* Corresponding author at: Flat Number 216, Indraprastha Apartments, Sector 12, Pocket 3, Dwarka, New Delhi, India.

E-mail address: mkanwat@gmail.com (M. Kanwat).

Fig. 1. Radiograph of the left hip showing a large mass at the inferior surface of the left femoral neck.

Fig. 2. CT scan showing a lobulated mass with extensive flocculent calcification with <1 cm space between the mass and lesser trochanter.

showed cortical and trabecular bone fragments covered by mature cartilaginous fragments, implying a diagnosis of osteochondroma with origin from soft tissue. [Fig. 4].

3. Discussion

Osteochondroma or exostosis is the most common primary bone tumor. These arise from the metaphysis of long bones most commonly from femur, tibia and humerus [1]. Extraskelatal osteochondromas are rare lesions with multiple postulated theories of origin. These can be broadly classified as paraarticular or those occurring away from the joint. It has been said that the connective tissue of the joint capsule or around it undergoes metaplasia to form these lesions [8]. Synovium has also been postulated as a possible origin [2]. Fracture of the osteochondroma are reported in the literature, but they were all acutely symptomatic and operated upon. It is very much possible, that, if the fracture is not operated, it can present like our case. This is a possibility but cannot be concluded. Most of these arise from synovium, with majority located in hands followed by feet [2].

Fig. 3. MRI showing a lobulated periarticular, intramuscular, non enhancing lesion isointense to the bone without any continuity to the adjacent femur.

Fig. 4. Mature bone fragments covered by mature cartilaginous fragments, implying a diagnosis of osteochondroma [H&E, 200X].

Paraarticular osteochondromas arise most commonly from the knee followed by ankle and feet [3,4]. Extraskelatal osteochondromas arising from other soft tissues have also been reported but are rarer. These have been reported in buttock, nape of neck and elbow [5–7,9]. Previously only one case has been reported in thigh, where it was located in the intermuscular plane between tensor fascia lata and glutei muscles [7]. Our case is second such report. In our case it was also situated in the intramuscular plane. The diagnosis was finally clinched on histopathology after excision.

The commonest differential diagnoses to be considered are myositis ossificans, tumoral calcinosis, synovial chondromatosis and chondrosarcoma. [Table 1] The patient responded to excision and was relieved of all symptoms.

Table 1
Differential diagnoses of Extraskelatal osteochondroma.

	Number Of Lesions	Location	Radiology	Histology
Myositis Ossificans	Solitary [5]	Away from joint [5]	Inhomogenous, egg shell calcification [5]	Zonal phenomenon with peripheral ossification [5]
Tumoral Calcinosis	Solitary/ Multiple	Away from joint	Devoid of bone, Layering when viewed by horizontal beam [5]	Calcification without bone formation
Synovial Chondromatosis	Multiple [5]	Involves joint/near the joint [5]	Multiple osteocartilagenous nodules	Hyaline cartilage bodies +/- calcification
Periosteal Chondrosarcoma [10]	Solitary	On the surface of bone, away from joint	Juxtacortical mass with irregular margins and cortical scalloping [10]	Increased cellularity and atypia of chondrocytes [10]
Calcified Synovial Sarcoma [5]	Solitary	Away from joint	Areas of peripheral calcification in a soft tissue mass [5]	Biphasic tumor composed of epithelial and spindle shaped cells along with calcification [10]
Extraskelatal Osteochondroma	Solitary [5]	Away from joint [5]	Solitary calcified lesion without periosteal reaction	Cartilaginous cap surrounding bone

4. Conclusion

Osteochondromas can at times be discontinuous from underlying bone with misleading radiology as was seen in our case. These should come in differential diagnosis whenever an ossified mass is seen in soft tissue. A knowledge about this rare lesion and its rare complications can help in accurate diagnosis and management. Can there be a possibility of such a lesion due to non-union of the fracture in the stalk of the osteochondroma? what is the natural progression of such a lesion? these questions need to be answered further.

Conflict of interest

None.

Acknowledgement

None.

References

- [1]. Peterson HA. Multiple hereditary osteochondromata. *Clin. Orthop. Relat. Res.* 1989;329:222–230.
- [2]. Gayle EL, Morrison WB, Carrino JA, Parsons TW, Liang CY, Stevenson A. Extraskelatal osteochondroma of the foot. *Skeletal Radiol.* 1999;28(10 October):594–598.
- [3]. Reith JD, Bauer TW, Joyce MJ. Paraarticular osteochondroma of the knee: report of 2 cases and review of literature. *Clin. Orthop.* 1997;334:225–232.
- [4]. Sakai H, Tamai K, Iwamoto A, Saotome K. Paraarticular chondroma and osteochondroma of the infrapatellar fat pad: a report of three cases. *Int. Orthop.* 1999;23:114–117.
- [5]. Lim SC, Kim YS, Moon YR. Extraskelatal osteochondroma of the buttock. *J. Korean Med. Sci.* 2003;18:127–130.
- [6]. Sood ID, Mudbhalkar NS. Extraskelatal osteochondroma: a case report. *Indian J. Pathol. Bacteriol.* 1967;10:72–76.
- [7]. Gulati Y, Maheshwari A, Sharma V, Mattoo R, Arora D, Gupta N. Extraskelatal osteochondroma of the thigh: a case report. *Acta Orthop. Belg.* 2005;71:115–118.
- [8]. Hagan PF, Schoenecker PL. Para-articular osteochondroma. *Am. J. Orthop.* 1995;24:65–67.
- [9]. Singh R, Sharma AK, Magu NK, Kaur KP, Sen R, Magu S. Extraskelatal osteochondroma in the nape of the neck: a case report. *J. Orthop. Surg.* 2006;14 (August 2):192–195 (Hong Kong).
- [10]. Ratcliff JR, Naqvi A, Roza G, Strauss JA, Damron TA. Soft tissue osteochondroma: case report and immunohistochemistry for parathyroid hormone-related protein. *Ann Diagn Pathol.* 2006;10:222–229.