Published in final edited form as:

Methods Enzymol. 2015 ; 562: 187–204. doi:10.1016/bs.mie.2015.04.004.

Elucidating complicated assembling systems in biology using size-and-shape analysis of sedimentation velocity data

Catherine T. Chaton^{*} and Andrew B. Herr[†]

^{*}Graduate Program in Molecular Genetics, Biochemistry & Microbiology, University of Cincinnati College of Medicine

[†]Division of Immunobiology and Center for Systems Immunology, Division of Infectious Diseases, Cincinnati Children's Hospital Medical Center, 3333 Burnet Avenue, MLC 7038, Cincinnati, OH 45229, USA

Abstract

Sedimentation velocity analytical ultracentrifugation (SV-AUC) has seen a resurgence in popularity as a technique for characterizing macromolecules and complexes in solution. SV-AUC is a particularly powerful tool for studying protein conformation, complex stoichiometry, and interacting systems in general. Deconvoluting velocity data to determine a sedimentation coefficient distribution $c(s)$ allows for the study of either individual proteins or multi-component mixtures. The standard $c(s)$ approach estimates molar masses of the sedimenting species based on determination of the frictional ratio (f/f_0) from boundary shapes. The frictional ratio in this case is a weight-averaged parameter, which can lead to distortion of mass estimates and loss of information when attempting to analyze mixtures of macromolecules with different shapes. A two-dimensional extension of the $c(s)$ analysis approach provides size-and-shape distributions that describe the data in terms of a sedimentation coefficient and frictional ratio grid. This allows for better resolution of species with very distinct shapes that may co-sediment and provides better molar mass determinations for multi-component mixtures. An example case is illustrated using globular and non-globular proteins of different masses with nearly identical sedimentation coefficients that could only be resolved using the size-and-shape distribution. Other applications of this analytical approach to complex biological systems are presented, focusing on proteins involved in the innate immune response to cytosolic microbial DNA.

Keywords

Size-and-shape analysis; sedimentation velocity; analytical ultracentrifugation; sedimentation coefficient; frictional ratio

Techniques for Studying Macromolecular Mixtures

Characterizing multi-component macromolecular systems in solution is important in many areas of biology and biotechnology. Many biological systems involve complicated

macromolecular assembly or aggregation phenomena; two examples from the field of innate immunity will be presented in this chapter. In biotechnology, quality control studies on biopharmaceutical products often need to resolve the bioactive species from potential contaminants and aggregated species. However, distinguishing the different components in a multi-component mixture, particularly those of widely varying shapes and sizes, can present significant challenges. Some techniques such as mass spectrometry that might be used are not solution-based approaches and require assumptions when extrapolating back to the aqueous sample. Other techniques such as native PAGE restrict analysis to macromolecular species that are small enough to enter the gel and require specialized running buffers.

Several solution-based techniques offer some advantages and the flexibility to analyze the macromolecule(s) of interest in defined buffers, which is often important both for biological function and for testing of biopharmaceuticals in their approved formulation state. Dynamic-light scattering (DLS) is one such technique for studying macromolecular molecules in solution. By measuring time-dependent fluctuations in light scattering intensity, it is possible to determine a diffusion coefficient and hydrodynamic radius (Cantor & Schimmel, 1980). For globular proteins, DLS can provide an estimate of the molecular weight, although without a direct determination of the diffusion or frictional coefficient, mass determinations are inaccurate for non-globular macromolecules. The DLS method has high sensitivity and requires very little sample volume. However, since the raw intensity of scattering is proportional to the square of the protein's molecular weight (i.e., for Rayleigh scattering, in which the particle size is small compared to the wavelength), large complexes and aggregates are over-weighted and contribute disproportionately to the overall signal. This increases the uncertainty when measuring relative concentrations of different species within a population if large complexes or aggregates are present. Another popular technique for analyzing macromolecular mixtures, size-exclusion (gel filtration) chromatography (SEC), can distinguish multiple species in solution. In general, SEC suffers from the same limitation as DLS when estimating the molecular weight of non-globular proteins. Combining data from SEC with experimentally-determined diffusion coefficients can provide accurate molecular weights for individual macromolecules (Ackers, 1967a; Ackers, 1967b), but this is not technically feasible for mixtures of macromolecular species with distinct diffusion and frictional coefficients.

Sedimentation velocity (SV) analytical ultracentrifugation (AUC) is a powerful technique for the study of macromolecules and nanoparticles in solution. This approach involves spinning a macromolecular sample in an ultracentrifuge and tracking the sedimentation rate of the macromolecule(s) through solution in real time, providing information on the size and shape of the sedimenting species. The macromolecular migration is independent of matrix interactions (unlike native PAGE or SEC) and it takes place at constant temperature under user-defined solution conditions. Unlike DLS, the signal observed in AUC is proportional to sample concentration, allowing accurate determination of the relative proportions of different species. Over the years, many different methods have been developed to analyze SV data, including van Holde-Weischet analysis (van Holde & Weischet, 1978); direct fitting to forms of the Lamm equation (Philo, 1997; Behlke & Ristau, 1997; Demeler & Saber, 1998); time-derivative approaches (Stafford, 1992; Philo, 2000; Schuck & Rossmanith, 2000; Stafford & Sherwood, 2004; Philo, 2006); and more recently, the c(s)

analysis approach that yields a sedimentation coefficient distribution (Schuck, MacPhee, & Howlett, 1998; Schuck, 2000). There are several good review articles that describe and compare these different analysis methods (Correia, 1998; Cole & Hansen, 1999; Howlett, Minton, & Rivas, 2006; Stafford, 2009; Zhao, Brautigam, Ghirlando, & Schuck, 2013). The $c(s)$ approach determines a sedimentation coefficient distribution for the sedimenting species and provides both hydrodynamic parameters and—particularly for single-component systems—quite accurate molecular weight determinations. However, there are some particular challenges when analyzing multi-component mixtures or complicated assembling systems. The Sedfit program developed by Schuck and colleagues now includes a two-dimensional extension of the $c(s)$ approach, allowing the study of complex mixtures of macromolecules by determining a size-and-shape distribution (Brown & Schuck, 2006). Other groups have also introduced similar two-dimensional approaches for analyzing velocity data (Brookes, Cao, & Demeler, 2010; Carney et al., 2011). In this chapter, we will briefly introduce the sedimentation coefficient distribution $c(s)$ and the size-and-shape distribution $c(s, f/f_0)$ and provide examples of how the new analytical approach can provide insight into multi-component mixtures or complex assembling systems in biology.

An Overview of the Standard Sedimentation Coefficient Distribution Approach

A typical sedimentation velocity experiment measures the concentration gradients formed by a sedimenting macromolecule driven by centrifugal force as it moves through a column of aqueous solvent. As the protein sediments, the boundary between protein and solvent moves from the meniscus (i.e., the inner boundary of the sample column) toward the outer wall of the sample cell. The rate of sedimentation and shape of the boundary give information about the macromolecule's size and shape via hydrodynamic properties such as the sedimentation, frictional and diffusion coefficients (Cantor & Schimmel, 1980; Laue & Stafford, 1999). These hydrodynamic properties are related by the Svedberg equation, shown in several equivalent forms:

$$\frac{s}{D} = \frac{\left(\frac{M_b}{f}\right)}{\left(\frac{RT}{f}\right)} = \frac{M_b}{RT} = \frac{M(1 - \bar{v}\rho)}{RT}$$

where s is the sedimentation coefficient, D is the diffusion coefficient, f is the frictional coefficient, M is the molar mass, M_b is the buoyant mass, and $\bar{v}$ is the partial specific volume of the macromolecule; ρ is the solvent density; R is the gas constant, and T is the absolute temperature. One can see that the sedimentation coefficient is proportional to the buoyant mass but inversely proportional to the frictional coefficient, and that knowledge of both s and f or s and D allow determination of the molar mass of the sedimenting macromolecule, after accounting for the partial specific volume of the macromolecule and the buffer density.

Sedimentation of a particle in a sector-shaped column within a centrifugal field is described by the Lamm equation (Lamm, 1929). Written as a partial differential, it is:

$$\frac{dc}{dt} = \frac{1}{r} \frac{d}{dr} \left[rD \frac{dc}{dr} - s\omega^2 r^2 c \right]$$

The concentration distribution $c(r,t)$ of the macromolecule as a function of radial position r and time t depends on the diffusion coefficient D , the strength of the centrifugal field $\omega^2 r$ and the sedimentation coefficient of the species. Measured in Svedberg units ($1 \text{ S} = 10^{-13} \text{ sec}$), the sedimentation coefficient is a proportionality constant that relates the velocity of the sedimenting particle to the applied force resulting from the centrifugal field. The sedimentation coefficient is dependent on both the size and shape of the particle.

In an idealized case where diffusion is neglected, the Lamm equation can be solved exactly. In this theoretical case, each sedimentation profile would have a sharp boundary created between the bulk macromolecules and the solution as macromolecules are driven toward the outer wall of the sample cell. In a more realistic experiment, diffusion acts to broaden the sedimentation boundary over time. This diffusional spread in the boundary is highly dependent on the frictional coefficient of the macromolecule; in other words, non-globular particles experience greater drag, which retards their rate of diffusion (and their rate of sedimentation, albeit to a lesser extent). Experimental factors such as radial dilution, non-uniformity in cell loading concentration, the formation of density gradients, and the accumulation of sedimented protein at the outside of the sample cell are several reasons that exact analytical solutions of the Lamm equation are not practical. Approximate analytical solutions do exist but are sensitive to experimental conditions (Philo, 1997; Behlke & Ristau, 1997). Another potential approach is to use numerical simulation to reproduce the natural progression of the sedimentation boundary (Demeler & Saber, 1998; Schuck et al., 1998; Stafford & Sherwood, 2004). Schuck and colleagues have developed the $c(s)$ analysis approach that directly models the boundary as a distribution of non-interacting particles and describes diffusion of the ensemble of sedimentation species (Schuck, MacPhee, & Howlett, 1998; Schuck, 2000; Dam & Schuck, 2004). They have shown that despite being based on equations describing the sedimentation behavior of non-interacting particles, the $c(s)$ analysis can also be applied to systems of interacting macromolecules that produce asymmetric boundaries according to Gilbert-Jenkins theory (Gilbert & Jenkins, 1959; Dam, Velikovsky, Mariuzza, Urbanke, & Schuck, 2005; Schuck, 2010).

Besides the sedimentation coefficient, which is the experimental parameter directly determined in an SV experiment, the major hydrodynamic property fitted for during $c(s)$ analysis is the frictional ratio f/f_0 . It is a measure of the frictional coefficient of the macromolecule compared to that of a compact, anhydrous, ideal sphere. Thus, the lower bound of this ratio is 1.0, corresponding to an ideal sphere; the frictional ratio increases for sedimenting macromolecules that are increasingly extended or non-globular. Because the diffusion rate is highly sensitive to the frictional coefficient, the shape of sedimentation boundaries can be directly fitted to determine the weight-averaged frictional ratio in the $c(s)$ analysis. Thus, for a single-component sample, it is possible to experimentally determine the values of the sedimentation coefficient s and the frictional coefficient f , allowing determination of the mass of the sedimenting species according to the Svedberg equation.

Since the frictional ratio determined in this way is a weight-averaged value, accurate mass determination is generally only possible for a single-component system unless additional species happen to have the same frictional ratio. As differences in the frictional ratio between components increase, the distortion in mass estimates will also increase. Despite this fact, the standard $c(s)$ approach is an excellent method for characterizing the number of species present and their relative concentrations in discrete macromolecular mixtures, even if accurate mass determinations are not feasible in all situations.

We will present an experimental situation that illustrates the limits of the standard $c(s)$ approach to distinguish heterogeneous species with similar sedimentation rates. We conducted sedimentation velocity experiments on two non-interacting proteins with very different shapes, each of which sediments near 2.0 S (Figures 1A–C). For a non-globular, highly-elongated protein we chose a construct from the Accumulation-Associated Protein (Aap), a cell-surface protein from *Staphylococcus epidermidis*. We have previously demonstrated that Aap is necessary for biofilm formation, as it mediates intercellular adhesion through zinc-induced self-assembly (Conrady, Brescia, Horii, Weiss, Hassett, & Herr, 2008). The C-terminal portion of Aap contains 5–17 tandem copies of the ‘B-repeat,’ a 128-amino acid sequence comprising two subdomains. The crystal structure of a minimal B-repeat construct, called Brpt1.5, revealed a highly elongated fold without a canonical hydrophobic core, with overall dimensions of 180 Å long by 20 Å wide (Figure 1D) (Conrady, Wilson, & Herr, 2013). Hen egg-white lysozyme (Figure 1E) was chosen as an example of a globular protein (Vaney, Maignan, Riès-Kautt, & Ducriux, 1996). In order to compare the sedimentation behavior of lysozyme and the Aap construct, each protein was sedimented individually, followed by a mixture of the two. The results will illustrate the challenges in resolving two proteins with different shapes but similar sedimentation coefficients.

When sedimented individually, the $c(s)$ distributions of lysozyme and the Aap construct are distinct but overlap (Figure 1F). The Aap construct sedimented with an $s_{20,w}$ (i.e., sedimentation coefficient corrected to standard conditions of 20° C in water) of 1.94 S, a frictional ratio of 2.41, and a resulting apparent molecular weight of 47.1 kDa (Figure 1A). The frictional ratio of the Aap construct is high, indicating a very elongated or non-globular protein, which is consistent with the crystal structure of the Brpt1.5 Aap construct (Figure 1D). Lysozyme sedimented with an $s_{20,w}$ of 1.77 S, a much lower frictional ratio of 1.21, and a lower apparent molecular weight of 14.5 kDa (Figure 1B). This frictional ratio is in the range typically observed for globular proteins (1.2 – 1.4). Since the buoyant mass is dependent on both s and f ($s = M_b/f$), it is clear that accurate determination of f/f_0 is essential to calculate the mass, since both proteins have nearly identical sedimentation coefficients.

The Aap construct and lysozyme were then mixed together in a 1:9 molar ratio, which corresponded to a slightly higher absorbance of lysozyme relative to the Aap construct at 235 nm (Figure 1C). Analysis of the mixture yielded a single peak in the $c(s)$ distribution at $s_{20,w} = 1.85$ S, with a weight-averaged frictional ratio of 1.62. Despite the acceptable residuals on all three experiments and RMSDs on all fits of less than 0.005, there is little evidence in the $c(s)$ distribution that would indicate the presence of two distinct species, despite the large difference in mass between the two constructs (Figure 1F). In order to

resolve these two species with very different shapes, a new analytical approach is required: determining the size-and-shape distribution $c(s, f/f_0)$ from the experimental velocity data.

Two Dimensional Size-and-Shape Analysis

As mentioned above, the standard $c(s)$ analysis approach uses the assumption that all sedimenting species can be described by a single frictional ratio, which is determined experimentally from the shape of the boundary. In short, non-globular or elongated particles with higher frictional coefficients experience greater drag, which reduces the rate of diffusion and therefore reduces boundary spreading. The use of a weight-averaged frictional ratio to estimate molar mass assumes that M scales with s according to a 2/3 power relationship, based on the Svedberg equation and the Stokes-Einstein relationship (Dam & Schuck, 2004). This assumption of increasing M with increasing s is appropriate when dealing with a single species, or with several species having similar frictional ratios. However, the assumption of a single frictional ratio obviously does not hold in the case of multiple heterogeneous species. Cases such as these can instead be analyzed using a relatively new approach that calculates a two-dimensional size-and-shape distribution describing the sedimentation behavior of a sample in terms of both sedimentation coefficient and frictional ratio distributions (Brown & Schuck, 2006). This $c(s, f/f_0)$ distribution is calculated on a grid of s and f/f_0 values, determining the best-fit value of f/f_0 for each s -value across the grid. The data are fitted to a modified form of the Lamm equation using finite element solutions and combined with Tikhonov-Philips or maximum entropy regularization to produce the most parsimonious $c(s, f/f_0)$ distribution that is consistent with the input data (Brown & Schuck, 2006). As with the $c(s)$ analysis, both time-invariant and radial-invariant noise is accounted for, improving the signal-to-noise ratio in the data. In some cases, f/f_0 may not be the hydrodynamic parameter of primary interest. This same computational approach allows for transformation into other two-dimensional distributions such as mass-Stokes radius distributions $c(M, R_s)$ or sedimentation coefficient-molar mass distributions $c(s, M)$. Furthermore, integration along the f/f_0 dimension can return the results to a one-dimensional $c(s)$ -equivalent distribution without the limitation of using a single weight-averaged frictional ratio. Referred to as the general $c(s, *)$ distribution, this transformation can serve as the final step in an analysis if diffusion information is not required or if the frictional ratio dimension is not well-determined.

The ability to resolve distinct frictional ratios for multiple species and therefore to more accurately assign molecular masses of multiple components adds a significant amount of valuable information. Importantly, calculation of these size-and-shape distributions is only marginally more computationally intensive than the standard $c(s)$ analysis, taking only minutes on a standard laptop. Experiments involving heterogeneous mixtures such as proteins of very different shapes, combinations of folded and intrinsically disordered proteins, or multiple assembling species in equilibrium are good candidates for the size-and-shape distribution approach. Returning to our test case data, we re-analyzed the same scans of the Aap protein and lysozyme mixture to determine the size-and-shape distribution $c(s, f/f_0)$. The $c(s, *)$ distribution (Figure 2A, black line) shows two peaks. Superimposed on this $c(s, *)$ distribution is a heat map illustrating a bird's-eye view of the resolved peaks in the size-and-shape distribution, plotted in terms of $s_{20,w}$ (the corrected sedimentation

coefficient) along the x-axis and f/f_0 along the y-axis. The same size-and-shape distribution is illustrated as a 3-D plot in Figure 2B, showing the clear separation of the two peaks along the f/f_0 axis, even though their sedimentation coefficient distributions almost overlap. Comparing the final $c(s,*)$ distribution of the mixture to the standard $c(s)$ distribution reveals additional information about the system and can resolve the second peak corresponding to the Aap construct (Figure 2C).

Case Study: Cyclic GMP-AMP Synthase Oligomerization

We present two case studies of proteins involved in cytosolic innate immune responses that undergo multiple assembly events upon activation. In both cases, use of the $c(s,f/f_0)$ analysis approach was essential to understand the system under study. The cGAS-STING pathway connects a cytosolic microbial DNA sensor to the production of type I interferons, forming a critical link in innate antimicrobial immunity (Shu, Li, & Li, 2014). The cyclic GMP-AMP synthase (cGAS), a 60 kDa dsDNA sensor, synthesizes cyclic 2',5' cGAMP from ATP and GTP upon activation by microbial dsDNA. cGAMP acts as a secondary messenger, facilitating the dimerization of STING (Ishikawa & Barber, 2008). In addition, STING can be activated directly by the cyclic dinucleotides c-di-GMP or c-di-AMP, both of which are conserved bacterial signaling molecules (Burdette, Monroe, Sotelo-Troha, Iwig, Eckert, Hyodo, et al., 2011). Upon activation by these second messengers, STING recruits tank-binding kinase (TBK)-1, an ER membrane serine/threonine protein kinase. TBK-1 can phosphorylate interferon regulatory factor (IRF)-3, leading to its assembly and translocation into the nucleus, where it stimulates transcription of interferon- β (Figure 3).

Several crystal structures of cGAS bound to dsDNA were published in 2013; the initial reports described an apparent 1:1 complex of cGAS:dsDNA (Civril, Deimling, de Oliveira Mann, Ablasser, Moldt, Witte et al., 2013; Gao, Ascano, Wu, Barchet, Gaffney, Zillinger, et al., 2013; Li, Shu, Yi, Chaton, Shelton, Diao, et al., 2013; Kranzusch, Lee, Berger, & Doudna, 2013; Kato, Ishii, Goto, Ishitani, Tokunaga, & Nureki, 2013; Zhang, Wu, Du, Xu, Sun, Chen, et al., 2014). The monomer's nucleotidyltransferase (NTase) fold (N-lobe) and five-helix bundle (C-lobe) were found to bind dsDNA down the length of the N-lobe. It was hypothesized that this allowed for a change in conformation of the active site buried in the cleft between both lobes. Our collaborators solved the cGAS:dsDNA structure as well and noticed the formation of a putative 2:2 dimer in the crystal (Li, Shu, Yi, Chaton, Shelton, Diao, et al., 2013). The N-lobes of each protomer face one another, creating a $\sim 1,330$ Å² patch of buried surface area which is mediated by 15 hydrogen bonds. Each protomer contains two distinct dsDNA-binding sites, both the originally-described dsDNA binding site (site A) next to the dimer interface, and a secondary site B on the opposite side of the dimer interface. Site B contributes 34–40% of the total buried surface area between cGAS and complexed dsDNA. Comparison of the free and bound cGAS structures revealed that the B-factors of the N-lobe dropped significantly when dsDNA was bound, suggesting that reducing flexibility of the NTase fold is necessary for formation of a functional active site.

In order to verify that the putative 2:2 cGAS:dsDNA dimer observed in the crystal could form in solution, we analyzed the complex by SV-AUC. cGAS alone sedimented as a monomer, but in the presence of dsDNA cGAS formed several higher-order oligomeric

species, as revealed by $c(s)$ analysis (Figure 4A). The cGAS dimer observed in the crystal revealed an important H-bonding interaction across the dimer interface mediated by K382; mutation of this residue to alanine prevented cGAS oligomerization, although the monomeric K382A mutant still was capable of binding dsDNA (Figure 4A). Importantly, the K382A mutant revealed a nearly complete loss of cGAS enzymatic activity, indicating that formation of the dsDNA-induced cGAS dimer is essential for catalysis. Mutation of residues at each DNA-binding site (K372E and R158E for site A and R342E and K335E for site B) also prevented cGAS oligomerization, showing the importance of DNA binding for dimer formation (Figure 4B).

The $c(s)$ distribution of wild-type cGAS bound to dsDNA indicated the presence of multiple species, not simply a dimeric 2:2 complex (Figure 4A). In order to delineate the species present in this sample, we determined the $c(s, f/f_0)$ distribution of cGAS in the presence of dsDNA (Figure 4C). While ~36% of the cGAS population formed a compact 2:2 complex, many other oligomeric species were observed. A more elongated complex with apparent 6:6 stoichiometry (25% of protein signal) was prominent along with several highly elongated, putative fiber-like species. Furthermore, the concentration regime over which cGAS assembly occurred was consistent with its affinity for dsDNA. cGAS samples at ~150 μM formed higher-order species, but dilution of the sample to ~30 μM led to dissociation of the oligomeric species. This agrees with the low affinity (K_D ~20 μM) observed for cGAS binding to dsDNA (Li et al., 2013).

Case Study: Interferon Regulatory Factor-3 Phosphorylation by Tank-binding Kinase-1

As mentioned above, cGAS activated by dsDNA-mediated oligomerization catalyzes the synthesis of the second messenger cGAMP, which binds to STING and triggers recruitment of the kinase TBK-1 (Figure 3). Our collaborators solved the crystal structure of TBK1 and analyzed its phosphorylation of IRF-3; we again used the $c(s, f/f_0)$ analysis of SV data to fully understand the effects of phosphorylation on IRF-3 assembly state (Shu, Sankaran, Chaton, Herr, Mishra, Peng, and Li, 2013). Once phosphorylated by TBK1 or I κ B kinase- ϵ (IKK ϵ), phospho-IRFs translocate to the nucleus and bind to CREB-binding protein (CBP) in order to initiate transcription of type I interferons (Fitzgerald, McWhirter, Faia, Rowe, Latz, Golenbock, et al., 2003; Lin, Heylbroeck, Pitha, & Hiscott, 1998). SEC analysis of *in vitro* phosphorylated human IRF-3 indicated that it migrated as a broad peak with a much higher apparent mass (~300 kDa) than the native monomeric form (~95 kDa). Sedimentation velocity analyses of native hIRF-3 revealed a single monomeric species with a frictional ratio of 1.5 (Figure 5A and C). Upon phosphorylation, the frictional ratio of the hIRF-3 monomer was lowered to near 1.0, indicating a conformational change to a compact structure (Figure 5B and D). In addition to conformational collapse of the primary species, several discrete higher order oligomers were observed. These included a potential tetramer with a frictional ratio above 3.0, as well as a series of highly-elongated fiber-like assemblies with apparent masses between a 13-mer and 32-mer. Further biochemical experiments showed that mutation of S86 to alanine prevented phosphorylation-dependent assembly of hIRF-3, and that an oligomeric form of phosphorylated hIRF-3 specifically bound to CBP

(most likely the dimeric form) (Shu et al., 2013). Similar results were found when IRF-3 was co-expressed with the kinase IKK ϵ , with IRF-3 forming monomers, dimers and higher-order oligomers (Takahasi, Horiuchi, Fujii, Nakamura, Noda, Yoneyama, et al., 2010). Though not yet fully understood, it is believed that the oligomeric forms of IRF-3 could play important roles in transcription activation due to higher avidity for DNA.

Conclusions

SV analysis of macromolecules in solution is a powerful technique that can provide a wealth of information about particle size, shape, assembly state, and can elucidate complex stoichiometry or conformational changes in solution. Deconvoluting the data into a $c(s)$ continuous sedimentation coefficient distribution is useful for determining the number of species in a sample and characterizing their relative size, but mass estimates are not accurate for mixtures of particles with varying frictional ratios. In contrast, the size-and-shape $c(s, f/f_0)$ distribution allows the determination of f/f_0 for each s -value on a two-dimensional grid. This improves the mass determination for multiple species and allows the resolution of multiple species of distinct shape, even those that sediment with similar rates.

There are certain limitations to the size-and-shape distribution approach that should be kept in mind. The $c(s)$ approach in general is based on equations for non-interacting particles, and some have raised concerns about the general applicability of this analysis method to interacting systems (Stafford, 2009); however, Schuck and colleagues have shown that $c(s)$ was able to accurately model sedimentation coefficients of interacting species under conditions of rapid equilibrium in a manner consistent with Gilbert-Jenkins theory (Dam et al., 2005; Schuck, 2010). Brown and Schuck do point out that the $c(s, f/f_0)$ distribution was not as successful as $c(s)$ at recreating the asymmetric boundaries from Gilbert-Jenkins theory for interacting systems, probably due to the slightly lower resolution of the $c(s, f/f_0)$ distribution (Brown & Schuck, 2006). Furthermore, the molar mass estimates for low-abundance species in the $c(s, f/f_0)$ distribution were not as accurate, and they suggest that $c(s)$ is better suited to the analysis of trace components (Brown & Schuck, 2006). Thus, one should use caution in interpreting mass estimates from size-and-shape distributions for interacting systems or for low-abundance species. Regardless, the use of the size-and-shape distribution will be very helpful in providing a greater level of detail than is otherwise possible for complicated multi-component solutions, particularly for species of widely varying shape. We believe that this type of analysis will find an increasing number of applications when studying biological systems that involve complicated assembly mechanisms or multi-component mixtures of macromolecules.

Acknowledgments

We thank Peter Schuck and Jim Cole for comments on the manuscript. This work was supported by funding from the National Institute of General Medical Sciences, R01 GM094363 (to ABH). Figure 4 was reprinted from *Immunity* **39**:1019–1031 (2013); X Li, C Shu, G Yi, CT Chaton, CL Shelton, J Diao, X Zuo, CC Kao, AB Herr, and P Li, “Cyclic GMP-AMP synthase is activated by double-stranded DNA-induced oligomerization” with permission from Elsevier. Figure 5 was reprinted from *Structure* **21**:1137–1148 (2013); C Shu, B Sankaran, CT Chaton, AB Herr, A Mishra, J Peng, and P Li, “Structural insights into the functions of TBK1 in innate antimicrobial immunity” with permission from Elsevier.

References

- Ackers GK. Molecular Sieve Studies of Interacting Protein Systems: 1. Equations for Transport of Associating Systems. *The Journal of Biological Chemistry*. 1967; 242:3026–34. [PubMed: 6027787]
- Ackers GK. A New Calibration Procedure for Gel Filtration Columns. *The Journal of Biological Chemistry*. 1967; 242:3237–38.
- Arthur KK, Kendrick BS, Gabrielson JP. Guidance to achieve accurate aggregate quantitation in biopharmaceuticals by SV-AUC. *Methods in Enzymology*. 2015; 562:477–500. [PubMed: 26412664]
- Behlke J, Ristau O. Molecular mass determination by sedimentation velocity experiments and direct fitting of the concentration profiles. *Biophysical Journal*. 1997; 72:428–434. [PubMed: 8994629]
- Brookes E, Cao W, Demeler B. A two-dimensional spectrum analysis for sedimentation velocity experiments of mixtures with heterogeneity in molecular weight and shape. *European Biophysics Journal*. 2010; 39:405–414. [PubMed: 19247646]
- Brown PH, Schuck P. Macromolecular Size-and-Shape Distributions by Sedimentation Velocity Analytical Ultracentrifugation. *Biophysical Journal*. 2006; 90:4651–61. [PubMed: 16565040]
- Burdette DL, Monroe K, Sotelo-Troha K, Iwig J, Eckert B, Hyodo M, Hayakawa Y, Vance R. STING is a direct innate immune sensor of cyclic di-GMP. *Nature*. 2011; 478:515–8. [PubMed: 21947006]
- Cantor, C., Schimmel, P. *Biophysical Chemistry Part II: Techniques for the study of biological structure and function*. New York, NY: W.H. Freeman and Company; 1980.
- Carney RP, Kim JY, Qian H, Jin R, Mehenni H, Stellacci F, et al. Determination of nanoparticle size distribution together with density or molecular weight by 2D analytical ultracentrifugation. *Nature Communications*. 2011; 2:335.
- Civril F, Deimling T, de Oliveira Mann C, Ablasser A, Moldt M, Witte G, et al. Structural mechanism of cytosolic DNA sensing by cGAS. *Nature*. 2013; 498:332–337. [PubMed: 23722159]
- Cole J, Hansen J. Analytical ultracentrifugation as a contemporary biomolecular research tool. *Journal of Biomolecular Technology*. 1999; 10:163–76.
- Conrady DG, Brescia CC, Horii K, Weiss AA, Hassett DJ, Herr AB. A zinc-dependent adhesion module is responsible for intercellular adhesion in staphylococcal biofilms. *Proc Natl Acad Sci U S A*. 2008; 105(49):19456–61. [PubMed: 19047636]
- Conrady DG, Wilson JJ, Herr AB. Structural basis for Zn²⁺-dependent intercellular adhesion in staphylococcal biofilms. *Proc Natl Acad Sci U S A*. 2013; 110(3):E202–11. [PubMed: 23277549]
- Correia JJ. Sedimentation velocity data analysis methods: what, when and why? *CHEMTRACTS - Biochemistry and Molecular Biology*. 1998; 11:944–8.
- Dam J, Schuck P. Calculating sedimentation coefficient distributions by direct modeling of sedimentation velocity concentration profiles. *Methods in Enzymology*. 2004; 384:185–212. [PubMed: 15081688]
- Dam J, Velikovskiy CA, Mariuzza RA, Urbanke C, Schuck P. Sedimentation velocity analysis of heterogeneous protein-protein interactions: Lamm equation modeling and sedimentation coefficient distributions c(s). *Biophysical Journal*. 2005; 89:619–34. [PubMed: 15863475]
- Demeler B, Saber H. Determination of molecular parameters by fitting sedimentation data to finite-element solutions of the Lamm equation. *Biophysical Journal*. 1998; 74(1):444–54. [PubMed: 9449345]
- Fitzgerald KA, McWhirter SM, Faia KL, Rowe DC, Latz E, Golenbock DT, Coyle AJ, Liao SM, Maniatis T. IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nature Immunology*. 2003; 4:491–496. [PubMed: 12692549]
- Gao P, Ascano M, Zillinger T, Wang W, Dai P, Serganov AA, Gaffney BL, Shuman S, Jones RA, Deng L, Hartmann G, Barchet W, Tuschl T, Patel DJ. Structure-function analysis of STING activation by c[G(2',5')pA(3',5')p] and targeting by antiviral DMXAA. *Cell*. 2013; 154:748–62. [PubMed: 23910378]
- Gilbert GA, Jenkins RC. Sedimentation and Electrophoresis of Interacting Substances. II. Asymptotic Boundary Shape for Two Substances Interacting Reversibly. *Proceedings of the Royal Society*. 1959; 253:420–37.

- Howlett GJ, Minton AP, Rivas G. Analytical Ultracentrifugation for the Study of Protein Association and Assembly. *Current Opinions in Chemical Biology*. 2006; 10:430–6.
- Ishikawa H, Barber GN. STING is an endoplasmic reticulum adaptor that facilitates innate immune signalling. *Nature*. 2008; 455:674–678. [PubMed: 18724357]
- Kato K, Ishii R, Goto E, Ishitani R, Tokunaga F, Nureki O. Structural and functional analyses of DNA-sensing and immune activation by human cGAS. *PLoS One*. 2013; 8:e76983. [PubMed: 24116191]
- Kranzusch PJ, Lee AS, Berger JM, Doudna JA. Structure of human cGAS reveals a conserved family of second-messenger enzymes in innate immunity. *Cell Reports*. 2013; 3:1362–1368. [PubMed: 23707061]
- Lamm O. Die differentialgleichung der ultrazentrifugierung. *Arkiv för Matematik, Astronomi, och Fysik*. 1929; 21B(2):1–4.
- Laue TM, Stafford WF. Modern Applications of Analytical Ultracentrifugation. *Annual Review of Biophysics and Biomolecular Structure*. 1999; 28:75–100.
- Li X, Shu C, Yi G, Chaton CT, Shelton CL, Diao J, Zuo X, Kao CC, Herr AB, Li P. Cyclic GMP-AMP Synthase Is Activated by Double-Stranded DNA-Induced Oligomerization. *Immunity*. 2013; 39:1019–31. [PubMed: 24332030]
- Lin R, Heylbroeck C, Pitha PM, Hiscott J. Virus-dependent phosphorylation of the IRF-3 transcription factor regulates nuclear translocation, transactivation potential, and proteasome-mediated degradation. *Molecular and Cellular Biology*. 1998; 18:2986–96. [PubMed: 9566918]
- Philo JS. An improved function for fitting sedimentation velocity data for low-molecular-weight solutes. *Biophysical Journal*. 1997; 72:435–44. [PubMed: 8994630]
- Philo JS. A method for directly fitting the time derivative of sedimentation velocity data and an alternative algorithm for calculating sedimentation coefficient distribution functions. *Analytical Biochemistry*. 2000; 279:151–63. [PubMed: 10706784]
- Philo JS. Improved methods for fitting sedimentation coefficient distributions derived by time-derivative techniques. *Analytical Biochemistry*. 2006; 354:238–46. [PubMed: 16730633]
- Schuck P. Size distribution analysis of macromolecules by sedimentation velocity ultracentrifugation and Lamm equation modeling. *Biophysical Journal*. 2000; 78:1606–19. [PubMed: 10692345]
- Schuck P. Sedimentation Patterns of Rapidly Reversible Protein Interactions. *Biophysical Journal*. 2010; 98:2005–13. [PubMed: 20441765]
- Schuck P, MacPhee CE, Howlett GJ. Determination of sedimentation coefficients for small peptides. *Biophysical Journal*. 1998; 74:466–74. [PubMed: 9449347]
- Schuck P, Rossmanith P. Determination of the sedimentation coefficient distribution by least-squares boundary modeling. *Biopolymers*. 2000; 54:328–41. [PubMed: 10935973]
- Shu C, Li X, Li P. The mechanism of double-stranded DNA sensing through the cGAS-STING pathway. *Cytokine Growth Factor Review*. 2014; 25:641–648.
- Shu C, Sankaran B, Chaton CT, Herr AB, Mishra A, Peng J, Li P. Structural Insights into the Functions of TBK1 in Innate Antimicrobial Immunity. *Structure*. 2013; 21:1137–48. [PubMed: 23746807]
- Stafford WF. Boundary analysis in sedimentation transport experiments: a procedure for obtaining sedimentation coefficient distributions using the time derivative of the concentration profile. *Analytical Biochemistry*. 1992; 203:295–301. [PubMed: 1416025]
- Stafford WF, Sherwood PJ. Analysis of heterologous interacting systems by sedimentation velocity: curve fitting algorithms for estimation of sedimentation coefficients, equilibrium and kinetic constants. *Biophysical Chemistry*. 2004; 108:231–43. [PubMed: 15043932]
- Stafford, WF. Protein-protein and ligand-protein interactions studied by analytical ultracentrifugation. In: Shriver, KW., editor. *Protein Structure, Stability, and Interactions*. Vol. 490. Humana Press; Totowa NJ: 2009. p. 83-113.
- Takahasi K, Horiuchi M, Fujii K, Nakamura S, Noda NN, Yoneyama M, Fujita T, Inagaki F. Ser386 phosphorylation of transcription factor IRF-3 induces dimerization and association with CBP/p300 without overall conformational change. *Genes to Cells*. 2010; 15:901–910. [PubMed: 20604809]
- Vaney MC, Maignan S, Riès-Kautt M, Ducruix A. High-resolution structure (1.33 Å) of a HEW lysozyme tetragonal crystal grown in the APCF apparatus. Data and structural comparison with a

crystal grown under microgravity from SpaceHab-01 mission. Acta Crystallographica Section D. 1996; 52:505–517.

van Holde KE, Weischedel WO. Boundary Analysis of Sedimentation Velocity Experiments with Monodisperse and Paucidisperse Solutes. Biopolymers. 1978; 17:1387–1403.

Zhang X, Wu J, Du F, Xu H, Sun L, Chen Z, et al. The cytosolic DNA sensor cGAS forms an oligomeric complex with DNA and undergoes switch-like conformational changes in the activation loop. Cell Reports. 2014; 6:421–430. [PubMed: 24462292]

Zhao H, Brautigam CA, Ghirlardo R, Schuck P. Current Methods in Sedimentation Velocity and Sedimentation Equilibrium Analytical Ultracentrifugation. Current Protocols in Protein Science. 2013; 20(Unit 20.12) <http://dx.doi.org/10.1002/0471140864.ps2012s71>.

Figure 1. Standard $c(s)$ analysis of test proteins

SV-AUC raw data for (A) The Aap repeat protein, (B) lysozyme, and (C) the ~1:9 molar mixture of Aap:lysozyme. (D) Crystal structures of a Brpt1.5 Aap construct (Conrady et al., 2013) (PDB 4FUN) and (E) hen egg-white lysozyme (Vaney et al., 1996) (PDB 193L). (F) The calculated $c(s)$ distributions for lysozyme, the highly elongated Aap protein, and the mixture. Despite a small shift in $s_{20,w}$, multiple species are not observed in the mixture.

Figure 2. Size-and-shape $c(s, f/f_0)$ analysis of test proteins

(A) The $c(s, f/f_0)$ distribution for the mixture of Aap and lysozyme is able to clearly resolve both species. Lysozyme appears at an $s_{20,w}$ of 1.84 and accounts for 90% of the loading signal. The highly-elongated Aap protein sediments with an $s_{20,w}$ of 1.98 and accounts for 9% of the loading signal. (B) 3D surface representation of the $c(s, f/f_0)$ distribution visualized with the lattice package in R. (C) A comparison showing the original $c(s)$ distribution calculated for the mixture of Aap and lysozyme overlaid on the $c(s,*)$ distribution. The presence of the minor species, the elongated Aap protein, is only visible after decoupling the scaling relationship between s and M implicit in the standard $c(s)$ analysis.

Figure 3. Schematic of innate immune signaling pathways in the cytosol
 Microbial dsDNA is recognized by the sensor protein cGAS, which upon activation catalyzes the synthesis of the secondary messenger 2',5' cGAMP. cGAMP binds to STING, activating recruitment of the kinase TBK-1. TBK-1 then phosphorylates IRF-3, which translocates to the nucleus and triggers expression of interferon-β. This figure was adapted from (Shu et al., 2014).

Figure 4. Sedimentation analysis of cGAS interacting with dsDNA

(A) $c(s)$ distributions of WT and K382A mutants of mouse cGAS in the presence and absence of dsDNA. WT cGAS forms multiple oligomeric species in the presence of dsDNA. The K382A mutant, which disrupts H-bonds at the cGAS dimer interface, forms a 1:1 complex with the DNA but was unable to oligomerize. (B) Normalized $c(s)$ distributions of cGAS mutants altered in the DNA-binding sites, including A-site mutants (K372E and R158E) and B site mutants (R342E and K335E), each sedimented in the presence of dsDNA. (C) Size-and-shape $c(s, f/f_0)$ analysis of WT cGAS:dsDNA demonstrates that multiple oligomeric forms of cGAS occur upon dsDNA binding. Both elongated and compact species are seen at a variety of possible stoichiometries, based on mass estimates from the $c(s, f/f_0)$ distribution. This figure was reprinted from (Li et al., 2013) with permission.

Figure 5. Size-and-shape analysis of hIRF-3 phosphorylated by TBK-1

The $c(s, f/f_0)$ distribution was determined for native IRF-3 (A) and phosphorylated IRF-3 (B). A number of distinct species appear upon phosphorylation, including compact monomer and dimer forms as well as elongated species. Two-dimensional size-and-shape distributions in (C) and (D) show major peaks and the putative oligomeric states based on predicted mass from the $c(s, f/f_0)$ analysis. This figure was reprinted from (Shu et al., 2013) with permission.