Published in final edited form as:

J Proteome Res. 2016 October 07; 15(10): 3656–3665. doi:10.1021/acs.jproteome.6b00468.

The impact of amidination on peptide fragmentation and identification in shotgun proteomics

Sujun Li[†], Aditi Dabir[‡], Santosh A. Misal[‡], Haixu Tang[†], Predrag Radivojac^{*,†}, and James P. Reilly^{*,‡}

School of Informatics and Computing, Indiana University, Bloomington, and Department of Chemistry, Indiana University, Bloomington

Abstract

Peptide amidination labeling using S-methyl thioacetimidate (SMTA) is investigated in an attempt to increase the number and types of peptides that can be detected in a bottom-up proteomics experiment. This derivatization method affects the basicity of lysine residues, and is shown here to significantly impact the idiosyncracies of peptide fragmentation and peptide detectability. The unique and highly reproducible fragmentation properties of SMTA-labeled peptides, such as the strong propensity for forming b₁ fragment ions, can be further exploited to modify the scoring of peptide-spectrum pairs and improve peptide identification. To this end, we have developed a supervised post-processing algorithm to exploit these characteristics of peptides labeled by SMTA. Our experiments show that, although the overall number of identifications are similar, the SMTA modification enabled the detection of 16-26% peptides not previously observed in comparable CID/HCD tandem mass spectrometry experiments without SMTA labeling.

Keywords

Mass spectrometry; Proteomics; Chemical modification; Peptide fragmentation; Amidination; Peptide identification; Fragment ions; Logistic regression

Introduction

Over the past decade, mass spectrometry has become the dominant platform in proteome research.^{1–5} Advances in technology and algorithm development have recently enabled deep proteome coverage of several model organisms with the current draft of the human proteome reaching 92%.⁶ Shotgun methods have played an important role in this field, although technical and statistical limitations still prevent complete coverage of identified protein sequences in a single experiment. This shortcoming compromises our ability to detect many biologically important events such as alternatively spliced products⁷ or post-translational modifications.⁸ Shotgun proteomics experiments are normally carried out in the following steps:⁹ proteins are extracted from lysed cells, they are enzymatically digested, and the resulting peptides are chromatographically separated and analyzed by mass spectrometry.

*To whom correspondence should be addressed.

[†]School of Informatics and Computing, Indiana University

[‡]Department of Chemistry, Indiana University

All of these steps still need to be perfected in order to improve protein amino-acid sequence coverage. For example, numerous papers have applied new enzyme digestion strategies for proteome analysis.^{10–12} The development of new particle sizes and coatings has led to improvements in chromatographic resolution^{13,14} and modifications in liquid chromatography solvents have led to enhanced electrospray ionization of peptides.^{15,16} Furthermore, peptides have been chemically modified to enable N- and C-terminal fragments to be distinguished and to modify ion fragmentation patterns.^{17,18}

Typically, for unmodified tryptic peptides, the collision-induced dissociation (CID) spectra are dominated by b- and y-type fragment ions. The latter tend to be more abundant than the former because b ions undergo secondary fragmentation more readily than y ions, thus creating small fragments that may not be detectable with the ion trap. Multiple strategies have been developed to alter the ion fragmentation patterns to improve peptide detectability by charge-remote or charge-induced processes.^{19–21} The locations of modified groups in peptides can directly influence the locations of protons and thus influence the fragment ion patterns; in addition, altering fragmentation by changing the basicity or acidity of residues may lead to different peptide identifications by mass spectrometry.¹⁸ Beyond the well-established database searching methods, computational researchers have incorporated this type of information into database searching and post-processing of search results.^{22–25}

We have previously studied labeling techniques and provided anecdotal evidence that peptide amidination leads to changes in peptide basicity thereby affecting their ionizability²⁶ and fragmentation.^{27,28} This is because N-terminal amidination leads to the formation of cyclic intermediates that enable peptide bond cleavage, stabilizing of otherwise unobservable b₁ ions and subsequent intense y_{N-1} ions.^{26–28} We have also systematically investigated peptide fragmentation and detectability.^{29–32} To follow up on that work, the present study focuses on quantifying the impact of amidination on peptide fragmentation and detectability of a large number of peptides. By exploiting the idiosyncrasies of peptides containing this label, we have then developed a new data interpretation algorithm to post-process peptide assignments from standard database search engines that ultimately lead to an improved diversity of identified peptides and better proteome coverage.

Methods

Sample preparation

Escherichia coli K12 MG1655 cells were grown at 37 °C in Luria Broth media for 16 hours. Cells were harvested by centrifugation at 7000 RPM for 10 min using a JA10 rotor. The bacterial pellet was resuspended in lysis buffer (25 ml Spedding buffer A, 8.75 μl 2-Mercaptoethanol and 3 protease inhibitor tablets) and lysed by a French press for three consecutive cycles at 10,000 PSI pressure at room temperature. Cell debris was removed by centrifugation with a JA20 rotor at 13000 RPM for 45 min. For each 100 μl of cell lysate, 200 μl glacial acetic acid and 33 μl of 1M MgCl₂ were added. The precipitated RNA was removed after centrifugation at 14000g on a table-top Eppendorf microcentrifuge. Excess acid was removed by filtration through a 3kDa Amicon filter several times until the pH reached 6. The pH was checked to be between 5 and 6 that is suitable for strong cation

exchange (SCX) chromatography. The protein concentration was determined by a Bradford assay using bovine serum albumin as a standard.

Strong cation exchange of cell lysate proteins

Approximately 250 μ g of whole cell lysate proteins were loaded onto a 4.6mm ID SCX chromatography column. The following mobile phases were used; Mobile phase A: 6M urea + 20mM acetic acid; Mobile phase B: 6M urea + 20mM acetic acid + 500mM NaCl; pH 5. Proteins were eluted with a 90-minute piecewise gradient that was optimized for cell lysate protein separation. Twenty fractions were collected using C4 trapping columns using previously described instrumentation.^{33,34} Each of the 20 fractions was desalted using 5% acetonitrile, 0.1% Trifluoroacetic acid (TFA) and then eluted using 90% acetonitrile, 0.1% TFA. The fractions were dried in a vacuum evaporator to remove the acid and acetonitrile. The samples were then resuspended in 25mM NH₄HCO₃ to prepare for digestion.

Trypsin digestion and amidination

SCX-fractionated proteins were reduced by adding 10mM dithiothreitol (DTT) and incubating for 1 hour at 60°C. The samples were allowed to cool to room temperature and alkylated using 10mM iodoacetamide. The reaction took place in the dark for 1 hour. Reduced and alkylated proteins were digested using trypsin with a protein-to-enzyme ratio of 50:1 for 18 to 20 hours at 37 °C. The resulting peptide samples were divided into two parts; one was amidinated and the other used as a control. For the amidination reaction, S-methyl thioacetamidate (SMTA) at a concentration of 43.4g/l was added to the peptides at a volume equal to the original volume of the peptide solution. The reaction was allowed to proceed for 1 hour at room temperature. During this time, an equal amount of SMTA solution was added to the reaction mixture every 15 minutes. After 1 hour the amidination reaction was stopped by adding 5% TFA and 5% ACN. Excess SMTA was removed and the samples were concentrated by Pierce PepClean C18 Spin columns. Peptides were eluted with 70% acetonitrile. Samples were dried in a vacuum evaporator, reconstituted in 25mM NH₄HCO₃ buffer with 0.1% formic acid, and mass analyzed. The control samples were also treated with 0.1% formic acid before mass analysis.

LC-MS/MS

One set of the SCX-separated protein fractions was digested and then analyzed with a Waters or Eksigent liquid chromatograph using a 1mm C18 reversed phase column with a 120 minute gradient. The eluent was electrosprayed into a ThermoFisher LTQ Orbitrap. By using its HCD capability, lower mass ion fragments such as b₁ ions could be detected that were not typically observable with the LTQ ion trap alone. Likewise, eight sets of amidinated and unlabeled digested fractions were run on an Eksigent capillary liquid chromatograph and ThermoFisher LTQ ion trap mass spectrometer. In these cases a 100 minute gradient was employed for each sample.

Database searches

Acquired mass spectrometer RAW files were converted to mgf files using the default parameters in ProteomicsTools.³⁵ Converted mgf files were searched by MSGF+³⁶ against

the *E. coli* reference proteome consisting of 4,306 sequences (Supplementary Information) and the often-observed contaminant proteins from the common Repository of Adventitious Proteins (<http://www.thegpm.org/crap>). In total, the database contained 4,306 *E. coli* proteins and 116 contamination proteins, as well as the 4,422 decoy reversed protein sequences to estimate false discovery rate by target-decoy approaches. For the LTQ ion trap data, the searches were carried out with a 1.5 Da precursor tolerance and semi-tryptic selectivity. Maximum charge was set at 5 and minimum charge at 1. Carbamidomethyl cysteine was set as a fixed modification while methionine oxidation, peptide N-terminal amidination (+41.027) and lysine amidination (+41.027) were all variable modifications. For the Orbitrap/HCD data, the precursor tolerance was set as 10ppm, the instrument type is 'High-res' LTQ and the fragment method is selected as 'HCD'. The other parameters were the same as with the LTQ instrument. It is worth noting that MSGF+ does not accept a "fixed" fragment ion tolerance. Instead, it utilizes mass errors in scoring; e.g., a peak with error 0.01 Da contributes higher score than another with error 0.2 Da. MSGF+ also selects an appropriate error model based on the "-inst" parameter. For the LTQ data, we used the default parameter of the instrument. For Orbitrap/HCD, we set the parameter as '-inst 1'.

Normalization method from spectral counting label-free quantification

We first obtained the raw spectral counts for each peptide across all samples. Assuming equal peptide abundances in each sample, we scaled the observed spectral count for each peptide in a sample by the sum of the spectral counts for all peptides detected in that sample. This information was used to interpret the relative detectability of amidinated peptides.

Classification-based post-processing

We developed a simple post-processing procedure to predict the probability that a PSM is a correct identification. The procedure involved training a logistic regression model based on the spectral and peptide features. The positive examples contained the top PSM for every spectrum whereas the negative examples contained the second best PSM. Both groups were selected from the top 10% of identifications, excluding those from the decoy database, in order to ensure that the top PSM candidates were correct. The predictor was then applied to top PSMs and the target-decoy approach was used to select the score threshold corresponding to the chosen FDR. The details of this workflow are shown in Figure 3A.

The features used to represent peptide-spectrum matches (PSMs) were exclusively related to the annotated fragmentation spectra. They included: (1) precursor error (in ppm units), (2) the charge of the matched spectrum, (3) a binary indicator of a b_1 ion match that was set to 1 if the intensity was higher than 5% of the maximum peak intensity, (4) b_1 ion match intensity relative to the maximum intensity, (5) a binary indicator of a b_2 ion match that was set to 1 if the intensity was higher than 5% of the maximum intensity, (6) b_2 ion match intensity, (7) a binary indicator of a y_{N-1} ion match that was set to 1 if the intensity was higher than 5% of the maximum intensity, (8) y_{N-1} ion match intensity, (9) b ion matches intensity other than b_1 and b_2 , (10) the fraction of total intensity of the b ion matches other than b_1 and b_2 , (11) the total intensity of the y ion matches other than y_{N-1} , (12) the fraction of total intensity of the y ion matches other than y_{N-1} . In total, there were twelve numerical features used for predictor development.

To estimate different feature combination effects, we compiled the above-mentioned twelve features into three different groups: *b₁-including*, *b₁-excluding* and *b₁-only*. The *b₁-including* feature set included all twelve features, the *b₁-excluding* feature set excluded *b₁*, *b₂* and *y_{N-1}* features and the *b₁-only* feature set only kept the *b₁* feature, specifically only consisted of *b₁* and *y_{N-1}*. The *b₁-including* and *b₁-excluding* combinations kept the precursor and charge features. The precursor and charge features were excluded from the *b₁-only* experiments.

Detectability of amidinated peptides

We developed another logistic regression classifier to study the signatures of peptides whose detectability is increased after SMTA labeling compared to their unlabeled counterparts. The features used to represent the identified peptides in this model were as follows: (1) a binary vector of length 20 representing the N-terminal amino acid, (2) a binary vector of length 20 representing the C-terminal amino acid, (3) a length-20 vector of amino acid compositions (relative frequencies), (4) molecular weight of the peptide sequence, (5) aromaticity of the peptide sequence,³⁷ (6) hydrophobicity value based on the KD scale,³⁸ (7) hydrophobicity value based on the HW scale,³⁹ (8) hydrophobicity value based on EM scale,⁴⁰ (9) instability index of the peptide sequence,⁴¹ (10) flexibility of peptide sequence,⁴² (11) isoelectric point of the peptide, (12) fraction of residues in peptides that tend to form helix (amino acids V, I, Y, F, W, L), turn (amino acids N, P, G, S) and sheet (amino acids E, M, A, L) secondary structure, (13) length of the peptide sequence. In total, we engineered 89 numerical features for the predictor development.

Results

Workflow overview

Figure 1 graphically displays the experimental protocol. Proteins were extracted from the whole cell lysate of *E. coli*, then fractionated into 20 traps by ion exchange chromatography in order to reduce protein complexity. Tryptic digestion of the contents of each trap was followed by a reversed-phase HPLC-MS analysis. Two types of mass spectrometers were employed to analyze the tryptic peptides. The first, an LTQ-Orbitrap with HCD ion fragmentation, enabled the detection of all fragment ions, including *b₁* ions. The second, a standalone LTQ ion trap, had a low-mass cutoff that prevented the observation of most *b₁* ions. The two types of instruments were utilized to explore the overall characteristics of SMTA labeling and its effect on peptide fragmentation.

Improving peptide identification by incorporating *b₁* ions

In previous work we demonstrated that peptide amidination can significantly affect the mass distribution of fragment ions generated by collisional activation.^{27,28} In fact, *b₁* and, to a lesser degree, *y_{N-1}* ions become dominant features in MS/MS spectra of labeled peptides. This is unusual since *b₁* ions are relatively unstable^{28,43,44} and are often not observed when unmodified peptide ions are fragmented. Their dominant appearance in the fragmentation of amidinated peptides has been explained by the creation of a cyclic intermediate involving the amidino modification.²⁷ This facilitates peptide backbone cleavage in a manner analogous to the effect that aspartic acid has in inducing charge-remote peptide

fragmentation.⁴⁵ A complication to exploiting this phenomenon is the low-mass cutoff of ion trap instruments that can obviate the observation of b_1 ions. Our previous solution to this problem was to employ a TOF mass analyzer that had no such limitation. In the present experiments we utilize an Orbitrap instrument with high-energy collisional dissociation (HCD) ion fragmentation to enable the detection of low b_1 ion masses. We are thus able to investigate how peptide ion fragment distributions in a large scale HCD experiment are affected by the amidination modification and whether b_1 ions can specifically be exploited to improve peptide identifications.

The numbers of identified spectra in labeled and unlabeled experiments were comparable. In orbitrap labeled experiments, we acquired 172,370 raw spectra. By using target-decoy approaches, based on 1% FDR filtering strategy, 17,575 PSMs were identified and 176 PSMs were matched to the decoy database with an MSGF+ E-value cutoff of $2.26 \cdot 10^{-8}$ (i.e., 1% FDR). In orbitrap unlabeled experiments, 171,664 raw spectra were recorded and 17,176 PSMs identified with 1% FDR with an E-value cutoff of $2.16 \cdot 10^{-8}$ (172 PSMs were matched to the decoy database). In terms of unique peptide identifications based on peptide sequence, not on charge states or PTMs, 2,408 unique peptides were identified in SMTA experiments, 2,847 were identified in unlabeled experiments, 1,683 peptides were identified in both experiments, while 725 peptides were identified in the SMTA sample that were not identified in the unlabeled sample (an increase of 25.5%).

Altered fragmentation pattern of amidinated peptides

Of the 17,575 peptide-spectrum matches (PSMs) identified in SMTA experiments, 15,516 (88.3%) were identified as N-terminally amidinated peptides. We focus on N-terminally amidinated peptides here because the b_1 ion is predominantly observed following N-terminal amidination. When annotating spectra of both amidinated and unlabeled samples, a b_1 ion assignment was considered to be credible only when the experimentally observed intensity was at least 1% of the maximum intensity in the spectrum. Among the 15,516 PSMs from the amidinated data, 14,126 (91.0%) PSMs included a matched b_1 ion. This fragment was often intense, accounting on average for 87.9% of the maximum peak intensity in the spectrum. In contrast, of the 17,176 PSMs from unlabeled peptide experiments, 17,007 (99.0%) were identified as non-amidinated and 169 (1.0%) were identified as N-terminally amidinated or lysine-amidinated. We attributed these 169 amidinated spectra to incorrect matches (Supplementary Information).

We also annotated other ions (b_2 , y_{N-1} , $y_{N/2}$) in order to compare their relative intensities, where $y_{N/2}$ indicates a middle y ion, that dominates the intensity in unmodified peptides. We demonstrate this significant difference in a violin plot (Figure 2). Violin plots are a combination of a box plot and a kernel density plot.⁴⁶ Figure 2 shows density of b_1 , b_2 , y_{N-1} , $y_{N/2}$ ions in amidinated and unlabeled peptides. Specifically, b_1 ions contribute on average 70.1% of the maximum fragment ion intensity in SMTA spectra. Conversely, only 2.3% of the intensity of unmodified spectra are annotated as b_1 ions. b_2 also shows an obvious difference: 4.6% of the intensity of SMTA spectra are annotated as b_2 , compared with 21.5% of the intensity in unlabeled spectra. It is noted here that this significant intensity difference in b_2 is the reason to exclude b_2 from the feature set *b_1 -excluding*. Because of the

ease of detecting b_1 ions, complementary y_{N-1} ions are also expected, and this is validated in this statistical data. y_{N-1} ions have average 10.8% intensity in SMTA labeled samples, while in unmodified samples they only have 4.5% average intensity. The average intensity of y_{N-2} ions is not significantly affected by amidination. In unlabeled peptides they have an average intensity of 28.5% while in amidinated peptides their average intensity is 24.3%. In conclusion, our fragmentation data demonstrate that the SMTA did not significantly alter the y series fragmentation ions, or the b series ion intensities except for b_1 , b_2 and y_{N-1} .

Reproducibility of amidinated peptides

We also investigated the reproducibility of fragmentation patterns in SMTA vs. SMTA, unmodified vs. unmodified, and SMTA vs. unmodified experiments. We annotated the spectra by b series, y series, b- H_2O series, y- H_2O series, b- NH_3 series and y- NH_3 series. For each unique peptide, we then obtained a set of fixed-length vectors of dimension $6(N-1)$, where N is the length of the peptide sequence, of annotated ion intensities (a fragment ion was assigned the intensity of 0 no experimental peak could be matched within the m/z tolerance). Based on these annotated fragment-ion vectors, we calculated the Pearson correlation coefficients between each pair of spectra identified as the same peptide. The average correlation between all replicated SMTA spectrum-pairs was 0.94 with standard error 0.0016 over 2,131 unique peptides that were identified by at least two spectra, whereas the correlation between replicated unmodified experiments, consistent with our previous observations,²⁹ was 0.92 with standard error 0.002 over 1,805 unique peptides. Using the set of 1,683 unique peptides that were identified in both SMTA and unmodified experiments, we found that the average correlation coefficient between the SMTA and unmodified experiments was only 0.45 with standard error 0.0055 (the reproducibility for a set of unrelated unmodified peptides of the same length was 0.42, whereas the corresponding reproducibility for the SMTA peptides was 0.54). The P-value for the difference in means between unmodified vs. unmodified and SMTA vs. unmodified groups was zero. In summary, the high reproducibility of SMTA spectra and a very low agreement between SMTA and unmodified spectra provide evidence that SMTA labeling results in altered yet reproducible peptide fragmentation patterns.

Peptide-spectrum match post-processing

Based on the tendency of amidinated peptides to form b_1 ions, we adopted a supervised approach, with features designed to incorporate b_1 ions into a post-processing procedure (Figure 3A). Prior to classifier development, regular database searching was performed that resulted in a ranked list of peptides for each experimental spectrum. The top PSMs from most confident 10% of identifications were treated as positive examples, while the second best PSMs from the corresponding identifications were selected as negative examples (all decoy hits were excluded). This method was designed to estimate the probability that a PSM reported by a database search engine is correct, for both SMTA and unlabeled experiments, for each feature set (b_1 -including, b_1 -excluding and b_1 -only; see Methods). The three classifiers, based on logistic regression, were used to better understand the importance of b_1 ions for classification.

Prediction accuracy

To evaluate classification models, we estimated the area under the Receiver Operating Characteristic (ROC) curve of each classifier through 5-fold cross-validation. The area under the ROC curves were estimated as 0.86 for b_1 -only, 0.96 for b_1 -including and 0.94 for b_1 -excluding for the SMTA experiments and 0.80 for b_1 -only, 0.95 for b_1 -including and 0.94 for b_1 -excluding for the unlabeled experiments. Details of the cross-validation results are shown in Supplementary Information.

Improving peptide identification by post-processing

After scoring each PSM, we set out to apply these scores in the post-processing procedure, as shown in Figure 3A. Given a particular threshold cutoff τ , we obtained two groups of peptides: one with the prediction score greater than or equal to the cutoff τ , while the other group has scores lower than the cutoff τ . Following this step, we applied a 1% FDR control using target-decoy approach on each of these two groups separately to obtain the final identifications. The underlying rationale for this procedure is that the group with higher prediction probability has a lower chance of having decoy identifications. In Figure 3B, we plotted the MSGF+'s $-\log(E\text{-value})$ as a function of the prediction score. Forward hits are depicted in black and the decoy hits are in red. Figure 3B shows that the density of decoy hits is very high in the area where the prediction score is low. As the prediction scores gets higher, the density of decoy hits becomes lower.

As shown in Figure 3C, the dependence of the number of identifications on the probability cutoff τ varies for different prediction models. Specifically, by using the b_1 -including model and the prediction threshold of $\tau = 0.4$, approximately 17.2% more PSMs are identified at 1% FDR. The other models, b_1 -excluding and b_1 -only, were only able to increase the number of identifications by approximately 12.0%. We also applied the same strategy on the unlabeled peptide experiments. The results show that b_1 -only model has no impact on the identification, while the other two (b_1 -including and b_1 -excluding) exhibit similar trends, approaching 12% more identified PSMs. As expected, by incorporating the dominant b_1 and b_1 -related features in the post-processing, identifications could be significantly improved in SMTA MS/MS experiments.

While the number of PSMs has been significantly improved by our post-processing approach, we also compared the unique peptides identified by the original searching (MSGF + results without post-processing, which will be referred to as "baseline") and prediction-based post-processing. Figure 4 depicts Venn diagrams that indicate the number of unique peptides in each of these four data sets: UnlabelO represents the baseline IDs (unique peptides) in the unlabeled dataset whereas UnlabelP represents the IDs from the unlabeled dataset after applying post-processing. Similarly, SMTAO and SMTAP represent the baseline and the post processing procedure applied to SMTA experiments, respectively. Figure 4A shows the number of unique peptides supported by at least one spectrum and Figure 4B shows the number of unique peptides supported by at least two spectra. After the post-processing of the SMTA data set, 227 unique peptides not seen in the unlabeled experiments were identified, 134 of which had at least two PSMs supporting each peptide. The number of identified unique peptides also increased after applying post-processing to

the unlabeled dataset, with 258 peptides supported by one spectrum and 174 peptides supported by at least two spectra in the unlabeled dataset. Thus, post-processing increased the number of unique peptides in both unlabeled and SMTA experiments, with a slightly higher percent improvement in the SMTA experiments (12.5% in SMTA vs. 9.6% in unlabeled), suggesting that incorporating the fragment ion information into the post-processing is beneficial.

Peptide identification in MS/MS spectra without b₁-ions

To investigate whether amidination improves peptide identification even when b₁ ions are not exploited, peptides were analyzed using an LTQ ion trap mass spectrometer. Whole-cell lysate protein mixtures digested by trypsin were separated into two batches. One batch followed the regular experimental procedure (as described in Methods), while the other was amidinated by SMTA. Both peptide batches were analyzed using LC-MS/MS, with CID-fragmentation in the ion trap. These pairs of experiments were repeated eight times, resulting in 16 different LTQ LC-MS/MS runs. Peptide identifications obtained in amidinated and unlabeled experiments were compared.

In total, 1,617,477 PSMs were identified from LTQ experiments, from which there were 1,621 decoy hits with an E-value cutoff at $1.1 \cdot 10^{-9}$; 196 unique peptides and 1,503 proteins were identified under 1% FDR at the PSM level. Analysis of *E. coli* whole-cell lysates showed that SMTA labeling enabled the identification of different peptides: 5,342 peptides from non-amidinated experiments, 4,068 peptides in SMTA experiments and 2,648 in both experiments. Amidination thus enabled the identification of 1,420 (26.6%) additional peptides in the cell lysates that were not identified without this modification. Since the LTQ instrument has lower mass resolution and thus gives less confident identifications, we considered peptides supported by at least two PSMs in order to estimate the increase in confidence by SMTA labeling. 4,032 unique peptides were identified (supported by two or more PSMs) in unlabeled experiments, 3,296 were identified in SMTA experiments and 2,648 unique peptides were identified in both experiments. 648 additional unique peptides were identified in the amidination experiments, corresponding to a 16.1% improvement. Hence, we conclude that amidination of tryptic peptides could alter both the fragmentation and the detectability of peptides, resulting in the identification of different peptides in MS/MS spectra.

Figure 5 further demonstrates the ability to detect different peptides through SMTA. The number of unique peptides identified in unlabeled experiments increases as more replicated experiments were conducted, but the rate of increase goes down gradually. On the other hand, the slope increases considerably when SMTA data are included. Specifically, after seven unlabeled experiments were combined, the 8th unlabeled experiment was only able to identify 3% more unique peptides, whereas one batch of SMTA experiment can identify 7% additional unique peptides. Based on the eight replicated experiments on unmodified peptides, we used a logarithmic function to forecast the incremental number of unique peptides with the increasing number of replicated experiments. The fitted function is expressed as $y = 1606.3 \cdot \ln(x) + 2225.6$ ($R^2 = 0.9934$) in which x represents the number of replicated experiments and y represents the total number of identified unique peptides. By

using this logarithmic function (Figure 5) to extrapolate additional increases in unlabeled experiments, it is evident that SMTA enhances peptide identification while the replicated unlabeled experiments approach saturation after a small number of experiments.

Quantity change in amidinated peptides

We observed that between 16.1% and 26.6% different peptides were identified in eight replicated LTQ experiments supported by either one or two spectra. This limited increase is reasonable because of the inability to detect b_1 ions in the LTQ instrument. In contrast, the Orbitrap instrument benefits from the uniqueness of b_1 ions. Nevertheless, even with the LTQ instrument's inability to detect b_1 ions, amidination still enabled more peptides to be detected. Altered ionization efficiency and modulation of the fragment distribution are the two sources of improvement. Moreover, the ionization efficiency alteration not only contributed to the unique peptide identifications, but also to the spectral count of detected peptides.

We further examined the peptides that exhibited significant quantitative changes with SMTA compared with unlabeled experiments. These results are displayed as a heatmap in Figure 6, where the horizontal axis represents 16 (eight unlabeled and eight amidinated) experiments and the vertical axis represents identified unique peptides. The red color represents higher quantity and the blue color represents lower quantity based on spectral counts. We first normalized the label-free spectral counts for each peptide (see Methods). Student's t-test was then used to check whether the quantities were significantly different in SMTA experiments and unlabeled experiments. The P-value of 0.05 was chosen as a cutoff to select the peptides, resulting in 212 peptides showing higher quantities in SMTA experiments and 119 peptides having higher quantities in unlabeled experiments (Figure 6). Although both experiments were loaded with equal amounts of proteins, different peptides show different quantitative characteristics. Hierarchical clustering visibly distinguishes the SMTA vs. unlabeled experiments through the quantities of identified peptides, indicating that amidination alters the detectability of some peptides. When multiple experiments are run, this process diversifies the set of identified peptides and prevents saturation related to repeated experiments.

We next investigated the characteristics of peptides that were differentially detected in the SMTA and unlabeled experiments. Based on these selected peptides, we trained a logistic regression classifier with the features as described in Methods to predict whether amidination increases or decreases their detected quantity. The area under the ROC curve of this classifier was 76% which indicates that this is a good classifier (see Supplementary Information for the performance of the classifier). We also examined the top ten features with greatest contribution to classification accuracy. As shown in Table 1, the N-terminal residues played a vital role in the classifier, with residues G, E, D and N having positive contributions and residues V and F having negative contributions. As expected, hydrophobicity-related characteristics of the peptides were important features for the classifier.³¹ Because the amidinated peptides are more basic than unlabeled peptides, this labeling technique results in altered detectability of peptides, depending on their amino acid composition and overall hydrophobicity.

Summary and Conclusions

Our prior experience led us to investigate the impact of S-methyl thioacetimidate (SMTA) labeling of peptides on protein identification in high-throughput shotgun proteomics. By introducing an amidino group into the N-termini of peptides, their fragmentation patterns and detectability have been significantly altered. We observed that b_1 ions are primarily and significantly enhanced, which enables accurate recognition of peptide N-terminal amino acids. Accordingly, the complementary y_{N-1} ions are enhanced to a lesser degree, while b_2 fragment ions are diminished. Even though b_1 and y_{N-1} ions are most significantly intensified in the MS/MS spectra of amidinated peptides, the intensities of $y_{N/2}$ fragment ions are hardly affected. This is a notable feature that can be exploited in targeted proteomics such as selective reaction monitoring (SRM) for protein quantification.

Based on the 16 replicated LTQ experiments in which b_1 ions were not detected, we found that SMTA modification not only enables the identification of additional peptides, but also significantly changes the fragmentation patterns of some peptides. We reasoned that the detectability of these peptides is altered by the increased basicity of their N-termini. Understanding the characteristics of the peptides with altered detectability could help to identify and quantify some peptides that are unlikely to be observed in experiments with unlabeled samples.

Finally, in this work, we also showed that integrating prior knowledge of amidinated peptide fragmentation into the computational post-processing can substantially improve peptide identification. Overall, we estimated that the post-processing procedure implemented in this study identifies 12.5% additional peptides in SMTA experiments that are unlikely to be identified otherwise.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH grant R01 GM103725.

References

1. Angel TE, Aryal UK, Hengel SM, Baker ES, Kelly RT, Robinson EW, Smith RD. Mass spectrometry-based proteomics: existing capabilities and future directions. *Chemical Society Reviews*. 2012; 41:3912–3928. [PubMed: 22498958]
2. Walther TC, Mann M. Mass spectrometry-based proteomics in cell biology. *The Journal of Cell Biology*. 2010; 190:491–500. [PubMed: 20733050]
3. Altelaar AM, Munoz J, Heck AJ. Next-generation proteomics: towards an integrative view of proteome dynamics. *Nature Reviews Genetics*. 2013; 14:35–48.
4. Mann M, Kulak NA, Nagaraj N, Cox J. The coming age of complete, accurate, and ubiquitous proteomes. *Molecular Cell*. 2013; 49:583–590. [PubMed: 23438854]
5. Richards AL, Merrill AE, Coon JJ. Proteome sequencing goes deep. *Current Opinion in Chemical Biology*. 2015; 24:11–17. [PubMed: 25461719]

6. Wilhelm M, et al. Mass-spectrometry-based draft of the human proteome. *Nature*. 2014; 509:582–587. [PubMed: 24870543]
7. Ramalho R, Li S, Radivojac P, Hahn MW. Proteomic evidence for in-frame and out-of-frame alternatively spliced isoforms in human and mouse. *IEEE/ACM Trans Comput Biol Bioinform*. 2015
8. Mann M, Jensen ON. Proteomic analysis of post-translational modifications. *Nat Biotechnol*. 2003; 21:255–261. [PubMed: 12610572]
9. Steen H, Mann M. The ABC's (and XYZ's) of peptide sequencing. *Nat Rev Mol Cell Biol*. 2004; 5:699–711. [PubMed: 15340378]
10. Choudhary G, Wu SL, Shieh P, Hancock WS. Multiple enzymatic digestion for enhanced sequence coverage of proteins in complex proteomic mixtures using capillary LC with ion trap MS/MS. *Journal of Proteome Research*. 2003; 2:59–67. [PubMed: 12643544]
11. Tran BQ, Hernandez C, Waridel P, Potts A, Barblan J, Lisacek F, Quadroni M. Addressing trypsin bias in large scale (phospho) proteome analysis by size exclusion chromatography and secondary digestion of large post-trypsin peptides. *Journal of Proteome Research*. 2010; 10:800–811. [PubMed: 21166477]
12. Rietschel B, Arrey TN, Meyer B, Bornemann S, Schuerken M, Karas M, Poetsch A. Elastase digests new ammunition for shotgun membrane proteomics. *Molecular & Cellular Proteomics*. 2009; 8:1029–1043. [PubMed: 19116210]
13. Nguyen DTT, Guilleme D, Rudaz S, Veuthey JL. Fast analysis in liquid chromatography using small particle size and high pressure. *Journal of Separation Science*. 2006; 29:1836–1848. [PubMed: 16970187]
14. Magrini L, Famiglini G, Palma P, Termopoli V, Cappiello A. Boosting the detection potential of liquid chromatography-electron ionization mass spectrometry using a ceramic coated ion source. *Journal of The American Society for Mass Spectrometry*. 2015:1–8.
15. Hahne H, Pachi F, Ruprecht B, Maier SK, Klaeger S, Helm D, Médard G, Wilm M, Lemeer S, Kuster B. DMSO enhances electrospray response, boosting sensitivity of proteomic experiments. *Nature Methods*. 2013; 10:989–991. [PubMed: 23975139]
16. Klammer AA, MacCoss MJ. Effects of modified digestion schemes on the identification of proteins from complex mixtures. *Journal of Proteome Research*. 2006; 5:695–700. [PubMed: 16512685]
17. Hennrich ML, Boersema PJ, van den Toorn H, Mischerikow N, Heck AJ, Mohammed S. Effect of chemical modifications on peptide fragmentation behavior upon electron transfer induced dissociation. *Analytical Chemistry*. 2009; 81:7814–7822. [PubMed: 19689115]
18. Keough T, Youngquist R, Lacey M. A method for high-sensitivity peptide sequencing using postsurface decay matrix-assisted laser desorption ionization mass spectrometry. *Proceedings of the National Academy of Sciences*. 1999; 96:7131–7136.
19. Paizs B, Suhai S. Fragmentation pathways of protonated peptides. *Mass Spectrometry Reviews*. 2005; 24:508–548. [PubMed: 15389847]
20. Harrison AG. To b or not to b: the ongoing saga of peptide b ions. *Mass Spectrometry Reviews*. 2009; 28:640–654. [PubMed: 19338048]
21. Laskin J, Yang Z, Lam C, Chu IK. Charge-remote fragmentation of odd-electron peptide ions. *Analytical Chemistry*. 2007; 79:6607–6614. [PubMed: 17676923]
22. Payne SH, Yau M, Smolka MB, Tanner S, Zhou H, Bafna V. Phosphorylation-specific MS/MS scoring for rapid and accurate phosphoproteome analysis. *Journal of Proteome Research*. 2008; 7:3373–3381. [PubMed: 18563926]
23. Savitski MM, Lemeer S, Boesche M, Lang M, Mathieson T, Bantscheff M, Kuster B. Confident phosphorylation site localization using the Mascot Delta Score. *Molecular & Cellular Proteomics*. 2011; 10:M110–003830.
24. Saeed F, Pisitkun T, Hoffert JD, Wang G, Gucuk M, Knepper MA. An efficient dynamic programming algorithm for phosphorylation site assignment of large-scale mass spectrometry data. *IEEE International Conference on Bioinformatics and Biomedicine Workshops*. 2012:618–625.

25. Li S, Arnold RJ, Tang H, Radivojac P. Improving phosphopeptide identification in shotgun proteomics by supervised filtering of peptide-spectrum matches. *Proceedings of the ACM Conference on Bioinformatics, Computational Biology and Biomedical Informatics*. 2013;316–323.
26. Beardsley RL, Reilly JP. Quantitation using enhanced signal tags: a technique for comparative proteomics. *Journal of Proteome Research*. 2003; 2:15–21. [PubMed: 12643539]
27. Beardsley RL, Reilly JP. Fragmentation of amidinated peptide ions. *Journal of the American Society for Mass Spectrometry*. 2004; 15:158–167.
28. Beardsley RL, Sharon LA, Reilly JP. Peptide de novo sequencing facilitated by a dual-labeling strategy. *Analytical Chemistry*. 2005; 77:6300–6309. [PubMed: 16194092]
29. Li S, Arnold RJ, Tang H, Radivojac P. On the accuracy and limits of peptide fragmentation spectrum prediction. *Analytical Chemistry*. 2011; 83:790–796. [PubMed: 21175207]
30. Ji C, Arnold RJ, Sokoloski KJ, Hardy RW, Tang H, Radivojac P. Extending the coverage of spectral libraries: A neighbor-based approach to predicting intensities of peptide fragmentation spectra. *Proteomics*. 2013; 13:756–765. [PubMed: 23303707]
31. Tang H, Arnold RJ, Alves P, Xun Z, Clemmer DE, Novotny MV, Reilly JP, Radivojac P. A computational approach toward label-free protein quantification using predicted peptide detectability. *Bioinformatics*. 2006; 22:e481–e488. [PubMed: 16873510]
32. Li YF, Arnold RJ, Tang H, Radivojac P. The importance of peptide detectability for protein identification, quantification, and experiment design in MS/MS proteomics. *Journal of Proteome Research*. 2010; 9:6288–6297.
33. Karty JA, Running WE, Reilly JP. Two dimensional liquid phase separations of proteins using online fractionation and concentration between chromatographic dimensions. *Journal of Chromatography B*. 2007; 847:103–113.
34. Liu X, Reilly JP. Correlating the chemical modification of Escherichia coli ribosomal proteins with crystal structure data. *Journal of proteome research*. 2009; 8:4466–4478. [PubMed: 19658437]
35. Sheng Q, Dai J, Wu Y, Tang H, Zeng R. BuildSummary: using a group-based approach to improve the sensitivity of peptide/protein identification in shotgun proteomics. *Journal of Proteome Research*. 2012; 11:1494–1502. [PubMed: 22217156]
36. Kim S, Pevzner PA. MS-GF+ makes progress towards a universal database search tool for proteomics. *Nature Communications*. 2014; 5
37. Lobry J, Gautier C. Hydrophobicity, expressivity and aromaticity are the major trends of amino-acid usage in 999 Escherichia coli chromosome-encoded genes. *Nucleic Acids Research*. 1994; 22:3174–3180. [PubMed: 8065933]
38. Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology*. 1982; 157:105–132. [PubMed: 7108955]
39. Hopp TP, Woods KR. Prediction of protein antigenic determinants from amino acid sequences. *Proceedings of the National Academy of Sciences*. 1981; 78:3824–3828.
40. Eisenberg D, Schwarz E, Komaromy M, Wall R. Analysis of membrane and surface protein sequences with the hydrophobic moment plot. *Journal of Molecular Biology*. 1984; 179:125–142. [PubMed: 6502707]
41. Guruprasad K, Reddy BB, Pandit MW. Correlation between stability of a protein and its dipeptide composition: a novel approach for predicting in vivo stability of a protein from its primary sequence. *Protein Engineering*. 1990; 4:155–161. [PubMed: 2075190]
42. Vihinen M, Torkkila E, Riikonen P. Accuracy of protein flexibility predictions. *Proteins: Structure, Function, and Bioinformatics*. 1994; 19:141–149.
43. Tu YP, Harrison AG. The b1 ion derived from methionine is a stable species. *Rapid Communications in Mass Spectrometry*. 1998; 12:849–851.
44. O'Hair RA, Reid GE. The search for stable gas phase b1 ions derived from aliphatic amino acids: a combined experimental and ab initio study. *Rapid Communications in Mass Spectrometry*. 2000; 14:1220–1225. [PubMed: 10918371]
45. Wysocki VH, Ross MM. Charge-remote fragmentation of gas-phase ions: mechanistic and energetic considerations in the dissociation of long-chain functionalized alkanes and alkenes. *International Journal of Mass Spectrometry and Ion Processes*. 1991; 104:179–211.

46. Hintze JL, Nelson RD. Violin plots: a box plot-density trace synergism. *The American Statistician*. 1998; 52:181–184.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Workflow overview. Proteins were extracted from whole cell lysate of *E. coli*, and fractionated into 20 traps by ion exchange chromatography. Tryptic digestion of the contents of each trap was followed by reversed phase HPLC-MS/MS analysis. Two types of mass spectrometers were employed to analyze the tryptic peptides.

Figure 2.

Violin plot of b_1 , b_2 , y_{N-1} , $y_{N/2}$ ions in amidinated and regular peptides. X-axis indicates the fragment ions and the samples, y-axis represents the intensity/maximum intensity ration.

Red color represents b_1 , blue color represent b_2 , green is y_{N-1} ion and black is the $y_{N/2}$.

Figure 3.

(A) Prediction procedure. (B) XYPlot of the Evaluated MSGF+ as a function of prediction scores. (C) The number of identified spectra (y-axis) at different threshold τ (x-axis) when the b₁-including, b₁-excluding and b₁-only models (represented by squares, triangles and circles, respectively) are used for the post-processing of SMTA (purple) and un-labeled (blue) experiments.

Figure 4.

Four-way Venn diagram of unique peptide identifications in unlabeled (UnlabelO and UnlabelP representing the IDs before and after post-processing, respectively) and the SMTA (SMTAO and SMTAP representing the IDs before and after post-processing, respectively) datasets. Panel *A* shows the number of unique peptides supported by at least one spectra, Panel *B* shows the number of unique peptides supported by at least two spectra.

Figure 5.

The number of identified unique peptides from eight unlabeled and eight SMTA experiments by MSGF+. The first eight points show the unique peptides identified in eight unlabeled experiments, while the following eight points show the incremental identification results from eight SMTA experiments. At point 16, 7196 were identified, whereas an estimated number of identified peptides after 16 unlabeled experiments is 6678, an estimated 7.7% increase.

Figure 6.

The heatmap of quantities of identified unique peptides in unlabeled and SMTA across 16 experiments. The x-axis is the 16 experiments (8 unlabeled and 8 SMTA) and the y-axis represents the identified unique peptides. The color reflects the normalized spectral counts quantity: blue indicates lower quantity and red indicates higher quantity.

Table 1

Regression coefficients of the top ten features contributing to the classifier of pep-tides with differentially detected quantities in SMTA and unlabeled datasets.

Feature	Regression coefficient
N-Terminal G	1.32
N-Terminal V	-1.08
N-Terminal E	1.08
N-Terminal F	-1.04
C-Terminal R	0.95
Grand Average of Hydropathy (GRAVY)	0.91
Hydrophobicity of Second C-terminus Amino Acid	-0.84
N-Terminal D	0.82
Hydrophobicity of First N-terminus Amino Acid	-0.72
N-Terminal N	0.71