12. Guirguis-Blake JM, Senger CA, Webber EM, Mularski RA, Whitlock EP. Screening for chronic obstructive pulmonary disease: evidence report and systematic review for the US Preventive Services Task Force. *JAMA* 2016;315:1378–1393.
13. Martinez FJ, O'Connor GT. Screening, case-finding, and outcomes for adults with unrecognized COPD. *JAMA* 2016;315:1343–1344.
14. Thorat Y, Salvi S, Kodgule R. Peak flow meter with a questionnaire and mini-spirometer to help detect asthma and COPD in real-life clinical practice: a cross-sectional study. *NPJ Prim Care Respir Med* 2017;27:32.

Copyright © 2018 by the American Thoracic Society

Cigarette Smoke Is an Endothelial Cell Toxin

To the Editor:

The report by Polverino and colleagues (1) provides a wonderful example of productive interdisciplinary collaboration. The authors connect lung and kidney endothelial cell damage and cigarette smoke in patients with chronic obstructive pulmonary disease and in smoke-exposed mice. This is an important study with far-reaching implications. Mechanistically, the authors point to generation of advanced glycation end products and receptor for advanced glycation end products activation triggered by oxidant stress, yet there are likely additional mechanisms that can be considered. Because the kidneys do not inhale, it is evident that it is the “circulating cigarette smoke” that is toxic to endothelial cells. For example, one component of cigarette smoke is the highly aggressive aldehyde acrolein (2–4), which can be measured in the blood and urine of smokers. A large number of experimental studies leave no doubt that cigarette smoke extract injures cultured endothelial cells. These findings have been frequently ignored or characterized as clinically irrelevant, yet they provide mechanistic insights and allow us to consider the hypothesis that emphysema is a vascular problem (5). If the notion that cigarette smoke cannot cross the alveolar–capillary barrier is dismissed, the fact that components of cigarette smoke are circulating has been firmly established. Although cigarette smoke extract *in vitro* kills endothelial cells directly, *in vivo* cigarette smoke components could, in susceptible individuals, trigger an immune response that includes the production of antiendothelial antibodies (6). Another implication of the findings published by Polverino and colleagues is that not only are the lung and kidney microvessels affected by cigarette smoke components but similarly the microcirculation of the skeletal muscles, the myocardium, and the retina are affected (1).

Interesting correlations, not provided by the authors, are those between albuminuria and the diffusing capacity of the lung for carbon monoxide and albuminuria and urinary isoprostane concentration (7). Last, this study suggests that the vascular chronic obstructive pulmonary disease phenotype might benefit from antioxidant treatments. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Norbert F. Voelkel, M.D.
University of Colorado
Denver, Colorado

References

1. Polverino F, Laicho-Contreras ME, Petersen H, Bijol V, Sholl LM, Choi ME, *et al*. A pilot study linking endothelial injury in lungs and kidneys in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:1464–1476.
2. Voelkel NF, Mizuno S, Yasuo M. Does drug-induced emphysema exist? *Eur Respir J* 2013;42:1464–1468.
3. Nana-Sinkam SP, Lee JD, Sotto-Santiago S, Stearman RS, Keith RL, Choudhury Q, *et al*. Prostacyclin prevents pulmonary endothelial cell apoptosis induced by cigarette smoke. *Am J Respir Crit Care Med* 2007;175:676–685.
4. Kitaguchi Y, Taraseviciene-Stewart L, Hanaoka M, Natarajan R, Kraskauskas D, Voelkel NF. Acrolein induces endoplasmic reticulum stress and causes airspace enlargement. *PLoS One* 2012;7:e38038.
5. Voelkel NF, Gomez-Arroyo J, Mizuno S. COPD/emphysema: the vascular story. *Pulm Circ* 2011;1:320–326.
6. Taraseviciene-Stewart L, Voelkel NF. Molecular pathogenesis of emphysema. *J Clin Invest* 2008;118:394–402.
7. Santini G, Mores N, Shohreh R, Valente S, Dabrowska M, Trové A, *et al*. Exhaled and non-exhaled non-invasive markers for assessment of respiratory inflammation in patients with stable COPD and healthy smokers. *J Breath Res* 2016;10:017102.

Copyright © 2018 by the American Thoracic Society

Reply to Voelkel

From the Authors:

We thank Dr. Voelkel for his complimentary comments acknowledging the importance of our study (1). As Dr. Voelkel correctly pointed out, the vascular chronic obstructive pulmonary disease (COPD) phenotype is a clinical entity that has been underrecognized. Initial studies from Dr. Voelkel's group provided the first clue that endothelial cell death is sufficient to induce airspace enlargement in rodents (2–4) and occurs in lungs of patients with COPD (5). Our studies of cigarette smoke (CS)-exposed mice and patients with COPD further support this hypothesis and demonstrate that endothelial injury also occurs in the kidneys in patients with COPD and smokers (1). Likely, CS injures endothelial cells in other tissues, including skeletal muscle and bones, and contributes to the comorbidities that are frequently detected in patients with COPD (6).

As Dr. Voelkel correctly points out, we focused on the contributions of the oxidative stress–advanced glycation end products (AGEs)–receptor for AGEs pathway in the pathogenesis of CS-induced endothelial injury. We investigated clues from studies of humans and experimental models of diabetes mellitus and hypertension (common comorbidities of COPD) that linked endothelial injury and microvascular complications of these diseases to

the oxidative stress–AGEs–receptor for AGEs pathway (7). It is likely that other important mechanisms contribute to CS-induced endothelial cell injury, which warrants investigation in the future.

Last, our study supports the notion that the vascular COPD phenotype might benefit from angiotensin-converting enzyme (ACE) inhibitor (ACEi) treatment. It is noteworthy in this respect that a population study recently reported that use of ACEi and angiotensin receptor blockers is associated with slower progression of emphysema assessed by chest computed tomography, particularly in former smokers (8). Treatment with ACEi for indications other than COPD is associated with decreased mortality in patients with COPD hospitalized for COPD exacerbation (9), reduced risk of pneumonia in patients with COPD (10), and a slower rate of FEV₁ decline in smokers without COPD (11). Our results indicate that randomized clinical trials should be conducted to assess the efficacy of ACEi in patients with COPD with versus without endothelial dysfunction (on the basis of the presence or absence of microalbuminuria [MAB]) to determine whether the vascular COPD phenotype benefits the most from ACEi therapy. Moreover, routine screening of patients with COPD for MAB may have utility in the phenotyping and management of patients with COPD. MAB should also be evaluated as a noninvasive and inexpensive biomarker to evaluate the responses of patients with COPD to treatment with ACEi. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Francesca Polverino, M.D., Ph.D.

Bartolome R. Celli, M.D.

Caroline A. Owen, M.D., Ph.D.

Brigham and Women's Hospital, Harvard Medical School
Boston, Massachusetts

and

Lovelace Respiratory Research Institute
Albuquerque, New Mexico

ORCID ID: 0000-0001-9686-5698 (F.P.).

References

1. Polverino F, Laicho-Contreras ME, Petersen H, Bijol V, Sholl LM, Choi ME, *et al*. A pilot study linking endothelial injury in lungs and kidneys in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:1464–1476.
2. Kasahara Y, Tudor RM, Taraseviciene-Stewart L, Le Cras TD, Abman S, Hirth PK, *et al*. Inhibition of VEGF receptors causes lung cell apoptosis and emphysema. *J Clin Invest* 2000;106:1311–1319.
3. Farkas L, Farkas D, Warburton D, Gauldie J, Shi W, Stampfli MR, *et al*. Cigarette smoke exposure aggravates air space enlargement and alveolar cell apoptosis in Smad3 knockout mice. *Am J Physiol Lung Cell Mol Physiol* 2011;301:L391–L401.
4. Tudor RM, Zhen L, Cho CY, Taraseviciene-Stewart L, Kasahara Y, Salvemini D, *et al*. Oxidative stress and apoptosis interact and cause emphysema due to vascular endothelial growth factor receptor blockade. *Am J Respir Cell Mol Biol* 2003;29:88–97.
5. Kasahara Y, Tudor RM, Cool CD, Lynch DA, Flores SC, Voelkel NF. Endothelial cell death and decreased expression of vascular endothelial growth factor and vascular endothelial growth factor receptor 2 in emphysema. *Am J Respir Crit Care Med* 2001;163:737–744.
6. Degens H, Gayan-Ramirez G, van Hees HW. Smoking-induced skeletal muscle dysfunction: from evidence to mechanisms. *Am J Respir Crit Care Med* 2015;191:620–625.
7. Ramasamy R, Yan SF, Schmidt AM. Receptor for AGE (RAGE): signaling mechanisms in the pathogenesis of diabetes and its complications. *Ann N Y Acad Sci* 2011;1243:88–102.
8. Parikh MA, Aaron CP, Hoffman EA, Schwartz JE, Madrigano J, Austin JHM, *et al*. The Multi-Ethnic Study of Atherosclerosis Lung Study. Angiotensin-converting inhibitors and angiotensin II receptor blockers and longitudinal change in percent emphysema on computed tomography. *Ann Am Thorac Soc* 2017;14:649–658.
9. Mortensen EM, Copeland LA, Pugh MJ, Restrepo MI, de Molina RM, Nakashima B, *et al*. Impact of statins and ACE inhibitors on mortality after COPD exacerbations. *Respir Res* 2009;10:45.
10. Kim J, Lee JK, Heo EY, Chung HS, Kim DK. The association of renin-angiotensin system blockades and pneumonia requiring admission in patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2016;11:2159–2166.
11. Petersen H, Sood A, Meek PM, Shen X, Cheng Y, Belinsky SA, *et al*. Rapid lung function decline in smokers is a risk factor for COPD and is attenuated by angiotensin-converting enzyme inhibitor use. *Chest* 2014;145:695–703.

Copyright © 2018 by the American Thoracic Society