Remarkably different results between two studies from North America on genomic mutations and sensitivity to DNA demethylating agents for myelodysplastic syndromes

Guiping Wang¹, Shanshan Guo¹, Huashi Xiao¹, Liang Zong^{1,2,3}, Tetsuya Asakawa⁴, Masanobu Abe⁵, Wenqing Hu⁶, Jiafu Ji²

¹Medical Research Center, Clinical Medical College of Yangzhou University (Northern Jiangsu People's Hospital), Yangzhou 225001, China; ²Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Center of Gastrointestinal Surgery, Peking University Cancer Hospital & Institute, Beijing 100142, China; ³Department of Gastrointestinal Surgery, Clinical Medical College of Yangzhou University (Northern Jiangsu People's Hospital), Yangzhou 225001, China; ⁴Department of Neurosurgery, Hamamatsu University School of Medicine, Handayama, Hamamatsu-city, Shizuoka 431-3192, Japan; ⁵Division for Health Service Promotion, University of Tokyo Hospital, Tokyo 113-003, Japan; ⁶Division for Health Service Promotion, Heji Hospital of Changzhi Medical College, Changzhi 046011, China

Correspondence to: Prof. Liang Zong. Clinical Medical College of Yangzhou University (Northern Jiangsu People's Hospital), Yangzhou 225001, China. Email: 250537471@qq.com.

Submitted Oct 04, 2017. Accepted for publication Nov 20, 2017.

doi: 10.21147/j.issn.1000-9604.2017.06.14

View this article at: <https://doi.org/10.21147/j.issn.1000-9604.2017.06.14>

Sekeres *et al.* (1) conducted a multicenter randomized, controlled trial to compare whether azacitidine-based combinations with lenalidomide or vorinostat produce superior overall response rates to azacitidine in the treatment of myelodysplastic syndromes (MDS). In that trial, 224 patients with higher-risk MDS and 53 with chronic myelomonocytic leukemia (CMML) were enrolled and randomly assigned to the “azacitidine” group, “azacitidine plus lenalidomide” group or “azacitidine plus vorinostat” group. The researchers found that patients with MDS treated with azacitidine-based combinations had similar response rate to azacitidine monotherapy. Using genomic mutation analysis, they found that the overall response rate to azacitidine-based treatment was higher for patients with mutations in *DNMT3A* and lower for those with mutations in *SRSF2*. Whereas in another study, Welch *et al.* enrolled 26 patients with MDS and 90 with acute myeloid leukemia (AML) who were treated with decitabine, and they found that patients with *TP53* mutations had a higher response rate, but not those with *DNMT3A* mutations (2). We propose that this big discrepancy in the conclusions between the two studies might have been caused by the presence of many co-interacting factors, e.g. study aims, DNA demethylating agents, treatment protocols, and patient sources. Still, given that both studies

focused on DNA-demethylating-agent-based therapy for MDS or leukemia, it seems that a direct comparison and discussion could be fruitful.

To easily understand the response rates for patients with and without mutations of *DNMT3A*, *SRSPF2*, and *TP53* obtained in the two studies, we summarized the studies' data in Table 1. First, both studies showed a relatively higher response rate for patients with *DNMT3A* mutations, but only the study by Sekeres *et al.* showed a difference that was statistically significant. Second, both studies clearly found that patients with *SRSF2* mutations showed a lower response rate. Third, the study by Welch *et al.* showed that patients with *TP53* mutations had a much higher response rate (100%) in comparison with those with wide-type *TP53* (41.0%), whereas these rates were 45.5% vs. 35.2%, respectively, in the study by Sekeres *et al.*

To explore the possible reason for the discrepancy between the two studies, we considered several possibilities, namely: 1) different DNA demethylating agents; 2) treatment strategies with single or combined agents; and 3) patients enriched for MDS or leukemia. However, the first two possibilities are unlikely to explain the discrepancy, because a single agent of decitabine was able to achieve a higher response rate in patients with wide-type *DNMT3A* or *TP53* mutations than azacitidine-based combination therapy. Further, azacitidine and decitabine are

Table 1 Response rates for patients with and without mutations of *DNMT3A*, *SRSPF2*, and *TP53*

Gene symbol	Studies (Ref)	Mutated response rate [n/N (%)]	WT response rate [n/N (%)]	P
<i>DNMT3A</i>	MDS+CMML (1)	8/12 (66.7)	34/101 (33.7)	0.025
	AML+MDS (2)	7/11 (63.6)	46/88 (52.3)	0.500
<i>SRSPF2</i>	MDS+CMML (1)	4/23 (17.4)	34/83 (41.0)	0.049
	AML+MDS (2)	2/11 (18.2)	29/49 (59.2)	0.020
<i>TP53</i>	MDS+CMML (1)	10/22 (45.5)	32/91 (35.2)	0.370
	AML+MDS (2)	21/21 (100)	32/78 (41.0)	<0.001

WT, wide-type.

antimetabolites with quite similar and unique mechanisms of action as epigenetic regulators. As for the third possibility, the patients in the studies by Sekeres *et al.* and Welch *et al.* were enriched for high-risk MDS and AML, respectively. Although Welch *et al.* showed that 12 AML and 9 MDS patients with *TP53* mutations all showed good responses to decitabine, the number of MDS patients included was small, and unfortunately their finding could not be validated in another trial with 224 higher-risk MDS patients by Sekeres *et al.*

Based on the above analysis, we might conclude that the sensitivity markers based on genomic mutations for MDS and AML might be different, although they share the same resistance marker. This finding should be further validated in the future.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

References

1. Sekeres MA, Othus M, List AF, et al. Randomized phase II study of azacitidine alone or in combination with lenalidomide or with vorinostat in higher-risk myelodysplastic syndromes and chronic myelomonocytic leukemia: North American Intergroup Study SWOG S1117. *J Clin Oncol* 2017;35:2745-53.
2. Welch JS, Petti AA, Miller CA, et al. *TP53* and decitabine in acute myeloid leukemia and myelodysplastic syndromes. *N Engl J Med* 2016;375:2023-36.

Cite this article as: Wang G, Guo S, Xiao H, Zong L, Asakawa T, Abe M, Hu W, Ji J. Remarkably different results between two studies from North America on genomic mutations and sensitivity to DNA demethylating agents for myelodysplastic syndromes. *Chin J Cancer Res* 2017;29(6):587-588. doi: 10.21147/j.issn.1000-9604.2017.06.14