duty of making a laborious, careful, and, as far as possible, unprejudiced examination of the evidence presented in relation to the subject, unless, indeed, its consideration were given up altogether, an alternative which could not for a moment be entertained. I have been compelled to arrive at conclusions somewhat at variance, perhaps, with my own preconceived notions; but I have the satisfaction of believing, that a fair and correct exposition of this much 'vexed question' has been presented, as the result of what has certainly been the most arduous portion of my labours in relation to this work. I have at least performed it with most honest intentions to eliminate the truth, as far as has been possible; whether altogether successfully or otherwise, time only, as it augments our experience, can determine."—(P. 275.)

We have no space left to consider the remaining chapters on "Urinary Abscess and Fistula," "Retention of Urine depending on Stricture," and on "Stricture of the Female Urethra," which are treated at considerable length, in a manner highly creditable to the author's talents and industry. Indeed we do think that Mr. Thomson has discussed all the points connected with stricture of the urethra with great ability, so that his treatise is one of much practical interest, and cannot fail to be useful to the profession. It amply justifies the College of Surgeons in awarding to him the Jacksonian prize.

N. W.

II. *The Pathology and Treatment of Leucorrhœa.* By W. TYLER SMITH, M.D. London: 1855.

In recent times there has, in the whole range of medicine, been no more fruitful source of polemical discussion than leucorrhœa. The reintroduction in the present century of the use of the speculum matricis into ordinary practice, revealed to obstetricians a new set of recondite lesions. The greatly increased frequency of physical examination of the female organs to which this gave rise, resulted in the proclamation, if not in the discovery, of various new disorders of these parts. The examinations necessary for their investigation were easy to men of the meanest capacity. The new field of practice was, as might be expected, entered upon with the greatest zeal and alacrity by a crowd of doctors. The cases were all in a class, having the greatest development of our common emotional feelings, but having also the peculiar weaknesses implied in the very name of *woman*. In such a field, it was not to be wondered at that enthusiasm should be engendered; it was much to be lamented that this should so soon have degenerated, in the case of many physicians, into great imprudence, and in not a few into gross charlatanry. Women, enervated by luxury, overpowered by hysteria, or weakened by disease—new doctors ascribing all sorts of complaints to disorder or disease of the womb—new diseases found to be alarmingly frequent, inflammation, ulceration, eruptions, hypertrophy, engorgement, displacement,

twistings, strictures—new instruments of alarming or imposing appearance, speculums, pessaries, probes, syringes, and a host of others—what could be expected? The name of the uterine mania began to be applied to the state of the public mind. Alarm was excited for the safety of female modesty, if not of female purity. The dogmas and conduct of the professional encouragers of this state of matters, became a subject of discussion in the medical journals. Like other errors, they could not bear the simple light. The discussions were often painful, if not disgusting, and often conducted in a bad spirit; but they have fortunately been, in a great degree, effectual in withering the exuberant proportions of the so-called uterine mania, and curbing the ardour of its professional abettors, who saw no difference, æsthetically, between the examination of the throat or the pulse and the examination of the womb, and who would have wished to assimilate, in freedom and familiarity, the conversation of the two sexes to that of individuals of the same sex.

In the midst of all this excitement, numerous physicians did not desert the strict path of duty, but, by careful observation and research, conducted in a spirit becoming the professors of a noble science and art, sought to establish the real nature and importance of the lesions found in *leucorrhœa*. These labours have been crowned with a great measure of success. It would be invidious to place the laurels on any individuals where so many have distinguished themselves, and where the exertions of the most successful have not always deserved unqualified praise. But the steps of progress in recent times may be best recognised in our country, by noticing, first, the general nosological descriptions of Sir Charles M. Clarke, then advancing to the observation of the abrasions and inflammations of the uterine neck, to which Dr. Henry Bennet has succeeded in attaching his name, and lastly, arriving at the minute, nay, microscopical account of these lesions, and appreciation of their nature, by Dr. Tyler Smith, in the work under our notice at present.

Just as these last and conclusive observations on *leucorrhœa* have been published, we find that another step in progress has been begun, and, as is not unnatural, it assumes, at first, a negative form. *Leucorrhœa* is now known and measured: it retains its importance as a frequent and debilitating disease; but it is discovered that in it we do not find, as was originally believed, the cause of the great mass of ordinary uterine ailments. Many of these remain after its cure, or after removal of the obvious lesions of the uterine neck. Many of these are found without, nearly as often as with these lesions; and attention is being more directed to the condition of the entire organ as the real cause of frequent aggravated cases of disease. But on this subject we must refrain from now entering.

Twelve chapters divide among themselves the whole work, but

the first five contain almost all the marrow. The last seven are, in some degree, wordy, not very closely reasoned, and not conclusive in their arguments.

In the first chapter, we have a minute account of the superficial layer of the mucous membrane, of the vagina, and os uteri. The whole of this tract is covered by squamous epithelium, beneath which are distributed numerous villi, which at the os uteri have a filiform shape, and are observed to contain vascular loops in their interior. Over the surface are distributed the openings of numerous mucous follicles, which are abundant in the lower part of the canal, but almost, if not altogether, disappear on the upper, and have not been observed on the os uteri. The second chapter describes the rugæ and columnar arrangement of the canal of the cervix, with its very numerous glandular follicles. The epithelium of the os uteri and the external portion of the cervix is, like that of the vagina, constantly squamous; the epithelium, just within the os uteri, is cylindrical, but not ciliated. The epithelium of the follicular surface of the canal of the cervix is cylindrical or dentated, like the epithelium just within the os. It is also ciliated low down in the cervix, but not at its very lowest part, and the ciliated character is continued into the cavity of the fundus uteri. The villi found in the upper portion of the cervix are covered by dentated epithelium, just as is the case with the villi of the lowest part of the cervix.

In the third chapter, we have an account of the healthy secretions of the genital passages:—

“The sebaceous follicles, or fat glands, of the vulva and external parts of generation, secrete an oily matter, which, when it is secreted in unusual quantities, or in persons not observing strict cleanliness, may be seen gathered between the folds of the nymphæ as white fatty matter. The secretion of the mucous glands of the ostium vaginæ is said to be connected with the sexual function, and to be increased under excitement. The mucus of the vaginal canal is not found in any considerable quantity in the healthy subject; it is only secreted in sufficient quantity to keep the mucous surface in a state of lubrication. It lies upon the mucous membrane as a milky fluid, containing quantities of small curdy points or masses, and consists of a transparent or semi-transparent plasma, containing an abundance of scaly epithelium and its debris. . . . The plasma of the vaginal mucus appears, when first secreted, to resemble the plasma of the cervical mucus, but it is less viscid and tenacious. It is only after it has lain a short time upon the vaginal surface that it becomes curdled. The vaginal mucus is, as *Donné* first remarked, distinctly acid; and it is to the effect of the acid coagulating the albumen of the mucus, and not to the presence of epithelium, that its curdled appearance is attributable. The margin of the cervical canal, and the limits of the villi, covered by squamous epithelium, and the commencement of the villi covered by cylinder epithelium, seem to mark the division between the acid and alkaline secretion, and is thus a confirmation of the rule laid down by *M. Donné*.”

“In the unimpregnated condition, when the cervix uteri is found perfectly healthy, little or no discharge is seen issuing from the cervical cavity; but when the labia uteri are separated, the canal of the cervix appears to be full of its peculiar secretion. In examinations after death, in cases in which the uterine organs are in a healthy condition, the mucous crypts and the canal of the cervix

are generally found filled with a clear viscid mucus, so as to entirely block up the passages from the vagina to the cavity of the fundus. . . . The mucus consists of myriads of mucous corpuscles, entangled in a transparent viscid plasma. The plasma is so tenacious, that the mucous corpuscles are found to be arranged in strings when placed under the microscope, and individual corpuscles are frequently seen to be elongated from the same cause."

In the fourth chapter, the author enters upon the subject of leucorrhœa. This he divides into two species. The first he calls epithelial. It is produced by the vaginal surface. The second he calls mucous. It is the product of the cervix uteri. He introduces this distinction by a comparison of the two sources of the discharge:—

"The lining membrane of the vagina approaches in organization to the skin. It is covered by a thick layer of scaly epithelium. It contains, in the greater part of its surface, few, if any, mucous follicles or glands. Its secretion is acid, consisting entirely of plasma and epithelium; and the chief object of the secretion is the lubrication of the surface on which it is formed. On the other hand, the lining of the canal of the cervix is a true mucous membrane. It is covered, in great part, by cylinder epithelium. It abounds with immense numbers of mucous follicles, having a special arrangement. It pours forth a true mucous secretion, alkaline in character, and consisting of mucous corpuscles and plasma, with little or no epithelium."

The mucous discharge is the more common and important. It is a glandular secretion:—

"Quantities of mucous corpuscles and oily particles, with particles of epithelium entangled in the viscid alkaline plasma which gives the mucus its clearness and consistence, are found. The clear mucus is seen at the os uteri, sometimes adhering to the os itself, at others extending through the vagina, and presenting at the os externum in the form of a string, and also lying upon the walls of the vagina, in the curdy or creamy state to which it is reduced by the action of the vaginal acid."

Sometimes the two kinds of discharge are mixed:—

"The secretion, in cases of vaginal leucorrhœa, generally consists entirely of epithelium, in every possible phase of development, mixed with acid mucous plasma. . . . When the vaginal form of leucorrhœa becomes very severe, the villi become affected; and not only is epithelium separated with extraordinary rapidity, but pus is formed upon the irritable sub-epithelial or villous surface, which, when mixed with the epithelial matter, can hardly be distinguished from the mucous corpuscles of the cervix mixed with scaly epithelium. In the second form of epithelial disorder, the epithelium is thrown off in large shreds or pieces, in which the pavement-like arrangement of the scales is perfectly preserved. These laminæ frequently have upon them marks of the rugæ of the vagina, and somewhat resemble the cuticle, in cases of acute desquamation of the surface of the body. . . . Sometimes, on making a specular examination in these cases, the whole surface of the vagina is seen covered with a white coating, which may be removed by a forceps, in membranous pieces of considerable extent and thickness."

The fifth chapter is one of the most interesting. It points out the ordinary sequence of changes in cases of leucorrhœa. First, the excessive discharge, with inflammation; subsequently, abrasion, ulceration, induration, and hypertrophy. Here Dr. Smith has reaped very great advantage from the use of the microscope,

which has enabled him to describe, with the most refined exactness, the anatomy of the abrasion and ulceration of the cervix. This employment of such a valuable instrument is unexceptionable and admirable. But in another part of his work, Dr. Smith wishes to found diagnostic signs upon the microscopic examination of discharges. He would indicate that the practitioner could tell the seat of disease by the discharges, and has attempted by figures (See pages 38, 63, 64, 69, &c.) to illustrate this method. But we are certain that this is an unwarrantable stretch, the result of an injudicious wish to make the microscope do what it, as yet at least, cannot. The cervical and vaginal discharges of disease, even if not mixed, could not be satisfactorily discriminated from one another. Even on Dr. Smith's own showing, healthy cervical mucus contains epithelial scales, derived from the vagina. This will go far, even in health, to assimilate the two discharges. In disease, the mucus of both parts will contain globules, which Dr. Smith, or any one else, would be puzzled to sort or divide into pus and mucous globules. These considerations alone, apart from others, send us back to simpler and readier means of ascertaining the site of disease of the genital passages.

We refrain from entering upon the remaining chapters, which are far from being so complete, so closely reasoned, or so honourable to their author, as those we have noticed. They have served to fill up the book, and make a goodly, handsome volume. The work contains little that is absolutely original, most of the points of novelty in it having been made out previously by Robert, Donné, Whitehead, Robin, Huguier, Bennet, and others. But the author deserves the greatest praise for having himself, as it were, re-discovered the whole, re-analysed the subject, and set it down with ability in a body of doctrine, which will long retain its place in our literature. Every obstetrician, who wishes to be *au courant*, must make himself master of the contents of this work.

J. M. Duncan

III. *On the Displacements of the Uterus.* By J. MATTHEWS DUNCAN, A.M., M.D., Lecturer on Midwifery, &c. Pp. 43. Edinburgh: Sutherland & Knox. 1854.

On the Statics of Pregnancy. By J. MATTHEWS DUNCAN, M.D., &c., &c. Pp. 20. Neil & Co. 1855.

THAT there is a fashion in physic, as well as in music and many other things, has been long ago asserted and believed; and many now go the length of saying that this extends to diseases, as well as to their mode of treatment. It is no doubt true, that for long we heard of little else but bilious disorders. The bile and the liver got the blame of all the morbid feelings incident to the human frame—of all the diseases to which the members of the faculty were