III.—*On Obscure Diseases of the Brain, and Disorders of the Mind; their Incipient Symptoms, Pathology, Diagnosis, Treatment, and Prophylaxis.* By FORBES WINSLOW, M.D., D.C.L., Oxon. London: John Churchill, 1860.

WE have previously taken occasion in these pages, to express our admiration of the high talents and keen powers of observation evinced in the writings of Dr. Forbes Winslow. The volume now before us will widely extend his reputation as a shrewd observer, a philosophical writer, and a sound practical physician. It forms merely the introduction to what we suppose will be Dr. Winslow's *magnum opus*, "On Softening and other types of Organic Disease of the Brain," but still of itself contains a vast fund of information, communicated in an extremely attractive style. The subjects treated of by Dr. Winslow are far too numerous and important to be noticed in detail, and we must therefore content ourselves by drawing attention to the general scope of the writer.

In the introductory chapter great stress, but not too great, is laid upon the importance of an early recognition of the incipient symptoms of brain disease, with a view to prompt treatment. "Upon investigating the history of the diseases of the encephalon, how frequently does the medical man discover that positive and unequivocal symptoms of brain affection have existed, and perhaps, during the early stage, been observed for months and in some cases for years, without exciting any apprehension on the part of the patient, his family, or his friends." This is undoubtedly true, and much to be deplored; but we doubt whether we can look for any change in this respect. It happens frequently, perhaps generally, that when insanity has been developed, the friends and relatives of a patient inform his medical attendant of little eccentricities, curious actions, sudden impulses, &c., which had for some time been observed, but which till now seemed to them inexplicable, or perhaps trivial. But the natural proneness to take a favourable view of the case, and to hope for an improvement, is so strong, that to any one conversant with the common springs of action, it does not seem surprising that these premonitory symptoms should, even when giving rise to anxiety, be concealed. Dr. Winslow says that when other deviations from organic conditions take place, they do not meet with the neglect which is accorded to the disturbances of the brain; but, of course, this is simply the result either of ignorance or of the desire to conceal what might attach, in the world's eye, a stigma to a family, or blight the prospects of an individual. While, therefore, it is greatly to be wished that more common sense and enlightened views should prevail on the importance of early recognizing cerebral disturbance, and taking

steps to check it, we are not sanguine that in this age there will be material improvement in this respect. We have rarely seen a case of insanity in which premonitions had not been observed; but the practical difficulty has always been, even when the disease was well established, to convince relatives of the necessity of prompt action.

Our author considers the incipient signs of insanity under six heads:—1. Morbid phenomena of intelligence. 2. Morbid states of motion. 3. Morbid conditions of sensation. 4. Morbid phenomena of the special senses. 5. Morbid phenomena of sleep and dreaming. 6. Morbid phenomena of organic or nutritive life. A seventh section is devoted to the general principles of pathology, treatment, and prophylaxis.

In treating of the morbid phenomena of intelligence, Dr. Winslow starts by affirming “that in every case of disease of the encephalon, particularly if the organic change or pressure be established in the vesicular matter, or in the membranes immediately investing the brain, a disordered or abnormal state of cerebro-psychical phenomena may in the incipient stage, on careful examination, be detected.” This section of the work is most interesting and valuable, and the reader will find numerous speculative points treated with great ability, and illustrated by well-selected examples.

In his third chapter our author points out the impossibility of defining insanity—an attempt at which has more than once been the means of throwing discredit on medical witnesses in courts of law. He dwells upon some of the analogies between insanity and dreaming, and discusses fully and clearly some of the speculative points which suggest themselves on this topic.

The fourth chapter gives in detail the confessions of a patient after recovery, a narrative which, in the pages of the *Psychological Journal*, attracted much attention, and which in many points of view is worthy of being permanently recorded. We have the state of the mind when recovering from an attack of insanity described and admirably illustrated and, in successive chapters, the stages of consciousness, exaltation, mental depression, and aberration are discussed. Five chapters are occupied by the consideration of the morbid phenomena of memory, which form such important features of insanity.

In the eighteenth chapter the morbid phenomena of motion are treated of.

Previous to any positive lesion of motility, general muscular debility is complained of. “As the affection of the brain involving a disordered state of the motor *force* advances, the patient’s feet slip on one side. He is observed frequently to stumble whilst walking, as if the ligaments of the ankle-joint were weakened or elongated. He cannot put his foot or leg forward without an *obviously conscious effort*. Succeeding this

general deficiency of muscular power, there is occasionally noticed a want of *local specific* motor strength in one of the limbs." "Thus, a gentleman who had previously manifested no symptom of decided illness, was observed frequently to drop his stick, as well as his umbrella, in the street. This was the first loss of motility observed for some weeks prior to an apparently sudden and acute attack of apoplexy, followed by paralysis."

A tremulous state of the muscular fibre is sometimes noticed in the incipient stages of cerebral disease, affecting the hands, arms, legs, or tongue; and irregular muscular action, as evidenced by crossing the legs, and such like movements, is another morbid phenomenon of motion. In considering convulsive action, Dr. Winslow divides the disease thus:—1. Epilepsy (with violent muscular movements). 2. Epilepsy (nocturnal in its character, and accompanied with slight muscular convulsion). 3. Epileptic vertigo (without muscular convulsions). The attacks of epilepsy occurring at night are often so slight as to escape serious attention, and may continue uncared-for till most serious results ensue. If this be the case, the physical and mental vigour are rapidly lost. Trousseau thinks "this mischievous form of epilepsy may continue for eight or ten years without any one, not even the patient himself, being aware of its existence. These are the principal diagnostic signs in such a case, viz., the biting of the tongue, and the involuntary emission of urine, especially in women. If the person who comes to consult you complains of waking with headache, if the lateral parts of the tongue are lacerated, and if you can ascertain that urine has been passed unconsciously, do not hesitate to declare that there has been a nocturnal attack of epilepsy. Moreover, in a very great number of cases, we may observe on the forehead, and especially below the eyes, myriads of petechiæ, the size of a pin's head, which are never produced under other circumstances." Epilepsy is preceded by a variety of symptoms—some affecting the motor power, others the sensory functions. In some the intellectual powers seem greatly exalted.

Affections of the tongue and muscles of the mouth are often observed in the early stage of paralysis, and in cases of general paralysis it is sometimes apparently hypertrophied. Perhaps one of the most striking and obvious precursory symptoms of organic disease of the brain, is the change which the hand-writing undergoes. Those who have been remarkable for a clear and exact caligraphy become slovenly, write unevenly, make mistakes and corrections. We have had occasion to observe this frequently, and can appreciate the value of the symptom. Indeed, the patient himself may be quite conscious of the change, and may even make plausible excuses for it. Paralysis agitans is another of the incipient morbid phenomena of the motor power. The morbid phenomena of speech, of sensation, of the special senses,

of sleep and dreaming, and of organic and nutritive life, are severally described.

Sleeplessness, it is well known, is a very common accompaniment of cerebral disease, and "in many cases a *certain forerunner of insanity*. "Cases of what may be termed idiopathic sleeplessness occasionally occur, in which the intellect is not (for a period) in the slightest degree disordered. Persons have been known to remain in this state of insomnia for several weeks—*never closing their eyes for five continuous* minutes in sleep. In one female patient the state of sleeplessness arose from a severe shock which she had received, consequent upon finding her husband, in the middle of the night, dead by her side, he having retired to bed apparently in excellent health. In these cases the patients are seldom heard to complain of the want of sleep. They appear to experience none of the usual sensations of bodily and mental fatigue, physical uneasiness, and discomfort, which follow ordinary states of partial and incomplete repose."

Dr. Winslow states that, while the pathological state of the brain may account for the sleeplessness so frequent in insanity, yet that, in many cases, the insomnia connected with mental derangement arises from a complete absorption, abstraction, concentration of the thoughts, and pre-occupation of the mind, in some terrible and fearful form of illusion, or frightful type of hallucination, that has firmly seized upon the morbid imagination.

The treatment of insanity is a matter of vital importance. "The cure and life of the patient, in all acute affections of the brain and disorders of the mind, depend—1. Upon the speedy detection of incipient symptoms; 2. Upon the accuracy of the diagnosis formed as to the nature of the cerebral affection; 3. Upon the immediate application of remedial treatment. The treatment of the early stage of insanity requires great delicacy, discrimination, and judgment. Under these circumstances, when the brain is in a morbid state of irritation, and the mind struggling between sanity and insanity, the person being conscious that his 'wits' are beginning 'to turn,' the medical attendant should proceed cautiously and discreetly in his examination. If the patient be led to believe from the conduct of the physician, or from anything which falls from him, that derangement of mind is suspected, the most painful and disastrous consequences in all probability will ensue. In the early stages of insanity the patient's suspicions are morbidly excited. Should he believe that he is imagined to be deranged, he will sometimes exhibit great violence and excitement. . . . When a medical man is called in to a case of this description, it is his duty first to direct his observations to the state of the general health. He will almost invariably detect either hepatic, gastric, cardiac, renal, or intestinal disorder, which may be irritating and sympathetically disordering the brain.

"It may occasionally be necessary to relieve the overloaded condition of the vessels of the brain. The patient often complains of severe headache, attended with an increase of temperature, for the relief of which the application of a few leeches, cold evaporating lotions, and ice to the head may be recommended. Great caution is, however, necessary in the use of depleting and antiphlogistic measures. Patients who have been injudiciously treated by such means, have often sunk into incurable melancholy. In *recent* attacks, occurring in young and plethoric subjects, when the symptoms are closely allied to inflammation of the brain, local bloodletting is often attended with the happiest results. . .

"In obscure and doubtful cases, tartrate of antimony will be found an excellent substitute for bleeding. Violent maniacal excitement, accompanied by every apparent indication of a high degree of cerebral congestion and inflammation, will often yield to the administration of this drug. The physician should begin with small doses, and gradually increase them, until the patient is able to take two or three grains without exciting actual vomiting. When bleeding is clearly inadmissible, cold applied to the head will be found not only to diminish vascular excitement, but to lessen powerfully the morbid sensitiveness of the cerebral organs. Should there, however, exist a tendency to active plethora and apoplexy, cold lotions and ice should be used with great caution. The prolonged hot bath, in conjunction with the cold *douche*, will often be found most efficacious in subduing maniacal excitement. The *douche* is to be used when the patient is in the hot bath.

"In the incipient, as well as in advanced stages of insanity, the generally overloaded and inactive state of the bowels should be relieved by means of purgatives. Much caution must be observed in the use of aperient medicine. Very frequently the whole surface of the mucous membrane of the intestinal canal is in a state of subacute inflammation. This condition acts sympathetically upon the brain and nervous system, and aggravates the mental irritation. When this morbid state of the intestines is present, the use of aperients should be preceded by the application of a few leeches (particularly if there be pain upon pressure), or counter-irritants to the neighbourhood of the abdominal affection. In other cases of insanity it will be necessary to exhibit drastic purgatives. Insanity has been known to yield to the steady and persevering use of cathartics. Hellebore had in ancient times the reputation of being a specific in cases of insanity. . . .

"In the treatment of incipient insanity, clearly unconnected with active head symptoms, there is no remedy which so effectually masters the disease as opium in one of its many formulæ. I am satisfied that a vast amount of mental derangement may be successfully treated in its early stage by the continuous and persevering administration of sedatives. When insanity is clearly

associated with a depressed condition of the vital powers, evidenced by a weak pulse, feeble action of the heart, and general anæmic state of the system, the exhibition of the hydrochlorate, acetate, or muriate of morphia, combined with iron and quinine, will, in a great majority of cases, be found to act like a charm in arresting the progress of the mental malady."

We have made this lengthy extract from Dr. Winslow's book with the view principally of showing that he does not concur in what we fear is the too common practice now in mental disorder—the expectant treatment. At the very time—if there be a time—when insanity is within the reach of medicinal treatment, how often is the opportunity neglected, not by relatives only, but by physicians, who allow precious moments to slip, and foster hopes of spontaneous amendment. Nay, we fear that in some of our asylums, where a wiser system should obtain, the patient is left very much without the chance even of benefiting by such means as those which experience has proved useful. Nor can we conceive any circumstances more likely to develope and confirm insanity, than those which surround the patient when first placed in seclusion. At once introduced into the company of a miscellaneous collection—it may be of imbeciles, maniacs, epileptics—he is left there to get well as best he can. Sometimes, indeed, a purgative course may be prescribed, and counter irritation be kept up, but beyond this no systematic course of treatment be resorted to, and maintained for a sufficient length of time to admit of beneficial results. Among other remedies than those mentioned which have been tried with advantage, are belladonna, conium, hydrocyanic acid, chloroform, Indian hemp, henbane, stramonium, and hops. Dr. Winslow has found the inhalation of chloroform sometimes very useful in cases of acute maniacal excitement.

In incipient general paralysis a persevering course of tonic treatment should be tried—iron, quinine, zinc, cod-liver oil, &c., while a seton or issue may be applied to the nape of the neck.

An important subject alluded to by our author is the necessity of watching with the utmost care the cerebral symptoms that follow all mechanical injuries to the head. It will be within the experience of most of our readers, that cases frequently occur in which cerebral symptoms manifest themselves long after the receipt of some, it may be very slight, injury of the head. Dr. Winslow gives some instances of this, which we may be allowed to supplement. An officer in the army, in the prime of youth, and of excellent constitution, was severely wounded at the storming of Badajoz. No bone was injured, but a musket-ball penetrated the scalp, and after running some way under it, passed out, without the bone having apparently been injured. He lay for some time insensible at the foot of the breach; was carried to his quarters; next day was quite well, and continued so for some years. By degrees traces of slight cerebral disturbance

were perceptible. These increased very slowly year by year, and ultimately, after a train of symptoms believed to depend upon ramollissement of the brain, and which included loss of motion of the lower limbs, inability to retain his urine, and loss of memory, he died exhausted. On inspection, in the neighbourhood of the vertex there was seen an irregular cicatrix extending for about one and a half or two inches in a transverse direction, more than a half being on the right side. The skull was a good deal thickened, principally in the frontal region, and its density was considerably increased. A large quantity of serum (from four to six ounces) issued from beneath the dura mater. On removing the skull-cap, the anterior lobes of the brain were found atrophied to a considerable extent, the dura mater lying in loose folds over them. The sub-arachnoid cellular tissue was greatly distended with serum, in some places to such an extent as to separate the convolutions, and assume the appearance of cysts. The substance of the brain was vascular, presenting numerous bloody points. There was great effusion in the lateral ventricles. The septum lucidum, a considerable portion of the roof of the left ventricle, and a less portion of the roof of the right, were found in a state of softening, yielding to the slightest touch. The surrounding parts of the brain were firmer than natural, and presented a clotted appearance from the existence of numerous punctiform extravasations. In this case, for many years the symptoms progressed so slowly as to be noticed only by the intimate friends of the patient. The density of the skull was doubtless the result of the increased action set up in the bone immediately after the injury was received. This officer survived the receipt of his wound thirty years, and it will not surprise the psychologist to learn that insanity occurred in one of his children.

In another case a gentleman, from twenty to twenty-five years of age, fell upon the deck of a ship which he commanded, and severely injured the skull. The trephine was employed, and a considerable portion of bone removed. He remained in excellent health for nearly thirty years. At the expiry of this time it was obvious to his friends that his memory was impaired. He gradually grew worse, was confined to bed, became deprived of the use of his limbs, and ultimately died, after a protracted illness, the duration of which could not be calculated, as the invasion and progress of the disease had been so insidious. All the medical men consulted were of opinion, that the symptoms were connected with the injury received in early life. On inspection, the whole skull was found so dense and thick as to resemble ivory. The site of the opening made by the trephine was not to be discovered, it was so completely obliterated; neither was the silver plate which all the patient's relatives believed to have been placed over the opening in the skull found. The dura

mater was ossified in numerous points, thickened throughout, and the falx was completely converted into bone. One other case may be cited, in which death followed injury of the head long after its receipt. A gentleman, having at a dinner party indulged too freely in the pleasures of the table, fell, and received a severe contusion upon the head. He remained insensible for some time, but ultimately recovered, and was not subjected to any active treatment. Years afterwards, his health became impaired. Uneasy feelings were complained of in the head, and ultimately, epilepsy having supervened, he died in a fit. On examination it was found that a fissure had taken place at the time of the fall, at the base of the brain; that nature had, in her attempts to cure the evil by increased action in the part, produced an increased deposit of bone; this increasing, and acting as a constant and progressing source of irritation, the epileptic fits became more numerous, and ultimately fatal. We have seen a boy suffer for long from head symptoms, in consequence of being struck on the head by a potato.

We have made no attempt in this notice to enter upon any analysis of Dr. Winslow's book, or to consider critically any of the views he has propounded. His work is one of a character which places it pretty much beyond the scope of a moderate review. It will be read by the physician from beginning to end with unflagging interest, and will well repay perusal. At the same time, we may be allowed to say that there are occasional symptoms of a carelessness which does not generally characterize Dr. Winslow's writings, and which, we presume, is the result of his constant occupations. There are frequent repetitions, and sometimes a neglect of arrangement, which we are confident will be amended in a future edition. That this volume will be read with avidity by other sections of the community than the medical, is certain. We are afraid that in their hands some of the striking facts contained in it may cause much distress and no little anxiety; but if it answers the great end of calling attention to the rational, which is the early treatment of insanity, it cannot fail to be productive of good. We look forward with interest to the appearance of the sequel, which will be doubtless worthy of the subject and of Dr. Winslow's high reputation.

IV.—*A Medico-legal Treatise on Malpractice and Medical Evidence, comprising the Elements of Medical Jurisprudence.*
By JOHN J. ELWELL, M.D., member of the Cleveland bar.
New York, 1860.

THE subject of malpractice has not received much attention from English writers on forensic medicine. Mr. Chitty, in the preface