Published in final edited form as:

Semin Cell Dev Biol. 2018 August ; 80: 13–16. doi:10.1016/j.semcdb.2017.07.038.

Wound redox gradients revisited

Philipp Niethammer

Cell Biology Program, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY 10065, USA

Abstract

Evidence emerges that redox gradients regulate morphogenesis, inflammation, regeneration, and healing of tissues. At the example of redox signaling during the zebrafish wound response, I briefly discuss current ideas on how such patterns might be sensed and spatially regulated to guide physiological processes over distances in animals.

Introduction

In the early 20th century, Charles Manning Child's "gradient theory" [1] proposed that metabolic redox patterns regulate morphogenesis. After the discovery of genes as drivers of tissue patterning, Manning's ideas became a side note of science history. To date, the question whether and how gradients of oxygen and its reactive metabolites regulate organismal biology is gaining renewed attention. Wound-induced redox patterns, which are conserved from plants to animals, provide an exemplary, post-developmental case for redox signaling on tissue scales [2]. Owing to its accessibility for intravital imaging approaches, transparent zebrafish larvae are an attractive system to study function and spatiotemporal regulation of these patterns during wound healing.

1. Wound redox gradients in zebrafish larvae

We became interested in biological redox patterns through our work on mitochondrial respiration in frog egg extracts [3,4]. In this system, mitochondria generate cytoplasmic redox gradients over tens to hundreds of microns. Such length scales are meaningful for spatial regulation of tissues, or unusually large cells, such as frog oocytes. The extract experiments inspired us to test whether redox gradients also occur in intact animals. By imaging the H₂O₂ biosensor HyPer in live zebrafish larvae, we found that tail fin tip amputation elicits a ~150–300 μ m wide H₂O₂ gradient that extends from the wound margin into the tissue (Figure 1A). The gradient is generated by the Ca²⁺-activated epithelial NADPH-oxidase Duox [5], which transfers electrons from cytoplasmic NADPH onto molecular oxygen outside of the cell. The generated superoxide is dismutated into H₂O₂ which diffuses within the extracellular space and enters cells [6]. Inhibition of Duox

Correspondence: Philipp Niethammer (niethamp@mskcc.org).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

suppresses the H_2O_2 gradient and wound recruitment of distant leukocytes [7], suggesting that Duox mediates long-range signaling.

2. Signaling through wound redox gradients

Understanding long-range communication through redox gradients demands addressing two separate questions: How are these patterns sensed, and how are they generated within tissues?

So far, two non-exclusive ideas have been put forward to explain how leukocytes sense wound redox patterns (Figure 1B): Directly, through H_2O_2 -mediated protein oxidation in leukocytes [8], or indirectly, through enzymatic oxidation of lipids that diffuse and trigger leukocyte chemotaxis through G-protein-coupled receptor signaling [9]. Direct H_2O_2 sensing has been reported to occur through Lyn—a membrane-bound non-receptor tyrosine kinase in leukocytes, which is activated by transient cysteine oxidation [8] (Figure 1B). Active, oxidized Lyn is required for leukocyte recruitment to zebrafish and drosophila wounds [8,10]. Transient oxidation of regulatory “thiol switches”, such as the cysteine 466 in zebrafish Lyn [8], constitutes one of the better studied redox signaling mechanisms to date. Proteins susceptible to this type of regulation often contain deprotonated, and thus more reactive thiols at functionally important sites [11,12]. Signaling specificity is thought to emerge from the higher oxidant reactivity of “regulatory thiols” as compared to other thiols in the proteome environment, and from the selective accessibility of regulatory thiols to a specific reactive oxygen species (ROS), for example H_2O_2 . Some of the most reactive cysteines are found in professional antioxidant peroxidases [13,14], or specialized H_2O_2 -sensitive transcription factors, such as the bacterial protein OxyR [15]. Their high reactivity has been exploited for H_2O_2 biosensor design through fusion with fluorescent domains. HyPer [16], which is based on OxyR, and Orp1-roGFP2 and roGFP2-Tsa2, which are based on yeast peroxidases [17,18], detect low, physiological amounts of H_2O_2 only because they effectively compete with cellular antioxidants. Judging from proteins whose H_2O_2 reactivity is known, redox sensitive, non-antioxidant proteins are oxidized far slower by H_2O_2 than professional antioxidants or OxyR. For instance, the rate constant of the redox-regulated tyrosine phosphatase 1B (PTP1B) for its reaction with H_2O_2 is ~7 times higher than that of free cysteine, but still 6 orders of magnitude lower than that of peroxiredoxin [19]. Accordingly, thiol switches of non-antioxidant proteins may only become oxidized when antioxidant activity is locally or temporally suppressed. Peroxiredoxins may be transiently depleted by high pulses of oxidants or locally inactivated by phosphorylation, for example, near membranes. This is thought to create discrete time windows or spatial nanodomains permissive for redox signaling to non-antioxidant proteins [20,21]. Whether, when and where antioxidants are inactivated during physiological wound redox signaling remains experimentally little addressed. To this end, we recently developed an imaged-based method that measures antioxidant reaction kinetics from time-lapse HyPer measurements in wounded zebrafish larvae [22]. Our data suggest that H_2O_2 inactivates antioxidants within ~30 μm of the wound margin. Interestingly, we found no evidence that the wound-induced H_2O_2 gradient breaches antioxidant barriers at larger distances. This suggests that Lyn must be exempted from general antioxidant competition for leukocytes to directly sense H_2O_2 from > 30 μm away. Future experiments should explicitly test this possibility.

Not all redox signaling mechanisms are antagonized by antioxidants: By breaking down H_2O_2 antioxidants themselves may transmit signals through redox chains [23,24]. Such “redox relay signaling” may contribute to wound chemotaxis of leukocytes in zebrafish (Figure 1B). Duox consumes O_2 and NADPH to generate H_2O_2 and NADP^+ . NADP^+ also accrues during H_2O_2 detoxification by thioredoxin and glutathione reductases of the antioxidant chain. Much attention has been focused on the role of H_2O_2 as diffusible redox signal. However, Duox-induced increases of cytoplasmic NADP^+ may be of regulatory importance, too. Biosensor imaging in zebrafish larvae suggests that the wound redox gradient is paralleled by a gradient of cytoplasmic NADP^+ , decreasing from the wound margin into the tissue [25]. Enzymatic reactions that require NADP^+ as cofactor may be sensitive to such a gradient. For example, the microsomal NADP^+ dehydrogenase 5-HEDH, uses NADP^+ to oxidize the hydroxyl group of the lipid mediator 5-HETE into a keto group [26–29]. This generates 5-KETE—an oxidized lipid product that is strongly chemotactic for neutrophils and eosinophils [30]. 5-KETE attracts neutrophils to zebrafish wounds [9]. Through mediating intracellular peroxiredoxin oxidation, diffusing H_2O_2 could propagate increased $\text{NADP}^+/\text{NADPH}$ ratios to neighboring cells to activate 5-HEDH and 5-KETE synthesis around epithelial wounds. Per this model, both H_2O_2 and 5-KETE diffusion would contribute to the overall range of chemotactic wound signaling. The model implies a novel role for NADPH oxidases as regulators of NADP^+ dependent metabolism (e.g., of inflammatory lipids), in addition to their well-known role as ROS generators. It will be intriguing to test the physiological relevance of this concept in the future.

3. Spatial regulation of wound redox gradients

Per the above considerations, H_2O_2 tissue diffusion plays a central, paracrine role, either through mediating oxidation of regulatory protein thiols or lipid oxidation via $\text{NADP}^+/\text{NADPH}$ ratio changes at distances. But how are H_2O_2 diffusion patterns regulated? Biophysical aspects of redox signaling have remained little studied, although they could help to quantitatively understand H_2O_2 's tissue signaling roles [31]. Like water, H_2O_2 diffuses fast. But its signaling range within the tissue is limited by sink reactions that constantly remove H_2O_2 from the extracellular space. Namely, fast extracellular H_2O_2 diffusion, is counterbalanced, at least in part, by aquaporin-mediated uptake into cells and peroxiredoxin-thioredoxin-mediated breakdown within them [22]. Whereas gradients generated by H_2O_2 diffusion alone are expected to be transient and spatially unstable, reaction-diffusion establishes H_2O_2 steady-state gradients of discrete length-scale that are robustly maintained in the tissues over tens of minutes, that is, presumably long enough to convey information on tissue injury over distances. This highlights another regulatory role of antioxidants as active mediators of spatial redox patterning.

Conclusion

A century ago, Charles Manning Child only had toxic live dyes, such as methylene blue, to study redox gradients, and little means to determine their function [32]. Today, we can harness a growing panel of genetically encoded biosensors and the power of genetics and pharmacology. Combined with quantitative thinking, these approaches may elevate tissue

redox patterns from the niche of fascinating phenomenology into the realm of mechanistic physiology.

Acknowledgments

I thank Tim Mitchison for his comments on the manuscript, and apologize to the colleagues whose research I did not cite in this comment due to space limitations. The work was supported by the NIH grant GM099970, and an American Asthma Foundation Scholar grant to Philipp Niethammer.

References

1. Child, CM. Individuality in organisms. Chicago: Chicago, Ill: The University of Chicago press; 1915.
2. Suzuki N, Mittler R. Reactive oxygen species-dependent wound responses in animals and plants. *Free Radic Biol Med*. 2012; 53:2269–2276. DOI: 10.1016/j.freeradbiomed.2012.10.538 [PubMed: 23085520]
3. Niethammer P, Kueh HY, Mitchison TJ. Spatial patterning of metabolism by mitochondria, oxygen, and energy sinks in a model cytoplasm. *Curr Biol*. 2008; 18:586–591. DOI: 10.1016/j.cub.2008.03.038 [PubMed: 18406136]
4. Kueh HY, Niethammer P, Mitchison TJ. Maintenance of mitochondrial oxygen homeostasis by cosubstrate compensation. *Biophys J*. 2013; 104:1338–1348. DOI: 10.1016/j.bpj.2013.01.030 [PubMed: 23528093]
5. Donkó A, Péterfi Z, Sum A, Leto T, Geiszt M. Dual oxidases. *Philos Trans R Soc Lond, B, Biol Sci*. 2005; 360:2301–2308. DOI: 10.1098/rstb.2005.1767 [PubMed: 16321800]
6. Enyedi B, Zana M, Donkó Á, Geiszt M. Spatial and temporal analysis of NADPH oxidase-generated hydrogen peroxide signals by novel fluorescent reporter proteins. *Antioxid Redox Signal*. 2013; 19:523–534. DOI: 10.1089/ars.2012.4594 [PubMed: 23121369]
7. Niethammer P, Grabher C, Look AT, Mitchison TJ. A tissue-scale gradient of hydrogen peroxide mediates rapid wound detection in zebrafish. *Nature*. 2009; 459:996–999. DOI: 10.1038/nature08119 [PubMed: 19494811]
8. Yoo SK, Starnes TW, Deng Q, Huttenlocher A. Lyn is a redox sensor that mediates leukocyte wound attraction in vivo. *Nature*. 2011; 480:109–112. DOI: 10.1038/nature10632 [PubMed: 22101434]
9. Enyedi B, Kala S, Nikolich-Zugich T, Niethammer P. Tissue damage detection by osmotic surveillance. *Nat Cell Biol*. 2013; 15:1123–1130. DOI: 10.1038/ncb2818 [PubMed: 23934216]
10. Evans IR, Rodrigues FSLM, Armitage EL, Wood W. Draper/CED-1 mediates an ancient damage response to control inflammatory blood cell migration in vivo. *Curr Biol*. 2015; 25:1606–1612. DOI: 10.1016/j.cub.2015.04.037 [PubMed: 26028435]
11. Poole LB. The basics of thiols and cysteines in redox biology and chemistry. *Free Radic Biol Med*. 2015; 80:148–157. DOI: 10.1016/j.freeradbiomed.2014.11.013 [PubMed: 25433365]
12. Brandes N, Schmitt S, Jakob U. Thiol-based redox switches in eukaryotic proteins. *Antioxid Redox Signal*. 2009; 11:997–1014. DOI: 10.1089/ars.2008.2285 [PubMed: 18999917]
13. Rhee SG, Kang SW, Chang TS, Jeong W, Kim K. Peroxiredoxin, a novel family of peroxidases. *IUBMB Life*. 2001; 52:35–41. DOI: 10.1080/15216540252774748 [PubMed: 11795591]
14. Toledano MB, Delaunay A, Monceau L, Tacnet F. Microbial H₂O₂ sensors as archetypical redox signaling modules. *Trends Biochem Sci*. 2004; 29:351–357. DOI: 10.1016/j.tibs.2004.05.005 [PubMed: 15236742]
15. Aslund F, Zheng M, Beckwith J, Storz G. Regulation of the OxyR transcription factor by hydrogen peroxide and the cellular thiol-disulfide status. *Proc Natl Acad Sci U S A*. 1999; 96:6161–6165. [PubMed: 10339558]
16. Belousov VV, Fradkov AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Tersikh AV, et al. Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nat Methods*. 2006; 3:281–286. DOI: 10.1038/nmeth866 [PubMed: 16554833]

17. Gutscher M, Sobotta MC, Wabnitz GH, Ballikaya S, Meyer AJ, Samstag Y, et al. Proximity-based protein thiol oxidation by H₂O₂-scavenging peroxidases. *J Biol Chem*. 2009; 284:31532–31540. DOI: 10.1074/jbc.M109.059246 [PubMed: 19755417]
18. Morgan B, Van Laer K, Owusu TNE, Ezeri a D, Pastor-Flores D, Amponsah PS, et al. Real-time monitoring of basal H₂O₂ levels with peroxiredoxin-based probes. *Nat Chem Biol*. 2016; 12:437–443. DOI: 10.1038/nchembio.2067 [PubMed: 27089028]
19. Winterbourn CC. The biological chemistry of hydrogen peroxide. *Meth Enzymol*. 2013; 528:3–25. DOI: 10.1016/B978-0-12-405881-1.00001-X [PubMed: 23849856]
20. Wood ZA, Poole LB, Karplus PA. Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling. *Science*. 2003; 300:650–653. DOI: 10.1126/science.1080405 [PubMed: 12714747]
21. Woo HA, Yim SH, Shin DH, Kang D, Yu D-Y, Rhee SG. Inactivation of peroxiredoxin I by phosphorylation allows localized H₂O₂ accumulation for cell signaling. *Cell*. 2010; 140:517–528. DOI: 10.1016/j.cell.2010.01.009 [PubMed: 20178744]
22. Jelcic M, Enyedi B, Xavier JB, Niethammer P. Image-Based Measurement of H₂O₂ Reaction-Diffusion in Wounded Zebrafish Larvae. *Biophys J*. 2017; 112:2011–2018. DOI: 10.1016/j.bpj.2017.03.021 [PubMed: 28494970]
23. Sobotta MC, Liou W, Stöcker S, Talwar D, Oehler M, Ruppert T, et al. Peroxiredoxin-2 and STAT3 form a redox relay for H₂O₂ signaling. *Nat Chem Biol*. 2015; 11:64–70. DOI: 10.1038/nchembio.1695 [PubMed: 25402766]
24. Travasso RDM, Sampaio Dos Aidos F, Bayani A, Abranches P, Salvador A. Localized redox relays as a privileged mode of cytoplasmic hydrogen peroxide signaling. *Redox Biol*. 2017; 12:233–245. DOI: 10.1016/j.redox.2017.01.003 [PubMed: 28279943]
25. Tao R, Zhao Y, Chu H, Wang A, Zhu J, Chen X, et al. Genetically encoded fluorescent sensors reveal dynamic regulation of NADPH metabolism. *Nat Methods*. 2017; 14:720–728. DOI: 10.1038/nmeth.4306 [PubMed: 28581494]
26. Patel P, Cossette C, Anumolu JR, Erlemann K-R, Grant GE, Rokach J, et al. Substrate selectivity of 5-hydroxyeicosanoid dehydrogenase and its inhibition by 5-hydroxy-Delta6-long-chain fatty acids. *J Pharmacol Exp Ther*. 2009; 329:335–341. DOI: 10.1124/jpet.108.143453 [PubMed: 19164464]
27. Erlemann K-R, Cossette C, Grant GE, Lee G-J, Patel P, Rokach J, et al. Regulation of 5-hydroxyeicosanoid dehydrogenase activity in monocytic cells. *Biochem J*. 2007; 403:157–165. DOI: 10.1042/BJ20061617 [PubMed: 17166093]
28. Erlemann K-R, Rokach J, Powell WS. Oxidative stress stimulates the synthesis of the eosinophil chemoattractant 5-oxo-6,8,11,14-eicosatetraenoic acid by inflammatory cells. *J Biol Chem*. 2004; 279:40376–40384. DOI: 10.1074/jbc.M401294200 [PubMed: 15234979]
29. Erlemann K-R, Cossette C, Gravel S, Lesimple A, Lee G-J, Saha G, et al. Airway epithelial cells synthesize the lipid mediator 5-oxo-E₂E in response to oxidative stress. *Free Radic Biol Med*. 2007; 42:654–664. DOI: 10.1016/j.freeradbiomed.2006.12.006 [PubMed: 17291989]
30. Powell WS, Rokach J. The eosinophil chemoattractant 5-oxo-E₂E and the OXE receptor. *Prog Lipid Res*. 2013; 52:651–665. DOI: 10.1016/j.plipres.2013.09.001 [PubMed: 24056189]
31. Weavers H, Liepe J, Sim A, Wood W, Martin P, Stumpf MPH. Systems analysis of the dynamic inflammatory response to tissue damage reveals spatiotemporal properties of the wound attractant gradient. *Curr Biol*. 2016; 26:1975–1989. DOI: 10.1016/j.cub.2016.06.012 [PubMed: 27426513]
32. Blackstone NW. Charles Manning Child (1869–1954): the past, present, and future of metabolic signaling. *J Exp Zool B Mol Dev Evol*. 2006; 306:1–7. DOI: 10.1002/jez.b.21085 [PubMed: 16353198]

Figure 1.

(A) Tail fin wounding of zebrafish larvae generates an H_2O_2 gradient that reaches from the wound site into the tissue. Shown, representative image of the H_2O_2 gradient at a zebrafish amputation wound obtained by intravital HyPer ratio-imaging. Red, high $[\text{H}_2\text{O}_2]$. Blue, low $[\text{H}_2\text{O}_2]$. Scale bar, 100 μm . (B) Hypothetical scheme of wound redox signaling through NADPH oxidases. NADPH oxidases may signal through increasing both intracellular NADP^+ and extracellular H_2O_2 . H_2O_2 detoxification through antioxidant enzymes also increases intracellular NADP^+ . NADP^+ stimulates lipid chemoattractant production (5-KETE) through an NADP^+ -dependent dehydrogenase (5-HEDH). Lyn oxidation (possibly within intracellular nanodomains) is also required for chemotactic wound signaling to leukocytes. Note the feed-forward loop structure of the system.