Published in final edited form as:

J Antibiot (Tokyo). 2018 February ; 71(2): 263–267. doi:10.1038/ja.2017.96.

Progress Toward the Total Synthesis of Hamigerans C and D: A Direct Approach to an Elaborated 6-7-5 Carbocyclic Core

Douglas C. Duquette¹, Dr. Thomas Jensen^{1,2}, and Brian M. Stoltz¹

¹The Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, 1200 East California Boulevard, MC 101-20, Pasadena, California 91125, United States

Abstract

The hamigeran family of natural products has been the target of numerous synthetic efforts due to its biological activity and interesting structural properties. Herein we disclose our efforts toward the synthesis of hamigerans C and D, unique among the initially isolated members due to their 6-7-5 carbocyclic core. Our approach directly targets this tricyclic motif by sequential Negishi and Heck coupling reactions, yielding an advanced intermediate with all necessary carbons and sufficient functionality poised for completion of the synthesis of these two natural products.

Introduction

The hamigeran family of natural products was isolated in 1999 from marine sponge *Hamigera tarangaensis* by Cambie and coworkers (Scheme 1).¹ While all four members of the family possess moderate cytotoxicity in P-388 leukemia cells (IC₅₀ = 31.6, 13.5, 16.0 and 8.0 μ M for hamigerans A, B, C and D, respectively), hamigeran B (**2**) displays 100% *in vitro* inhibition against both Herpes and Polio viruses.

Due to this attractive biological activity and the synthetically challenging 6-6-5 tricyclic structure bearing three consecutive stereocenters on the congested A-ring, hamigeran B has been the target of continual efforts of many synthetic organic chemists. Hamigeran B (along with hamigeran A and several analogs) was first synthesized in 2003 by Nicolaou and coworkers^{2,3} and since has succumbed to a number of syntheses,^{2,3,4,5,6,7,8,9,10,11,12,13,14} including epimeric and formal syntheses. In contrast, at the outset of our studies no synthesis of hamigeran C or hamigeran D (**3** and **4**), which share a unique 6-7-5 tricyclic core, had been reported. This dearth in syntheses is presumably due to the relative scarcity of methods for the synthesis of seven-membered rings as compared to their six-membered counterparts. Hamigeran D has been synthesized recently by Gao and coworkers via a biomimetic skeletal rearrangement from a 6-6-5 tricycle,¹⁵ and thus our strategy of directly targeting the seven-membered ring structure remains unique (*vide infra*).

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

²Current address: Lundbeck Pharmaceuticals, Copenhagen, Denmark

The Stoltz group has developed a method for the asymmetric synthesis of γ -quaternary acylcyclopentenones (Scheme 2, **10**) by a retro-aldol/aldol condensation sequence resulting in a two-carbon contraction of vinylogous esters of cycloheptenone (**13**).¹⁶ The versatility of this intermediate and the excellent scalability of this sequence inspired us to pursue the synthesis of natural products with cores containing densely-functionalized cyclopentanes with quaternary carbon stereocenters, making hamigerans C and D ideal targets.

Results and Discussion

Studies toward the syntheses of hamigerans C and D were initiated during our investigations of the asymmetric synthesis of γ -quaternary acylcyclopentenones.¹⁶ From acylcyclopentene **10**, we were able to assemble the 6-7-5 carbocyclic core of hamigerans C and D (Scheme 2, **18**) in good overall yield over four synthetic steps, featuring two sequential Heck cross-coupling reactions.

Notably lacking on tricycle **18**, however, is substitution on the aromatic C-ring and any functional handle on the benzannulated cycloheptyl B-ring of the natural products. The latter is due to the (*E*)-selectivity in the coupling of **10** with *o*-iodophenol. The resulting olefin must be hydrogenated in order to perform the second Heck coupling, resulting in loss of the otherwise useful olefin moiety. As such, we envisioned a retrosynthetic strategy that would allow us to perform the first cross-coupling reaction to yield a (*Z*)-olefin with an appropriately substituted aromatic ring (Scheme 1, **7**). Such a strategy would allow us to obtain a tricyclic intermediate (**6**) with all of the necessary functional handles in place to advance toward the total syntheses of both of these natural products.

Our first synthetic efforts were directed toward synthesis of Negishi coupling partners **8** and **9**. We desired a differentially protected orcinol derivative, as one phenol group would need to be deprotected and converted to an aryl triflate for the planned Heck cross-coupling, while the other phenol would need to be robustly protected as a methyl ether to be revealed late in the synthesis as the phenol present in hamigeran C or the phenol precursor to the oxazine moiety of Hamigeran D. Synthesis of the *O*-methyl-*O'*-tetrahydro-2-pyranylorcinol **8** was accomplished by dimethylation and mono-demethylation of orcinol (**19**),¹⁷ followed protection as a THP ether in 64% yield over three steps (Scheme 3).

The synthesis of (*Z*)-vinyl iodide **9** was accomplished by a novel sequence from acylcyclopentene **10**. Following the established procedure for ketal protection of the enone,¹⁶ the resulting diene could be selectively ozonolyzed to yield aldehyde **21** in 71% yield over 2 steps. However, yields obtained using the (*Z*)-iodomethylenation procedure reported by Stork¹⁸ proved unreliable upon several repeated experiments (Table 1).

While mass-balance for this transformation was consistent, varying amounts of starting material contaminated by deprotection side-products were reisolated, with yields of the desired vinyl iodide **9** ranging from 33–67% (Table 1, entry 1). Moreover, the 5:1 ratio of *Z*:*E* isomers (as determined by integration of the terminal vinyl proton in the ¹H NMR spectrum) could not be separated by chromatography through silica gel, alumina, or silver nitrate-impregnated silica gel with a wide variety of eluent combinations. As unreacted

starting material appeared to be a major cause of loss for the yield in this transformation, we considered that, given the quaternary center proximal to the aldehyde of **21**, nucleophilic approach by the Wittig reagent was extremely hindered. We reasoned that the Wittig reagent was instead acting as a base, deprotonating the α -position of **21** to yield an unreactive enolate.

Raising the temperature of addition in this reaction to 40 °C and maintaining that temperature throughout the reaction yielded a consistent boost in yield (66–87%) without any detriment to the (*Z*)-selectivity (Table 1, entry 2). Reaction at 30 °C also appeared to be an improvement, though longer reaction times resulted in increased appearance of side products resulting from deprotection of the ketal (entry 3). By increasing the temperature to 50 °C and performing the reaction in benzene, we were able to achieve a good yield, but noted a dramatic decrease in selectivity from 5:1 to 1.3:1 in favor of the (*Z*)-olefin. (entry 4).

Satisfied with the improved yields obtained by performing the reaction with an addition temperature of 40 °C, we sought to improve the (*Z*)-selectivity by modifying the nature of the Wittig reagent. We imagined that modification of the aggregation state of the deprotonated Wittig salt by modifying the counterion or base used or by chelation of the counterion by hexamethylphosphoramide (HMPA), could benefit the selectivity of the reaction.

Unfortunately, substitution of sodium hexamethyldisilylamide (HMDS) for lithium or potassium HMDS (Table 2, entries 2 and 4) both resulted in a decrease in selectivity for the desired (*Z*)-olefin, as did addition of HMPA (entry 3) or the use of sodium *t*-butoxide (entry 5). Moreover, tuning of the sterics of the Wittig reagent itself by substituting (iodomethyl)triphenylphosphonium iodide for the less bulky (bromomethyl)triphenylphosphonium bromide (entry 6) or (iodomethyl)tri-*n*-butylphosphonium iodide (entry 7) also resulted in a decrease in selectivity. To our surprise, the more sterically-hindered (iodomethyl)tri-*o*-tolylphosphonium iodide gave a modest selectivity of 0.9:1 in favor of the undesired (*E*)-olefin (entry 8). As such, we returned to our initial conditions and proceeded with our synthetic studies.

With the proposed Negishi cross-coupling partners in hand, we began investigations of this reaction. We anticipated being able to affect a selective deprotonation of **8** *ortho*- to the two ether substituents with *n*-butyllithium, followed by transmetalation with zinc(II) chloride to afford the Negishi coupling partner for vinyl iodide **9** (Scheme 3).

Gratifyingly, this reaction proceeded in good yield to afford **22** as a single product with respect to the deprotonation of **8**, though still as a mixture of olefin isomers. Removal of both acetal and THP protecting groups proceeded smoothly using Montmorillonite K-10 clay in methanol at 50 °C, the crude product of which was >95% pure by ¹H NMR and could be used directly in the following step. We also found that triflation of the intermediate phenol could be affected using trifluoromethanesulfonyl anhydride and DMAP over a slightly longer reaction time but without loss of yield when compared to the more costly Comins' reagent (*N*-(5-chloro-2-pyridyl)bis(trifluoromethanesulfonimide)), furnishing the desired aryl triflate **7** in 55–62% yield over three steps from Negishi coupling partners **8** and

9. At this point, the minor olefin isomer had also been removed over the course of purification and isolation steps, or had otherwise isomerized to the major isomer due to exposure to light and/or heat.

Aryl triflate **7** was subjected to conditions developed for the model Heck reaction (Scheme 2), modified to be run in a microwave reactor, providing tricycle **6** in 88% yield on 0.12 mmol scale (Scheme 3). However, upon scale up, cleavage of the triflate caused a significant decrease in yield, and a large amount of an undesired acylated side-product was obtained. Further improvements to this Negishi-Heck sequence, including attempts at performing the steps in a one-pot cascade fashion using Pd and Ni catalysis, have been the subject to ongoing research in our group.

With functionalized tricyclic core of hamigerans C and D (**6**) in hand, we sought to pursue the selective functionalization of the disubstituted olefin. A number of dihydroxylation and epoxidation conditions including AD-mix, stoichiometric osmium tetroxide, and *m*-CPBA all failed to provide the desired product. Epoxidation of the disubstituted alkene **6** was finally affected in a mixture of aqueous bleach and dichloromethane with (*R,R*)-Mn(salen) epoxidation catalyst **23** to afford epoxide **24** in 89% yield and a 7.5–10:1 ratio of inconsequential diastereomers (Scheme 3).

Treatment of **24** with catalytic gold(III) chloride in dry acetone¹⁹ directly yielded protected diol **25** in 74% yield as a single diastereomer, although the stereochemistry of the B-ring substituents has eluded definitive determination. The final carbon of hamigerans C and D could be installed by submitting acetal **25** to Wittig conditions, yielding desired compound **26**. Diene **26** represents the most advanced intermediate we have synthesized to date, and completion of the synthesis of hamigerans C and D requires hydrogenation of the diene moiety, deprotection and oxidation of the B-ring diol, bromination and demethylation of the aromatic C-ring. Synthesis of hamigeran C would require installation of the acetoxy group on the B-ring, and completion of hamigeran D would occur by completion of the oxazine D-ring.

In conclusion, we have achieved the synthesis of the 6-7-5 tricyclic core of hamigerans C and D via a unique strategy of directly targeting synthesis of the seven-membered ring via sequential Negishi and Heck cross-coupling reactions. Our targeted synthesis is made catalytically enantioselective by application of our asymmetric alkylation/ring-contraction strategy for synthesis of chiral acylcyclopentenones. Our advanced intermediates possess the requisite carbons and functionality for elaboration to both of these targets, and research into the remaining steps is underway in our group.

For experimental procedures, please see accompanying supporting information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Dedicated to Professor K. C. Nicolaou for his tremendous scientific contributions to the total synthesis of highly complex and biologically important natural products. The authors wish to thank NIH-NIGMS (R01GM080269) for supporting this research. D.C.D. would like to thank the NSF (predoctoral research fellowship, No. DGE-1144469) and Caltech for financial support. Dr. David VanderVelde (Caltech) is thanked for aid in NMR structural determination. Dr. Allen Hong and Dr. Scott Virgil are thanked for helpful discussions.

References

1. Wellington KD, Cambie RC, Rutledge PS, Bergquist PR. Chemistry of Sponges 19 Novel Bioactive Metabolites from *Hamigera tarangaensis*. *J Nat Prod*. 2000; 63:79–85. [PubMed: 10650083]
2. Nicolaou KC, Gray D, Tae J. Total Synthesis of Hamigerans: Part 1. Development of Synthetic Technology for the Construction of Benzannulated Polycyclic Systems by the Intramolecular Trapping of Photogenerated Hydroxy-*o*-quinodimethanes and Synthesis of Key Building Blocks. *Angew Chem Int Ed*. 2001; 40:3675–3678.
3. Nicolaou KC, Gray D, Tae J. Total Synthesis of Hamigerans: Part 2. Implementation of the Intramolecular Diels-Alder Trapping of Photochemically Generated Hydroxy-*o*-quinodimethanes; Strategy and Completion of the Synthesis. *Angew Chem Int Ed*. 2001; 40:3679–3683.
4. Mehta G, Shinde HM. Enantiospecific total synthesis of 6-*epi*-(–)-hamigeran B. Intramolecular Heck reaction in a sterically constrained environment. *Tetrahedron Lett*. 2003; 44:7049–7053.
5. Clive DLJ, Wang J. Synthesis of (±)-Hamigeran B, (–)-Hamigeran B and (±)-1-*epi*-Hamigeran B: Use of Bulky Silyl Groups to Protect a Benzylic Carbon–Oxygen Bond from Hydrogenolysis. *J Org Chem*. 2004; 69:2773–2784. [PubMed: 15074927]
6. Sperry JB, Wright DL. Synthesis of the hamigeran skeleton through an electro-oxidative coupling reaction. *Tetrahedron Lett*. 2005; 46:411–414.
7. Trost BM, Pissot-Soldermann C, Chen I. A Short and Concise Asymmetric Synthesis of Hamigeran B. *Chem Eur J*. 2005; 11:951–959. [PubMed: 15612053]
8. Taber DF, Tian W. Synthesis of (–)-Hamigeran B. *J Org Chem*. 2008; 73:7560–7564. [PubMed: 18771326]
9. Miesch L, Welsch T, Rietsch V, Miesch M. Intramolecular Alkynylogous Mukaiyama Aldol Reaction Starting from Bicyclic Alkanones Tethered to Alkynyl Esters: Formal Total Synthesis of (±)-Hamigeran B. *Chem Eur J*. 2009; 15:4394–4401. [PubMed: 19283820]
10. Cai Z, Harmata M. Studies Directed toward the Synthesis of Hamigeran B: A Catalytic Oxidative Cyclization. *Org Lett*. 2010; 12:5668–5670. [PubMed: 21080691]
11. Lau SYW. Concise and Protective Group-Free Synthesis of (±)-Hamigeran B and (±)-4-Bromohamigeran B. *Org Lett*. 2011; 13:347–349. [PubMed: 21126051]
12. Mukherjee H, McDougal NT, Virgil S, Stoltz BM. A Catalytic, Asymmetric Formal Synthesis of (+)-Hamigeran B. *Org Lett*. 2011; 13:825–827. [PubMed: 21271716]
13. Lin H, et al. Enantioselective Approach to (–)-Hamigeran B and (–)-4-Bromohamigeran B via Catalytic Asymmetric Hydrogenation of Racemic Ketone To Assemble the Chiral Core Framework. *Org Lett*. 2016; 18:1434–1437. [PubMed: 26943071]
14. Jiang B, Li MM, Xing P, Huang ZG. A Concise Formal Synthesis of (–)-Hamigeran B. *Org Lett*. 2013; 15:871–873. [PubMed: 23379694]
15. Li X, Xue D, Wang C, Gao S. Total Synthesis of the Hamigerans. *Angew Chem Int Ed*. 2016; 55:9942–9946.
16. Hong AY, et al. Ring-Contraction Strategy for the Practical, Scalable, Catalytic Asymmetric Synthesis of Versatile γ -Quaternary Acylcyclopentenones. *Angew Chem Int Ed*. 2011; 50:2756–2760.
17. Mirington RN, Feutrell GI. Orcinol Monomethyl Ether. *Org Synth*. 1973; 53:90–93.
18. Stork G, Zhao K. A stereoselective synthesis of (Z)-1-iodo-1-alkenes. *Tetrahedron Lett*. 1989; 30:2173–2174.
19. Balamurugan R, Kothapalli RB, Thota GK. Gold-Catalysed Activation of Epoxides: Application in the Synthesis of Bicyclic Ketals. *Eur J Org Chem*. 2011; 2011:1557–1569.

Scheme 1.

Structures of first four identified members of the hamigeran family of natural products, and retrosynthetic analysis of hamigerans C and D.

Scheme 2.

Previously achieved methods for the synthesis of chiral acylcyclopentenones and elaboration to the carbocyclic core of Hamigerans C and D. a) Pd₂(pmdba)₃ (1.25 mol %), (S)-*t*-BuPHOX (**12**, 3.13 mol %), toluene, 30 °C, 93% yield, 87% ee. b) LiAlH₄, Et₂O, 0 °C; 10% aq HCl, 81%. c) LiOH, CF₃CH₂OH, THF, 60 °C, 91%. d) Pd(OAc)₂ (10 mol %), tetrabutylammonium iodide (TBAI), NEt₃, *o*-iodophenol, DMF, 100 °C, 90%. e) Rh(PPh₃)₃Cl (10 mol %), H₂ (1 atm), toluene, 23 °C, 95%. f) Comins' Reagent, 4-dimethylaminopyridine (DMAP), CH₂Cl₂, 78%. g) Herrmann-Beller catalyst (**17**, 10 mol %), Bu₄NOAc, DMA, 115 °C, 77%.

Scheme 3.

Synthetic efforts toward the total synthesis of hamigerans C and D. a) Me_2SO_4 , K_2CO_3 , refluxing acetone, 95%. b) NaH , EtSH , refluxing DMF, 84%. c) dihydropyran (DHP), p -toluenesulfonic acid, THF, 81%. d) neopentyl glycol, pyridinium p -toluenesulfonate (PPTS), toluene, 140°C , 73%. e) O_3 , CH_2Cl_2 , -78°C ; PPh_3 , $-78^\circ\text{C} \rightarrow 23^\circ\text{C}$, 97%. f) $\text{ICH}_2\text{PPh}_3\text{I}$, sodium bis(trimethylsilyl)amide (NaHMDS), THF, 40°C 66–87%, 5:1 *Z:E* ratio by ^1H NMR. g) $n\text{-BuLi}$ (2.5 M in hexanes), THF, $0^\circ\text{C} \rightarrow 23^\circ\text{C}$; ZnCl_2 ; vinyl iodide **9**, $\text{PdCl}_2(\text{PPh}_3)_2$, 85–95%. h) Montmorillonite K-10, MeOH, 50°C . i) TiF_2O , DMAP, CH_2Cl_2 , 23°C , 65% yield over two steps. j) Herrmann–Beller catalyst (**17**, 10 mol %), Bu_4NOAc , DMA, 120°C (μwaves), 88% (0.12 mmol scale) or 54% (2.0 mmol scale). k) $(R,R)\text{-Mn(salen)}$ catalyst (**23**, 15 mol %), NaOCl (aq, 0.54 M, pH 11.2), CH_2Cl_2 , 23°C , 89%, 7.5–10:1 d.r. by ^1H NMR analysis. l) AuCl_3 (20 mol %), acetone, 23°C , 74%, single diastereomer by ^1H NMR analysis. m) MePPh_3Br , $n\text{-BuLi}$ (2.06 M in hexanes), benzene, $23^\circ\text{C} \rightarrow 90^\circ\text{C}$, 67%.

Table 1

Temperature and solvent optimization studies for yield and selectivity in olefination reaction.

Entry	Solvent	Temperature	Z:E Ratio	Yield
1	THF	0 → 50 °C	5:1	33–67%
2	THF	40 °C	5:1	66–87%
3	THF	30 °C	5:1	63%
4	Benzene	50 °C	1.3:1	85%

Table 2

Phosphonium ylide and base optimization studies for selectivity in olefination reaction.

Entry	R	Base	X	Z:E Ratio
1	Ph	NaHMDS	I	5:1
2	Ph	LiHMDS	I	2.5:1
3	Ph	LiHMDS (5:1 THF:HMPA)	I	1.5:1
4	Ph	KHMDS	I	3.3:1
5	Ph	NaOt-Bu	I	2.3:1
6	Ph	NaHMDS	Br	1.4:1
7	<i>n</i> -Bu	NaHMDS	I	2.0:1
8	<i>o</i> -Tol	NaHMDS	I	0.9:1