BNOS 2017 Abstracts: Oral Presentations

LOCAL DELIVERY OF IRINOTECAN TO RECURRENT GLIOBLASTOMA AT REOPERATION OFFERS A SAFE THERAPEUTIC ADVANTAGE OVER SYSTEMIC DELIVERY.

Prof Garth Cruickshank, Mr Desire Ngoga, Dr Allah Detta, Prof Andy Lewis, Dr Rebecca Holden, Mr Olufawe Fayaye; University of Birmingham and Queen Elizabeth Hospital

Direct drug delivery to a brain tumours offers the surety of access, together with the potential to realise at least therapeutic concentrations. With previously used systemic drugs toxicity and removal mechanisms are avoided. With new formulations drug activation and duration can be explored for low local toxicity, and tumour impact with sequential imaging. In this early study Irinotecan a drug with known efficacy against glioma but with limiting systemic toxicity has been incorporated into biodegradable hydrogel microspheres and injected into the post-surgical cavity wall in patients with recurrent glioblastoma. 10 patients with focally recurrent GBM were recruited with 9 undergoing injection with up to 3mls of microbeads in alginate suspension in up to 60 ~ 8mm injection tracks after surgical resection (100mg). Patients were assessed for immediate (72hrs) toxicity with clinical assessment and imaging. Pharmacokinetic determination of Irinotecan and SN 38 active metabolite allowed confirmation of activation and comparator with known plasma toxicity profiles. Follow-up was planned for a minimum of 6/12 for SAE but continued to allow overall survival estimation. No overt SAE were determined for this group of patients with imaging and steroid use indicating less local swelling and wound healing issues than have been demonstrated for Carmustine wafers despite early offloading. Pharmacokinetic measures (SN38 plasma curves) indicate a much higher activation of Irinotecan (>90%) than expected but shorter period of exposure. Survival curves suggest clinical benefit with 4/9 patient living longer than 8 months warranting further exploration of this safe approach

RISK FACTORS FOR DEVELOPING POST-OPERATIVE SEIZURES FOLLOWING MENINGIOMA RESECTION

Ms Arousa Ali, Mr Ananyo Bagchi, Dr Samantha Mills, Dr David Giraldo, Mr Emmanuel Chavredakis, Mr Andrew Brodbelt, Mr Michael Jenkinson; University of Liverpool

BACKGROUND: Meningiomas are benign primary tumours treated by surgical resection. Following surgery patients may develop postoperative seizures. The aim of this study was to investigate the risk factors associated with developing postoperative seizures and determine whether prophylactic antiepileptic drugs (AED) should be prescribed. **METHODS:** Retrospective analysis of patients undergoing surgical resection of meningioma between 2010–2015. The following clinical factors were assessed: tumour location, grade, Simpson resection, post-operative haemorrhage/infection and use of AED prophylaxis. MRI parameters were assessed including: tumour volume, adjacent parenchymal T2 signal change, midline shift. Logistical regression was performed. **RESULTS:** 285 patients were identified with a median age of 60 years (range: 16–83 years). 68 (24%) had pre-operative seizures of which 62 were on AEDs and post-operatively 48 (17%) had seizures. 217 (76%) were seizure naïve, 19 were given prophylactic AEDs and post-operatively 5 had seizures. Of the 198 patients who did not receive prophylactic AED, 43 had post-operative seizures (22%). Prophylactic AED did not reduce the risk of seizures compared to no use (OR = 1.85, p = 0.260). Clinical risk factors for developing post-operative seizures were non-skull base location (n = 207, OR = 0.35, p = 0.010) and presence of post-operative haemorrhage/infection (n = 25, OR = 1.69, p = 0.002). MRI factors associated with developing post-operative seizures were presence of T2 signal intensity change greater than 66% of tumour volume (n = 81, OR = 4.71, p = 0.000). **CONCLUSION:** Risk factors associated with postoperative seizure development were the non-skull base tumour location and post-operative haemorrhage/infection. The role of prophylactic AED could not be determined in this study due to insufficient sample size. Randomized control trials are required.

DIFFUSION-WEIGHTED MRI MAY SUPPORT THE IDENTIFICATION OF NON-ENHANCING IDH WILDTYPE ASTROCYTOMA

Dr Stefanie Thust, Dr Sotirios Bisdas, Dr Jeremy Rees, Dr Harpreet Hyare, Prof Sebastian Brandner, Dr Sara Hassnein, Prof H Rolf Jager, Dr Almas Siddiqui, Dr Laura Mancini; National Hospital for Neurology and Neurosurgery, UCLH

AIM: To determine if apparent diffusion coefficient (ADC) measurements permit the distinction of non-enhancing IDH mutant (IDHmut) from IDH

wildtype (IDHwt) astrocytoma. **BACKGROUND:** Lack of Gadolinium contrast enhancement represents a typical feature of low grade astrocytomas, but is increasingly considered an unreliable predictor of non-aggressive behaviour. The latest World Health Organisation (WHO) classification recognises a distinct subgroup of rapidly progressive gliomas (IDHwt astrocytoma) without histopathological grade IV characteristics. On molecular grounds, IDHwt astrocytomas constitute malignant neoplasms and are suspected to represent early glioblastomas (GBM). Their histology may be matched by initially innocuous imaging appearances, which could lead to misdiagnosis if a 'watch and wait' approach is adopted. **METHODS:** From a cohort of 58 WHO grade II/III astrocytomas (34 IDHwt/24 IDHmut), 22 non-enhancing tumours were selected. Manually obtained T2w volumetric segmentation masks were spatially registered (FSL) to ADC maps derived from standard DWI (b0-b1000). A quantitative analysis of diffusion values was performed blinded to molecular data and compared (Student's t-test) for both groups. **RESULTS:** 13 IDHwt (6 WHO II/7 WHO III) and 9 IDHmut (6 WHO II/3 WHO III) astrocytomas were analysed. Maximum, mean and minimum ADC values for the IDHwt group were significantly lower (p < 0.01) than for the IDHmut group. **CONCLUSION:** ADC values derived from clinical DWI may reveal the aggressive nature of non-enhancing IDHwt astrocytomas, whereby mean ADC measurements are obtainable on most hospital imaging systems. Further study is required to determine, if a threshold value can be defined and to consider potential influences from WHO grade.

HIGH-RESOLUTION RNA-SEQ PROFILING OF GLIOBLASTOMA RESIDUAL INVASIVE CELLS ISOLATED BY A NOVEL 5-ALA FLUORESCENCE BASED METHOD

Mr Jonathan Rowlinson, Dr Maria de los Angeles Estevez-Cebrero, Dr Anbarasu Lourdasamy, Dr Ruman Rahman, Dr Stuart Smith; The University of Nottingham

INTRODUCTION: Malignant brain cancer Glioblastoma (GBM) kills 3500 people per year in the UK with an average survival of 6 months. The intra-tumour heterogeneity of GBM has meant tailored therapies based on the genetic mutations within the cancer cells have so far failed to replicate in vitro promise in clinical trials. Previous research has focused on the core of the tumour; however this is removed by surgery and doesn't reflect the residual cancer cells left in the patient's brain. The aim of this study is to generate new potential targets from looking at the expression of those residual cancer cells left behind after surgery that ultimately lead to tumour recurrence. **METHOD:** 5-aminolevulinic acid (5-ALA) is a clinically used drug that makes cancer cells glow (fluoresce) pink, allowing more complete removal of brain tumours. 10 Tumour samples were dissociated and fluorescent activated cell sorting (FACS) was used to separate the residual fluorescent cancer cells (1%) from normal brain at the tumour's edge. Samples from 3 distinct tumour regions, core, rim and invasive margin without FACS sorting from the same patients were also included for comparison to the FACS samples. Gene expression was analysed by RNA-seq and validated by qPCR. **RESULTS:** High-resolution RNA-seq profiles were generated for distinct tumour regions and the fluorescently isolated invasive cell population, allowing potential targetable abnormalities to be identified and validated. **CONCLUSION:** This study shows that pure tumour cells can be separated from normal cells using fluorescence generated from tumour metabolism of 5-ALA. Further optimisation of FACS is required to increase RNA yield from FACS sorted samples. Our approach gives hope that we can interrogate the true residual disease, and for the first time gain insight into the source of tumour recurrence.

PSYCHOMETRIC EVALUATION OF THE CAREGIVER NEEDS SCREEN IN NEURO-ONCOLOGY FAMILY CAREGIVERS

Dr Florian Boele, Dr Lauren Terhorst, Ms Jennifer Prince, Dr Heidi Donovan, Mr Jason Weimer, Dr Frank Lieberman, Dr Jan Drappatz, Dr Paula Sherwood; University of Leeds

INTRODUCTION: The informal care demands of brain tumour patients include various issues associated with neurological and cognitive symptoms. Caregiver needs questionnaires developed for cancer populations do not include these disease-specific symptoms. Therefore, we have developed the neuro-oncology Caregiver Needs Screen (CNS) and evaluated its psychometric properties. **METHODS:** In-depth qualitative interviews with caregivers were used to generate items, which were reviewed by experts to add or collapse items as necessary. The resulting 32-item instrument was tested in a sample of 122 neuro-oncology caregivers, along measures of depression (Center for Epidemiological Studies – Depression scale), anxiety (Profile of Mood States – Anxiety), caregiver burden (Caregiver Reaction Assessment) and mastery (Mastery Scale). Exploratory factor analysis was used to determine internal structure. Internal consistency reliability and construct validity were assessed. **RESULTS:** Six subscales were identified (neurologic symptoms; oncologic symptoms; personal communication; communicating with health care providers; resources;

caregiver health). Internal consistency for the subscales ranged between $\alpha = .653$ and $\alpha = .886$. A floor effect was found for the personal communication subscale only. Convergent validity was verified by moderate to strong correlations between measures of caregiver wellbeing and CNS scale scores. **DISCUSSION:** Findings provide preliminary evidence of reliability and validity for the CNS as a measure of neuro-oncology caregiver distress as a result of common care issues. This instrument can prove useful in research setting as well as in clinical practice when assessing caregivers' needs for supportive care.

UNDERSTANDING MISSED OPPORTUNITIES FOR MORE TIMELY DIAGNOSIS OF BRAIN CANCER – WHAT CAN WE LEARN FROM THE BRACED QUALITATIVE INTERVIEW STUDY WITH ADULT GLIOMA PATIENTS?

Ms Clarissa Penfold, Dr Alexis J Joannides, Dr William Sage, Mr Andrew Brodbelt, Mr Michael D Jenkinson, Mr Stephen Price, Dr Colin Watts, Mrs Joyce Bell, Mrs Margaret Johnson, Prof Willie Hamilton, Dr Fiona M Walter; University of Cambridge

BACKGROUND: BRACED (The BRAin Cancer Early Detection study) is a qualitative study of patient perspectives on factors affecting timely diagnosis of primary brain tumours. Patient, doctor and health-care system-related factors can all contribute towards prolonged time to cancer diagnosis, impacting on patient experience, morbidity and mortality. **AIM:** To develop a richer understanding of adult glioma patients' experiences and attributions of symptoms, and their routes to diagnosis. **METHODS:** In-depth qualitative interviews with 20 patients with early analysis completed; a second phase of interviews is underway with another 20 patients, for a detailed exploration of early emerging themes. **RESULTS:** First phase participants were 14 men and 6 women, aged 22 - 76 (median 63) years. Participants reported varied routes and times to diagnosis, often involving more than one healthcare professional. Factors contributing to prolonged times to diagnosis were explored in analysis of patient narratives relating to initial assessment, diagnostic test performance and interpretation, and follow-up and coordination. There was evidence of challenges in presenting multiple and subtle symptoms in the context of a routine GP appointment. Some participants felt that their diagnosis may have been delayed in primary care by non-urgent referral to specialist care or 'false' reassurance following normal test results. Prolonged time to diagnosis was also suspected where: no CT or MRI scan was performed on emergency presentation despite being suggested by a clinician; a neurologist was called but did not see the patient prior to discharge; delays in reporting scan results occurred; or there was an apparent absence of 'safety-netting' advice to support patients in their ongoing appraisal of symptoms. **DISCUSSION:** Early findings identify a number of factors that may contribute to prolonged time to diagnosis after presentation, which could be minimised by improving guidance for both patients and clinicians.

INTEGRATED SPECT-MRI FOR ENHANCED STRATIFICATION AND EARLY TREATMENT RESPONSE ASSESSMENT IN GLIOMA

Prof Susan Short, Prof Brian Hutton, Dr Luisa Ottobri, Dr Irene De Francesco, Prof Carlo Fiorini; University of Leeds

BACKGROUND: Personalised treatment based on individual tumour biology tracked over time is an important goal to improve outcomes in glioma. Imaging biomarkers have the potential to permit individualization by defining heterogeneous biology in 3D space and tracking response. To optimize application of this approach, biological readouts need to be aligned with high quality anatomical information. The INSERT consortium seeks to address this unmet need using simultaneous multi-tracer SPECT and MRI. **AIM:** Design and build a multi-modality imaging tool to allow simultaneous SPECT-MR assessment and read-out of multiple tumour biology biomarkers in patients with glioma. **METHODS:** A fully MRI compatible SPECT imaging system was designed for i) pre-clinical and ii) clinical applications. In both systems the SPECT imaging frame is inserted to the bore of a standard MRI scanner and can be moved between scanners. Novel components of the system include: unique MR compatible slit-slat collimator design, hand-built in precision-cut tungsten, high-resolution detection modules with customized SiPM readout, housed in a robust, 3D-printed unit that can be aligned with MR imaging isocentre to scan the whole brain. To develop optimized applications of this system, pre-clinical experiments were undertaken to assess the utility of novel SPECT tracers with MR imaging in orthotopic (U251) glioma models. **RESULTS:** The pre-clinical and clinical INSERT systems have been constructed and early feasibility testing is ongoing. In orthotopic models, SPECT combined with MRI is a promising approach to early response assessment that is highly correlated with bioluminescence imaging, autoradiography, PET and histology. A protocol for early assessment of the same biomarkers in glioma patients is agreed and will recruit in 2017. **CONCLUSION:** There remains a significant unmet need for better biological readouts of tumour response in glioma. The application of multi-modality imaging with the advanced technology and practical advantages of INSERT delivers novel opportunities in this setting.

PROGNOSTIC STRATIFICATION OF IDH WILD TYPE LOWER GRADE GLIOMAS

Prof Ho Keung Ng; The Chinese University of Hong Kong

IDH wild-type astrocytoma is described as a provisional entity within the new WHO classification. Some groups believe that IDH wild-type lower-grade gliomas, when interrogated for other biomarkers, will mostly turn out to be glioblastomas. We hypothesize that not all IDH wild-type lower-grade gliomas have very poor outcomes and the group can be sub-stratified prognostically. 718 adult WHO Grade II and III gliomas from our hospitals were reviewed and tested for IDH1/2 mutations. 166 IDH wild-type cases were identified for further studies and tested for EGFR and MYB amplification, mutation of histone H3F3A, TERT promoter and BRAF. EGFR amplification, BRAF and H3F3A mutations were observed in 13.3%, 8.6% and 9.1% respectively, in a mutually exclusive pattern in IDH wild-type lower-grade gliomas. TERTp mutations were detected in 26.4% of cases. Favorable outcome was observed in tumors with oligodendroglia phenotype, grade II histology and total resection. Independent adverse prognostic values of older age, non-total resection, grade III histology, EGFR amplification and H3F3A mutation were confirmed by multivariate analysis. Tumors were further classified into "molecularly" high grade (harboring EGFR, H3F3A or TERTp) (median OS = 1.2 yrs) and lower grade (lacking all of the three) (median OS = 7.6 yrs) with independent prognostic relevance. The most favorable survival was noted in molecularly lower-grade gliomas with MYB amplification. In summary, adult IDH wild-type lower-grade gliomas are prognostically heterogeneous and do not have uniformly poor prognosis. Clinical information and additional markers including MYB, EGFR, TERTp and H3F3A should be examined to delineate bad and good prognostic groups.

MINIMISING LENGTH OF STAY FOLLOWING BRAIN TUMOUR SURGERY: BENEFITS TO PATIENTS, PROVIDERS AND THE NHS

Mr Paul Grundy; University Hospital Southampton

INTRODUCTION: Length of stay (LOS) following surgery is highly variable between providers. Techniques can be employed to minimise LOS to the benefit of patients, providers and the NHS. Data is now available to benchmark and to provide an opportunity for aspiring to best practice. **METHODS:** We introduced measures to reduce LOS. We explored data from Hospital Episode Statistics published for providers by the National Neurosurgical Audit Program (NNAP) and Get It Right First Time (GIRFT) in 2016/7. **RESULTS:** In our unit the most frequent pathway for surgery for brain tumours was as elective admissions: 75% of patients with high-grade gliomas; 81% with metastases and 85% with benign tumours were admitted for surgery in an elective manner. Median LOS for intrinsic tumours was 1 day (versus national 6 days, NNAP). Mean LOS (vs national) was 2.5 (6.4) days for HGG; 2.9 (6.5) days for metastases; 4.7 (9.2) days for benign tumours; all the lowest in UK. Despite this, readmission rates were within the normal range at 20% (18%) for HGG; 21% (20%) for metastases; 15% (14%) for benign tumours. Re-operation rates over 1 year were also in normal range: 11% (12%) HGG; 9% (11%) for metastases; 12% (11%) for benign tumours. Elective cranial mortality was 0.87 (0.94). **DISCUSSION:** Adopting several strategies to minimise LOS has proven very successful and resulted in excellent benchmarked data. This has proven popular with patients, has improved efficiency at our trust, reduced cost, cancellations and waiting times and has more widespread implications across the NHS. Short stay neurosurgery for brain tumours is safe and does not significantly increase readmission, re-operation or mortality rates. Techniques employed to deliver this will be discussed. **CONCLUSION:** All providers of neurosurgery should use national data to improve practice in the future, including engaging with peers and utilising strategies proven to be successful elsewhere.

A HUMAN IPS CELL-BASED MODEL OF MEDULLOBLASTOMA DEMONSTRATES CO-OPERATIVITY BETWEEN SHH SIGNALLING AND MUTATION IN AN EPIGENETIC MODIFIER

Mr Jignesh Tailor, Dr Miller Huang, Prof Austin Smith, Prof William Weiss; St George's University Hospital, London

OBJECTIVE: Medulloblastoma (MB), the most common malignant pediatric brain tumor, is comprised of four subgroups (WNT, SHH, Group 3, Group 4). While recurrent single nucleotide variations is rare in MB, mutation of epigenetic modifiers is frequent, suggesting a mechanism for tumorigenesis. Genetically engineered mouse models (GEMMs) recapitulate some aspects of MB, but fail to replicate the epigenetics characteristic of human MB. Therefore, we aimed to develop the first human stem cell based model of MB using iPSC derived from Gorlin syndrome (GS), a disease that predisposes patients to SHH MB due to heterozygosity for PTCH1. **Methods:** We differentiated iPSC from healthy (WT) or GS patients toward hindbrain neuroepithelial stem (NES) cells, a presumed cell of origin of MB. We then electroporated the NES cells with RNA-guided FokI Nucleases (RFN, CRISPR/dCas9-FokI) targeting GSE1 (formerly KIAA0182), a gene encoding an epigenetic modifier