It may be proper to mention, that, since 1843, there has been published at Berlin a journal devoted to the subject of the diseases of children. The first volume contains various instructive essays and memoirs.*

ART. II.—*A Practical Treatise on the Domestic Management and most important Diseases of Advanced Life. With an Appendix, containing a series of Cases illustrative of a New and Successful Mode of treating Lumbago and other forms of Chronic Rheumatism, Sciatica, and other Neuralgic Affections, and certain forms of Paralysis.* By GEORGE E. DAY, M.D., Fellow of the Royal College of Physicians, and Physician to the Western General Dispensary. London, 1849. 8vo, pp. 342.

WHILE the characters of diseases are much modified by the state of the human frame in infancy and childhood, they receive an influence not less real and considerable from the progress of advancing years and the approach and presence of old age. We have seen, that not only do particular forms of disease prevail most frequently in the early period of life, but the diseases which do prevail at that period receive, as it were, a particular impression from states of the vascular and nervous system, which are predominant during the years of infancy and childhood. Again, certain forms of disease prevail more after puberty and during adolescence than previous to this period; and certain textures and organs are then unquestionably more liable to be affected by the morbid element than others, and than at either previous or subsequent periods of life. During the period of manhood, or after the body has attained its complete growth, diseases assume a character not less peculiar to that time, and different in certain features from those which prevail at other periods of life. The morbid element evinces its effects in different textures and organs, and in different and characteristic modes. At length, old age approaches,—in some early, in others at later periods, but to all, though at different times, with inevitable certainty; and in its train come not only maladies of a peculiar character, but all diseases receive from the state of the system, then either approaching or established, a modification which is so characteristic, that it has long attracted, in a general manner, the observation of physicians and the attention of philosophers.

* Journal fur Kinderkrankheiten, herausgegeben von Behrend und Hildebrand. I. Band, 6 Heften. Berlin, 1843.

It was with the elder pathological writers, as Boerhaave, Gaubius, Nietzki, and Caldani, a favourite doctrine, that in the human frame there were implanted seminal principles of disease (*seminia morbi*), which were at all times present in the fluids and solids of the body, and the slow and imperceptible action of which, effected in various organs those changes, which rendered these organs progressively less fit for performing their functions, and at length totally unfit to perform them. It was understood, of course, that the power and influence of these morbid principles were more or less aided and augmented in different individuals, by external morbid agents, and external powers of various kinds; by climate, the weather, and physical agents in general, different forms of food and drink, occupations, the operation of the passions, and all similar causes. To these, however, was assigned only a concurrent and accessory power, not one quite independent; and it was always believed and taught, that the primary causes of disease, whether dynamic or organic, were seated, fixed, and inherent in the frame of the human body itself.

This idea has never been wholly abandoned, either by physicians, by philosophers, or even by the vulgar. It may have assumed a new aspect; it may have been designated by a new denomination; but the object signified has been always more or less present to the imagination. Has this idea a real existence, or rather, it ought to be said, has the thing signified, the internal morbid principle, a real existence? It is perhaps idle to enter at present into the inquiry, which is more a metaphysical and psychological, than a pathological one. Neither, were we inclined, do our limits permit to enter into the discussion of such a subject. We merely observe, that various writers on pathology, since the beginning of the present century, have more or less distinctly reproduced the idea and doctrine of their predecessors. Kurt Sprengel, for instance, Professor at Halle, one of the great authorities on these points, thirty years ago delivers the doctrine in a form sufficiently distinct and intelligible:—"The internal condition or diathesis, or susceptibility we call that disposition of the body about to become diseased, by which it results, that either actual disease takes place, or may probably take place. First, therefore, is the capacity to be affected by disease universal and natural to the whole human race, and which must be taken into consideration, however active external agents may be. Poison, the sword, and fire-arms or artillery, inflict diseases on those who are in apparently perfect health, or, at most, not tending to disease. But the susceptibility to be affected by disease is less entitled to consideration, than that internal condition, which implies a certain individual proclivity to diseases.*

* Curtii Sprengel Institutiones Medicæ, Pathologiæ Generalis, Volumen Primum, § 24. Diathesis seu interna Conditio, p. 40. Mediolani, 1817, 8vo.

About ten years after the period when these sentences were the second time published to the medical world of Germany, Philip Charles Hartmann, Professor in the University of Vienna, expressed his views of the same subject, in the following manner.

“§ 595.—If the human body were adequate either to repel or to overcome all those agents, which threaten its health, it would give access neither to hurtful force nor to disease. But as this immunity is not enjoyed by beings living on the earth, which possess only limited powers of resistance and subjugation, the result is, that these bodies are able to subject external agents only of a certain degree and kind. The living body, accordingly, when exposed to the influence of external agents, necessarily yielding to this influence, may be made to deviate from the normal condition, and be subjected to disease, in consequence of the efforts which it makes to resist the influence of these agents.

“§ 596. That condition of the human body, by virtue of which its vital properties may be by inimical agents excited to abnormal action, is named the *Disposition to Disease*; and those external agents, which produce this effect, are denominated *hurtful powers*. Further, man possesses a twofold disposition to disease; one *general*, the other *special*. The general disposition to disease is that which affects the whole human race, since it is rooted in the nature of the human frame itself. The special disposition is that which is proper to certain individuals, and derives its origin from particular relations and modes, by which life is regulated in each individual.”*

From other sources, especially the *Elements of Analytical Pathology*, by Bufalini,† similar passages might be easily adduced, tending to show, that pathological speculators and physicians have at all times assumed the existence of some internal morbid principle or condition in the human body. This, however, it is unnecessary further to do. It is sufficient to mention the circumstance, to show that this morbid principle or element, at all times existing, and probably operating more or less steadily, however imperceptibly, constitutes that essence or condition upon which, extending through all the periods of life, external causes and agents act in inducing diseases, and which at different periods of life being acted upon, or itself re-acting in different modes, impresses on the diseases of each period their peculiar characters.

* Ph. Caroli Hartmann, M. D., et P. P. O. in *Scientiarum Universitate Vindobonensi, Theoria Morbi, Seu Pathologia Generalis Prælectionibus Academicis Accommodata, Editio Altera Emendatior*. Vindobonæ, 1828.

† *Fondamenti di Patologia Analitica* di Maurizio Bufalini Cesenate, Med. in Patria, Già Assistente et Lettore Straordinario alla Cattedra di Clinica Medica, nell' Università di Bologna, Attuale Prof. O. de Medicina Teorico-Pratica, Nell' Università di Urbino, &c. Edizione Terza, Tomo i. Pesaro, 1828. Tomo ii., 1830.

It has been seen that, though the list of diseases to which infant and early life is exposed, is very large and serious, yet some diseases there are, which more particularly predominate at that period of existence. Thus, affections of the nervous system predominate most in infancy, and next to these, disorders of the alimentary canal. Bichat has observed that diseases of the thoracic organs, that is, the organs of respiration and circulation, predominate most after puberty, and between that period and the time that marks the termination of manhood; that in old age, disorders of the intestinal tube and assistant chylopoietic viscera, with those of the urinary organs, begin to predominate; and that, as life continues to advance, again various affections of the nervous system begin to make their appearance. This, of course, is merely such a general view as might occur to a man of observation placed in a large hospital, and seeing daily and weekly many persons labouring under different forms of disease.

Any inquiry relating to the diseases of old age, involves, at least, two questions. The *first* is, of any given number of individuals coming into the world at the same time, what proportion shall live, how many or how few shall be permitted, to see and endure what is denominated old age? The *second* is, what are the maladies most likely to affect that period of life, and what is the comparative degree of their influence in shortening its duration?

As to the first question, the documents and facts collected by the Reports of the Registrar-General, as well as the disquisitions on what is called the mean duration of human life, afford the means of throwing upon it some light, and understanding it more accurately than has hitherto been done. Of 1000 persons, male and female, born in this country, at any given time, at the age of eleven years 702 are alive, and at the age of twelve years, 699; showing that in the course of twelve years, 300 of the thousand have been cut off. At the age of thirty-one years, 598 are left, showing that another hundred persons have in the course of nineteen years been added to the three hundred swept away by disease. At about the age of forty-five years and six months, five hundred are left; showing, that in the course of fourteen years and a half longer, one-half of the whole number of persons born have been carried to the tomb. At the end of sixty-six years and about three months, ($66\frac{1}{4}$) the number of survivors is reduced to three hundred (300). At $72\frac{1}{2}$ years, only two hundred (200) are alive. At seventy-nine and a half, only 100 are alive; and at the age of ninety-six years, all the males have departed, and the sole representative of the thousand persons is one decrepit old woman, deaf, generally blind, and enduring with great difficulty the weight of her advanced age and infirmities.

Between fifty-two years of age and sixty, which may be regard-

ed as the beginning of old age, between 451 and 380 persons are left to look forward to the hope of adding from ten to fifteen, or at most twenty years, to their measure of life. At fifty-two and fifty-three, the probable future duration of life is about nineteen years. At sixty, it is fourteen years for both sexes; and as life advances to three-score and ten, and beyond that period, it rapidly descends to nine, eight, six, and five years. It is further to be observed, that all the survivors cannot live these periods. It is only a certain proportion of them, who have, among others, a chance of, at this time of life, adding five, six, or eight years to their previous existence.*

For giving an answer to the second question, not many or extensive data are furnished. There are, nevertheless, some means, especially those furnished by the weekly tables of mortality for London, in which the ages below five, between fifteen and sixty, and above sixty, have their respective numbers of deaths by different diseases specified. From a tabular view of these for the years 1845 and 1846, the following results may be deduced.

The mortality by diseases of the nervous system, which is very large below the fourth year, undergoes a great and striking diminution above that age, till the fifth year, when it is slightly increased. It continues, however, very low till the 35th year, when it begins to be increased, and continues increasing till the seventieth year, after which it is greatly diminished.

The mortality by diseases of the organs of respiration, is in the aggregate of the first five years of life, very little, that is, about one-twentieth, below that by diseases of the nervous system. But it is differently distributed during the four first years. It is during the first year not quite one-half of the whole mortality of the first five years; and in the ensuing two years it is quite as much as in the first year. In the fifth year it is greater than either in the third or fourth year. Between five and ten it is greatly reduced; but at fifteen it is again increased. Between the age of fifteen, and onwards to that of sixty years, the mortality from this source is at its greatest height; and this mortality is chiefly caused by pulmonary consumption in the first fifteen years, and by different forms of pneumonia and *bronchitis* during the last thirty years of the period.* This class of diseases forms indeed the largest and most copious stream of mortality to the human race; and it is so much more destructive, that it carries off the largest proportion of human beings during the period of adolescence and manhood.

Zymotic diseases, or those the origin of which depends on the presence and operation of a poisonous agent, either in the atmo-

* Sixth Annual Report of the Registrar-General, p. 520.

† Under five years, 2,305; above five years, 5,574; between fifteen and sixty years, 4,241. Between sixty and ninety-five, 821. This is for males only.

sphere or evolved from the earth, but most usually the former, form the next great destroyers of the human race. These prevail most numerous, and cause the largest amount of mortality, in children below four years. An increase takes place in the fifth year; and after this, a considerable diminution till the twentieth year, when an increase is again observed; after this there is a slight diminution; but, nevertheless, the mortality is considerable. The mortality between the ages of ten and sixty is caused chiefly by typhus fever.

The diseases most prevalent and most mortal after the age of fifty and fifty-five, are disorders of the nervous system, as apoplexy, palsy, and similar affections of the brain; disorders of the organs of respiration, as bronchitis, and different forms of pneumonia; disorders of the organs of circulation, as the heart and arteries, and occasionally the veins; certain disorders of the secreting glands, as the liver and kidneys; disorders of the digestive organs; and the not very well defined head and title of old age.

What, it may be asked, is death by old age? In the Report of the Metropolitan mortality for 1845, 1,111 males and 1,848 females are represented to have been the prey of this anomalous and undefined agent. In that for 1846, 842 males and 1,399 females are stated to have died from this cause. In the one year, therefore, 2,959 persons, and in the other, only 2,241, died of old age, making a total difference of 718 persons. The difference is great, and not to be accounted for by the natural and regular workings of old age upon the human frame. The difference depends, we are satisfied, upon improvement in diagnosis; upon more care being taken to ascertain the exact causes of death. The reason, why we think so, is to be found in the fact, that there is not a proportionate difference in the total mortality for these two years.

These deaths take place in persons aged, or supposed to be aged, above sixty; and the great majority of deaths take place between the ages of sixty-five and eighty-five.

The truth is, that of these numbers, approaching in the year 1845 to three thousand persons, not probably one hundred really died of old age, strictly speaking. Many of those who are believed and reported to die of old age, die of positive diseases, which doubtless acquire a peculiar character at that period of life. Thus, pneumonia and bronchitis destroy not a few; pneumonia especially in a latent form, and *bronchitis* proceeding to emphysema and different forms of asthma. It has been ascertained repeatedly, that aged persons in public institutions, hospitals, and asylums for the aged, supposed to die of advanced age, have actually been destroyed by pneumonia in a latent form, sometimes advancing slowly, sometimes more rapidly, but always present so

as to destroy the structure and functions of the lungs as a breathing organ.

Again, diseases of the heart, sometimes alone, sometimes with disease of the lungs, destroy many of those supposed and stated to die of old age. Ossification and coalescence of the valves, contraction of the apertures, ossification of the coronary arteries, atrophy and flaccidity of the heart, *steatosis* or fatty transformation of the heart, all tend to produce death in the aged, and all have produced death, which by ignorant or indifferent reporters has been ascribed to old age.

About the age of thirty-seven, or previous to that period in some, but in general about and after that period, the arteries begin in many persons to be affected by the waxy, steatomatous or fatty deposit. This is followed by, or accompanied with, the calcareous deposit; and in this manner, and by these newly-secreted substances, the arteries of the whole body, but especially those of the chest, abdomen, and brain, may be so penetrated, transformed, and rendered brittle, that they can no longer perform their functions as transmissible tubes. As these changes proceed, increasing as the individuals advance in life, they give rise, or may under certain circumstances give rise, to various diseases in the brain, in the chest, and in the abdomen. The arteries of the brain thus affected, cause many of the diseases of that organ peculiar to advanced life, as softening, hemorrhage, palsy, and other disorders.

Inflammation, though occurring in the veins at all ages, is liable, after fifty and fifty-five, to attack spontaneously these canals, and thus to prove a cause of death, which, we have no doubt, has often escaped notice. Several cases of inflammation of the iliac and pelvic veins occurring in persons above the age of fifty-five and sixty years, we have seen prove fatal, either of themselves or by their secondary effects on the lungs, in which these lesions tend to cause the deposition of clots of blood within the pulmonary veins. It is not a matter of doubt, that this state, as a cause of death, has often escaped notice, and that deaths in consequence of the cause now mentioned, have not unfrequently been set down to the head of debility, old age, and similar causes.

Diseases of the kidneys, and the urinary organs in general, have also been overlooked; and the persons dying of them have been supposed and represented to have died of old age.

Diseases of the pancreas have also caused death, and in the same manner have been overlooked and ascribed to old age.

On all the questions thus arising, Dr Day has endeavoured in the present volume to furnish correct information, and if not to remove doubts and uncertainties, at least to diminish their number.

The work consists of thirteen chapters.

In the first chapter, which is general and introductory, the author gives a short view of the most important changes taking place in the system in advanced life. The second chapter is devoted to the subject of the means of preserving the health in declining life. The third chapter consists of general observations on the medical treatment of advanced life; the fourth is devoted to climacteric disease; and the fifth, to senile marasmus and wasting.

In the sixth chapter, Dr Day takes a general view of the diseases most fatal to persons in advanced life. In the seventh he considers the diseases affecting the organs of respiration; in the eighth, those affecting the nervous system; in the ninth, those of the alimentary canal and its appendages; in the tenth, diseases of the heart; in the eleventh, diseases affecting the urinary and generative functions; in the twelfth, diseases of the skin, ulcers on the legs, and senile gangrene; and in the thirteenth, the subjects of gout and rheumatism in their most usual forms.

The years of declining life are, according to Dr Day, naturally divisible into the following epochs:—

1st, Declining age, extending in females to about the fifty-second year, and in males to about the sixtieth year.

2d, Advanced age or incipient old age, extending in females from fifty-three to about sixty-five, and in males from sixty to seventy.

3d, Mature or ripe old age, dating from the periods already specified, and extending to about seventy-five in females, and eighty in males.

4th, Decrepitude, or second infancy, constituting in those whose measure of existence is so far prolonged, the last epoch of human life.

During the first of these periods, the feelings, disappointments, and anxieties of life exhibit their effects on the internal organs, as well as on the external appearance, in a more distinct manner and more forcibly than in earlier life. In consequence of the circulation becoming more languid, venous congestions and visceral obstructions, with the various diseases thence arising, become frequent. Piles, apoplexy and palsy, diseases of the liver, kidneys, and bladder, structural changes in the heart, dropsy, chronic disorders of the organs of respiration, gout and mental derangement, are not unfrequently developed about this period. These affections are common to both sexes. About the same time, however, in females, various changes are liable to take place either in the sexual organs, or in the system at large in connection with changes in the sexual organs. Disorders of the uterus, ovaries, and breasts are liable to ensue at, or about this time.

During the second period and the beginning of the third, these changes are liable to increase in force, if not in number. At the

close of the third period, the decision of the teeth causes great changes in the organs of assimilation. The osteo-steatomatous transformation of the arteries disposes to apoplexy and palsy, cerebral softening, and mortification of the extremities; and the state of the kidneys and bladder, as well as the tendency to morbid changes in the urine, contribute greatly to impair general health and strength.

In the last period, general decay of the system is manifest from a concurrence of causes. Obscure and latent inflammatory attacks may terminate life.

Among changes taking place in advanced life, affecting both the vital and mechanical properties of textures, next to those already noticed in the arteries, are the ossification of various cartilaginous structures; for instance, the thyroid and cricoid cartilages, affecting the voice and respiration; and the cartilages of the ribs, by which thoracic respiration is greatly impaired and limited, and that function is merely performed by the diaphragm and abdominal muscles. Fragility of the bones is another change of great moment and influence in advanced life.

Physiologists have usually considered it as an established fact, that the frequency of the action of the heart is diminished as life advances. This Dr Day represents to be not the case. Among 562 healthy females, whose mean age is 73 years, he finds that the average number of pulsations is a fraction above 79 in the minute; and among 197 healthy men of the mean age of 68, the beats are 72.5.

It is not our intention to enter on the often-repeated subject of the diet and general management requisite for those advanced in life. We believe that the proper course, what to avoid and what to choose, is well known, whether it be followed or not. In the present state of the question, it is a matter more for the moralist than the physician. Every elderly lady and elderly gentleman knows quite well how she or he ought to live; and if they do not live in that manner, it is because they will not. Self-denial is a thing, which is little known, or if known, it is not practised except in workhouses. If any one, nevertheless, wishes information, this chapter is recommended to especial attention.

About wine, tea, coffee, and other liquors, physicians always become eloquent, and sometimes speak not a little nonsense. Dr Day cautions against excess; and thinks that, to those who have not indulged in the use of wine, it may be well to say, "Drink no longer water; but use a little wine for thy stomach's sake, and thine often infirmities." The wisdom of advising this as a dietetic or an oinochrestic direction may be greatly doubted. It seems almost as if it were one of those texts by which the comfortable dean, with reddish nose, and a decanter of tawny port

before him, in the snug old-fashioned parlour of the deanery, encouraged himself to take his daily dose of two ample pints. Surely it must occur to any educated man reading the Epistles, that they were written for more elevated and nobler purposes than instructing men whether they should drink wine, or refrain from that generous but somewhat fiery liquor; and that St Paul, though an excellent apostle and theologian, might have made but an indifferent physician. The evil is, that it is exceedingly difficult to keep men, or even ladies, within the strict letter of the law as to the use of wine; and, where it is permitted to take a little, every one thinks that the little that he or she takes, is really adhering to the precept. So often, however, have we seen and known this text quoted, and made to justify what it never was intended to do, that a more dangerous one we think hardly exists; and we must say, that it is safest to confine the precepts in the Epistles, so far as they are intelligible, to their legitimate and proper uses.

In the whole of these directions regarding the use of wine for the aged and the invalid, there is some danger and not a little folly. One physician recommends for the aged invalid Madeira, another Amontillado, and a third Madre di Xeres; while a fourth tells his patients that port or tent, or some other vinous compound, is the grand remedy for his decaying tabernacle. What advantage have several of these forms of vinous liquor, except that they are expensive, and consequently unattainable by the humble invalid? It appears to us, that all are only different forms of poisonous articles, the use of which has been rendered necessary by the preposterous and pernicious habits of society. A man begins to take wine in early life; and, as he continues it as long as he can, without receiving any intimation of its bad effects, he is then given to understand, that he has done himself and his health irreparable injury, but that if he stops, he is sure to be made much worse, and probably to die. Would it not be much better, for all persons to be taught, in early life, that wine is an indulgence both unnecessary and hurtful, and as the time must come, when its use must be given up, it is best and safest not to begin it at all? Why should any habit, which is admitted on all hands to be bad, be either commenced, encouraged, or carried on?

But why is the wine necessary? Oh, says some sagacious person, it enables a man to digest his food, and is a pleasant tonic. If either of these statements were facts, then it might be reasonable to say, that the moderate use of wine is beneficial. In a large proportion of cases, however, these statements are not facts. Whenever wine is believed or felt to be necessary after taking food, it may be safely asserted, that food too much in quantity and improper in quality has been taken; and the same is still more

decidedly true regarding spirits. First, after taking as much food, and eating as great a variety of articles as might serve for two, if not for three meals, wine or diluted spirits, and sometimes undiluted spirits, are taken, in order to enable the stomach, it is said, to perform its duty; in other words, to obscure, stifle, and extinguish all those uneasy sensations which it is made to feel from being taxed far beyond its powers; to render the individual insensible to those salutary warnings, which, in the natural state, are intended to inform him, that he has committed several serious errors in diet. Then, after two or three hours more, when other uneasy feelings begin to arise, he endeavours to allay and stifle them by the use of strong astringent liquors; and lastly, perhaps after committing another error in diet, he has recourse to some brandy and water, and then retires to a disturbed and restless sleep, with a stomach loaded to distension with food little or not at all digested.

Such a course of constant offence to a delicate, membranous, vascular, and nervous organ, cannot proceed long without causing serious injury both to its functions and its structure, and through that to other organs and the system at large. And then the poor victim of all this self-delusion is told that he must not give up his evil courses; for the end of that man would be worse than the beginning. If this really be the fact, patients should be warned in time, while they have not yet proceeded irrecoverable lengths in this career of gastronomic stimulation,—that it is easier and safer not to begin, than, after having begun, to check this course. On the other hand, we must say this, that we have not witnessed, from a gradual and judiciously-managed stoppage of the rations in the persons now referred to, those dreaded effects on which physicians love to expatiate. The subject is, after all, an unprofitable one, for nobody listens to lectures on temperance and abstinence; and if any one does, he forms resolutions which are made only to be broken.

The observations on the uses of medicine in the management of the general health and diseases of the aged, are judicious, and show that the author has carefully considered this delicate subject.

We have always been skeptical as to the existence of climacteric disease, and doubtful whether the true causes of the several forms of this state have been justly assigned. At the same time, as we have more than once considered this subject, and as we press not our views on others, we have only to say, that Dr Day gives, after Sir Henry Hallford, an interesting description of the causes and phenomena of this condition of the system.

On the diseases most fatal to persons advanced in life, Dr Day communicates several useful pieces of information. During the five years between 1843 and 1847 inclusive, he finds that the

total number of deaths among persons aged sixty years and upwards, occurring in the metropolis, amounts to 53,048. Of these, 15,136, or about two-sevenths, are recorded as having died from the effects of old age. Dr Day agrees with most physicians in thinking that if this class of deaths were ascertained by inspection, the number would be very much diminished. Taking the numbers of both, however, as they stand, 47,902 instances of death by actual disease are left. These are distributed in the following manner :—

Death is ascribed to disease of the respiratory organs in 12,598 instances ; to diseases of the nervous system in 6947 instances ; to diseases of the digestive organs in 3141 instances ; and to diseases of the circulating system in 2841 cases.

Besides these, there are recorded 1076 cases of diarrhoea, 748 of influenza, mostly during the close of 1847, and 417 of erysipelas.

The following table shows the comparative frequency of the causes of death at and after the age of 60 years. Among 1000 persons who have attained that age, there die of old age 285·3

Diseases of the Organs of Respiration.	{	Bronchitis	79·3	}	237·3		
		Asthma	62·4				
		Consumption	35·7				
		Pneumonia.....	27·1				
		Hydrothorax	10·4				
Diseases of the Nervous System.	{	Other diseases	22·6	}	130·9		
		Apoplexy	53·0				
		Palsy	51·2				
Diseases of the Digestive Organs				}	59·2		
Diseases of the Circulating System.	{	Other diseases.....	26·7				
		Diseases of the Heart...	51·3				
		Pericarditis	1·3				
Aneurism				}	53·5		
Diarrhoea						20·3	
Influenza						14·2	
Erysipelas.....						7·8	
Other diseases						808·7	
						191·3	
						1000	

The residual 191·3 marked as other diseases, consist in a great measure of typhus and dropsy. These the author does not include in the table, because typhus is made to include all cases of continued fever, and because dropsy is a symptom, and may arise from different primary diseases.

It may be further observed, that the rather large head of asthma—62·4 in 1000,—is usually little entitled to be, in all instances, regarded as a primary and independent disease, and in all probability includes various diseases of the lung, as chronic bronchitis, emphysema, and several diseases of the heart, valvular disease, and various forms of dilatation. Hydrothorax, in like manner, which figures at the rate of about one per cent., is very probably symptomatic either of disease of the heart or disease of the lungs. Of the class, other diseases, under those of the respiratory organs and of the nervous system, it is not easy to know what to make. Most assuredly they owe their large proportion and present position to imperfect and careless diagnosis, and probably to the ignorance and indifference of several of the reporters.

Pneumonia, in different forms, is, as has been already stated, the most frequently fatal disease of the aged. This indeed does not appear from the London tables of mortality, in which *bronchitis* is the most frequently fatal disease among those of the organs of respiration affecting the aged. But the fact is abundantly established by the researches of physicians in hospitals in other countries. In the inspection of the bodies of 390 persons at the Bicetre in Paris, the ages of whom ranged between 60 and 90 years, M. Prus found that death was the effect of disease of the respiratory organs in 149 cases; and that these were distributed in the following manner:—

Pneumonia, 77, of whom 6 terminated in abscess of the lung.	
Pleurisy, ...	26
Tubercular consumption, ...	18
Asthma, ...	10
Bronchitis, ...	8
Pulmonary congestion, ...	4
Asphyxia from meteorism dependent on indigestion, ...	3

The remaining four were, isolated cases of comparatively rare disease.

Thus, among the entire number of 390 bodies inspected, it appears that in 149 instances, or nearly five-thirteenths, death was caused by disease in the organs of respiration; that among the instances of disease of the respiratory organs, more than one-half were cases of pneumonia; and among the entire number of bodies examined, one-fifth were examples of pneumonia. Dr Day therefore draws a conclusion, which to us has been long familiar, namely, that pneumonia in some form is the most usual cause of death to the aged among the diseases affecting the organs of respiration, and that very probably it is the most frequent cause of death among all diseases at an advanced period of life.

Its most common causes are chronic bronchorrhœa or previously-

existing chronic catarrh, the cold season of the year, especially cold frosty nights, and, in certain circumstances, long confinement to the recumbent position in bed. The latter, nevertheless, is rather a cause of hypostatic congestion than of genuine pneumonia. Pneumonia occurs also in the gouty and rheumatic, and those who live rather freely. This was the disease which proved fatal to old Parr; and in his case, it was induced by transition from a coarse spare mode of living, to one in which he lived well, and drank strong wines and other fermented liquors.

Dr Day is rather averse to general blood-letting in the treatment of this malady in the aged, and thinks that local depletions are more eligible and more safe. Antimonials, diaphoretics, and counter-irritants form the principal remedial means.

Bronchitis is liable to terminate in *bronchorrhœa*, which is very common in certain individuals, especially of gouty and rheumatic diathesis. It consists in a profuse secretion of mucous matter from the bronchial membrane, which, at first inflamed, becomes afterwards relaxed. The disease must then be regarded as one of inordinate secretion and debility.

The best remedies are, first, ipecacuanha in emetic doses, then in nauseating doses; then the inhalation of some of the gum-resin vapours; vapour and sulphur baths; dry cupping; gentle laxatives; meadow-saffron in the gouty; and a combination of one grain of blue pill, one-eighth of a grain of tartar emetic, and two or three grains of extract of conium, three times daily, then twice daily, and finally, once daily.

Bronchorrhœa is one of the maladies which give rise to the peculiar symptom called asthma.

Of asthma, Dr Day describes the urinous and gouty forms. These are closely allied. In the urinous asthma, the symptoms are produced chiefly by the presence either of urea, or rather of urate of ammonia in the blood, and causing morbid states in various textures and organs, and morbid phenomena in almost all the functions. The gouty asthma is merely one of the forms of misplaced gout.

Stramonium is spoken of as a useful remedy; and medicated cigarettes compounded according to the formula in the subjoined note,* are represented to be much and beneficially used on the Continent.

* Picked leaves of Belladonna	.	.	6 grains.
Hyoscyamus, (leaves, we suppose),	.	.	3 grains.
Stramonium,	.	.	3 grains.]
Phellandrium aquaticum,	.	.	1 grain.
Watery extract of opium,	.	.	$\frac{1}{4}$ of a grain.
Cherry laurel water,	.	.	sufficiency.

The leaves, previously stripped of nerves, to be chopped and mixed; the opium to be dissolved in the cherry laurel water, and the whole to be mixed.

Attention to diet, regulation of the bowels and the non-naturals, in general, are the great points of cure.

All that we require to notice as to influenza is, that Schönbein, the discoverer of gun-cotton, thinks that epidemic influenza depends on the presence of *ozone*, an imponderable agent, in the atmosphere. This product is supposed to be the result of atmospheric electricity; but nothing is hitherto ascertained regarding its properties, except that it may be prepared in the laboratory of the chemist, that it is an intensely oxidizing agent, and that its presence may be demonstrated by chemical tests.

From the Tables for the mortality of the metropolis, it results, that during the five years extending from the beginning of 1843 to the close of 1847, 6,947 deaths, in persons at and above the age of sixty, arose from diseases of the nervous system; and that of these, 2,812 are ascribed to apoplexy, and 2,806 to palsy, leaving 1,329 cases to cerebral softening, tumours, and diseases of the membranes. Dr Day expresses himself severely against the classification; assigning as the reason;—that the statement that palsy caused death to 2,806 persons conveys no correct information whatever. It may be presumed, that of this number, several, or even many deaths, were caused by cerebral softening. In many instances also, death seemingly produced by affections of the brain, depends, in truth, upon disease of the liver, of the kidneys, or of the arteries. Some more careful distinction ought to be made to render these tables of any utility, either pathological or practical.

Of the next subjects, meningeal apoplexy, cerebral softening, *meningitis*, and the different forms of mental derangement, we have so often spoken in detail in recent volumes, that it is unnecessary to do more than mention the names.

The frequency of diseases of the alimentary canal may be understood from the numerical proportion in the table already given. The comparative frequency of each disorder, and of disorders of different divisions of the tube and its appendages, there are no means of exactly ascertaining. The only accurate facts are obtained from the report or memoir by Prus, already referred to, in which it results, that among 390 persons dead above 60 years from various diseases, in only 49 cases was death caused by diseases of the alimentary canal and its peritoneal covering, and in 8 instances from disease of the liver and gall bladder. In these 57 cases, death was caused in 27 instances by *enteritis*; in 10, by cancer of the stomach; in 4, by gastro-enteritis; in 3, by dysentery; in 3, by hepatitis; and, in 10, by miscellaneous disorders.

Dyspepsia, in different forms and with different symptoms, is a common but not a fatal affection in advanced life. It may, however, produce other disorders, as urinary and renal affections, gout

and rheumatism; and it may, during periods of epidemic influence, pave the way for diarrhœa and dysentery and cholera. Thus, in the summer of 1846, many aged persons habitually dyspeptic were cut off by diarrhœa. During the prevalence of the cholera epidemic, which is not yet completed, many persons above fifty, habitually dyspeptic, were also attacked and destroyed. And, lastly, it is not unusual for persons who have had dysentery abroad, and in miasmatic situations, to be attacked and destroyed by this disease after the age of sixty years.

The present author thinks that the most frequent forms of the disorder in the aged are acute atonic dyspepsia, and that described by Dr Tweedie J. Todd as follicular gastric dyspepsia. In the latter case, the muciparous follicles are in a state of chronic irritation and congestion.

Diseases of the rectum are not uncommon at an advanced period of life. Such are hemorrhoids; inflammation of the rectum (*proctitis*); contraction of the rectum; follicular disease; cancer of the rectum or of the colon, mostly tubercular; and occasionally abscess around the bowel.

The affections of the heart may take place at any period of life; yet a certain order of them is greatly more common at advanced periods than previously. All the affections which depend on, or are connected with, steatomatous and osseous deposit, are peculiarly the appanage of advanced life. *Angina pectoris* is very seldom met with before the age of 50 years. After that it is a frequent cause of death. Valvular disease, in like manner, is more common after 60 years than previous to that age. But it is a common observation, confirmed by all experience, that the symptoms produced at that period of life, are rarely so well marked as when lesions of this kind appear at earlier periods; and hence aged persons more frequently are cut off suddenly and unexpectedly than younger individuals labouring under these disorders.

The several orifices of the heart become enlarged as life advances, but at different rates. Hasse ascertained, that while the progressive enlargement of the two auriculo-ventricular orifices is tolerably uniform, the orifices of the aorta and pulmonary artery increase equally till the meridian of life; and that the aortic orifice enlarges more rapidly in advanced life than that of the pulmonary artery, so that, in aged persons, the pulmonary orifice is narrower than the aortic. This comparative contraction of the pulmonary artery, in relation to the aorta, is one among other causes of the venous plethora or congested state of the veins, which is almost constantly observed in advanced life. The valves, also, become inadequate, generally in consequence of chronic inflammation rendering them rigid, and shortening or contracting their texture.

On the subject of diseased states of the urine and the urinary organs, on affections of the skin, and on gout and rheumatism, it is unnecessary to enter, as these subjects have been frequently and fully considered in treating of various other works.

It is only necessary to say, that Dr Day has produced a work containing much instructive information on the diseases of aged persons and the methods of treatment, medical and domestic, most suited to the removal and alleviation of these diseases. He has been at considerable pains in collecting from various sources whatever facts might elucidate either the pathology, the etiology, or the treatment of these diseases. The volume cannot fail to be read with advantage; and we shall not further anticipate the interest of the reader.

ART. III.—*Elements of Anatomy*. By JONES QUAIN, M.D. Fifth Edition. Edited by Richard Quain, F.R.S., and William Sharpey, M.D., F.R.S., Professors of Anatomy and Physiology in University College, London. In two vols., Part iii., forming the conclusion of volume second. London, 1848. 8vo, Pp. 879—1363.

THIS third part concludes the new edition of the work of Dr Jones Quain. It is only necessary to say, that the work now, when entirely before the public, fully realizes the opinion which was given in this Journal on the first two portions. It may be said to be, in a great degree, a new work. This character is particularly applicable to the Treatise on General Anatomy, which occupies in the present edition three hundred and seven pages. It is manifest that the author, Dr Sharpey, has with this part taken very great pains; and that, from personal observation and that of the most skilful microscopical observers, he has presented a very correct and faithful view of the present state of knowledge in the minute anatomy of the different tissues.

The Descriptive Anatomy in the present portion is accomplished with not less ability. It embraces the anatomy of part of the nervous system, the anatomy of the eye, the ear, the alimentary canal, the organs of respiration, the urinary organs, and the reproductive organs, with an account of the anatomy of the parts concerned in hernia and in lithotomy.

We have no hesitation in recommending this treatise on anatomy as the most complete on that subject in the English language; and the only one, perhaps, in any language, which brings the state of knowledge forward to the most recent discoveries.