which he formerly enjoyed. This moral disease penetrates deeper into life than can be well conceived. It saddens the hearts of thousands who, once happy and useful members of society, are now lost in the depths of darkness and despair.

But this great evil should be considered as national, and the preservation of the public health and morals not only as an act of humanity and Christian charity, but one specially of public economy; for, without taking into account the loss of labour, which must be very great, we are to look to the heavy expenses incurred during sickness, the consequences of numerous deaths, involving the support of so many widows and orphans, who must fall on the public, either on the parish or on private charity; and, lastly, the immense cost to the country for the maintenance, the prosecution, and punishment of criminals.

Under these circumstances it will be evident that, by carrying out these improvements, not only will a great saving be effected to the public; but that, by affording sufficient employment, suitable education and religious instruction, and the suppression of many of the public houses and beer-shops,—those great sources of moral and physical corruption and ruin,—a vast change would speedily be apparent in the well-being and happiness of the great mass of the people; for we cannot admit that moral or physical degradation, with all its humiliating and distressing consequences, belongs to any class of society; as it is well known that the most exalted state of moral feeling and courage is to be found amongst the poorest of our race, unless the faculties have become, by long-continued hardship and misery, incapable of sustaining the moral energy.

By thus devising and carrying into effect this system of reformation and improvement among the lost orders of the lower classes, man would be seen to emerge from this depraved and neglected state, and to rise to that condition originally destined for him by his all-wise and merciful Creator.

ART. III.—*The Brain the Sole Centre of the Human Nervous System.* By EDWIN LEE, Member of several of the principal European Medical and Chirurgical Societies. (Read before the Royal Society, May 1848).

THIS paper was written for the Royal Medical and Chirurgical Society, the subject appearing to the author one eminently calculated to elicit useful and interesting discussion upon points of great practical importance. The committee, however, thought

proper to decide that the paper was inadmissible ; it was therefore offered to the Royal Society.

The following brief considerations are submitted to the Society in consequence of the opinion which I have always entertained, that the inferences which have been deduced from experiments upon the inferior animals are less applicable than has been supposed to man and the higher mammalia, in whom different anatomical and physiological conditions are seen to prevail. That this is a point the decision of which is of great practical importance, is sufficiently obvious ; and having already expressed these opinions in a work, of which a new edition is preparing, I am desirous of previously having the subject discussed in the Society, in order if possible to meet any objections that may be advanced against it and the corroborative opinions which I shall adduce.

It is well known that in the lowest class of the animal creation, vitality is pretty equally diffused, so that subdivision does not occasion destruction of life ; and this is also the case somewhat higher in the scale, where the rudiments of a distinct nervous system are perceptible, as in the tape-worm. In proportion, however, as we ascend to the more perfect animals, the vital powers tend to become more centralized and dependent upon the integrity of particular organs. Thus, in reptiles and frogs, which possess a brain and heart, though the vital phenomena may continue to be manifested after the destruction of these organs, this will be but for a short time, and the prolonged continuance of life is impossible. The more complete dependance of life in the higher animals, upon these organs, need not here be dwelt upon ; even slight lesion of them being mostly destructive.

From the two great divisions of structural difference in the nervous system,—viz. the granular or vesicular grey substance, and the fibrous white matter,—the difference of function has been inferred with every appearance of truth, both from experiments and from pathological observations, viz. the former being more immediately connected with the generation of nervous power ; the latter serving solely for its conduction. In inferior animals, where the brain forms but a small portion of the nervous system, re-active effects may be produced by stimulation for a longer or shorter period after its removal ; this organ being, however, necessary in all cases in originating actions, and in the perception of sensations. But in the higher mammalia, and especially in man, where the grey matter is for the most part agglomerated in the brain, being but in very minute proportions elsewhere (the spinal cord or the ganglia), these reactive effects do not take place, or very speedily cease, when the influence of this organ is no longer exerted ; and, consequently, the inferences which have been drawn from obser-

vation of phenomena manifested in animals in which a totally different condition of the nervous system exists, do not appear to me to be confirmed by the arguments, which have been brought forward in proof of the independence of separate portions of the human nervous system. Bichat, as is well known, considered the sympathetic as constituting a distinct system from the cerebro-spinal, and each individual ganglion as a central focus of organic life, analogous to the great and only centre of animal life, the brain. He also considered that each nerve was in great measure independent of the ganglion whence it proceeded; and even that each portion of such nerve was independent of the rest, forming in itself a distinct focus of nervous influence. Kolliker, likewise, in common with some other modern physiologists, has recently advocated the opinion that each isolated ganglion, with its connected nerves, is the seat of a special activity constituting a simple functional whole; but this view does not seem to be in accordance with physiological and pathological facts; and the opinion of Galen, by whom, says Mr Solly, in his work on the Brain, the ganglions of the sympathetic nerves were supposed to act as buttresses in order to strengthen them as they recede from their reputed origin, seems to me to approach nearer to the truth. Willis also regarded the ganglionic nerves as an appendage to the cerebro-spinal system; and on viewing the question in its more comprehensive aspect, most persons would, I think, arrive at the conclusion, that, at least in man, these separate portions are essentially dependent upon the brain for the accurate performance of their functions. Mr Longet remarks, with reference to the supposed independence of the sympathetic,—“Facts are far from confirming the opinion, that each sympathetic ganglion is to be considered as a small centre acting independently of the cerebro-spinal axis. The acts of sensibility and contractility termed organic are but very imperfectly manifested, when the nerves of the cerebro-spinal system which converge to their ganglia, have been cut. The imperfect movements which persist for a certain time, take place even after the section of the nerves which arise from the ganglia, to be distributed to the viscera, as the heart and intestines, and last *until the entire exhaustion of the power diffused in the minute nervous fibrils.*” This position is well exemplified by the observation of Henle, which has been verified by Kolliker, that in rabbits, dogs, and horses, a fresh piece of intestine cut off close to the mesentery, and chemically or mechanically irritated, contracts only circularly at the irritated points. If, however, the mesentery with its nerves and ganglia be taken out with the intestine, and the part be then irritated, the contraction extends at the same time over a greater space; and if the nerves of the intestine are allowed to remain united to the spinal cord, the movement extends over the whole alimentary canal.

Professor Muller likewise observes upon this question,—“The moveable parts supplied by the sympathetic (heart and bowels) are, to a certain extent, independent of the brain and spinal cord. These organs must, however, also be regarded as the source of the functions of the sympathetic, *if its power is not to be exhausted.*”* “It is well known that the action of the heart varies from passion, and torpor of the abdominal viscera ensues on paralysis of the cord.” “From these facts we may infer, that the sympathetic is charged from the central parts of the nervous system as sources of the nervous principle, and that being once laden it retains its charge, continuing to resist nervous power, even when the more distant supply is suspended, and is renewed only after a time. If, however, the brain and cord lose the capability of being the source of the nervous principle, exhaustion ensues, whence the debility and scarcely-perceptible pulse at the close of acute diseases.”

Hence may be seen the reason of the uninterrupted persistence of the organic functions during sleep, as also why impressions made upon internal organs are so seldom transmitted to the sensorium. “This system,” says another French physiologist, “specially consecrated to the nutritive and visceral functions, is but seldom affected by exterior impressions. It is acted upon by organs upon which it reacts, but this takes place quietly and obscurely. This depends upon the weakness of the conducting property. The ganglia of this system disseminated along its course impede the too free communications; and thus violent commotions of the cerebro-spinal system (the passions, &c.) are requisite for the influence to be propagated to, and to affect the sympathetic; and very sudden and intense disorder of the viscera must occur before their sensations can be transmitted to the percipient organ. It would appear that each ganglion constitutes a particular focus or centre, in which the operations, of a purely local innervation, are arrested and perfected. From this results the advantage, that the quiet and equilibrium of the vegetative functions are more easily preserved, less oscillation is to be apprehended, and there is more stability and continuity in the functional phenomena.”†

The same arguments against the non-independence of the sympathetic, apply with equal force to that of the spinal cord, and consequently to the theory of Dr M. Hall; according to which, certain muscular movements termed reflex, depend upon a particular class of nerves, of which the spinal cord is the central organ, and which are independent of the brain; these movements, in experiments upon animals, not being producible when the spinal cord is removed. In contending for the originality of his views, Dr Hall says;—“Has it been stated in any work, ancient or modern, that

* The italics are mine.—E. L. † Duges, Physiologie Comparée.

the deglutition of water by the pharynx, the exclusion of carbonic acid by the larynx, the retention of the urine or fæces by the sphincters, are *exclusively functions of the spinal marrow, and of a peculiar system of excitor and motor nerves, of which it is the centre and axis?*" It may be true that this view had not been previously taken by others, and, as I think, with reason; for, upon a closer investigation, it will be found that the influence of the brain is exerted in these actions, though it may not be so clearly manifested as in the case of the more strictly voluntary movements. To take, as an illustration, the act of deglutition;—according to Dr M. Hall, the prehension of food is a voluntary action, and cannot be performed without the concurrence of the brain; but the contraction of the fauces and pharynx upon the morsel is a reflex action, and swallowing may take place even after the brain has been removed. The *sphincter ani* remaining contracted after decapitation, in one of the lower animals, the lower portion of the spinal cord being uninjured, but becoming relaxed when this portion is destroyed, affords another illustration of the reflex theory. In making these deductions, however, it would seem that the distinction has not been sufficiently kept in view between the results of these experiments, and those which would ensue upon analogous ones being made upon animals higher in the scale of creation, in which, as has been already observed, the portions of the nervous system tend more to a common centre; and in the higher mammalia, the inability to produce muscular reactions after removal of the brain, and especially after decapitation, clearly evidences the dependence of the use of the nervous system upon this organ. If a morsel of food could be placed in the pharynx of one of this class immediately after removal of the brain, there it would remain.

"It results from the preceding observations," says M. Duges, "that the cerebral lobes are the chief centres of innervation, and that the spinal cord is rather destined to conduct than to elaborate the impressions and mental determinations; it is, however, not incapable of all action of centrality, if the term may be allowed; and this action will be so much the more apparent and easily produced in proportion as the preponderance of the brain is less, and as the spinal cord presents a greater analogy with the anterior ganglia of the invertebrata. This analogy is seen in the tubercles near the origin of nerves in serpents and lizards, the tail of which, when separated from the body, moves actively; and the decapitation of a turtle, of a lizard, a serpent, or of a frog, still admits of life being prolonged in the mutilated body for several days; the animal is agitated, draws up its limbs when pinched, leaps or moves when strongly excited. Legallois observed similar effects in young decapitated rabbits, in which the circulation was arti-

ficially kept up after ligature of the carotids. Analogous phenomena have been observed in the anencephalous fœtus. Birds hold in this respect an intermediate position between the mammalia and reptiles."

In the former editions of my work on nervous disorders, before having had cognizance of the works whence the preceding extracts are taken, I expressed my opinion as opposed to the theory of the independent action of the spinal cord, as also of its isolated portions, which some have deduced from the phenomena occurring after its section in the inferior animals, in these terms; viz. "that the nervous power retained in the spinal cord for the immediate actions of the parts which it supplies must soon be exhausted if communication be cut off from the source of nervous energy." This view of the question, I have since perceived, is the one taken by some distinguished physiologists, to whose opinions I will briefly advert.

It has been ascertained that nerves do not terminate, as was formerly supposed, in the parts which they supply, but are reflected in loopings, and that the filaments of which a nerve is composed, though bound together in the same sheath, yet pass separately into the spinal cord, and, according to some physiologists (Valentin), on to the brain, forming loopings also in the cerebral hemispheres, coming in contact in their passage along the cord with its grey matter; and this view appears to be corroborated by the phenomena of disease. Muller remarks that "the existence of distinct paralysis of sensation and of motion goes to prove that the sensitive and motor filaments pass up separately to the brain;" and again, "we must also consider that it depends upon the brain how many muscles of the trunk are to be moved in each action; whence it would seem to be a necessary deduction, that the primitive fibrils of the nervous trunks which enter into the cord do not there unite, but run parallel with each other to the brain, in order to convey to it local sensations, and to receive from it isolated excitations for the movements of parts."

On the other hand, Sir C. Bell, Mr Grainger, and Dr M. Hall, distinguished in the anterior as well as in the posterior roots of the spinal nerves, filaments passing directly into, and terminating in, the grey matter, the others proceeding directly to the brain; and Stilling has more recently given in his work a representation of longitudinal and transverse fibres ending in the cord. Professor Carus, however, referring to these contradictory opinions, observes that, although these nervous filaments do intertwine in a complex manner with the grey matter while in the cord, yet that their true terminal loopings are in the brain. "The spinal cord," he adds, "must therefore be considered as the common trunk of all the nerves of the body, and differs from a nerve only in the circum-

stance of the considerable admixture of grey substance. My investigations accord with those of Valentin; those of the other physiologists do not take account of the nervous loopings, and are not founded upon such accurate microscopic observations."

"As the radii of the globes converge towards a centre, and as gravitation attracts bodies in this direction, so in like manner do the primitive filaments of the nerves of the spinal cord converge to the brain, in order there, and there only, to form their internal or central loopings, the external or peripheric ones being diffused throughout the organism. The same distinguished physiologist further observes, with reference to the reflex actions,—“herein nothing more takes place in the central nervous system than what so commonly and universally occurs in the sympathetic system, viz. that the stream of sensitive innervation in the nervous filaments, instead of proceeding from the peripheric loopings direct to the central loopings, passes in its course through grey substance, in which sensation is concentrated, and, by reacting, produces muscular contractions.

"Thus we may readily perceive the reason why, wherever grey matter is in intimate contact with nervous filaments, the transmission of innervation in the latter must be modified; and it cannot be otherwise than that, by such intermediate operation, the action of the central grey masses on the primitive filaments will necessarily be to a certain extent lessened or altered. We see that in proportion as nervous filaments pass through a greater quantity of grey matter, so much the more must their sensitive as well as their reacting powers be obscured. The nerves of the lower extremity consequently produce more obtuse sensations, and less rapidity and delicacy of movement, than those of the upper extremities, and especially of the head, or those which are concerned in speech; for it is clear that the former pass through very much more not central grey matter than the latter. Thus the nerves of the three higher senses manifest an extraordinary peculiar power, inasmuch as the course between their periphery and the central vesicular mass is short and direct, without passing through any intermediate grey matter. The ganglia of the posterior roots of the spinal nerves likewise serve to check a too powerful stream of sensitive innervation."

The following passage confirms the views already expressed relative to the sympathetic ganglia:—"The nerves of the sympathetic follow throughout the same law as those of the cerebral system; and it is because their filaments pass through one or more masses of grey substance before reaching the brain, that the sensations transmitted to this organ, as also its reactions are more obscure. Hence, vegetative life pursues undisturbed its peculiar course. When sensations and reactive movements are necessary

for vegetative life, they are found to be everywhere supplied through the medium of the sympathetic without the higher powers of the central system being drawn into direct co-operation, and consequently the actions of organic life become perceptible to consciousness only under certain circumstances (such as increased activity of the organs, the alterations produced by disease), and are felt so much the more powerfully from their being unusual. Hence, on the other hand, the influence of the mind upon vegetative life is so considerable, and thus we may properly estimate the opening and closing of the sphincters, the peristaltic movements of the intestines, that of the absorbents, &c. which always require the unconscious co-operation of the cerebro-spinal nerves.

“ The nervous system in general, and the central portion especially, can only be considered as a whole, which, originally small and simple, becomes afterwards larger and more complicated. To entertain the idea of a nervous system as existing piecemeal, and subsequently uniting into a whole, would be much the same as if one were to imagine that, in our bodies, the trunk and extremities existed separately, and afterwards grew together. It is highly interesting, with reference to this point, to take a glance at animal life. The brain is small in proportion as the spinal cord is large. The more grey substance there is in the cord, the less there are of primitive filaments, so much the more will these reflex actions be manifested. If the body of an eel be cut into four portions, each one will move for a long period upon any slight excitation; whereas the decapitated trunk of an adult mammal immediately lies still, or exhibits contractions only upon irritating the cord itself. On this account, also, the reflex movements are so much the more strongly manifested in proportion as the animal is young, because the grey matter preponderates in the cord. In the fœtus, the sympathetic nerve, and especially the chief ganglia, are proportionally large.”*

These opinions are corroborated by the high authority of Professor Cruveilhier, who says,—“ The independence of different portions of the spinal cord of one another; the independence of the cord itself of the brain, which has been pretty generally admitted of late,—seems to me to be a serious physiological error, founded upon ingenious experiments. The theory of the ancients, who looked upon the spinal marrow as a great bundle of nerves destined to respond to all the nerves of the economy, in order to transmit definitively these impressions to the brain, or to receive from it the voluntary or organic impulses, is much more in accordance with facts and with the great anatomical law of the continuity of the nervous system.”

MM. Foville and Pinel Grandchamp remark that, “ As nume-

* Physiologie.

rous observations demonstrate, that diseases of the cerebrum and cerebellum, co-existing with integrity of the spinal cord, destroy, pervert, and alter the sensation and motion of parts supplied by the spinal nerves, we must infer that the cord, like the nerves, depends upon a central focus of action, which must be referred to the cerebrum and cerebellum."

"Two rabbits," says Dr M. Hall (endeavouring to show that the tone of muscular fibre depends upon the cord), "were taken and their heads removed. The spinal marrow of one was cautiously destroyed by a sharp instrument, and its limbs were perfectly lax; whereas in the other, where the spinal marrow had not been touched, they retained a certain degree of firmness and elasticity. The difference was most obvious. *On the following day*, the limbs of both animals *were equally rigid*.* The same objection which has been made with reference to reflex action, depending solely upon the cord, also applies to muscular tonicity."

"Exposed muscles," says Professor Muller, "on being irritated, contract without either the centripetal or centrifugal nervous action being requisite; but when the muscles are covered with a skin, they must derive their stimulus to contraction, in the first place, from excitation of their sensitive covering, which induces centripetal action of the sensorial nerves, and centrifugal motor excitation *from the brain*. In this way is produced contraction of the larynx and air-passages from acid gases, or cough from irritation of these parts. It is the same with respect to the sphincters of the anus and bladder. These muscles cannot be excited directly by the presence of the egesta, but these substances act on the sensitive nerves of the membrane, and excite the spinal cord, which being *constantly charged with motor power*, reacts upon them. Hence, after lesion of the cord, their contraction ceases."

M. Flourens also remarks on this point,—“The lower end of a divided nerve may still continue to excite contractions, but only when it is irritated. The nerve has then an action proper to it, but this action must be brought into play by external excitation. The nerve is therefore only a subordinate agent. It is the same with respect to the spinal cord. On its being separated from the brain, all spontaneous movement of the body immediately ceases; it still retains, however, a certain degree of action, and an external irritant may bring this action into play. Like the nerve, it has therefore an action proper to it; but, like the nerve, it has neither spontaneity nor primordality of action. The spinal cord is consequently likewise only a subordinate agent. This spontaneity of action arises *solely from the brain*.”

These quotations may suffice to show the true relation which the spinal cord holds to the brain, as also the error of a too hasty

* The italics are mine.

generalization of the results of experiments upon cold-blooded animals. Thus, when the body of a viper was observed, after decapitation, to continue to direct its course to a hole in the wall to which it was accustomed to retreat; and also when some birds are seen, immediately after decapitation, to continue to run or even to fly to a short distance,—it is by no means to be inferred that these movements are performed independently of the brain, but these instances merely prove the persistence, for a brief period, of an impulse already given. Thus, also, in soldiers tired by a forced march, even while asleep, the legs may continue the movements so frequently repeated, and carry them on unconsciously, and, as it were, mechanically.

“Many movements,” observes Cabanis, “take place in the animal economy without consciousness (*a l'insu du moi*), but, nevertheless, through the medium of the sensitive organ. In all those inferior animals which survive the destruction of their brain, the muscular parts, isolated from the sensitive centre, still exhibit, during a longer or shorter period, movements which are maintained entirely by the influence of a sensibility which may be termed posthumous.”

In the young of some warm-blooded animals, which, after decapitation, exhibit reactions to external impressions, vitality remains in the trunk so much the longer in proportion as the animal is near the time of its birth. Legallois found that in newly-born rabbits under these circumstances, sensation and motion were retained for a quarter of an hour; whereas in those a month old, these signs of vitality scarcely continued two minutes. The same principle also applies to the instances of anencephalous infants, which have existed for a short time after birth, breathing, and performing some of the instinctive actions, which cannot be continued without a brain; and life consequently soon becomes extinct, being retained longer in those cases where the *medulla oblongata* is present. That neither this part, however, nor even the spinal cord, is essential for the manifestation of these phenomena, is shown by the cases in which this portion of the nervous system is likewise absent. M. Gerdy, in a recent physiological work, states that MM. Fauvel and Mery each exhibited an infant born without either brain or *spinal cord*, but which, nevertheless, lived for several hours. That of Mery lived twenty-one hours, and even took some nourishment. As regards also the sensations and spasmodic contractions of paralyzed limbs on being irritated, which have been adduced as purely reflex actions of the cord, there is no doubt in my mind that the influence of the brain is still excited, though there may not be consciousness of its operation, as is the case when paralyzed limbs are convulsed in epileptic attacks. It would, however, occupy

too much time to consider upon the present occasion the various practical bearings of this question. I will therefore conclude by submitting the following deductions:—

1. That in man and the higher mammalia, the brain is the sole centre of the nervous system and the source of power.

2. That the grey matter of the spinal cord and the ganglia of the sympathetic are to be regarded as reservoirs of this power for the immediate actions of the parts supplied, which speedily becomes exhausted if communication with the brain be cut off, and also as modifiers of impressions transmitted from the periphery to the brain.

3. That the automatic and instructive actions are not performed independently of the concurrence of the brain, though its influence is less direct and manifest than in the case of voluntary movements.

4. That the brain is the organ principally implicated in several disorders which have been too exclusively referred to the spinal cord.

ART. IV.—*Practical Remarks on the Endemic Fever of Ceylon.*

By J. C. CAMERON, M. D., Staff-Surgeon of the 2d Class.
(Communicated to the Editor by the Director-General of the Army Medical Department.)

THE following details and conclusions are drawn from the experience gathered at the bed-side, during nearly thirteen years' service in this colony, which has led me to abandon the use of bleeding and salivation as the chief remedies in the form of tropical fever prevalent here, and to substitute in their stead the free use of quinine, without any regard to exacerbations, remissions, or local complications.

This practice, although very analogous to that recommended by Clark and Lind in the last century, is, I believe, opposed to the doctrines of the modern schools, as I have generally found my newly-arrived brethren exceedingly surprised by it, and afraid that the worst results would follow the use of quinine during the hot stage of fever. A single trial has, however, generally convinced them of its efficacy in this island. And, in fact, the practice may be considered almost universal in the colony, so that I am induced to submit the results for the information of the profession, in the hopes, considering the general resemblance of tropical disease, that what has been found so great an improvement here, may, upon trial, prove successful