Published in final edited form as:

Schizophr Res. 2018 June ; 196: 29–34. doi:10.1016/j.schres.2017.07.017.

Thought, Language, and Communication Deficits and Association with Everyday Functional Outcomes Among Community-Dwelling Middle-Aged and Older Adults with Schizophrenia

Anjana Muralidharan^{1,2}, Anastasia Finch¹, Christopher R. Bowie^{3,4}, and Philip D. Harvey^{5,6}

¹VA Capitol Healthcare Network, Mental Illness Research Education and Clinical Center, 10 N. Greene Street, Baltimore, MD, 21201, USA

²Department of Psychiatry, University of Maryland School of Medicine, 655 W. Baltimore Street, Baltimore, MD, 21201, USA

³Department of Psychology, Department of Psychiatry, Queen's University, Kingston, ON, K7L3N6, Canada

⁴Center for Addiction and Mental Health, Toronto, ON, Canada

⁵Department of Psychiatry, Leonard M. Miller School of Medicine, 1600 NW 10th Avenue, Miami, FL, 33136, USA

⁶Research Service, Bruce W. Carter VA Medical Center, 1201 NW 16th Street, Miami, FL, 33125, USA

Abstract

Corresponding author: Anjana Muralidharan, Ph.D., VA Capitol Healthcare Network, Mental Illness Research Education and Clinical Center, 10 N. Greene Street, Attn: MIRECC (Annex Bldg.), Baltimore, MD 21201, Phone: 1-410-637-1872, Fax: 1-410-637-1880, anjana.muralidharan2@va.gov.

Role of the Funding Source

This research was funded by NIMH grant MH 63116 to Dr. Harvey and by the U.S. Department of Veterans Affairs VISN 3 MIRECC. The funding sources had no role in the study design, in the collection, analysis and interpretation of data, in the writing of the report, or in the decision to submit the article for publication.

Contributors

Dr. Harvey obtained the research funding and designed the overall study with Dr. Bowie. Dr. Bowie supervised data collection. Dr. Muralidharan conceptualized the research question, conducted data analyses, and oversaw the writing of the manuscript. All four authors contributed to the writing of this manuscript. All authors contributed to and have approved the final manuscript.

Conflict of Interest

Dr. Muralidharan and Dr. Finch have no conflicts of interest to report with regard to this work.

Dr. Harvey has received consulting fees or travel reimbursements from Allergan, Boehringer Ingelheim, Lundbeck Pharma, Minerva Pharma, Otsuka Digital Health, Sanofi Pharma, Sunovion Pharma, and Takeda Pharma during the past year. He has a research grant from Takeda and from the Stanley Medical Research Foundation.

Dr. Bowie has received consulting fees from Boehringer Ingelheim, Lundbeck Pharma, Otsuka Digital Health, and Takeda Pharma. He has received grant support from Lundbeck, Pfizer, and Takeda.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Older adults with schizophrenia experience poorer community integration and social functioning compared to same-age peers with no mental health disorders; these individuals are at elevated risk for functional decline and early institutionalization in long-term care facilities. Deficits in thought, language, and communication (TLC; that is, thought disorder and alogia) are core features of schizophrenia and may worsen with age; however, little research focuses on the functional sequelae of these impairments among older adults with schizophrenia. The present study aimed to examine the relationships among age, TLC deficits, and functional outcomes in a sample of community-dwelling middle-aged and older adults with schizophrenia (N=245; ages 40–85). Participants completed assessments of symptoms, neurocognition, TLC deficits, and functional outcomes. Two different categories of TLC deficits were examined: verbal underproductivity (i.e., alogia) and disconnected speech. Regression analyses, controlling for gender, age, Veteran status, smoking status, cognitive impairment, and symptom severity, found that disconnected speech predicted occupational functioning, while verbal underproductivity predicted capacity to communicate skillfully in semi-structured social situations, as well as community functioning across interpersonal, occupational, and everyday living domains. Exploratory mediation analyses found significant indirect effects of age, through TLC deficits, on certain functional outcomes. Targeted training to improve TLC deficits, especially verbal underproductivity, among older adults with schizophrenia could have downstream effects on community functioning, improving outcomes for a vulnerable group.

Keywords

older adults; schizophrenia; community functioning; functional capacity; thought disorder; alogia

1. Introduction

Community-dwelling older adults with schizophrenia experience poorer community integration and social functioning when compared to their same-age peers, including elevated rates of unemployment, underperformance in everyday activities, and social isolation (Abdallah et al., 2009; Madieto et al., 2012; Meesters et al., 2010). Moreover, older adults with schizophrenia are vastly overrepresented in nursing homes (Miller et al., 2006; Miller et al., 2007) and are at increased risk for nursing home placement as early as their forties (Andrews et al., 2009). This indicates that, although there is great heterogeneity in outcomes as individuals with schizophrenia age (Jeste et al., 2011), for some, aging may be accompanied by faster and more severe functional decline than that experienced by their same-age peers without psychiatric disorders. Further elucidation of the factors which contribute to poor functional outcomes in community-dwelling older adults with schizophrenia is necessary to inform targeted interventions for this group.

While cognitive decline accounts for worsening of functional outcomes in older adults with schizophrenia in some cases (Reichenberg et al., 2014), research indicates that only a subset of these individuals exhibit substantial cognitive decline with age (Harvey et al., 2010; Thompson et al., 2013). Additionally, while considerable research has examined the relationships among neurocognition, symptom severity, and functional outcomes in adults

with schizophrenia (e.g., Bowie et al., 2006), much less attention has focused on one of the cardinal features of the illness: impairments in thought, language, and communication.

Thought, language, and communication (TLC) deficits, also referred to as formal thought disorder and alogia, are common in schizophrenia, and are associated with impairments in everyday functioning (Racenstein et al., 1999; Bowie et al., 2011). TLC deficits are thought to fall into two categories: “positive” thought disorder, characterized by the presence of disorganized, circumstantial, or tangential speech, and “negative” thought disorder (i.e., alogia), characterized by an absence of speech production (Andreasen, 1986). Previous research with chronically institutionalized older patients with schizophrenia, both “positive” TLC deficits (hereafter referred to as disconnected speech) and “negative” TLC deficits (hereafter referred to as verbal underproductivity) were associated with deficits in social outcomes, but not with deficits in other functional domains (e.g., activities of daily living, recreation; Bowie & Harvey, 2008). Additionally, verbal underproductivity worsened with age, especially among the older-old and oldest-old patients (ages 74 and up; Bowie et al., 2005). Importantly, the relationships among age, TLC deficits, and functional outcomes have not been examined in community-dwelling older adults with schizophrenia, for whom adaptive functioning may be much more dependent on the ability to skillfully communicate.

Previous studies on TLC deficits and functional outcomes in schizophrenia have generally relied on consumer or informant reports of everyday functioning (Racenstein et al., 1999; Bowie & Harvey, 2008). One limitation of this approach is that real-world functional performance may be influenced by factors beyond a consumer’s communication abilities, such as socioenvironmental constraints. Therefore, it is unknown whether TLC deficits are associated with poorer functional outcomes because they directly influence a consumer’s *ability* to successfully communicate (i.e., a consumer cannot communicate his/her thoughts or needs when given the opportunity) or because that consumer has *fewer opportunities* to communicate (i.e., TLC deficits lead to increased social isolation and fewer opportunities for interaction). To address this, performance-based measures of functioning, which prompt participants to demonstrate social and occupational skills through in-vivo tasks, may be used. These measures capture *functional capacity*, or their *ability* to accomplish activities of daily living under optimal conditions (Bowie et al., 2006). The only study to examine the association of TLC deficits with functional capacity in adults with schizophrenia, found that both verbal underproductivity and disconnected speech were associated with performance on in-vivo, social skills based role plays (Bowie et al., 2011). These associations have not been examined in an older adult sample of individuals with schizophrenia.

The present study aimed to further elucidate the association of TLC deficits and functional outcomes among community-dwelling middle-aged and older adults with schizophrenia. We aimed to examine, adjusting for demographic factors, cognitive impairment, and symptom severity, whether verbal underproductivity and disconnected speech would be associated with poorer clinician-rated community functioning across occupational, everyday living, and interpersonal domains. Secondly, we aimed to examine, again adjusting for demographic factors, cognitive impairment, and symptom severity, whether verbal underproductivity and disconnected speech would be associated with poorer performance on laboratory social skills tasks and functional tasks of communication. Exploratory analyses tested whether

TLC deficits mediated relationships between age and functional outcomes, to explain the greater functional deficits seen in older people with schizophrenia.

2. Methods

2.1. Participants

This study presents analyses of baseline data from a large longitudinal study, which assessed the course of cognition and functioning in a sample of community-dwelling adults with schizophrenia (see Bowie et al., 2006 for more details on methodology). The present analyses utilized data from all participants with a schizophrenia diagnosis, ages 40 and up, who completed baseline assessments for the study ($n = 245$; see Table 1). Participants were recruited through academic, state, and VA outpatient treatment programs, and were required to meet criteria for a primary DSM-IV diagnosis of schizophrenia or schizoaffective disorder, as assessed through the Comprehensive Assessment of Symptoms and History (Andreasen et al., 1992). Additionally, individuals needed to exhibit evidence of current active illness at baseline assessment, as evidenced by an inpatient hospitalization or emergency room visit for psychosis within the last two years, or at least moderate levels of positive symptoms. Exclusion criteria were the presence of medical illnesses that could impact cognitive functioning, and a Mini-Mental Status Examination (MMSE; Folstein et al., 1975) score below 18.

2.2. Procedures

All study procedures were approved by the appropriate institutional review boards and participants completed written informed consent prior to participating in any study activities. Eligible participants completed assessments of cognition, functional and communication skills, and symptoms. All assessments were completed by trained raters. Participants' case managers completed ratings of community functioning within one week of the assessments.

2.3. Measures

Demographic information and smoking status were ascertained from the participant. Height, weight, and presence of diabetes and heart disease were ascertained by chart review when available.

2.3.1 Symptom Severity—The Positive and Negative Syndrome Scale (PANSS; Kay, 1991) was used as a measure of clinical symptom severity. The PANSS is comprised of 30 individual items that are scored using seven point Likert scales (1= absent, 7= extreme) and was rated through a structured interview with the participant and collateral information (i.e., chart review and clinician informant). The interrater reliability for this measure was high (Bowie & Harvey, 2008). The PANSS Positive and PANSS Negative scale scores were calculated excluding three individual items, due to their high overlap with TLC: conceptual disorganization, lack of spontaneity and flow of conversation, and social withdrawal (Bowie & Harvey, 2008). The widely used and well-validated Beck Depression Inventory (BDI; Beck, Steer, & Brown, 1986) was used as a measure of depression symptom severity.

2.3.2. Global Cognition—Global cognitive functioning was assessed through comprehensive neuropsychological assessment. Age-based norms were used to produce standard scores, and these scores were combined to create a cognitive composite score which included the following domains: constructional praxis, attention and concentration, executive functioning, and verbal learning, memory, and fluency. For a detailed description of the cognitive composite see Bowie et al. (2006).

2.3.3. TLC Deficits—The Thought, Language, and Communications (TLC) Scale is an 18-item interviewer-rated measure of thought disorder, based on participant speech and communication during unscripted conversation (Andreasen, 1986). The TLC scale has been successfully used in previous research to measure two subtypes of TLC deficits: disconnected speech (TLC_DS) and verbal underproductivity (TLC_VU; Bowie et al., 2005; Bowie & Harvey, 2008). TLC_DS is calculated as a mean of a six items representing loose or incoherent speech (e.g., tangentiality, circumstantiality), while TLC_VU was defined by the poverty of speech item. This use of the TLC measure has demonstrated adequate-to-good interrater reliability (Bowie et al., 2005; Bowie & Harvey, 2008; Harvey et al., 1997).

2.3.4. Performance-Based Measures of Functional Capacity—Performance-based functional and social skills were measured using the UCSD Performance-Based Skills Assessment (UPSA; Patterson et al., 2001) and the Social Skills Performance Assessment (SSPA; Patterson et al., 2001). The UPSA was created to assess functional skills among older outpatients with serious mental illness in five domains: communication, finance, transportation, planning recreational activities, and household chores (Bowie et al., 2006). For the purposes of this study, the communication domain (UPSA communication) was utilized as a measure of functional communication abilities. Participants completed a series of task-oriented role plays (e.g., making an emergency phone call) assessing their ability to successfully complete each communication task, producing a communication subscale score, with higher scores indicating better performance.

The SSPA involves a series of role plays performed between the participant and the interviewer. Role plays are less structured and involve unscripted communication with the interviewer in two different “scenes”: greeting a new neighbor and negotiating with a landlord. These scored role plays follow a practice scene where the participant has to make plans with a friend. Role plays are coded on a scale from 1 to 5 (1=low; 5=high) on the following items: interest, fluency, clarity, focus, affect, grooming, negotiation ability, submission, overall argument/overall conversation, and social appropriateness. The mean score (SSPA mean) represents a global estimate of social skill level for each participant. The SSPA has good test-retest and interrater reliability (Patterson et al., 2001).

2.3.5. Community Functioning—Community functioning was assessed through the Specific Level of Function Scale (SLOF; Schneider & Struening, 1983). This measure has high interrater reliability, internal consistency, and factorial validity (Bowie et al., 2006; Harvey et al., 2011). The SLOF contains 43 items rated on a 5-point Likert scale. Ratings are based on informant report of an individual’s performance across six functional domains. The relevant domains for the present analyses were: interpersonal relationships (SLOF interpersonal; e.g., forming friendships), daily activities (SLOF activities; e.g., shopping,

taking medication, handling personal finances), and work skills (SLOF work; e.g., has employable skills). Case managers who had at least a 'very good' (score 4 or 5 on a 5 point scale) knowledge of the participants rated these items within one week of the assessment and they were unaware of any of the other assessment results.

2.4. Data Analysis

First, multiple linear regression analyses were performed to test our primary hypotheses, with the following measures as outcome variables: SLOF work, SLOF activities, SLOF interpersonal, UPSA communication, and SSPA mean. In each regression, gender, Veteran status, and smoking status were entered simultaneously in the first step. Then, age, global cognitive impairment, depression severity, positive symptom severity, and negative symptom severity were entered simultaneously in the next step. Then, TLC_DS and TLC_VU were entered in the final step as predictor variables. Cumulative R-squared statistics were generated to examine the relative contribution of each step in the model. Finally, for each functional outcome which had at least one TLC measure as a significant predictor, an exploratory mediation analysis was conducted using the PROCESS macro for SPSS. For each mediation analysis, gender, Veteran status, smoking status, global cognitive impairment, depression symptom severity, positive symptom severity, and negative symptom severity were entered as covariates in the model, with age as the predictor and TLC_VU and/or TLC_DS as mediators, depending on which TLC measures were significant predictors in the regression model for that functional outcome. PROCESS uses a bootstrapping, regression-based procedure to estimate the magnitude and significance of the indirect effect (Hayes, 2013); in this analysis, if the confidence interval does not include zero, the indirect effect is assumed to be significant. Number of bootstrap samples was set to 5,000.

3. Results

Demographics and descriptive statistics are presented in Table 1. Age ranged from 40 to 85 years. All participants were receiving antipsychotic and/or other psychotropic medication(s). Among participants for whom chart data was available (~70%), the majority of participants were overweight or obese (mean BMI = 30.34 +/- 7.05), 14.2% had Type I or II diabetes, and 5.9% had heart disease. Missing data was less than 10% for all variables in the analyses except the SLOF scale, which was missing for approximately 20% of participants, typically due to lack of availability of or responsiveness from a case manager with sufficient knowledge of the participant. All available data was used for each analysis, resulting in different sample sizes for each analysis. Bivariate correlations among variables are displayed in Table 2; because a number of variables exhibited non-normality, especially the TLC and SLOF scales, Spearman's rho correlations are reported. Older age was associated with worse performance on the UPSA and the SSPA, more disconnected speech, and poorer performance on work and everyday activities. Higher performance on cognitive tests was associated with less verbal underproductivity, better work and everyday activity performance, and better scores on the UPSA and SSPA.

Results from regression analyses are displayed in Table 3. All analyses controlled for gender, Veteran status, smoking status, age, global cognitive impairment, depression symptom severity, positive symptom severity, and negative symptom severity. TLC_DS and TLC_VU were significantly associated with SLOF work, explaining 4.7% additional variance, collectively. TLC_VU was also significantly associated with SLOF activities (explaining 4.8% additional variance), SLOF interpersonal (explaining 4.8% additional variance), and SSPA mean (explaining 3.4% additional variance). All regression models met assumptions of approximately normal distribution of residuals, with no evidence of problematic multicollinearity or autocorrelation.

Exploratory mediation analyses indicated that, controlling for gender, Veteran status, smoking status, age, global cognitive impairment, depression symptom severity, positive symptom severity, and negative symptom severity, TLC_VU mediated the relationships between age and SLOF interpersonal (95% CI = $-.050, -.001$). TLC_DS mediated the relationship between age and SLOF work (95% CI = $-.065, -.002$). The TLC measures were not significant mediators in any other mediation analysis.

4. Discussion

Older adults with schizophrenia were previously reported to exhibit age-related changes in cognition, communication, and everyday functioning. In this sample of community-dwelling middle-aged and older adults with schizophrenia, TLC deficits were associated with poorer functioning across occupational, interpersonal, and everyday living domains, accounting for between 3.4% and 4.8% of the variance in community functioning. This is in contrast to previous work with chronically institutionalized older adults with schizophrenia, which found that TLC deficits were solely associated with social functioning deficits (Bowie & Harvey, 2008). It may be that in a sample that consisted of a chronically institutionalized population, occupational and self-care functioning was generally limited in terms of opportunities; this lack of variance may have precluded significant associations between TLC deficits and functional outcomes. Alternatively, functional outcomes may be less dependent on communication skills for older adults with schizophrenia in long-term care facilities, who have access to structure and support in performing activities of daily living and recreation. In contrast, adaptive functioning for older adults with schizophrenia in the community requires proactive engagement and seeking out of opportunities, which is highly dependent on communication skills. Community-dwelling older adults with schizophrenia could benefit from targeted treatment aimed at improving TLC deficits.

Findings suggest that disconnected speech and verbal underproductivity are associated with functional outcomes in different ways. Disconnected speech was associated with deficits in work skills, but not with deficits in other areas of community functioning. Verbal underproductivity, on the other hand, was associated with deficits across multiple domains of functioning. It makes intuitive sense that reduced verbal output would be associated with decreased success in interpersonal relationships, posing a barrier to initiating and maintaining conversations and friendships; this was in fact, one of the functional domains in which verbal underproductivity accounted for the most variance (4.8%). In addition, verbal underproductivity was associated with poorer performance in semi-structured social

situations (accounting for 3.4% of the variance explained), but not with social skills tasks with specific instructions. Therefore, with structure, older adults with schizophrenia may be able to overcome verbal underproductivity to accomplish specific tasks or activities. However, verbal underproductivity poses a barrier to connecting with others or expressing one's needs in less structured interactions. Verbal underproductivity was also associated with poorer work skills and performance of daily activities; these associations are more difficult to explain. Poverty of speech in and of itself should not prevent an individual from successfully completing work tasks without supervision, for example, or engaging in daily activities such as taking care of household chores or managing his/her personal finances. It may be that verbal underproductivity is associated with general social isolation, which leads to decreased opportunities to engage in activities of daily living and demonstrate functional skills.

Exploratory analyses examined whether TLC deficits mediated the link between older age and poorer functional outcomes. Findings indicated that disconnected speech mediated the link between age and poorer work functioning. In addition, the effect of age on interpersonal functioning was mediated through verbal underproductivity. Given that older age is associated with greater social isolation in the general population (Cattan et al., 2005), and with increased verbal underproductivity with age among adults with schizophrenia (Bowie et al., 2005), the interaction of these factors could lead to even greater social isolation and poorer functional outcomes. Worsening in TLC deficits with age could contribute to functional decline among older adults with schizophrenia. These individuals may benefit from skills training with a focus on decreasing disconnected speech, which could improve performance on work tasks, and increasing verbal output in unstructured social situations, which could have downstream effects across functional domains.

The present study utilized a large sample of an understudied population; namely, community-dwelling middle-aged and older adults with schizophrenia, including older-old individuals in their 70's and 80's. In addition the present study utilized rigorous clinical assessment procedures, and ratings of community functioning were made by independent observers who were not involved in other study assessments. While these are considerable strengths, there are also limitations of note. This is a post-hoc analysis on data which were not collected to test the hypotheses put forth here; thus these findings need to be confirmed in a prospective hypothesis testing study. Data analyses included multiple comparisons, increasing risk for Type I error. In addition, participant scores on the TLC measure were generally low, indicating generally low levels of thought disorder and alogia in the sample. At the same time, participant scores on the community functioning measure were generally high, indicating a relatively high-functioning sample. This restriction of range may have led to an underestimate of the association between TLC deficits and functional outcomes. Furthermore, the sample was predominantly male and restricted to individuals with active illness; thus, the findings may not apply to largely female groups or to asymptomatic individuals with schizophrenia. In addition, we were unable to account for the effects of psychotropic medications on TLC deficits or functional outcomes. We also did not have a measure of family or social support so could not examine the impact of these variables on outcomes. Finally, given that this is a cross-sectional study, causal conclusions regarding the associations among the variables cannot be made. Future studies should examine the impact

of TLC deficits on functioning in a sample with greater variance in TLC deficits, and could utilize a longitudinal research design.

In conclusion, in a sample of community-dwelling middle-aged and older adults with schizophrenia, TLC deficits were associated with poorer community functioning across domains, even when controlling for demographic factors, symptom severity, and cognitive impairment. Disconnected speech was associated with deficits in task-oriented social skills and occupational functioning, while verbal underproductivity was associated with poorer performance in unstructured social interactions, and poorer community functioning across occupational, interpersonal, and recreational domains. Targeted training to improve these deficits among older adults with schizophrenia could improve social skills performance and have downstream effects on community functioning, improving functional outcomes for a vulnerable and complex group.

Acknowledgments

This research was funded by NIMH grant MH 63116 to Dr. Harvey and by the U.S. Department of Veterans Affairs VISN 3 MIRECC. This manuscript is result of work supported with resources and the use of facilities at the U.S. Department of Veterans Affairs VISN 5 MIRECC. All authors who contributed to this paper are listed as authors. No professional medical writer was involved in any portion of the preparation of the manuscript. This work reflects the authors' personal views and in no way represents the official view of the Department of Veterans Affairs or the U.S. Government.

References

- Abdallah C, Cohen CI, Sanchez-Almira M, Reyes P, Ramirez P. Community integration and associated factors among older adults with schizophrenia. *Psychiatr Serv.* 2009; 60(12):1642–1648. [PubMed: 19952155]
- Andreasen NC. Scale for the assessment of thought, language, and communication (TLC). *Schizophr Bull.* 1986; 12(3):473. [PubMed: 3764363]
- Andreasen NC, Flaum M, Arndt S. The Comprehensive Assessment of Symptoms and History (CASH): an instrument for assessing diagnosis and psychopathology. *Arch Gen Psychiatry.* 1992; 49(8):615–623. [PubMed: 1637251]
- Andrews AO, Bartels SJ, Xie H, Peacock WJ. Increased risk of nursing home admission among middle aged and older adults with schizophrenia. *Am J Geriatr Psychiatry.* 2009; 17:697–705. [PubMed: 19625787]
- Beck, A., Steer, R., Brown, GK. *The Beck Depression Inventory.* 2. The Psychological Corporation; San Antonio: 1986.
- Bowie CR, Gupta M, Holshausen K. Disconnected and underproductive speech in schizophrenia: Unique relationships across multiple indicators of social functioning. *Schizophr Res.* 2011; 131:152–156. [PubMed: 21561739]
- Bowie CR, Harvey PD. Communication abnormalities predict functional outcomes in chronic schizophrenia: differential associations with social and adaptive functions. *Schizophr Res.* 2008; 103(1):240–247. [PubMed: 18571378]
- Bowie CR, Reichenberg A, Patterson TL, Heaton RK, Harvey PD. Determinants of real-world functional performance in schizophrenia subjects: correlations with cognition, functional capacity, and symptoms. *Am J Psychiatry.* 2006; 163(3):418–425. [PubMed: 16513862]
- Bowie CR, Tsapelas I, Friedman J, Parrella M, White L, Harvey PD. The longitudinal course of thought disorder in geriatric patients with chronic schizophrenia. *Am J Psychiatry.* 2005; 162(4): 793–795. [PubMed: 15800156]
- Cattan M, White M, Bond J, Learmouth A. Preventing social isolation and loneliness among older people: a systematic review of health promotion interventions. *Ageing Soc.* 2005; 25(1):41–67.

- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975; 12(3):189–198. [PubMed: 1202204]
- Harvey PD, Lombardi J, Leibman M, Parrella M. Age-related differences in formal thought disorder in chronically hospitalized schizophrenic patients: a cross-sectional study across nine decades. *Am J Psychiatry.* 1997; 154(2):205. [PubMed: 9016269]
- Harvey PD, Raykov T, Twamley EW, Vella L, Heaton RK, Patterson TL. Validating the measurement of real-world functional outcomes: Phase I results of the VALERO Study. *Am J Psychiatry.* 2011; 168:1195–1201. [PubMed: 21572166]
- Harvey PD, Reichenberg A, Bowie CR, Patterson TL, Heaton RK. The course of neuropsychological performance and functional capacity in older patients with schizophrenia: Influences of previous history of long-term institutional stay. *Biol Psychiat.* 2010; 67:933–939. [PubMed: 20202624]
- Hayes, AF. *Introduction to Mediation, Moderation, And Conditional Process Analysis: A Regression-Based Approach.* Guilford Press; New York: 2013.
- Jeste DV, Wolkowitz OM, Palmer BW. Divergent Trajectories of Physical, Cognitive, and Psychosocial Aging in Schizophrenia. *Schizophr Bull.* 2011; 37(3):451–455. [PubMed: 21505111]
- Kay, SR. *Positive and Negative Syndromes in Schizophrenia Assessment and Research.* Brunner/Mazel Inc; New York: 1991.
- Madiedo CJ, Garcia-Aracena EF, Ryu HH, Cohen CI. Community integration in older adults with schizophrenia on 4-year follow-up. *Am J Geriatr Psychiatry.* 2012; 20(3):91–92.
- Meesters PD, Stek ML, Comijs HC, de Haan L, Patterson TL, Eikelenboom P, Beekman AT. Social functioning among older community-dwelling patients with schizophrenia: a review. *Am J Geriatr Psychiatry.* 2010; 18(10):862–878. [PubMed: 20808137]
- Miller EA, Rosenheck RA. Mental illness and use of home care nationally in the U.S. Department of Veterans Affairs. *Am J Geriatr Psychiatry.* 2007; 15:1046–1056. [PubMed: 18056823]
- Miller EA, Rosenheck RA. Risk of nursing home admission in association with mental illness nationally in the Department Of Veterans Affairs. *Med Care.* 2006; 44:343–351. [PubMed: 16565635]
- Patterson TL, Goldman S, McKibbin CL, Hughs T, Jeste DV. UCSD Performance-Based Skills Assessment: development of a new measure of everyday functioning for severely mentally ill adults. *Schizophr Bull.* 2001; 27(2):235–245. [PubMed: 11354591]
- Patterson TL, Moscona S, McKibbin CL, Davidson K, Jeste DV. Social skills performance assessment among older patients with schizophrenia. *Schizophr Res.* 2001; 48(2):351–360. [PubMed: 11295387]
- Racenstein JM, Penn D, Harrow M, Schleser R. Thought disorder and psychosocial functioning in schizophrenia: The concurrent and predictive relationships. *J Nerv Ment Dis.* 1999; 187:281–289. [PubMed: 10348082]
- Reichenberg A, Feo C, Prestia D, Bowie CR, Patterson TL, Harvey PD. The course and correlates of everyday functioning in schizophrenia. *Schizophr Res Cogn.* 2014; 1:47–52.
- Schneider LC, Struening EL. SLOF: A behavioral rating scale for assessing the mentally ill. *Soc Work Res Abstr.* 1983; 19(3):9–21. [PubMed: 10264257]
- Thompson WK, Savla GN, Vahia IV, Depp CA, O'Hara R, Jeste DV, Palmer BW. Characterizing trajectories of cognitive functioning in older adults with schizophrenia: does method matter. *Schizophr Res.* 2013; 14:90–96.

Table 1**Demographics and Descriptive Statistics (N=245)**

Variable	N (%)	
Gender		
Male	178 (72.7%)	
Female	64 (26.1%)	
Race/Ethnicity		
White	128 (52.2%)	
African-American	71 (29.0%)	
Other	16 (6.5%)	
Marital Status		
Never married	124 (50.6%)	
Veteran Status		
Yes	94 (38.4%)	
No	151 (61.6%)	
Smokes Cigarettes		
Yes	147 (60.0%)	
No	92 (37.6%)	
Missing	6 (2.4%)	
	M (SD)	Range
Age (years)	56.01 (9.07)	40.00 – 85.00
Education (Years)	12.75 (0.16)	5.00 – 20.00
BDI Total	11.90 (10.11)	0.00 – 50.00
PANSS Positive ^a	11.81 (4.51)	6.00 – 26.00
PANSS Negative ^a	9.88 (3.64)	5.00 – 23.00
Cognitive composite ^b	-1.48 (0.99)	-3.94 – 0.48
TLC_DS ^c	0.27 (0.37)	0.00 – 1.60
TLC_VU ^d	0.39 (0.66)	0.00 – 2.00
SLOF Work ^e	24.17 (5.64)	6.00 – 30.00
SLOF Activities ^e	49.66 (7.28)	24.00 – 55.00
SLOF Interpersonal ^e	29.73 (5.37)	14.00 – 35.00
UPSA Communication ^f	16.77 (5.26)	0.00 – 25.00
SSPA Mean ^g	3.83 (0.75)	1.36 – 5.00

Notes. BDI = Beck Depression Inventory; PANSS = Positive and Negative Syndrome Scale; TLC_DS = Thought, Language, and Communication Scale, Disconnected Speech; TLC_VU = Thought, Language, and Communication Scale, Verbal Underproductivity; SLOF = Specific Levels of Functioning scale; UPSA = University of California San Diego Performance-Based Skills Assessment; SSPA = Social Skills Performance Assessment.

^aPANSS scores do not include conceptual disorganization, lack of spontaneity and flow of conversation, and passive/apathetic social withdrawal items. Higher scores reflect more severe psychopathology.

^bMean of z-scores across a battery of neurocognitive measures.

^cScores can range from 0 to 4, with higher scores reflecting more extreme disconnections in speech.

^dScores can range from 0 to 4, with higher scores reflecting higher verbal underproductivity.

^eSLOF work scores can range from 6 to 30, SLOF Activities scores can range from 11 to 55, SLOF Interpersonal scores can range from 7 to 35. For each domain, lower scores indicate more assistance needed to perform that skill.

^fScores can range from 1 to 25 with lower scores reflecting more impairment in functioning.

^gScores can range from 1 to 5 with lower scores reflecting more difficulties with social skills.

Table 2

Spearman's Rho Bivariate Correlations.

	1	2	3	4	5	6	7	8	9	10	11
1. Age											
2. BDI Total	-.165*										
3. PANSS Positive	-.233**	.214**									
4. PANSS Negative	.030	.135*	.265**								
5. Cognitive composite	-.004	-.048	-.049	-.212*							
6. TLC_VU	.100	.052	-.099	.378**	-.245**						
7. TLC_DS	.341**	-.025	.043	.171**	-.097	-.006					
8. SLOF Work	-.304*	-.080	.108	-.144*	.233**	-.274**	-.279**				
9. SLOF Activities	-.394**	-.033	.162*	-.193**	.377**	-.387**	-.284**	.683**			
10. SLOF Interpersonal	-.114	-.125	.044	-.345**	.131	-.369**	-.216**	.472**	.411**		
11. UPSA Communication	-.254**	-.024	-.046	-.185**	.481**	-.184**	-.282**	.299**	.366**	.142	
12. SSPA Mean	-.172**	.012	.026	-.345**	.360**	-.375**	-.139*	.361**	.386**	.253**	.326**

Notes. BDI = Beck Depression Inventory; PANSS = Positive and Negative Syndrome Scale; TLC_VU = Thought, Language, and Communication Scale, Verbal Underproductivity; TLC_DS = Thought, Language, and Communication Scale, Disconnected Speech; SLOF = Specific Levels of Functioning scale; UPSA = University of California San Diego Performance-Based Skills Assessment; SSPA = Social Skills Performance Assessment

All available data were used for each correlation, resulting in different sample sizes for each correlation, ranging from n= 184 to n=241.

* p < .05;

** p < .01

Table 3

Multiple Regressions: TLC Deficits Predicting Functional Outcomes

Step	β	t	p	R ²	R ² change
<u>SLOF Work (n = 189)</u>					
1.					
	Gender *	.133	2.014	.046	.293
	Veteran status	-.030	-0.408	.683	
	Smoking status	-.109	-1.679	.095	
2.	Age **	-.312	-4.215	<.001	.170
	Cognitive composite	.109	1.642	.102	
	BDI Total	-.098	-1.472	.143	
	PANSS Positive	.103	1.393	.165	
	PANSS Negative	.005	.062	.950	
3.	TLC_DS *	-.205	-2.784	.006	.047
	TLC_VU *	-.182	-2.435	.016	
<u>SLOF Activities (n = 186)</u>					
1.					
	Gender	.012	.191	.849	.420
	Veteran status	-.050	-.743	.459	
	Smoking status	-.002	-.033	.974	
2.	Age **	-.370	-5.461	<.001	.293
	Cognitive composite **	.240	3.965	<.001	
	BDI Total	-.050	-.821	.413	
	PANSS Positive	.154	2.269	.024	
	PANSS Negative	-.004	-.061	.952	
3.	TLC_DS	-.093	-1.387	.167	.048
	TLC_VU **	-.253	-3.709	<.001	
<u>SLOF Interpersonal (n = 189)</u>					
1.					
	Gender	.107	1.558	.121	.234
	Veteran status	.037	.492	.624	
	Smoking status	.012	.178	.859	
2.	Age	-.056	-.724	.470	.166

Step	β	t	p	R ²	R ² change
	Cognitive composite	-.032	-.466	.642	
	BDI Total *	-.138	-2.003	.047	
	PANSS Positive	.143	1.849	.066	
	PANSS Negative *	-.244	-3.062	.003	
3.	TLC_DS	-.082	-1.066	.288	.048
	TLC_VU *	-.255	-3.281	.001	
	<u>UPSA Communication (n = 224)</u>				
1.	Gender	-.012	-.214	.831	.368
	Veteran status	-.084	-1.333	.184	
	Smoking status	-.013	-.236	.813	
2.	Age **	-.262	-4.081	<.001	.313
	Cognitive composite **	.456	7.940	<.001	
	BDI Total	-.008	-.137	.891	
	PANSS Positive	-.067	-1.037	.301	
	PANSS Negative	-.026	-.400	.689	
3.	TLC_DS	-.117	-1.842	.067	.010
	TLC_VU	-.021	-.327	.744	
	<u>SSPA Mean (n = 222)</u>				
1.	Gender	.086	1.485	.139	.358
	Veteran status	-.029	-.451	.652	
	Smoking status *	-.185	-3.259	.001	
2.	Age **	-.269	-4.150	<.001	.277
	Cognitive composite **	.268	4.610	<.001	
	BDI Total	.030	.512	.609	
	PANSS Positive	-.011	-.170	.865	
	PANSS Negative *	-.194	-2.898	.004	
3.	TLC_DS	.020	.309	.758	.034
	TLC_VU *	-.212	-3.243	.001	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Notes: SLOF = Specific Levels of Functioning scale; BDI = Beck Depression Inventory; PANSS = Positive and Negative Syndrome Scale; TLC_VU = Thought, Language, and Communication Scale, Verbal Underproductivity; TLC_DS = Thought, Language, and Communication Scale, Disconnected Speech; UPSA = University of California San Diego Performance-Based Skills Assessment; SSPA = Social Skills Performance Assessment.

* p < .05;
** p < .01
*** p < .001