Published in final edited form as:

Appetite. 2017 August 01; 115: 45–53. doi:10.1016/j.appet.2016.11.015.

Wanting and liking: Separable components in problematic eating behavior?

Sarah E. Polk, Erica M. Schulte, Celina R. Furman, and Ashley N. Gearhardt*

University of Michigan, 530 Church St., Ann Arbor, MI 48103, USA

Abstract

Some individuals may have an addictive-like response to certain foods, possibly contributing to problematic eating. Highly processed foods, with added fats and/or refined carbohydrates, are suggested to be most associated with addictive-like eating. The incentive sensitization theory suggests that wanting (e.g. craving) may drive compulsive drug use rather than liking (e.g. enjoyment), but it is unknown whether highly processed foods elicit similar wanting and liking patterns as drugs of abuse, or whether individual differences exist. The current study examines the association of highly processed foods with craving and liking, and whether these relationships differ by food addiction symptomology, cognitive restraint, or body mass index (BMI). Participants ($n = 216$) reported craving and liking for 35 foods and completed the *Yale Food Addiction Scale* (YFAS) and *Three Factor Eating Questionnaire* (TFEQ). Highly processed foods were craved more overall. Craving of highly processed foods was predicted negatively by restraint and positively by YFAS score. Liking of highly processed foods was predicted negatively by restraint and positively by BMI. In conclusion, craving and liking appear distinct with respect to highly processed foods, and may be influenced by addictive-like eating, cognitive restraint, and BMI. This suggests that the incentive sensitization framework may also be relevant for problematic food consumption, especially for individuals reporting food addiction symptoms.

Keywords

Craving; Liking; Obesity; Restraint; Food addiction

1. Introduction

The prevalence of obesity and eating-related problems worldwide is a matter of scientific and public interest. In the past 30 years, global overweight and obesity rates have risen by 27.5% for adults and 47.1% for children, and the number of overweight and obese individuals is now over 2 billion (The GBD 2013 Obesity Collaboration, 2014). Overweight and obesity are estimated to be a greater health burden than smoking and problem drinking (Jia & Lubetkin, 2010; Sturm, 2002), and are associated with many preventable chronic diseases, such as cardiovascular disease, high blood pressure, type-2 diabetes, and some cancers. In addition to the current threat, the rates of overweight and obesity are rising; in the past 40 years, the increase in mean BMI has accelerated, and global prevalence is now

*Corresponding author. 2268 East Hall, 530 Church St. Ann Arbor, MI 48103, USA. agearhar@umich.edu (A.N. Gearhardt).

twice as high as in 1980 (Malik, Willett, & Hu, 2013; Stevens et al., 2012). In the United States, the country with the highest rates of overweight and obesity, it is estimated that by 2030, 65 million more adults will become obese (Wang, McPherson, Marsh, Gortmaker, & Brown, 2011). These trends have facilitated the need for more in-depth research on obesity, particularly on potential mechanisms of overeating.

1.1. Evidence for “food addiction”

There is growing evidence that certain foods, particularly highly processed foods (i.e. foods high in added refined carbohydrates and/or fat), may trigger biological and behavioral mechanisms in a manner akin to drugs of abuse. In animal studies, it has been shown that rats undergo neurobiological changes in the reward system, such as the down-regulation of dopamine receptors, in response to intermittent sugar binges or heightened consumption of high-sugar, high-fat foods (Avena, Bocarsly, Rada, Kim, & Hoebel, 2008; Avena, Rada, & Hoebel, 2009; Johnson & Kenny, 2010). Further, rats demonstrate elevated motivation to seek out these high-fat, high-sugar foods and experience symptoms of withdrawal (e.g. teeth chattering) when sugar is removed from the diet (Avena, Long, & Hoebel, 2005; Colantuoni et al., 2002, 2001; Lenoir, Serre, Cantin, & Ahmed, 2007).

In humans, the *Yale Food Addiction Scale* (YFAS; Gearhardt, Corbin, & Brownell, 2009; Gearhardt, Corbin, & Brownell, 2016) is currently the only validated tool to operationalize behavioral indicators of addictive-like eating. The measure applies the diagnostic criteria for substance-related and addictive disorders (American Psychiatric Association, 2000; American Psychiatric Association, 2013) to the consumption of highly palatable foods, such as losing control over consumption, or continued use despite negative consequences. Individuals who endorse many behavioral indicators of addictive-like eating on the YFAS show similar biological (Davis et al., 2013; Gearhardt et al., 2011) and behavioral characteristics (Gearhardt, White, Masheb, & Grilo, 2013; Gearhardt et al., 2012, 2011; Meule, Lutz, Voge, & Kübler, 2012; Murphy, Stojcek, & MacKillop, 2014; Pivarunas & Conner, 2015) as individuals with substance-use disorders, for example elevated impulsivity, reward dysfunction, and emotion dysregulation. Individuals with high YFAS scores are more likely to have increased body mass index (BMI), more frequent food binges, and stronger cravings for certain foods (Pursey, Stanwell, Gearhardt, Collins, & Burrows, 2014).

In particular, highly processed foods with added fat and refined carbohydrates, such as pizza, chocolate, cake, and cookies, have been shown to be more associated with problematic, addictive-like eating than foods in their natural form, such as nuts, fruit, and vegetables (Curtis & Davis, 2014; Schulte, Avena, & Gearhardt, 2015). Highly processed foods are also more frequently consumed among individuals reporting YFAS indicators of food addiction (Pursey, Collins, Stanwell, & Burrows, 2015). Additionally, strong cravings (Gilhooly et al., 2007; Ifland et al., 2009; Weingarten & Elston, 1991; White & Grilo, 2005), stress-triggered consumption (Epel, Lapidus, McEwen, & Brownell, 2001; Oliver & Wardle, 1999; Oliver, Wardle, & Gibson, 2000; Zellner et al., 2006), loss of control (Arnold, Kenardy, & Agras, 1992; Vanderlinden, Dalle Grave, Vandereycken, & Noorduyn, 2001; Waters, Hill, & Waller, 1999), and binge eating (Rosen, Leitenberg, Fisher, & Khazam, 1986; Vanderlinden et al., 2001; Yanovski et al., 1992) are all more strongly associated with highly processed foods

than less processed foods. Finally, neuroimaging studies suggest that cues for highly processed foods activate the reward-related systems in humans in a similar manner as cues for drugs of abuse (Tang, Fellows, Small, & Dagher, 2012; Volkow, Wang, Fowler, Tomasi, & Baler, 2011).

1.2. Craving and liking

Given that an addictive-like process may underlie problematic eating behavior for some individuals, mechanisms relevant to addictive disorders may also contribute to “food addiction.” The incentive sensitization theory (Robinson & Berridge, 1993, 2000) is a preeminent framework in the field of addiction research, which posits that “wanting” (e.g. craving, desire, motivation) of the substance, more than “liking” (e.g. enjoyment, pleasure) the effects of the substance, indicator of compulsive and problematic substance use. In other words, in the context of addiction, liking may stay relatively stable, or even decrease, as wanting increases. Thus, it may be especially important to target the biological, behavioral, and psychological underpinnings of wanting, rather than liking, in addiction-focused prevention and intervention efforts. This framework is also applicable to food reward (Berridge, 1996, 2009), and indeed wanting and liking of sucrose have been separated in animal models using pharmaceutical manipulations and hyper-dopaminergic knockout mice, resulting in increased wanting but stable or decreased liking (Berridge & Valenstein, 1991; Peciña, Cagniard, Berridge, Aldridge, & Zhuang, 2003; Smith & Berridge, 2005; Wyvell & Berridge, 2000). Wanting also appears to increase with prolonged sucrose consumption (Avena et al., 2005). While the role of incentive sensitization in humans is less clear, there is research supporting this framework for drug consumption (see review by Leyton, 2007), and the separation of wanting and liking in humans in response to food cues and consumption can be shown both in behavioral studies (Finlayson, King, & Blundell, 2007; Finlayson, King, & Blundell, 2008) and neuroimaging studies (Born et al., 2011; Jiang, Soussignan, Schall, & Royet, 2015).

In the context of food addiction, however, the partitioning of wanting and liking in humans has not been extensively investigated. There is evidence that food addiction is associated with elevated craving for food, which corresponds closely to wanting (Robinson & Berridge, 1993). For example, Meule and Kübler (2012) showed that higher scores on the *YFAS* do predict greater food cravings, but the authors did not distinguish between food types (i.e. high-sugar, high-fat). Similarly, Davis, Levitan, Kaplan, Kennedy, and Carter (2014) showed that a food addiction diagnosis by the *YFAS* predicted greater snack food craving, but this study used a limited range of foods (i.e. candy bars, cookies, potato chips). There is limited research on the association between both craving and liking of highly processed foods and the *YFAS*. Gearhardt, Rizk, and Treat (2014) examined how symptoms of addictive-like eating, as measured by the *YFAS*, may be associated with craving and liking of various foods. While this study was unable to decouple craving and liking, it did show that both were increased for highly processed foods, those which have been most implicated in addictive-like responses (Schulte et al., 2015). However, there were also some limitations to this research, such as the inclusion of only overweight/obese females, which may have constrained the ranges of craving and liking, as well as the range of *YFAS* scores (Gearhardt, Boswell, & White, 2014).

1.3. Cognitive restraint

Restraint may also be associated with differences in craving and liking for food. Although restraint is a key construct in the field of eating (Heatherton, Polivy, & Herman, 1990; Herman & Mack, 1975; Ruderman, 1986), it is still unclear exactly what measures of restraint are assessing. Restraint was originally conceptualized as a biological construct with restrained individuals simply eating less than an individual pre-determined set point (Nisbett, 1972), but measures of restraint have not been consistently associated with either BMI or overall caloric intake (Herman & Mack, 1975; Snoek, van Strien, Janssens, & Egnels, 2008; Stice, Fisher, & Lowe, 2004). More recently, restraint has been proposed to be a cognitive process that reflects a conscious desire to restrict actual food intake (Laessle, Tuschl, Kotthaus, & Prike, 1989). Despite this desire to limit consumption, restrained eaters are more prone to overeat following a stressor or a caloric pre-load, suggesting that cognitive restraint is associated with disinhibited eating (Fedoroff, Polivy, & Herman, 1997; Mills & Palandra, 2008; Ruderman, 1985; Wardle, Steptoe, Oliver, & Lipsey, 2000). However, it's unclear whether cognitive restraint precedes disinhibited eating (Herman & Polivy, 1990; Howard & Porzelius, 1999; Polivy & Herman, 1985), or if cognitive restraint develops as a compensatory mechanism following episodes of disinhibited eating (Grilo & Masheb, 2000; Hilbert et al., 2014; Reas & Grilo, 2007; Spurrell, Wilfley, Tanofsky, & Brownell, 1997). Although understanding the casual role of cognitive restraint in eating problems calls for more longitudinal research, another approach to investigating cognitive restraint is to evaluate whether it shows similar or distinct associations with craving and liking for foods, relative to other eating-related constructs (e.g. food addiction, BMI). To date, there is mixed evidence regarding associations between craving and liking and cognitive restraint. Some studies have found positive associations between craving and cognitive restraint (Fedoroff, Polivy, & Herman, 2003; Hill, Weaver, & Blundell, 1991; Massey & Hill, 2012; Polivy, Coleman, & Herman, 2005), while others have found no associations (Gearhardt, Rizk et al., 2014; Hill et al., 1991; Jauregui-Lobera, Bolaños-Ríos, Valero, & Ruiz Prieto, 2012; Rodin, Mancuso, Granger, & Nelbach, 1991; Weingarten & Elston, 1991). Regarding liking, positive associations (Fedoroff et al., 1997), negative associations (Keskitalo et al., 2008), and no associations with cognitive restraint (Lemmens et al., 2010; Lähteenmäki & Tuorila, 1995) have been found. Thus, the association of cognitive restraint with craving and liking is not fully understood.

1.4. Body Mass Index

Elevated BMI may also be associated with differences in craving and liking. Some studies have found that higher BMI is positively associated craving, particular for high-fat foods (Franken & Muris, 2005; Rissanen et al., 2002; Rodin et al., 1991; White, Whisenhunt, Williamson, Greenway, & Netemeyer, 2002). However, Gearhardt, Rizk et al. (2014) found that in a sample of all overweight/obese individuals, higher BMI was associated with lower craving for fatty foods. The association of BMI with food liking has also not always been consistent; a number of studies have found positive associations between BMI and liking of salty, sweet, and high-fat foods (Cox, Hendrie, & Carty, 2016; Deglaire et al., 2015; Drewnowski, Kurth, Holden-Wiltse, & Saari, 1992; Laurent-Jaccard, De Matteis, Hofstetter, & Schutz, 1994; Mela & Sacchetti, 1991). However, other studies suggest that higher BMI is associated with less liking of certain food types, such as sweet, fatty and salty food (Cox et

al., 1998; Gearhardt, Rizk et al., 2014), and yet another study showed no differences in liking between lean and obese participants (Cox, Perry, Moore, Vallis, & Mela, 1999). Thus, more exploration is needed to understand the association between BMI and craving of liking.

1.5. The current study

In the current study, we aim to further investigate the incentive sensitization model by examining the associations of food craving and liking with food addiction, cognitive restraint, and BMI in a community sample of men and women that range from underweight to obese. Utilizing hierarchical linear modeling, we will examine 1) the association between craving and liking for highly processed versus minimally processed foods at level one, and 2) whether individual characteristics (food addiction symptoms, cognitive restraint, BMI) alter these associations at level two. As highly processed foods (e.g. ice cream, pizza, French fries) relative to minimally processed foods (e.g. nuts, fruits, vegetables) are more strongly implicated in addictive-like eating (Schulte et al., 2015), we predict that craving, more so than liking, will overall be higher for highly processed foods. Further, as the dissociation of wanting and liking is thought to be a feature of addictive processes, we predict that food addiction symptoms will be more strongly associated with food craving than food liking. Finally, we hypothesize that BMI and cognitive restraint will not show this pattern, as they are not measures of addiction.

2. Method and materials

2.1. Participants

Participants ($n = 216$) from the United States were recruited through Amazon Mechanical Turk (MTurk) and completed all measures remotely via online questionnaires. While samples from MTurk are not nationally representative, the participant pool is large and demographically diverse, comparable to traditional convenience samples (Paolacci & Chandler, 2014). Participants were offered \$0.75 for the completion of 20-min task, approximately \$2.25 per hour, higher than the median hourly rate of \$1.38 for MTurk studies (Horton & Chilton, 2010). Participants were excluded from analyses if they provided incorrect answers to catch questions (“Who was the first president of the United States?,” “Have you ever had a fatal heart attack while watching TV?,” “What is $2 + 2$?,” “ $n = 6$ ”), provided impossible data (e.g. height 6’1”, weight 21 lbs.; $n = 3$), or identified with a gender other than male or female or did not provide this information ($n = 4$). The University of Michigan Health and Behavioral Sciences Institutional Review Board approved the current study and written informed consent was obtained from all participants.

Participants were asked to self-report demographic information. Participants’ age ranged from 19 to 75 years old ($M = 38.15$, $SD = 14.02$), 61.6% were female ($n = 133$), and reported ethnicity varied (80.1% Caucasian/White, 3.2% Asian/Pacific Islander, 5.1% Hispanic, 4.6% African-American, 0.5% Arab, 1.4% reported Other, and 5.1% Biracial/Multiracial). BMI was determined by self-reported height and weight. Height and weight may be under-reported and thus bias BMI (Connor Gorber, Tremblay, Moher, & Gorber, 2007; Taylor et al., 2006), an adjustment algorithm was developed by Connor Gorber,

Shields, Tremblay, and McDowell (2008) based on a nationally representative Canadian sample. Analyses of the current study were conducted with both unadjusted and adjusted BMI, and no differences were seen. The adjusted BMI variable is used in the current study. Participants fell into all weight categories ($M = 27.98$, $SD = 6.28$): 2.8% underweight (min = 16.64; $n = 6$), 43.1% normal weight ($n = 93$), 28.7% overweight ($n = 62$), and 25.4% obese (max = 51.95; $n = 55$). The distribution of BMI in this sample was positively skewed (1.07, $SE = 0.17$), thus analyses were performed with the log-transformed BMI data (skewness = 0.47, $SE = 0.17$). No differences were seen, thus for clarity of interpretation, the reported results reflect the non-transformed BMI variable.

2.2. Procedures and assessment measures

2.2.1. Three-Factor Eating Questionnaire—Cognitive restraint was measured using the cognitive restraint of eating subscale of the *Three-Factor Eating Questionnaire* (*TFEQ*), which has excellent internal consistency and test-retest validity (Stunkard & Messick, 1985). Though there are multiple measures of cognitive restraint, we chose to use the *TFEQ*, as it has been suggested as the best measure of cognitive restraint and an appropriate measure of actual restriction of food intake (French, Jeffery, & Wing, 1994; Laessle et al., 1989). The *TFEQ* cognitive restraint scale is comprised of 21 items: 12 true/false statements, eight 4-point scales (i.e. “Never” to “Always”, “Not at all” to “Very much”), and one 6-point scale (0–5). In the current study, scores on the *TFEQ* cognitive restraint scale ranged from 10 to 18 ($M = 13.26$, $SD = 2.10$). Scores were normally distributed.

2.2.2. Yale Food Addiction Scale—The *Yale Food Addiction Scale* (*YFAS*) examines addictive-like eating behavior using the DSM-IV-TR diagnostic criteria for substance dependence. It has been shown to have adequate reliability (Gearhardt et al., 2009) and adequate internal consistency in a diverse sample (Gearhardt, Boswell et al., 2014). The *YFAS* contains 25 items to assess behavioral indicators of addictive-like eating and clinical impairment/distress. The measure can be scored to produce a symptom count, ranging from 0 to 7, or a dichotomous diagnosis indicating whether the individual meets criteria for a diagnosis of food addiction (three or more symptoms and clinical impairment/distress). The current study utilized the symptom count method of scoring the *YFAS*, and participants’ symptoms ranged from 0 to 7 ($M = 1.93$, $SD = 1.56$). In this sample, the *YFAS* symptom count distribution was positively skewed (1.11, $SE = 0.17$), thus *YFAS* symptom count data were log-transformed after adding 1, as symptom count could be 0, and analyses were run with the log-transformed data (skewness = 0.07, $SE = 0.166$). No differences were seen, thus the reported results reflect the non-transformed *YFAS* symptom count variable.

2.2.3. Craving and liking—Craving and liking were assessed for 35 nutritionally diverse foods, systematically selected based on processing, fat, sodium, sugar, carbohydrate, protein, and fiber content. Foods were categorized as processed if they contained added fat and/or refined carbohydrates (e.g. sugar, flour). Eighteen of the selected foods were highly processed, (e.g. chocolate, cake, chips, cheeseburger, pizza, French fries, cookies), and seventeen were minimally processed (e.g. nuts, chicken breast, apple, egg, broccoli).

General Labeled Magnitude Scales (gLMS) were developed to assess craving and liking, based on practices by Bartoshuk et al. (2004) and Kalva, Sims, Puentes, Snyder, and Bartoshuk (2014).

Labeled scales (e.g., Visual Analog Scales) are susceptible to inconsistencies due to differences in experience of participants, for example, someone who has experienced child labor may say that is the most intense pain they had ever endured, while a child might say a skinned knee is the highest pain ever experienced. The *gLMS* aims to “stretch” the labeled scale, such that all ratings are made with the participant’s minimum and maximum experience intensity in mind, which allows for ratings that are not confounded by individual differences to be compared across participants (Bartoshuk et al., 2004).

Craving was defined for the subject as an intense desire or want for something. For *gLMS* training, the participant was asked to think of the strongest craving they had ever experienced of any kind as an anchor for the top of the scale. Then, subjects rated their craving for each of the 35 foods on a scale from 0 (“No craving at all”) to 100 (“Strongest craving you’ve ever experienced of any kind”).

Liking was measured in a similar manner. For *gLMS* training on the hedonic scale, participants were first asked to think of the thing in the world they dislike the most to anchor the bottom of the scale and the thing in the world they like the most to anchor the top. Participants then rated their liking for each of the 35 foods on a scale from –100 (“Greatest dislike”) to 100 (“Greatest like”).

The order in which the craving and liking questions were presented to participants was evenly counterbalanced, such that some participants rated craving first and others reported liking first.

3. Results

3.1. Craving

Hierarchical linear modeling (Raudenbush & Bryk, 2002) with robust standard errors was used to examine associations between processing (dichotomous, such that unprocessed foods were coded as 0, and processed foods were coded as 1) and individual differences (food addiction symptoms, cognitive restraint, BMI) with craving reports for each food. Data were analyzed using HLM7 (Raudenbush, Bryk, & Congdon, 2011) with two-level regression analyses, with participants’ reports of how much they craved the 35 foods set as the outcome, and nutritional components of individual foods at level one, nested within the 216 participants’ characteristics at level two. By using this multi-level analysis, we were able to investigate both the association of food attributes with reported craving, and the influence of individual characteristics on the association of food attributes and craving. That is, we aimed to understand if and how individual characteristics (*YFAS* symptom count, cognitive restraint, BMI) changed the association between processed (versus unprocessed) foods and craving. A significance level of $p < 0.05$ was used in this study. There were no missing cases.

Processing had main effects on the reported craving of a food, as seen in the level-one equation. The intercept for the level-one equation (β_0) represents the model-predicted participant-reported craving for an unprocessed food. The partial slope can be interpreted as the influence of processing (β_1) on reported craving (Table 1).

Level-One Equation for Processing as a Predictor of a Reported Craving of a Food:

$$Craving_{ij} = \beta_{0j} + \beta_{1j} * (Processing_{ij}) + r_{ij}$$

Chi-square tests revealed significant variation across participants reported craving focusing on processing, $\chi^2(212) = 246.83$, $p = 0.050$. Thus, food-specific predictors of the intercept and processing were examined and both treated as random effects. Within level-one regression analyses, processing was seen as a large, positive predictor for craving in the average individual ($d = 0.711$). That is, processed foods on average were given higher craving ratings ($M = 39.04$) than unprocessed foods ($M = 35.58$).

YFAS symptom count (centered), cognitive restraint (centered), and adjusted BMI (centered) were then entered into this model as level-two predictors to examine whether individual characteristics alter the association between processing and craving ratings. Though there were significant correlations between *YFAS* symptom count and BMI ($r = 0.233$), and *YFAS* symptom count and cognitive restraint ($r = 0.148$), these correlations were not strong enough to warrant multicollinearity concerns. For level-two equations, the intercept indicates the average value of the relevant level-one variable, assuming mean values for all level-two parameters. For example, γ_{10} indicates the average impact of an unprocessed food on craving ratings for a participant with average scores on *YFAS* symptom count, cognitive restraint, and BMI. The partial slopes within the level-two equations represent the association between a one-unit increase in a level-two predictor and the effect of the level-one predictor on craving. For example, the regression coefficient for the effect of *YFAS* symptom count (γ_{11}) represents the model-predicted impact of a one-unit increase in *YFAS* symptom count on the association between processing and craving.

Level-Two Equations for Participant-Specific Predictors of Level-One Parameters:

$$\begin{aligned}\beta_{0j} &= \gamma_{00} + \gamma_{01} * (YFAS\ Symptom\ Count_j) + \gamma_{02} * (Cognitive\ Restraint_j) + \gamma_{03} * (BMI_j) + u_{0j} \\ \beta_{1j} &= \gamma_{10} + \gamma_{11} * (YFAS\ Symptom\ Count_j) + \gamma_{12} * (Cognitive\ Restraint_j) + \gamma_{13} * (BMI_j) + u_{1j}\end{aligned}$$

An average participant, defined as a participant with mean values on all level-two parameters, reported an average craving of 37.33, on a scale from 0 (“No craving at all”) to 100 (“Strongest craving every experienced of any kind”), for an unprocessed food item (y_{00}).

With respect to individual characteristics, two patterns emerged. *YFAS* symptom count was a small, positive predictor of the association between reported craving and processing ($d = 0.297$); as *YFAS* symptom count increased, craving for processed foods increased compared to craving for unprocessed foods. Cognitive restraint was a moderate, negative predictor of

the association between reported craving and processing ($d = -0.323$), meaning that as cognitive restraint increased, craving for processed foods decreased moderately compared to craving for unprocessed foods. BMI did not significantly alter the association between craving and processing.

3.2. Liking

Hierarchical linear modeling with robust standard errors was also used to examine associations between processing and individual differences with liking reports for each food.

The intercept for the level-one equation (β_0) represents the model-predicted participant-reported liking for an unprocessed food. The partial slope can be interpreted as the influence of processing (β_1) on reported liking (Table 2).

Level-One Equation for Processing as a Predictor of a Reported Craving of a Food:

$$Liking_{ij} = \beta_{0j} + \beta_{1j} * (Processing_{ij}) + r_{ij}$$

Chi-square tests revealed significant variation across participants' reported liking based on level of processing, $\chi^2(212) = 252.56, p = 0.029$. Therefore, food-specific predictors of the intercept and processing were examined and both treated as random effects. On average, processing was not significantly associated with liking ratings ($p = 0.091$).

YFAS symptom count, cognitive restraint, and BMI were entered into the model as level-two predictors to examine changes in the association between processing and liking ratings based on participant-specific characteristics.

Level-Two Equations for Participant-Specific Predictors of Level-One Parameters.

$$\begin{aligned}\beta_{0j} &= \gamma_{00} + \gamma_{01} * (YFAS\ Symptom\ Count_j) + \gamma_{02} * (Cognitive\ Restraint_j) + \gamma_{03} * (BMI_j) + u_{0j} \\ \beta_{1j} &= \gamma_{10} + \gamma_{11} * (YFAS\ Symptom\ Count_j) + \gamma_{12} * (Cognitive\ Restraint_j) + \gamma_{13} * (BMI_j) + u_{1j}\end{aligned}$$

An average participant reported a mean liking of 32.25, on a scale from -100 ("Greatest dislike") to 100 ("Greatest like"), for an unprocessed food item (γ_{00}). For unprocessed foods, YFAS symptom count, cognitive restraint, and BMI did not alter the association between processing and liking.

Two patterns emerged during this analysis. Cognitive restraint was a moderate, negative predictor of the association between reported liking and processing ($d = -0.330$); as cognitive restraint increased, liking for processed foods decreased moderately compared to liking for unprocessed foods. BMI was a small, positive predictor of the association between reported liking and processing ($d = 0.282$); as BMI increased, liking for processed foods increased slightly compared to liking for unprocessed foods. YFAS symptom count did not significantly impact liking ratings for processed foods.

4. Discussion

This study examined the effects of individual differences (food addiction symptomology, cognitive restraint, and BMI) on craving and liking of highly processed versus minimally processed foods. In the current sample, craving was elevated for highly processed foods on average, and this association increased with *YFAS* symptom count. The association between liking and processing was weaker, however, only trending toward significance ($p = 0.091$). Individuals with higher cognitive restraint indicated lower craving and liking for highly processed foods. Finally, for individuals with elevated BMI, craving of highly processed foods remained stable and liking increased.

4.1. Food processing

In the present study, highly processed foods were generally related to elevated craving, but not significantly associated with elevated liking. Studies have consistently found associations between craving and food processing (Gilhooly et al., 2007; Ifland et al., 2009; Weingarten & Elston, 1991; White & Grilo, 2005; White et al., 2002); the positive association between craving and processing in this study corroborates these previous findings. However, the finding that elevated liking was not significantly associated with processing was surprising, given that it is generally assumed that processed foods are liked more than non-processed foods. Viewing highly processed foods seems to elicit reward responses (Simmons, Martin, & Barsalou, 2005), and foods high in sugar and fat have been associated with higher self-reported liking in some studies (Cox et al., 2016; Finlayson et al., 2007; Warwick & Schiffman, 1990). However, there is also prior research that is consistent with the current findings; one study found no difference in liking of processed versus non-processed foods (Cox et al., 1999), and another found a negative association (Cox et al., 1998). Further research should continue investigating the contribution of food processing to altered craving and liking of certain foods in the general population.

4.2. Addictive-like eating

The association between highly processed foods and increased craving was significantly stronger for individuals who exhibited indicators of food addiction measured by the *YFAS*, relative to individuals without food addiction symptomology. Additionally, endorsement of addictive-like eating indicators on the *YFAS* was not related to liking of highly processed foods. This pattern of elevated craving, but not liking, suggests that the incentive sensitization framework of addictive disorders (Robinson & Berridge, 1993, 2000) may be relevant for problematic consumption of highly processed foods, particularly for individuals endorsing *YFAS* indicators of addictive-like eating. Further, *YFAS* food addiction indicators were not related to craving or liking for minimally processed foods. In substance-use disorders, there is an interaction between the addictive substance and an individual's propensity to develop an addictive-like response (Everitt & Robbins, 2005; Koob & Le Moal, 2005; Volkow & Morales, 2015). Similarly, the present results support a substance-based, food addiction framework, where highly processed foods appear to interact with individual characteristics (e.g. addictive-like eating) to differentially activate addictive mechanisms (e.g. craving). As such, it may be that highly processed foods have a central

role in activating an addictive-like process and contributing to problematic eating behavior in certain individuals, though future research is needed.

The present findings are inconsistent with the only previous study examining food craving and liking and their associations with *YFAS* symptoms of food addiction. Using a similar paradigm as the one in the current study, Gearhardt, Rizk et al. (2014) were not able to separate craving and liking in the overall sample or among individuals with elevated reports of *YFAS* food addiction symptomology. However, their sample may have had limited generalizability, given the exclusion of men and healthy weight individuals, and the limited range of reported food addiction symptoms. Thus, the present sample may have been more sufficiently powered to detect the observed patterns of craving and liking, as well as individual differences. Another advantage to the current study was the use of the *gLMS* (Bartoshuk et al., 2004; Kalva et al., 2014), which may be more suitable than a 7-point Likert scale for differentiating individuals' food experiences, especially across groups.

In light of the current findings, it is important to note that food liking is dynamic, changing over time, and may decrease as individuals eat more rewarding foods (Burger & Stice, 2012). The incentive sensitization framework clarifies that rewarding substances (i.e. drugs) are often initially consumed for the pleasurable effects, but during a transition to addiction, these pleasurable effects are diminished, and the substances are reportedly liked less (Robinson & Berridge, 2008). Some evidence has also been presented supporting a similar pattern of change in food liking; Burger and Stice (2012) found that individuals who more frequently consumed ice cream exhibited reduced reward-region (e.g. striatum) responsivity, while reported craving of ice cream increased. Thus the frequency of consumption of certain foods may influence the relationship between craving and liking over time.

4.3. Cognitive restraint

Individuals reporting elevated cognitive restraint exhibited a different pattern of craving and liking than persons with greater reported indicators of addictive-like eating. In the current sample, increased cognitive restraint, as measured by the *TFEQ* (e.g. successful limitation of food consumption; Stunkard & Messick, 1985), was associated with lower levels of craving and liking for processed foods. However, this contrasts with previous research in healthy participants, which found positive associations between craving/liking and cognitive restraint (Komatsu & Aoyama, 2014; Polivy et al., 2005). One possibility for these discrepant findings is that the association between cognitive restraint and craving/liking may differ based on exposure to other factors. For example, pre-meal/post-meal consumption, high stress, and exposure to food cues or caloric preloading are all known to affect restrained eaters differently than unrestrained eaters (Born et al., 2011; Fedoroff et al., 1997; Rotenberg & Flood, 2000; Wardle et al., 2000). Thus, it is possible that different associations between cognitive restraint and food craving/liking would emerge in response to exposure to other conditions. Future studies are needed to examine the effects of certain states on craving and liking in individuals exhibiting higher levels of cognitive restraint.

4.4. Body Mass Index

For highly processed foods, the associations of craving and liking with BMI were different than their associations with *YFAS* symptom count or cognitive restraint. For individuals with higher BMIs, craving ratings did not differ, but liking ratings were more positive. Thus, the current study suggests that liking may be more strongly associated with BMI than craving for highly processed foods. The present finding that BMI was not related to craving of highly processed foods was unexpected, given evidence in prior work for increased craving associated with higher BMI (Franken & Muris, 2005; Rodin et al., 1991; White et al., 2002). However, as *YFAS* symptom count was included in the model with BMI in the current study, it is possible that an addictive-like eating phenotype is more strongly associated with elevated craving than BMI. However, higher BMI appears to be more closely related to liking than food addiction symptoms. Thus, individuals with elevated BMI who do not endorse food addiction symptoms may benefit more from interventions that target food liking, rather than craving.

4.5. Limitations

The current study used a cross-sectional design, thus only correlational relationships between the variables may be inferred. Future research is needed to explore how these relationships may develop, and how well these patterns can predict eating behavior. The current participants were recruited from MTurk, samples from which can be used in the place of traditional convenience samples (Paolacci & Chandler, 2014), however it is not nationally representative, thus generalizations should be made cautiously. Within the current sample, Hispanic and African-American populations were underrepresented, and Caucasian/White and multi-/biracial populations and females were overrepresented (United States QuickFacts, 2015). All measures were collected through self-report, thus it will be important to examine the behavioral and biological correlates of craving and liking and individual differences. Finally, participants were not asked to weigh themselves before self-reporting on their weight, which may have led to inaccurate reports. Future research would benefit from the direct measurement of height and weight.

5. Conclusion

Craving and liking of certain foods and food types, especially highly processed foods, may contribute to problematic eating behaviors. This study investigated changes in craving and liking associated with food addiction, cognitive restraint, and BMI as individual factors, and how these associations may change for highly processed foods versus minimally processed foods. Different patterns emerged for all three individual factors, adding to the evidence suggesting that they are separate mechanisms, each of which affect eating behaviors differently. This study was also able to disentangle craving and liking of food within a wide range of participants. Most notably, increased reports of *YFAS* indicators of food addiction were positively associated with craving, but not liking, for highly processed foods. This provides evidence that the incentive sensitization model may be relevant for “food addiction” and elevated craving for highly processed foods may contribute to compulsive consumption in a similar manner as in compulsive drug use. Further, the current findings support previous research that suggests highly processed foods may be most implicated in

addictive-like eating behavior (Schulte et al., 2015), which may warrant the term “food addiction” to be refined to “highly processed food addiction.”

Acknowledgments

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 42000. text rev.<http://dx.doi.org/10.1176/appi.books.9780890423349>
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 52013. <http://dx.doi.org/10.1176/appi.books.9780890425596>
- Arnow B, Kenardy J, Agras WS. Binge eating among the obese: A descriptive study. *Journal of Behavioral Medicine*. 1992; 15:155–170. <http://dx.doi.org/10.1002/eat.20352>. [PubMed: 1583679]
- Avena NM, Bocarsly ME, Rada P, Kim A, Hoebel BG. After daily bingeing on a sucrose solution, food deprivation induces anxiety and accumbens dopamine/acetylcholine imbalance. *Physiology & Behavior*. 2008; 94:309–315. <http://dx.doi.org/10.1016/j.physbeh.2008.01.008>. [PubMed: 18325546]
- Avena NM, Long KA, Hoebel BG. Sugar-dependent rats show enhanced responding for sugar after abstinence: Evidence of a sugar deprivation effect. *Physiology & Behavior*. 2005; 84:359–362. <http://dx.doi.org/10.1016/j.physbeh.2008.01.008>. [PubMed: 15763572]
- Avena NM, Rada P, Hoebel BG. Sugar and fat bingeing have notable differences in addictive-like behavior. *The Journal of Nutrition*. 2009; 139:623–628. <http://dx.doi.org/10.3945/jn.108.097584>. [PubMed: 19176748]
- Bartoshuk LM, Duffy VB, Green BG, Hoffman HJ, Ko CW, Lucchina LA, ... Weiffenbach JM. Valid across-group comparisons with labeled scales: The gLMS versus magnitude matching. *Physiology & Behavior*. 2004; 82:109–114. <http://dx.doi.org/10.1016/j.physbeh.2004.02.033>. [PubMed: 15234598]
- Berridge KC. Food reward: Brain substrates of wanting and liking. *Neuroscience Biobehavioral Review*. 1996; 20:1–25. [http://dx.doi.org/10.1016/0149-7634\(95\)00033-B](http://dx.doi.org/10.1016/0149-7634(95)00033-B).
- Berridge KC. ‘Liking’ and ‘wanting’ food rewards: Brain substrates and roles in eating disorders. *Physiology & Behavior*. 2009; 97:537–550. <http://dx.doi.org/10.1016/j.physbeh.2009.02.044>. [PubMed: 19336238]
- Berridge KC, Valenstein ES. What psychological process mediates feeding evoked by electrical stimulation of the lateral hypothalamus. *Behavioral Neuroscience*. 1991; 105:3–14. <http://dx.doi.org/10.1037/0735-7044.105.1.3>. [PubMed: 2025391]
- Born JM, Lemmens SG, Martens MJ, Formisano E, Goebel R, Westerterp-Plantenga MS. Differences between liking and wanting signals in the human brain and relations with cognitive dietary restraint and body mass index. *American Journal of Clinical Nutrition*. 2011; 94:392–403. <http://dx.doi.org/10.3945/ajcn.111.012161>. [PubMed: 21653801]
- Burger KS, Stice E. Frequent ice cream consumption is associated with reduced striatal response to receipt of an ice cream-based milkshake. *American Journal of Clinical Nutrition*. 2012; 95:810–817. <http://dx.doi.org/10.3945/ajcn.111.027003>. [PubMed: 22338036]
- Colantuoni C, Rada P, McCarthy J, Patten C, Avena NM, Chadeayne A, et al. Evidence that intermittent, excessive sugar intake causes endogenous opioid dependence. *Obesity Research*. 2002; 10:478–488. <http://dx.doi.org/10.1038/oby.2002.66>. [PubMed: 12055324]
- Colantuoni C, Schwenker J, McCarthy J, Rada P, Ladenheim B, Cadet JL, et al. Excessive sugar intake alters binding to dopamine and mu-opioid receptors in the brain. *Neuroreport*. 2001; 12:3549–3552. <http://dx.doi.org/10.1097/00001756-200111160-00035>. [PubMed: 11733709]

- Connor Gorber S, Shields M, Tremblay MS, McDowell I. The feasibility of establishing correction factors to adjust self-reported estimates of obesity. *Health Reports*. 2008; 19:71–82.
- Connor Gorber S, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: A systematic review. *Obesity Review*. 2007; 8:307–326. <http://dx.doi.org/10.1111/j.1467-789X.2007.00347.x>.
- Cox DN, Hendrie GA, Carty D. Sensitivity, hedonics and preferences for basic tastes and fat amongst adults and children of differing weight status: A comprehensive review. *Food Quality and Preference*. 2016; 48:359–367. <http://dx.doi.org/10.1016/j.foodqual.2015.01.006>.
- Cox DN, Perry L, Moore PB, Vallis L, Mela DJ. Sensory and hedonic associations with macronutrient and energy intakes of lean and obese consumers. *International Journal of Obesity & Related Metabolic Disorders*. 1999; 23:403–411. <http://dx.doi.org/10.1038/sj.ijo.0800836>. [PubMed: 10340819]
- Cox DN, Van Galen M, Hedderley D, Perry L, Moore PB, Mela DJ. Sensory and hedonic judgments of common foods by lean consumers and consumers with obesity. *Obesity Research*. 1998; 6:438–447. <http://dx.doi.org/10.1002/j.1550-8528.1998.tb00376.x>. [PubMed: 9845234]
- Curtis C, Davis C. A qualitative study of binge eating and obesity from an addiction perspective. *Eat Disord*. 2014; 22:19–32. <http://dx.doi.org/10.1080/10640266.2014.857515>. [PubMed: 24365525]
- Davis C, Levitan RD, Kaplan AS, Kennedy JL, Carter JC. Food cravings, appetite, and snack-food consumption in response to a psychomotor stimulant drug: The moderating effect of “food-addiction”. *Frontiers in Psychology*. 2014; 5:1–8. <http://dx.doi.org/10.3389/fpsyg.2014.00403>. [PubMed: 24474945]
- Davis C, Loxton NJ, Levitan RD, Kaplan AS, Carter JC, Kennedy JL. Food addiction and its association with a dopaminergic multilocus genetic profile. *Physiology & Behavior*. 2013; 149:63–69. <http://dx.doi.org/10.1016/j.physbeh.2013.05.014>.
- Deglaire A, Méjean C, Castetbon K, Kesse-Guyot E, Hercberg S, Schlich P. Associations between weight status and liking scores for sweet, salt and fat according to the gender in adults (The Nutrinet-Santé study). *European Journal of Clinical Nutrition*. 2015; 69:40–46. <http://dx.doi.org/10.1038/ejcn.2014.139>. [PubMed: 25074389]
- Drewnowski A, Kurth C, Holden-Wiltse J, Saari J. Food preferences in human obesity: Carbohydrates versus fats. *Appetite*. 1992; 18:207–221. [http://dx.doi.org/10.1016/0195-6663\(92\)90198-F](http://dx.doi.org/10.1016/0195-6663(92)90198-F). [PubMed: 1510463]
- Epel E, Lapidus R, McEwen B, Brownell K. Stress may add bite to appetite in women: A laboratory study of stress-induced cortisol and eating behavior. *Psychoneuroendocrinology*. 2001; 1:37–49. [http://dx.doi.org/10.1016/S0306-4530\(00\)00035-4](http://dx.doi.org/10.1016/S0306-4530(00)00035-4).
- Everitt BJ, Robbins TW. Neural systems of reinforcement for drug addiction: From actions to habits to compulsion. *Nature Neuroscience*. 2005; 8:1481–1489. <http://dx.doi.org/10.1038/nn1579>. [PubMed: 16251991]
- Fedoroff IC, Polivy J, Herman CP. The effect of pre-exposure to food cues on the eating behavior of restrained and unrestrained eaters. *Appetite*. 1997; 28:33–47. <http://dx.doi.org/10.1006/appe.1996.0057>. [PubMed: 9134093]
- Fedoroff IC, Polivy J, Herman CP. The specificity of restrained versus unrestrained eaters’ responses to food cues: General desire to eat, or craving for the cued food? *Appetite*. 2003; 41:7–13. [http://dx.doi.org/10.1016/S0195-6663\(03\)00026-6](http://dx.doi.org/10.1016/S0195-6663(03)00026-6). [PubMed: 12880616]
- Finlayson G, King N, Blundell JE. Liking vs. wanting food: Importance for human appetite control and weight regulation. *Neuroscience and Biobehavioral Reviews*. 2007; 31:987–1002. <http://dx.doi.org/10.1016/j.neubiorev.2007.03.004>. [PubMed: 17559933]
- Finlayson G, King N, Blundell JE. The role of implicit wanting in relation to explicit liking and wanting for food: Implications for appetite control. *Appetite*. 2008; 50:120–127. <http://dx.doi.org/10.1016/j.appet.2007.06.007>. [PubMed: 17655972]
- Franken IHA, Muris P. Individual differences in reward sensitivity are related to food craving and relative body weight in healthy women. *Appetite*. 2005; 45:198–201. <http://dx.doi.org/10.1016/j.appet.2005.04.004>. [PubMed: 15949869]
- French SA, Jeffery RW, Wing RR. Food intake and physical activity: A comparison of three measures of dieting. *Addictive Behaviors*. 1994; 19:401–409. [PubMed: 7992675]

- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., ... Gakidou, E. The GBD 2013 Obesity Collaboration. Global, regional and national prevalence of overweight and obesity in children and adults 1980–2013: A systematic analysis; Lancet (London, England). 2014. p. 384 [http://dx.doi.org/10.1016/S0140-6736\(14\)60460-8](http://dx.doi.org/10.1016/S0140-6736(14)60460-8)
- Gearhardt AN, Boswell RG, White MA. The association of “food addiction” with disordered eating and body mass index. *Eating Behaviors*. 2014; 15:427–433. <http://dx.doi.org/10.1016/j.eatbeh.2014.05.001>. [PubMed: 25064294]
- Gearhardt AN, Corbin WR, Brownell KD. Preliminary validation of the Yale food addiction scale. *Appetite*. 2009; 52:430–436. <http://dx.doi.org/10.1016/j.appet.2008.12.003>. [PubMed: 19121351]
- Gearhardt AN, Corbin WR, Brownell KD. Development of the Yale food addiction scale version 2.0. *Psychology of Addictive Behaviors*. 2016; 30:113–121. <http://dx.doi.org/10.1037/adb0000136>. [PubMed: 26866783]
- Gearhardt AN, Rizk MT, Treat TA. The association of food characteristics and individual differences with ratings of craving and liking. *Appetite*. 2014; 79:166–173. <http://dx.doi.org/10.1016/j.appet.2014.04.013>. [PubMed: 24768936]
- Gearhardt AN, White MA, Masheb RM, Grilo CM. An examination of food addiction in a racially diverse sample of obese patients with binge eating disorder in primary care settings. *Comprehensive Psychiatry*. 2013; 54:500–505. <http://dx.doi.org/10.1016/j.comppsy.2012.12.009>. [PubMed: 23332551]
- Gearhardt AN, White MA, Masheb RM, Morgan PT, Crosby RD, Grilo CM. An examination of the food addiction construct in obese patients with binge eating disorder. *International Journal of Eating Disorders*. 2012; 45:657–663. <http://dx.doi.org/10.1002/eat.20957>. [PubMed: 22684991]
- Gearhardt AN, Yokum S, Orr PT, Stice E, William R, Brownell KD. Neural correlates of food addiction. *Archives of General Psychiatry*. 2011; 68:808–816. <http://dx.doi.org/10.1001/archgenpsychiatry.2011.32>. [PubMed: 21464344]
- Gilhooly CH, Das SK, Golden JK, McCrory MA, Dallal GE, Saltzman E, et al. Food cravings and energy regulation: The characteristics of craved foods and their relationship with eating behaviors and weight change during 6 months of dietary energy restriction. *International Journal of Obesity*. 2007; 31:1849–1858. <http://dx.doi.org/10.1038/sj.ijo.0803672>. [PubMed: 17593902]
- Grilo CM, Masheb RM. Onset of dieting vs binge eating in outpatients with binge eating disorder. *International Journal of Obesity*. 2000; 24:404–409. <http://dx.doi.org/10.1038/sj.ijo.0801171>. [PubMed: 10805495]
- Heatherton TF, Polivy J, Herman CP. Dietary restraint: Some current findings and speculations. *Psychology of Addictive Behavior*. 1990; 4:100–106. <http://dx.doi.org/10.1037/h0080580>.
- Herman CP, Mack D. Restrained and unrestrained eating. *Journal of Personality*. 1975; 43:647–666. <http://dx.doi.org/10.1111/j.1467-6494.1975.tb00727.x>. [PubMed: 1206453]
- Herman CP, Polivy J. From dietary restraint to binge eating: Attaching causes to effects. *Appetite*. 1990; 14:123–125. [http://dx.doi.org/10.1016/0195-6663\(90\)90009-W](http://dx.doi.org/10.1016/0195-6663(90)90009-W). [PubMed: 2337336]
- Hilbert A, Pike K, Goldschmidt A, Wilfley D, Fairburn C, Dohm F, et al. Risk factors across the eating disorders. *Psychiatry Research*. 2014; 220:500–506. <http://dx.doi.org/10.1016/j.psychres.2014.05.054>. [PubMed: 25103674]
- Hill AJ, Weaver CF, Blundell JE. Food craving, dietary restraint and mood. *Appetite*. 1991; 17:187–197. [http://dx.doi.org/10.1016/0195-6663\(91\)90021-J](http://dx.doi.org/10.1016/0195-6663(91)90021-J). [PubMed: 1799281]
- Horton, JJ., Chilton, LB. The labor economics of paid crowdsourcing. *Proceedings of the 11th ACM Conference on Electronic Commerce*; 2010.
- Howard CE, Porzelius LK. The role of dieting in binge eating disorder: Etiology and treatment implications. *Clinical Psychology Review*. 1999; 19:25–44. [http://dx.doi.org/10.1016/S0272-7358\(98\)00009-9](http://dx.doi.org/10.1016/S0272-7358(98)00009-9). [PubMed: 9987582]
- Ifland JR, Preuss HG, Marcus MT, Rourke KM, Taylor WC, Burau K, et al. Refined food addiction: A classic substance use disorder. *Medical Hypotheses*. 2009; 72:518–526. <http://dx.doi.org/10.1016/j.mehy.2008.11.035>. [PubMed: 19223127]
- Jáuregui-Lobera I, Bolaños-Ríos P, Valero E, Ruiz Prieto I. Induction of food craving experience: The role of mental imagery, dietary restraint, mood and coping strategies. *Nutricion hospitalaria*. 2012; 27:1928–1935. <http://dx.doi.org/10.3305/nh.2012.27.6.6043>.

- Jia H, Lubetkin EI. Trends in quality-adjusted life-years lost contributed by smoking and obesity. *American Journal of Preventive Medicine*. 2010; 38:138–144. <http://dx.doi.org/10.1016/j.amepre.2009.09.043>. [PubMed: 20117569]
- Jiang T, Soussignan R, Schall B, Royet JP. Reward for food odors: An fMRI study of liking and wanting as a function of metabolic state and BMI. *Social Cognitive and Affective Neuroscience*. 2015; 10:561–568. <http://dx.doi.org/10.1093/scan/nsu086>. [PubMed: 24948157]
- Johnson PM, Kenny PJ. Dopamine D2 receptors in addiction-like reward dysfunction and compulsive eating in obese rats. *Nature Neuroscience*. 2010; 13:635–641. <http://dx.doi.org/10.1038/nn.2519>. [PubMed: 20348917]
- Kalva JJ, Sims CA, Puentes LA, Snyder DJ, Bartoshuk LM. Comparison of the hedonic general labeled magnitude scale with the hedonic 9-Point scale. *Journal of Food Science*. 2014; 79:238–245. <http://dx.doi.org/10.1111/1750-3841.12342>.
- Keskitalo K, Tuorila H, Spector TD, Cherkas LF, Knaapila A, Kaprio J, ... Perola M. The three-factor eating questionnaire, body mass index, and responses to sweet and salty fatty foods: A twin study of genetic and environmental associations. *The American Journal of Clinical Nutrition*. 2008; 88:263–271. [PubMed: 18689360]
- Komatsu S, Aoyama K. Food craving and its relationship with restriction and liking in Japanese females. *Foods*. 2014; 3:208–216. <http://dx.doi.org/10.3390/foods3020208>. [PubMed: 28234314]
- Koob GF, Le Moal M. Plasticity of reward neurocircuitry and the ‘dark side’ of drug addiction. *Nature Neuroscience*. 2005; 8:1442–1444. <http://dx.doi.org/10.1038/nn1105-1442>. [PubMed: 16251985]
- Laessle RG, Tuschl RJ, Kotthaus BC, Prike KM. A comparison of the validity of three scales for the assessment of dietary restraint. *Journal of Abnormal Psychology*. 1989; 98:504–507. <http://dx.doi.org/10.1037/0021-843X.98.4.504>. [PubMed: 2592686]
- Lähteenmäki L, Tuorila H. Three-factor eating questionnaire and the use and liking of sweet and fat among dieters. *Physiology & Behavior*. 1995; 57:81–88. [http://dx.doi.org/10.1016/0031-9384\(94\)00210-V](http://dx.doi.org/10.1016/0031-9384(94)00210-V). [PubMed: 7878129]
- Laurent-Jaccard A, De Matteis L, Hofstetter JR, Schutz Y. Are food preferences influenced by body mass index, age, sex, and tobacco? *Schweizerische Medizinische Wochenschrift*. 1994; 124:2039–2041. [PubMed: 7973538]
- Lemmens SG, Born JM, Rutters F, Schoffelen PF, Wouters L, Westerterp-Plantenga MS. Dietary restraint and control over “wanting” following consumption of “forbidden” food. *Obesity (Silver Spring)*. 2010; 18:1926–1931. <http://dx.doi.org/10.1038/oby.2010.36>. [PubMed: 20186140]
- Lenoir M, Serre F, Cantin L, Ahmed SH. Intense sweetness surpasses cocaine reward. *PLoS ONE*. 2007; 2:e698. <http://dx.doi.org/10.1371/journal.pone.0000698>. [PubMed: 17668074]
- Leyton M. Conditioned and sensitized responses to stimulant drugs in humans. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2007; 31:1601–1613. <http://dx.doi.org/10.1016/j.pnpbp.2007.08.027>. [PubMed: 17888557]
- Malik VS, Willet WC, Hu FB. Global obesity: Trends, risk factors and policy implications. *Nature Reviews Endocrinology*. 2013; 9:13–27. <http://dx.doi.org/10.1038/nrendo.2012.199>.
- Massey A, Hill AJ. Dieting and food craving. A descriptive, quasi-prospective study. *Appetite*. 2012; 58:781–785. <http://dx.doi.org/10.1016/j.appet.2012.01.020>. [PubMed: 22306437]
- Mela DJ, Sacchetti DA. Sensory preferences for fats: Relationships with diet and body composition. *American Journal of Clinical Nutrition*. 1991; 53:908–915. [PubMed: 2008871]
- Meule A, Kübler A. Food cravings in food addiction: The distinct role of positive reinforcement. *Eating Behaviors*. 2012; 13:252–255. <http://dx.doi.org/10.1016/j.eatbeh.2012.02.001>. [PubMed: 22664405]
- Meule A, Lutz A, Vogele C, Kübler A. Women with elevated food addiction symptoms show accelerated reactions, but no impaired inhibitory control, in response to pictures of high-calorie food-cues. *Eating Behaviors*. 2012; 13:423–428. <http://dx.doi.org/10.1016/j.eatbeh.2012.08.001>. [PubMed: 23121803]
- Mills JS, Palandra A. Perceived caloric content of a preload and disinhibition among restrained eaters. *Appetite*. 2008; 50:240–245. <http://dx.doi.org/10.1016/j.appet.2007.07.007>. [PubMed: 17888542]

- Murphy C, Stojcek MK, MacKillop J. Interrelationships among impulsive personality traits, food addiction and body mass index. *Appetite*. 2014; 73:45–50. <http://dx.doi.org/10.1016/j.appet.2013.10.008>. [PubMed: 24511618]
- Nisbett RE. Hunger, obesity, and the ventromedial hypothalamus. *Psychological Review*. 1972; 79:433–453. <http://dx.doi.org/10.1037/h0033519>. [PubMed: 4564223]
- Oliver G, Wardle J. Perceived effects of stress on food choice. *Physiology & Behavior*. 1999; 66:511–515. [http://dx.doi.org/10.1016/S0031-9384\(98\)00322-9](http://dx.doi.org/10.1016/S0031-9384(98)00322-9). [PubMed: 10357442]
- Oliver G, Wardle J, Gibson EL. Stress and food choice: A laboratory study. *Psychosomatic Medicine*. 2000; 62:853–865. <http://dx.doi.org/10.1097/00006842-200011000-00016>. [PubMed: 11139006]
- Paolacci G, Chandler J. Inside the turk: Understanding mechanical turk as a participant pool. *Current Directions in Psychological Science*. 2014; 23:184–188. <http://dx.doi.org/10.1177/0963721414531598>.
- Peciña S, Cagniard B, Berridge KC, Aldridge JW, Zhuang X. Hyperdopaminergic mutant mice have higher “Wanting” but not “liking” for sweet rewards. *The Journal of Neuroscience*. 2003; 23:9395–9402. <http://dx.doi.org/10.1097/00008877-200409000-00094>. [PubMed: 14561867]
- Pivarunas B, Conner BT. Impulsivity and emotion dysregulation as predictors of food addiction. *Eating Behaviors*. 2015; 19:9–14. <http://dx.doi.org/10.1016/j.eatbeh.2015.06.007>. [PubMed: 26164390]
- Polivy J, Coleman J, Herman CP. The effect of deprivation on food cravings and eating behavior in restrained and unrestrained eaters. *International Journal of Eating Disorders*. 2005; 38:301–309. <http://dx.doi.org/10.1002/eat.20195>. [PubMed: 16261600]
- Polivy J, Herman CP. Dieting and bingeing: A causal analysis. *American Psychologist*. 1985; 40:193–201. <http://dx.doi.org/10.1037/0003-066X.40.2.193>. [PubMed: 3857016]
- Pursey KM, Collins CE, Stanwell PM, Burrows TL. Foods and dietary profiles associated with ‘food addiction’ in young adults. *Addictive Behaviors Reports*. 2015; 2:41–48. <http://dx.doi.org/10.1016/j.abrep.2015.05.007>.
- Pursey KM, Stanwell P, Gearhardt AN, Collins CE, Burrows TL. The prevalence of food addiction as assessed by the Yale food addiction scale: A systematic review. *Nutrients*. 2014; 6:4552–4590. <http://dx.doi.org/10.3390/nu6104552>. [PubMed: 25338274]
- Raudenbush, SW., Bryk, AS. Hierarchical linear models. 2. SAGE Publications, Inc; 2002.
- Raudenbush, SW., Bryk, AS., Congdon, R. Hlm 7.00 for windows [computer software]. Skokie, IL: Scientific Software International, Inc; 2011.
- Reas DL, Grilo CM. Timing and sequence of the onset of overweight, dieting, and binge eating in overweight patients with binge eating disorder. *International Journal of Eating Disorders*. 2007; 40:165–170. <http://dx.doi.org/10.1002/eat.20353>. [PubMed: 17089414]
- Rissanen A, Hakala P, Lissner L, Mattlar CE, Koskenvuo M, Rönnemaa T. Acquired preference especially for dietary fat and obesity: A study of weight-discordant monozygotic twin pairs. *International Journal of Obesity*. 2002; 26:973–977. <http://dx.doi.org/10.1038/sj.ijo.0802014>. [PubMed: 12080452]
- Robinson TE, Berridge KC. The neural basis of drug craving: An incentive-sensitization theory of addiction. *Brain Research Reviews*. 1993; 18:247–291. [http://dx.doi.org/10.1016/0165-0173\(93\)90013-P](http://dx.doi.org/10.1016/0165-0173(93)90013-P). [PubMed: 8401595]
- Robinson TE, Berridge KC. The psychology and neurobiology of addiction: An incentive-sensitization view. *Addiction*. 2000; 95:S91–S117. <http://dx.doi.org/10.1080/09652140050111681>. [PubMed: 11002906]
- Robinson TE, Berridge KC. The incentive sensitization theory of addiction: Some current issues. *Philosophical Transactions of the Royal Society B*. 2008; 363:3137–3146. <http://dx.doi.org/10.1098/rstb.2008.0093>.
- Rodin J, Mancuso J, Granger J, Nelbach E. Food cravings in relation to body mass index, restraint and estradiol levels: A repeated measures study in healthy women. *Appetite*. 1991; 17:177–185. [http://dx.doi.org/10.1016/0195-6663\(91\)90020-S](http://dx.doi.org/10.1016/0195-6663(91)90020-S). [PubMed: 1799280]
- Rosen JC, Leitenberg H, Fisher C, Khazam C. Binge-Eating episodes in bulimia nervosa: The amount and type of food consumed. *International Journal of Eating Disorders*. 1986; 5:255–267. [http://dx.doi.org/10.1002/1098-108X\(198602\)5:2<255::AID-EAT2260050206>3.0.CO;2-D](http://dx.doi.org/10.1002/1098-108X(198602)5:2<255::AID-EAT2260050206>3.0.CO;2-D).

- Rotenberg KJ, Flood D. Dietary restraint, attributional styles for eating, and preloading effects. *Eating Behavior*. 2000; 1:63–78. [http://dx.doi.org/10.1016/S1471-0153\(00\)00005-2](http://dx.doi.org/10.1016/S1471-0153(00)00005-2).
- Ruderman AJ. Dysphoric mood and overeating: A test of restraint theory's disinhibition hypothesis. *Journal of Abnormal Psychology*. 1985; 94:78–85. <http://dx.doi.org/10.1037/0021-843X.94.1.78>. [PubMed: 3980858]
- Ruderman AJ. Dietary restraint: A theoretical and empirical review. *Psychological Bulletin*. 1986; 99:247–262. <http://dx.doi.org/10.1037/0033-2909.99.2.247>. [PubMed: 3515384]
- Schulte EM, Avena NM, Gearhardt AN. Which foods may be addictive? The roles of processing, fat content, and glycemic load. *PLoS ONE*. 2015; 10:e0117959. <http://dx.doi.org/10.1371/journal.pone.0117959>. [PubMed: 25692302]
- Simmons WK, Martin A, Barsalou LW. Pictures of appetizing foods activate gustatory cortices for taste and reward. *Cerebral Cortex*. 2005; 15:1602–1608. <http://dx.doi.org/10.1093/cercor/bhi038>. [PubMed: 15703257]
- Smith KS, Berridge KC. The ventral pallidum and hedonic reward: Neurochemical maps of sucrose “liking” and food intake. *The Journal of Neuroscience*. 2005; 25:8637–8649. <http://dx.doi.org/10.1523/jneurosci.1902-05.2005>. [PubMed: 16177031]
- Snoek HM, van Strien T, Janssens JMAM, Egnels RCME. Restrained eating and BMI: A longitudinal study among adolescents. *Health Psychology*. 2008; 27:753–759. <http://dx.doi.org/10.1037/0278-6133.27.6.753>. [PubMed: 19025271]
- Spurrell EB, Wilfley DE, Tanofsky MB, Brownell KD. Age of onset for binge eating: Are there different pathways to binge eating? *International Journal of Eating Disorders*. 1997; 21:55–65. [http://dx.doi.org/10.1002/\(SICI\)1098-108X\(199701\)21:1<55::AID-EAT7>3.0.CO;2-](http://dx.doi.org/10.1002/(SICI)1098-108X(199701)21:1<55::AID-EAT7>3.0.CO;2-) [PubMed: 8986518]
- Stevens GA, Singh GM, Lu Y, Danaei G, Lin JK, Ginucane MM, ... Ezzati M. National, regional, and global trends in adult overweight and obesity prevalences. *Population Health Metrics*. 2012; 10:22–37. <http://dx.doi.org/10.1186/1478-7954-10-22>. [PubMed: 23167948]
- Stice E, Fisher M, Lowe MR. Are dietary restraint scales valid measures of acute dietary restriction? Unobtrusive observational data suggest not. *Psychological Assessment*. 2004; 16:51–59. <http://dx.doi.org/10.1037/1040-3590.16.1.51>. [PubMed: 15023092]
- Stunkard AJ, Messick S. The three-factor eating questionnaire to measure dietary restraint, disinhibition and hunger. *Journal of Psychosomatic Research*. 1985; 29:71–83. [PubMed: 3981480]
- Sturm R. The effects of obesity, smoking, and drinking on medical problems and costs. *Health Affairs*. 2002; 21:245–253. <http://dx.doi.org/10.1377/hlthaff.21.2.245>. [PubMed: 11900166]
- Tang DW, Fellows LK, Small DM, Dagher A. Food and drug cues activate similar brain regions: A meta-analysis of functional MRI studies. *Physiology & Behavior*. 2012; 106:317–324. <http://dx.doi.org/10.1016/j.physbeh.2012.03.009>. [PubMed: 22450260]
- Taylor AW, Dal Grande E, Gill TK, Chittleborough CR, Wilson DH, Adams RJ, ... Ruffin RE. How valid are self-reported height and weight? A comparison between CATI self-report and clinic measurements using a large cohort study. *Australian and New Zealand Journal of Public Health*. 2006; 30:238–246. <http://dx.doi.org/10.1111/j.1467-842X.2006.tb00864.x>. [PubMed: 16800200]
- United States QuickFacts. U.S. Census bureau; Retrieved June 1, 2016, from <https://www.census.gov/quickfacts/table/PST045215/00>
- Vanderlinden J, Dalle Grave R, Vandereycken W, Noorduin C. Which factors do provoke binge-eating? An exploratory study in female students. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*. 2001; 2:79–83. <http://dx.doi.org/10.1007/BF03325086>.
- Volkow ND, Morales M. The brain on drugs: From reward to addiction. *Cell*. 2015; 162:712–725. <http://dx.doi.org/10.1016/j.cell.2015.07.046>. [PubMed: 26276628]
- Volkow ND, Wang GJ, Fowler JS, Tomasi D, Baler R. Food and drug reward overlapping circuits in human obesity and addiction. *Current Topics in Behavioral Neuroscience*. 2011; 11:1–24. http://dx.doi.org/10.1007/7854_2011_169.
- Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M. Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet*. 2011; 378:815–825. [http://dx.doi.org/10.1016/S0140-6736\(11\)60814-3](http://dx.doi.org/10.1016/S0140-6736(11)60814-3). [PubMed: 21872750]

- Wardle J, Steptoe A, Oliver G, Lipsey Z. Stress, dietary restraint and food intake. *Journal of Psychosomatic Research*. 2000; 48:195–202. [http://dx.doi.org/10.1016/S0022-3999\(00\)00076-3](http://dx.doi.org/10.1016/S0022-3999(00)00076-3). [PubMed: 10719137]
- Warwick ZS, Schiffman SS. Sensory evaluations of fat-sucrose and fat-salt mixtures: Relationship to age and weight status. *Physiology & Behavior*. 1990; 48:633–636. [http://dx.doi.org/10.1016/0031-9384\(90\)90202-F](http://dx.doi.org/10.1016/0031-9384(90)90202-F). [PubMed: 2082362]
- Waters A, Hill A, Waller G. Internal and external antecedents of binge eating episodes in a group of women with bulimia nervosa. *International Journal of Eating Disorders*. 1999; 29:10–16. [http://dx.doi.org/10.1002/1098-108X\(200101\)29:1<17::AID-EAT3>3.0.CO;2-R](http://dx.doi.org/10.1002/1098-108X(200101)29:1<17::AID-EAT3>3.0.CO;2-R).
- Weingarten HP, Elston D. Food cravings in a college population. *Appetite*. 1991; 17:167–175. [http://dx.doi.org/10.1016/0195-6663\(91\)90019-O](http://dx.doi.org/10.1016/0195-6663(91)90019-O). [PubMed: 1799279]
- White MA, Grilo CM. Psychometric properties of the Food Craving Inventory among obese patients with binge eating disorder. *Eating Behaviors*. 2005; 6:239–245. <http://dx.doi.org/10.1016/j.eatbeh.2005.01.001>. [PubMed: 15854870]
- White MA, Whisenhunt BL, Williamson DA, Greenway FA, Netemeyer RG. Development and validation of the food-craving inventory. *Obesity Research*. 2002; 10:107–114. <http://dx.doi.org/10.1038/oby.2002.17>. [PubMed: 11836456]
- Wyvell CL, Berridge KC. Intra-accumbens amphetamine increases the conditioned incentive salience of sucrose reward: Enhancement of reward “wanting” without enhanced “liking” or response reinforcement. *The Journal of Neuroscience*. 2000; 20:8122–8130. [PubMed: 11050134]
- Yanovski SZ, Leet M, Yanovski JA, Flood MN, Gold P, Kissileff H, et al. Food selection and intake of obese women binge-eating disorder. *American Journal of Clinical Nutrition*. 1992; 56:975–980. [PubMed: 1442665]
- Zellner DA, Loaiza S, Gonzalez Z, Pita J, Morales J, Pecora D, et al. Food selection changes under stress. *Physiology & Behavior*. 2006; 87:789–793. <http://dx.doi.org/10.1016/j.physbeh.2006.01.014>. [PubMed: 16519909]

Table 1

Hierarchical lineal model with craving as outcome variable.

Parameter	Estimate	Std Error	t-Value	df	p-Value	d-Value
For Intercept, β_0						
Intercept, γ_{00}	37.335	1.036	36.038	212	<0.001	4.950
YFAS symptom count, γ_{01}	1.278	0.772	1.656	212	0.099	0.227
Cognitive restraint, γ_{02}	0.209	0.535	0.391	212	0.696	0.054
BMI, γ_{03}	0.233	0.161	1.452	212	0.148	0.199
For Processing slope, β_1						
Intercept, γ_{10}	3.501	0.677	5.174	212	<0.001	0.711
YFAS symptom count, γ_{11}	0.899	0.416	2.164	212	0.032	0.297
Cognitive restraint, γ_{12}	-0.748	0.317	-2.355	212	0.019	-0.323
BMI, γ_{13}	0.026	0.097	0.267	212	0.789	0.037

Note. BMI = Body Mass Index; YFAS = Yale Food Addiction Scale.

Table 2

Hierarchical linear model with liking as outcome variable.

Parameter	Estimate	Std Error	t-Value	df	p-Value	d-Value
For Intercept, β_0						
Intercept, γ_{00}	32.251	1.285	25.088	212	<0.001	3.446
YFAS symptom count, γ_{01}	0.657	0.880	0.747	212	0.456	0.103
Cognitive restraint, γ_{02}	0.506	0.644	0.786	212	0.433	0.108
BMI, γ_{03}	0.299	0.171	1.749	212	0.082	0.240
For Processing slope, β_1						
Intercept, γ_{10}	1.883	1.108	1.699	212	0.091	0.233
YFAS symptom count, γ_{11}	0.436	0.648	0.673	212	0.502	0.092
Cognitive restraint, γ_{12}	-1.246	0.518	-2.404	212	0.017	-0.330
BMI, γ_{13}	0.332	0.162	2.051	212	0.041	0.282

Note. BMI = Body Mass Index; YFAS = Yale Food Addiction Scale.