Published in final edited form as:

Drug Alcohol Depend. 2017 September 01; 178: 39–42. doi:10.1016/j.drugalcdep.2017.04.027.

Expectancy of impairment attenuates marijuana-induced risk taking

Rachel L Gunn^{3,4}, Linda Skalski^{2,3}, and Jane Metrik^{2,1,3}

¹Center for Alcohol and Addiction Studies, Brown University School of Public Health, Providence, RI, 02903

²Providence VA Medical Center, Providence, RI, 02908

³Department of Psychiatry and Human Behavior, Alpert Medical School of Brown University, Providence, RI, 02912

⁴Department of Psychological and Brain Sciences, Indiana University-Bloomington, Bloomington, IN, 47405

Abstract

Background—Marijuana use has been associated with increased risk-taking and impulsive behavior. While pharmacologic effects of marijuana can lead to inhibitory impairment, expectancy of potential impairment may result in compensatory behavioral response by decreasing impulsive decisions and risky behaviors. With the increases in marijuana use and related problems, a better understanding of the individual characteristics associated with marijuana intoxication and risky behavior is needed. This study examined the role of impairment expectancies in marijuana's acute effects on behavioral measures of impulsivity and risk-taking.

Methods—Participants (N=136) were regular marijuana users. A balanced placebo design (BPD) was used crossing marijuana administration (i.e., 0% Tetrahydrocannabinol (THC) vs. 2.8% THC) with stimulus expectancy (i.e., Told Placebo vs. Told THC). Marijuana outcome expectancies were measured by self-report and dependent measures including a number of behavioral impulsivity tasks and the balloon analogue risk task (BART).

Results—Among participants who received THC, higher expectancies for cognitive-behavioral impairment (CBI) were related to lower risk-taking on the BART. Among those who received placebo, there was no association between CBI expectancies and BART performance. CBI

Correspondence: Jane Metrik, PhD, Center for Alcohol and Addiction Studies, Brown University, Box G-S121-4, Providence, RI 02912., Jane_Metrik@brown.edu.

Contributors: Dr. Metrik designed the project that collected the data for the current study. Ms. Gunn conducted literature searches and provided summaries of previous research studies under the guidance of Dr. Metrik. Ms. Gunn conducted the analyses under the guidance of Dr. Metrik. Ms. Gunn wrote the first draft of the manuscript with significant contribution from Dr. Skalski and all authors contributed to and have approved the final manuscript.

Conflict of Interest: All of the listed authors declare that they have no conflicts of interests.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

expectancies did not moderate drug effect on the BART and drug or stimulus expectancy effects on impulsivity measures.

Conclusions—Results provide initial evidence that expectancies of greater impairment are associated with compensatory behavior on a risk-taking task under acute marijuana intoxication. Future studies should examine the role of impairment expectancies on risk behaviors of substantial public health concern such as driving while under the influence of marijuana.

Keywords

marijuana; expectancies; BART; balanced placebo design; risk-taking

1. Introduction

Ample evidence suggests that marijuana use is associated with increased likelihood of risk-taking behaviors, including intoxicated driving (Aston et al., 2016; Johnson et al., 2012), automobile accidents (Li et al., 2012; Ramaekers et al., 2004), and risky sexual behaviors (Simons et al., 2010). Laboratory studies suggest that acute marijuana administration is associated with poor inhibitory control on the Stop-signal task (Hart et al., 2001; McDonald et al., 2003; Metrik et al., 2012; Ramaekers et al., 2006) and the Stroop Task (Hooker and Jones, 1987; Metrik et al., 2012). Findings with marijuana's acute effects on risky decision-making are mixed, some studies showing increased risky decision-making on the Iowa Gambling Task (Lane et al., 2005), and others showing no effect (Ramaekers et al., 2006; Vadhan et al., 2007). Additionally, administration of marijuana is not acutely associated with increased impulsive decision-making during the delay discounting task (McDonald et al., 2003; Metrik et al., 2012).

Few studies have separated the effects of the drug from expectancies, which may also contribute to impulsive and risk-taking behaviors. Balanced placebo design (BPD) (Marlatt and Rohsenow, 1980) allows for the examination of the expectancy that marijuana was smoked independently from the pharmacological effect of delta-9-tetrahydrocannabinol (THC). This 2×2 factorial design crosses drug administration (THC or placebo) with instructions that THC was smoked (i.e., stimulus expectancy: Told THC or Told placebo) (Metrik et al., 2009).

Expectancy theory has helped elucidate the cognitive and motivational mechanisms for marijuana and other substance use (Abrams and Niaura 1987; Goldman and Brown, 1987; Schafer and Brown, 1991). Research and theory suggests that when stimulus expectancies are activated, outcome expectancies (the expected behavioral and affective response to the drug) are in turn initiated (Kirsch and Sapirstein, 1999; Metrik and Rohsenow, 2013). Outcome expectancies are strong determinants of marijuana use in youth and adults (Aarons, et al., 2001; Gaher and Simons, 2007; Neighbors, et al., 2008; Simons, et al., 2009). More specifically, expectancies of cognitive-behavioral impairment (CBI) are inversely related to adolescent lifetime marijuana use and past 3-month use in adults (Kristjansson, et al., 2012). Further, negative marijuana outcome expectancies are related to lower likelihood of risk-taking behavior, such as driving while intoxicated (Arterberry et al., 2013; Aston et al., 2016). In a BPD study, stimulus expectancy effects have decreased impulsive decision-

making on a delay discounting task and increased perception of sexual risk among women, consistent with a compensatory effect (Metrik et al., 2012).

Few studies have examined the potential moderating effects of outcome expectancies on marijuana's pharmacological and stimulus expectancy effects. Metrik and colleagues (2011) found more salient CBI expectancies led to more anxiety under acute marijuana intoxication. In alcohol (Testa et al., 2006; Vogel-Sprott and Fillmore, 1999) and nicotine studies (Juliano and Brandon, 2002), negative outcome expectancies moderated pharmacological effects on behavioral tasks and negative affect. Overall, evidence suggests positive outcome expectancies amplify or increase the likelihood of behavior, whereas negative outcome expectancies result in reduction in behavior or compensation for the negative effects (Metrik and Rohsenow, 2013). No studies thus far have examined the effect of marijuana CBI expectancies on behavioral impulsivity and risk-taking behavior under acute administration. The present study is a secondary analysis of the Metrik et al. (2012) data testing the hypothesis that higher CBI expectancies would be associated with lower levels of impulsivity and risk taking among those who received THC and were told THC relative to participants without any THC or expectancy of THC.

2. Methods

2.1 Participants

Participants met the following inclusion criteria: native English speakers, 18 to 30 years of age, marijuana use at least once a week in the past month and at least ten times in the past 6 months, and self-reported ability to abstain from marijuana for 24 h without withdrawal. Exclusion criteria were: history of substance abuse treatment and intent to quit or receive treatment for marijuana abuse; use of other illicit drugs; pregnancy; nursing; past month affective disorder or history of panic attacks, psychotic, or suicidal state assessed by psychiatric interview; alcohol dependence; contraindicated medical issues by physical exam; 20+ tobacco cigarettes a day; and prior knowledge about the study procedures or contact with participants. The final sample ($N = 136$) had a mean age of 21.4 ($SD = 3.1$), was 64.7% female, 65.4% Caucasian non-Hispanic, and averaged 41.2% ($SD = 24.4$) marijuana use days at baseline. Further detail on demographic and substance use characteristics of the sample are presented in Metrik et al. (2012).

2.2 Procedures

Full details of procedures used in the current study have been previously outlined (Metrik et al., 2012). At baseline, participants completed questionnaires and impulsivity and risk taking tasks to provide within-subjects control for repeated measures. An alveolar carbon-monoxide (CO) of < 6 ppm was used to confirm no recent (past 12 hours) smoking (Cooper and Haney, 2009; Metrik et al., 2012) with a Bedfont Scientific Smokelyzer®. Zero breath alcohol concentration was verified with an Alco-Sensor IV (Intoximeters, Inc., St Louis, MO., USA). Participants were then randomized to one of the four experimental BPD conditions: Told THC/Received THC, Told THC/Received Placebo, Told Placebo/Received THC, and Told Placebo/Received Placebo. At the second session, participants were instructed by a research assistant about which cigarette (THC or placebo) they were assigned

to smoke (see Metrik et al., 2009 for full details of the instructional set manipulation procedures). Marijuana cigarettes (placebo or 2.8% THC) were provided by the National Institute on Drug Abuse, rolled at both ends, humidified, and smoked according to the standardized paced puffing procedure (Foltin, et al., 1987). Participants were fully debriefed about the deception after completing the study. A previous report of the data examined here revealed successful manipulation of drug and expectancy effects (Metrik et al., 2012).

2.3 Measures

2.3.1 Baseline outcome expectancies—Beliefs about possible cognitive-behavioral consequences of smoking marijuana were assessed with the 10-item CBI subscale of the *Marijuana Effect Expectancy Questionnaire* (MEEQ; Schafer and Brown, 1991). The MEEQ is scored on a 5-point Likert scale.

2.3.2. Impulsive disinhibition—The *Stroop Color-Word task* (Stroop 1935) instructed participants to press as quickly as possible the designated key on the keyboard first in response to the color of a symbol string (e.g., XXXX) and then the color of the color-incongruent word. The primary dependent variable was response latency in milliseconds on color-incongruent trials. The *Stop Signal task* (Logan et al., 1997) measures inhibition of a prepotent response with two concurrent tasks: (1) the go task is a choice reaction-time task that requires participants to rapidly discriminate two symbols (maximum presentation 1,250 ms) and (2) the stop task involves presentation of a tone (75ms, 1,000 Hz) that signals one to inhibit the response to the go task. The time in milliseconds required for the participant to stop the go response (stop signal reaction time, SSRT) was the primary dependent measure.

2.3.3. Impulsive decision-making—The *Delay Discounting Questionnaire* (DDQ; Richards et al., 1999) uses a computerized adjusting amount procedure to measure discounting of delayed monetary reinforcers (hypothetical choices between \$10 available after a delay or a smaller amount available immediately). The primary dependent variable was area under the curve connecting indifference points and the x-axis, from 0.0 (steepest discounting) to 1.0 (no discounting) (Myerson et al., 2001). The *Experiential Discounting task* (EDT; Reynolds and Schiffbauer 2004) also assesses delayed-discounting. During a block of trials, participants chose between a standard amount (\$0.30) that was probabilistic and an adjusting amount of money (initially set to \$0.15) that was always delivered immediately and was certain. Participants were told that if they chose the standard 30 cents, the next adjusting choice would increase, while choosing the immediate amount caused the next adjusting choice to decrease. The primary dependent variable was the indifference point (i.e., an equal number of choices to both the standard and immediate options). The *Balloon Analogue Risk Task*, automatic response version (BART; Lejuez et al., 2002; Pleskac et al., 2008), is a computerized behavioral measure of risk taking. Thirty balloon images were presented one at a time with participants designating the number of pumps to have each balloon inflated. Each pump earned 1 cent, but money was lost if the balloon exploded (at 64 pumps on average). The primary dependent variable was the average number of pumps across trials.

2.4. Data Analysis Plan

ANOVA was conducted to test for differences in CBI expectancies between drug and stimulus expectancy manipulation groups at baseline. BPD groups did not differ on demographics, substance use characteristics, or baseline impulsivity measures in prior analyses (Metrik et al. 2012). Multiple regression analyses were used to examine the hypothesis Marijuana Expectancies and Risk Taking 7 that CBI expectancies would moderate drug manipulation and stimulus expectancy effects on outcomes. Five separate models were run for each of the five outcome measures (Stroop, Stop-signal, BART, EDT, and DDQ). In each model, baseline task performance and task administration order were entered at step one as covariates, main effects of drug manipulation, stimulus expectancy and CBI expectancy MEEQ subscale were entered at step two, and two interactions of CBI expectancy with drug manipulation and stimulus expectancy were entered in the third and final step to test for moderation. Drug by stimulus expectancy interactions were not significant across all measures in prior analyses (Metrik et al., 2012).

3. Results

ANOVA revealed no significant difference in CBI expectancies at baseline between drug manipulation, $F(1,134) = 2.78$, $p = .10$ nor stimulus expectancy groups, $F(1,134) = .904$, $p = .34$. Multiple regression revealed non-significant interactions of CBI expectancies with drug manipulation or stimulus expectancy effects on the Stroop, Stop-Signal, EDT, or DDQ tasks after smoking. On the BART task, there was a significant interaction of CBI expectancies with drug manipulation, $B = -.190$, $SE = .08$, $p = .02$, $sr^2 = .04$, but not with stimulus expectancies, $B = -.091$, $SE = .08$, $p = .27$, $sr^2 = .01$ (Table 1). Post-hoc analyses without the CBI expectancy by stimulus expectancy interaction term in the model revealed higher expectancies of CBI were associated with less balloon pumps on the BART (lower levels of risk-taking) among participants who received THC, $B = -.16$, $SE = .08$, $p < .05$, $sr^2 = .06$. Among those who received placebo, there was no significant association between CBI expectancies and pumps on the BART, $B = .09$, $SE = .07$, $p = .22$, $sr^2 = .02$ (Figure 1).

4. Discussion

This study examined whether marijuana outcome expectancies of CBI would moderate the pharmacological and stimulus expectancy effects of marijuana on measures of impulsivity and risk-taking. Results suggested that CBI expectancies moderated the pharmacological, but not stimulus expectancy, effect of marijuana on the BART task. Among participants who received THC, higher expectancies for impairment were related to lower risk-taking (less balloon pumps). However, among those who received placebo, there was no association between impairment expectancies and performance on the BART task.

Consistent with previous research examining the relationship between negatively-valenced expectancies and behavior, results suggest a compensatory behavioral response for individuals who expect marijuana will impair thoughts and behaviors (Metrik and Rohsenow, 2013). These results build on previous findings on compensatory behavior observed in Metrik et al., 2012, in which THC stimulus expectancy reduced risky decision-making. Additionally, in alcohol studies, impairment expectancies among social drinkers are

associated with less impairment on a stop-signal task (Fillmore and Vogel-Sprott, 1996). This response was also evident in an alcohol and caffeine study, in which social drinkers who were told that caffeine would counteract the effect of alcohol showed more impairment than those subjects who were not given this expectancy (Fillmore, et al., 2002). Together with the current findings, this suggests that expecting impairment from a substance may operate as a protective factor against impulsive and risky behaviors when under the influence of that substance. Research also suggests negative marijuana outcome expectancies are associated with compensatory behaviors while driving (i.e., reduction in speed) (Bates and Blakely, 1999). However, this study also found that this increased false perceptions of safety, highlighting the potential negative consequences of these expectancies.

Somewhat surprisingly, CBI expectancies did not moderate drug or stimulus expectancy effects on behavioral impulsivity measures. It is possible that marijuana expectancies are not as relevant to more automatic processes in impulsivity such as behavioral disinhibition. Indeed, there was no effect of stimulus expectancy on the Stroop or the Stop Signal tasks in the parent study (Metrik et al., 2012). Contrary to our expectation, CBI expectancies did not modulate marijuana's effects on delay-discounting measures, which assess impulsive decision-making involving a more deliberate process of evaluation of outcomes.

5. Conclusions

The findings provide further evidence towards the value of carefully targeting outcome expectancies for intervention. Results suggest that expectancies about the impairing effects of marijuana lead to lower levels of risk-taking among users when under the influence of the drug. The present study also suggests that this moderating effect may be unique to risk-taking behavior, given impairment expectancies did not moderate drug or expectancy effects on other measures of impulsive decision making or disinhibition. Given the BART was the only measure of risk-taking in the present study, it is unknown whether such compensatory action would be observed with marijuana's acute effects on other measures of risk-taking. Future controlled studies should investigate the potential moderating role of impairment expectancies on other measures that are of significant public health concern, such as driving under the influence of marijuana.

Acknowledgments

Role of Funding Sources: This study was supported by a grant R01 DA021403 from the National Institute on Drug Abuse (NIDA) to Jane Metrik. The views expressed in this article do not necessarily reflect the position or policy of the Department of Veteran Affairs.

The authors gratefully acknowledge Dr. James Harper, III, Amy Mochel, Suzanne Sales, and Timothy Souza for their contribution to the project.

References

- Aarons GA, Brown SA, Stice E, Coe MT. Psychometric evaluation of the marijuana and stimulant effect expectancy questionnaires for adolescents. *Addict Behav.* 2001; 26:219–236. [PubMed: 11316378]
- Abrams DB, Niaura RS. Social learning theory. *Psychol Theor Drink Alcohol.* 1987; 1:131–178.

- Arterberry BJ, Treloar HR, Smith AE, Martens MP, Pedersen SL, McCarthy DM. Marijuana use, driving, and related cognitions. *Psych Addict Behav.* 2013; 27:854.
- Aston ER, Merrill JE, McCarthy DM, Metrik J. Risk factors for driving after and during marijuana use. *J Stud Alcohol Drugs.* 2016; 77:309–316. [PubMed: 26997189]
- Bates MN, Blakeley TA. Role of cannabis in motor vehicle crashes. *Epidemiol Rev.* 1999; 21:222–232. [PubMed: 10682259]
- Cooper ZD, Haney M. Comparison of subjective, pharmacokinetic, and physiological effects of marijuana smoked as joints and blunts. *Drug Alcohol Depend.* 2009; 103:107–113. [PubMed: 19443132]
- Fillmore MT, Roach EL, Rice JT. Does caffeine counteract alcohol-induced impairment? The ironic effects of expectancy. *J Stud Alcohol.* 2002; 63:745–754. [PubMed: 12529075]
- Fillmore MT, Vogel-Sprott M. Evidence that expectancies mediate behavioral impairment under alcohol. *J Stud Alcohol.* 1996; 57:598–603. [PubMed: 8913990]
- Foltin RW, Fischman MW, Pedrosa JJ, Pearlson GD. Marijuana and cocaine interactions in humans: Cardiovascular consequences. *Pharmacol Biochem Behav.* 1987; 28:459–464. [PubMed: 2829241]
- Gaher RM, Simons JS. Evaluations and expectancies of alcohol and marijuana problems among college students. *Psychol Addict Behav.* 2007; 21:545. [PubMed: 18072837]
- Goldman, MS., Brown, SA., Christiansen, BA. Expectancy theory-thinking about drinking. In: Blane, Leonard, KE., editors. *Psychological theories of drinking and alcoholism.* Guilford Publications; New York, NY: 1987. p. 181
- Hart CL, Van Gorp W, Haney M, Foltin RW, Fischman MW. Effects of acute smoked marijuana on complex cognitive performance. *Neuropsychopharmacol.* 2001; 25:757–765.
- Hooker WD, Jones RT. Increased susceptibility to memory intrusions and the Stroop interference effect during acute marijuana intoxication. *Psychopharmacol.* 1987; 91:20–24.
- Johnson MB, Kelley-Baker T, Voas RB, Lacey JH. The prevalence of cannabis-involved driving in California. *Drug Alcohol Depend.* 2012; 123:105–109. [PubMed: 22101027]
- Juliano LM, Brandon TH. Effects of nicotine dose, instructional set, and outcome expectancies on the subjective effects of smoking in the presence of a stressor. *J Abnorm Psych.* 2002; 111:88–97.
- Kirsch, I., Sapirstein, G. Listening to Prozac but hearing placebo: A meta-analysis of antidepressant medications. In: Kirsch, I., editor. *How Expectancies Shape Experience.* American Psychological Association; Washington, DC: 1999. p. 303-320.
- Kristjansson SD, Agrawal A, Lynskey MT, Chassin LA. Marijuana expectancies and relationships with adolescent and adult marijuana use. *Drug Alcohol Depend.* 2012; 126:102–110. [PubMed: 22682980]
- Lane SD, Cherek DR, Tcheremissine OV, Lieving LM, Pietras CJ. Acute marijuana effects on human risk taking. *Neuropsychopharmacol.* 2005; 30:800–809.
- Lejuez CW, Read JP, Kahler CW, Richards JB, Ramsey SE, Stuart GL, Strong DR, Brown RA. Evaluation of a behavioral measure of risk taking: The Balloon Analogue Risk Task (BART). *J Exp Psych Appl.* 2002; 8:75.
- Li MC, Brady JE, DiMaggio CJ, Lusardi AR, Tzong KY, Li G. Marijuana use and motor vehicle crashes. *Epidemiol Rev.* 2012; 34:65–72. [PubMed: 21976636]
- Logan GD, Schachar RJ, Tannock R. Impulsivity and inhibitory control. *Psychol Sci.* 1997; 8:60–64.
- Marlatt GA, Rohsenow DJ. Cognitive processes in alcohol use: Expectancy and the balanced placebo design. *Advances in substance abuse: Behav Biol Res.* 1980; 1:159–199.
- McDonald J, Schleifer L, Richards JB, de Wit H. Effects of THC on behavioral measures of impulsivity in humans. *Neuropsychopharmacol.* 2003; 28:1356–1365.
- Metrik J, Kahler CW, McGeary JE, Monti PM, Rohsenow DJ. Acute effects of marijuana smoking on negative and positive affect. *J Cogn Psychother.* 2011; 25:31–46.
- Metrik J, Kahler CW, Reynolds B, McGeary JE, Monti PM, Haney M, de Wit H, Rohsenow DJ. Balanced placebo design with marijuana: pharmacological and expectancy effects on impulsivity and risk taking. *Psychopharmacol.* 2012; 223:489–499.
- Metrik, J., Rohsenow, DJ. *Addiction. The Wiley-Blackwell Handbook of Addiction Psychopharmacology*; 2013. Understanding The Role Of Substance Expectancies; p. 459-487.

- Metrik J, Rohsenow DJ, Monti PM, McGeary J, Cook TA, de Wit H, Haney M, Kahler CW. Effectiveness of a marijuana expectancy manipulation: Piloting the balanced-placebo design for marijuana. *Exp Clin Psychopharmacol*. 2009; 17:217. [PubMed: 19653787]
- Myerson J, Green L, Warusawitharana M. Area under the curve as a measure of discounting. *J Exp Anal Behav*. 2001; 76:235–243. [PubMed: 11599641]
- Neighbors C, Geisner IM, Lee CM. Perceived marijuana norms and social expectancies among entering college student marijuana users. *Psych Addict Behav*. 2008; 22:433.
- Pleskac TJ, Wallsten TS, Wang P, Lejuez CW. Development of an automatic response mode to improve the clinical utility of sequential risk-taking tasks. *Exp Clin Psychopharmacol*. 2008; 16:555. [PubMed: 19086776]
- Ramaekers JG, Berghaus G, van Laar M, Drummer OH. Dose related risk of motor vehicle crashes after cannabis use. *Drug Alcohol Depend*. 2004; 73:109–119. [PubMed: 14725950]
- Ramaekers JG, Kauert G, van Ruitenbeek P, Theunissen EL, Schneider E, Moeller MR. High-potency marijuana impairs executive function and inhibitory motor control. *Neuropsychopharmacol*. 2006; 31:2296–2303.
- Reynolds B, Schiffbauer R. Measuring state changes in human delay discounting: An experiential discounting task. *Behav Process*. 2004; 67:343–356.
- Richards JB, Zhang L, Mitchell SH, Wit H. Delay or probability discounting in a model of impulsive behavior: Effect of alcohol. *J Exp Anal, Behav*. 1999; 71:121–143. [PubMed: 10220927]
- Schafer J, Brown SA. Marijuana and cocaine effect expectancies and drug use patterns. *J Consult Clin Psych*. 1991; 59:558.
- Simons JS, Dvorak RD, Lau-Barraco C. Behavioral inhibition and activation systems: Differences in substance use expectancy organization and activation in memory. *Psych Addict Behav*. 2009; 23:315–328.
- Simons JS, Maisto SA, Wray TB. Sexual risk taking among young adult dual alcohol and marijuana users. *Addict Behav*. 2010; 35:533–536. [PubMed: 20071102]
- Stroop JR. Studies of interference in serial verbal reactions. *J Exp Psych*. 1935; 18:643.
- Testa M, Fillmore MT, Norris J, Abbey A, Curtin JJ, Leonard KE, Mariano KA, Thomas MC, Nomensen KJ, George WH, Livingston JA, Saenz C, Buck PO, Zawacki T, Parkhill MR, Jacques AJ, Hayman LW Jr, Van Zile-Tamsen C. Understanding alcohol expectancy effects: Revisiting the placebo condition. *Alcohol Clin Exp Res*. 2006; 30:339–348. [PubMed: 16441283]
- Vadhan NP, Hart CL, van Gorp WG, Gunderson EW, Haney M, Foltin RW. Acute effects of smoked marijuana on decision making, as assessed by a modified gambling task, in experienced marijuana users. *J Clin Exp Neuropsychol*. 2007; 29:357–364. [PubMed: 17497559]
- Vogel-Sprott, M., Fillmore, MT. American Psychological Association. Washington, DC: 1999. Expectancy and Behavioral Effects of Socially Used Drugs.

Highlights

- Expectancy of marijuana's impairment attenuated marijuana's effect on risk taking
- Under Tetrahydrocannabinol (THC), higher impairment expectancies were related to lower risk taking
- Under placebo, this association was not present
- Impairment expectancies predict compensatory behavior under the influence of THC

Figure 1. Interaction of CBI Expectancies and Drug Effect on BART

Note. Interaction effect from post-hoc regression analysis examining Marijuana Cognitive-Behavioral Impairment Expectancies (centered) by Received (THC/Placebo) on Average number of pumps on the BART post-smoking. Variance displayed using 95% confidence intervals.

Table 1
Step-wise Multiple Regression Predicting post-smoking BART performance

Predictor	β	S.E.	t	P	sr^2
Step 1					
BART(BL)	.78	.05	207.61	.00	.59
Task order	.04	.05	.60	.44	.00
CBI Expect	-.05	.05	.94	.33	.01
Step 2					
BART(BL)	.79	.06	203.81	.00	.61
Task order	.04	.06	.64	.43	.01
CBI Expect	-.05	.06	.68	.41	.01
Told (THC/Placebo)	-.03	.06	.27	.60	.00
Received (THC/Placebo)	.03	.06	.37	.54	.00
Step 3					
BART(BL)	.78	.05	207.54	.00	.62
Task order	.05	.05	.98	.32	.01
CBI Expect	.17	.10	2.77	.10	.02
Told (THC/Placebo)	-.05	.06	.70	.41	.01
Received (THC/Placebo)	.03	.06	.30	.59	.00
CBI $\times$ Received	-.19	.08	5.4	.02	.04
CBI $\times$ Told	-.09	.08	1.21	.27	.01

Note. BART(BL) = Average number of pumps across balloon trials on the BART at baseline (centered) (Mean= 60.23, SD = 11.47); CBI Expect= Cognitive-Behavioral Impairment Expectancies (centered) (M = 2.73, SD = .72); Told (coded 0 = placebo, 1 = THC); and Received (coded 0 = placebo, 1 = THC); CBI $\times$ Received= CBI Expectancies by drug (Received THC or placebo) interaction term, CBI $\times$ Told = CBI Expectancies by stimulus expectancy (Told THC or placebo) interaction term. sr^2 = squared semipartial correlation.