ART. V.—*On the Action of certain Inorganic Compounds, when introduced directly into the Blood.* By JAMES BLAKE, Esq., M. R. C. S. L., F. R. M. C. S. [Read before the Royal Society, January 21 and 28, 1841.]

IN a paper on the actions of poison, which I presented to the Royal Society in April 1839, and which was read on the 13th of June in the same year, I incidentally alluded to some experiments I had performed on the action of salts, when injected into the veins. On that occasion I merely noticed these experiments, on account of their having enabled me to determine the time required by the blood, to pass from one part of the system to another.

The results of my experiments on the introduction of saline substances into the vascular system, were communicated to the "Academie des Sciences" at Paris in the same year. (This memoir was published in the "Archives Generales de Medecine," November 1839.) Since this period, a further investigation of the subject has afforded me many new facts, which I now propose to lay before the Society.

At the time I wrote my former memoirs, I had only investigated the actions of the substances I had experimented with, in relation to their action on the heart, and the changes to which they appeared to give rise in the lungs. I had, in fact, divided these substances into two classes, one of which included those which appeared to produce death by destroying the irritability of the heart, whilst the other contained those substances, which prove fatal by arresting the pulmonary circulation. Entirely occupied by these two phenomena, I had noticed but casually the action which the salts of soda and potash appeared to exert on the general capillary circulation. The experiments I have now to bring forward will prove, that the effects that these and some other substances produce on the general capillary circulation, form an important element in the investigation of their general action on the animal economy. But besides those phenomena which depend on the action of these salts on the vascular system, there are others to which I had not at all directed my attention, viz. the effect which these substances appear to give rise to, by their contact with the nervous and muscular tissues. In the present memoir I have carefully recorded the effects produced by each substance on the different tissues to which I have now alluded, at least where these effects have been of a nature to be readily appreciated. I have, in fact, endeavoured to investigate the properties of the living tissues, in the same manner as the chemist would investigate the nature

of an inorganic compound, that is, by observing the phenomena it presents, when brought into relation with other substances.

I am aware that, in the present state of physiology, such a step may appear rash, and not likely to lead to any positive results ; but I know of no surer means that we can command of becoming better acquainted with the complicated phenomena which constitute life, than by introducing certain definite modifications in the circumstances under which these phenomena occur ; and I think there can be no question, that the direct introduction of inorganic substances into the blood, affords us one of the most certain means of producing definite and appreciable changes in these circumstances. It is, however, equally true, that the present imperfect state of chemistry leaves us completely in the dark as to the nature of the perturbations which are thus produced, depriving us at the outset of every hope of arriving at any general conclusions, and confining our researches to the mere observation of facts. It is these considerations that have induced me to lay before the Society the following memoir, although I am fully aware that the experiments I have performed, still leave many voids to be filled up. The vast extent of the subject, however, involving, as it does, some of the highest physiological problems, precludes the hope that I should ever have arrived at a point, at which I might consider these researches as complete.

Before proceeding to the detail of the experiments, I shall shortly describe the manner in which they have been performed. The phenomena connected with the vascular system have been observed by the aid of the hæmadynamometer,—an instrument invented by M. Poiseuille, for ascertaining the pressure that the blood exerts on the parietes of the vessels. Had it not been for the almost mathematical precision which this valuable instrument enables us to introduce in researches on the circulation, I certainly should never have ascertained many of the facts which I have now to bring forward. A description of this instrument will be found in Todd's *Cyclopædia of Anatomy and Physiology*, Art. "Circulation." In ascertaining the pressure of the blood in the arterial system, this instrument has always been connected with the femoral artery. When the substance experimented on has been injected into the veins, it is the external jugular that has always been selected for this operation ; when into the arteries, the injecting tube has been introduced either into the right or left axillary artery, the point of the tube looking towards the heart, so that any fluid injected through it should be forced through the subclavian artery into the aorta, and thus become mixed with the blood circulating through the body. In all operations requiring the use of the hæmadynamometer, it is necessary that the animal should be confined, in order to command the requisite degree of

steadiness. In these circumstances, it is impossible to observe with accuracy the general symptoms that are produced. I have, therefore, always devoted a distinct experiment for the purpose of ascertaining these symptoms, a tube being inserted into the jugular vein, and the animal being placed on the floor, and allowed full liberty, the moment after the substance to be experimented with had been introduced into the vein. In these circumstances, any general symptoms that were produced would be readily detected. The whole of these experiments have been performed on dogs; they have been chosen as nearly as possible of the same size and strength: the general weight was from 12 to 16 pounds. The substances I have used in these experiments have been the sulphate and nitrate of magnesia; sulphate of zinc; sulphate of copper; nitrate of lime, and chloride of calcium; chloride of strontium, and nitrate of strontia; chloride of barium, and nitrate of baryta; acetate of lead; nitrate and acetate of silver; chloride of sodium, and nitrate and sulphate of soda; nitrate and hydrochlorate of ammonia; nitrate, sulphate, biarseniate, and bicarbonate of potassa, and the alkali itself.

The order in which I propose stating the experiments I have performed with these substances, is that in which they are here arranged. This arrangement may be considered in some respects a natural one; for each class of salts, from magnesia to soda, will be found connected by its physiological action more closely with the salt that more immediately precedes and follows it, than with any other.

The fact which I stated in the memoir presented to the Academy of Sciences at Paris, (see Archives Generales de Med. Nov. 1839), that all the salts of the same base exert an analogous action when introduced directly into the blood, a fact, the correctness of which has been proved by a far more extended series of experiments, will dispense with my stating the action of more than one salt of each base.

Magnesia.—When a solution of a salt of this base is injected into the veins, its first effect, if it be not too strong, is to quicken the action of the heart; and it also exerts a powerful influence on the character of the pulse, as is shown by the change which takes place in the oscillations of the column of mercury, indicating the pressure in the arterial system. Six grains of sulphate of magnesia, dissolved in five drachms of water, when injected into the veins, caused the pulsations of the heart to become quicker ten seconds after the injection, or at the moment when the blood containing the salt was circulated over the heart; at the same time, the oscillations of mercury in the hæmadynamometer became much diminished. The pressure in the arterial system was, at the commencement of the experiment, equal to a column of mercury, varying from five to seven inches and a half, at each pulsation of the

heart ; but a few seconds after the injection, the column of mercury was nearly stationary at six inches, the oscillations not being more than two or three-tenths of an inch, instead of two inches and a half, as they were before the injection. Two minutes after the injection, the effects of the salt had quite gone off, the action of the heart being just the same as it was before the introduction of the salt. Five minutes after the former experiment, a solution containing sixteen grains of the salt was injected into the veins, and produced analogous effects on the heart in about ten seconds ; thirty seconds after the injection, the pressure in the arterial system became lower, proving that some change had taken place either in the quantity of blood sent into the arteries, or in the facility with which it was discharged through the systematic capillaries ; for it can only be by a modification of one or the other of these elements that any change can take place in the pressure in the arterial system. When injected in larger quantities, this salt suddenly arrests the action of the heart. A solution containing a drachm of the salt to six drachms of the water, arrested the action of the heart eight seconds after its introduction into the veins. The inspection of the thorax immediately after death, discovered the heart motionless, although exposed half a minute after respiration had ceased. On opening the pericardium, slight movements were observed on that part of the parietes of the ventricle corresponding to the interventricular septum. The auricles were quite motionless. The left cavities of the heart were filled with blood, of a bright scarlet colour ; that in the right cavity was dark. The presence of arterialized blood in the left cavities of the heart, affords a certain proof that the movements of this organ had stopped, before respiration had become arrested.

When injected into the arteries, the sulphate of magnesia does not appear to have any influence on the passage of the blood through the general capillaries. A tube being introduced into the right axillary artery, a solution containing sixteen grains of the salt was injected into the artery. No immediate change was observed in the pressure in the arterial system, which could not have been the case, had any change taken place in the passage of the blood through the general capillaries. The immediate effect on the column of mercury in the hæmadynamometer cannot be carefully noted, owing to the violent cries and efforts of the animal, which produce great oscillations in the mercury ; the pressure certainly was not augmented ; and in about a minute, when the struggles of the animal had ceased, the column of mercury was found to have diminished about one inch. Five minutes after the above experiment, a second solution, containing thirty grains of the salt, was introduced into the artery. The phenomena that followed were analogous to those just described, except that the oscillations in the column of mercury were rendered less. When a much

larger quantity of this salt is injected, death appears to follow, owing to its action on the nervous system; as it appears not to arrive at the heart in a sufficiently concentrated state to arrest its pulsations.

The general symptoms that follow the introduction of this substance into the veins depend, as might be supposed, on the quantities used. A solution, containing eight grains of the salt, when injected into the veins, caused slight dyspnœa ten seconds after its introduction; in twelve seconds, the animal fell down, as if suddenly paralysed, but soon rose again, and did not appear to suffer any inconvenience. Two minutes after the injection, vomiting took place. After a few minutes, a solution containing fourteen grains was injected; the animal was evidently affected in ten seconds; in twelve seconds, it fell on its side; the legs extended, and appeared totally deprived of all power to move; the muscles seemed even to have lost their tonicity, as the limbs remained in the position in which they were placed; there were not the slightest spasms or convulsions, nor did the animal appear to suffer, although perfectly sensible. After lying in this state for ten minutes, the animal got up and walked, but very unsteadily. The action of the heart was arrested on introducing into the veins a solution containing thirty-five grains of the salt; the heart became paralysed, and the animal fell about twelve seconds after the injection, stretched its legs, and was dead in about thirty seconds,—not more than two or three respiratory movements taking place after the action of the heart had ceased. On opening the thorax, the heart was found in the same state as in the case above related, where this salt was injected into the veins. The presence of this salt in the blood did not interfere with its coagulation. The lungs were rather more scarlet than natural, but their texture was not altered.

Before proceeding to relate the experiments that have been performed with the salts of zinc, I would offer a few remarks on the facts which have been just described. The effect produced by small quantities of the salts of magnesia on the heart, is, as I have stated, to quicken the pulsations of that organ, and also to cause a marked difference in the character of the pulse, as shown by the diminutions of the oscillations of the column of mercury indicating the pressure in the arterial system. This alteration in the character of the pulse may depend on two causes;—either that the ventricular systole requires a longer time for its completion, the blood being thus forced into the aorta more gradually;—or it might be due to an increase in the elasticity of the parietes of the arteries. It is to the first of these causes that I am inclined to attribute the phenomenon in question, although the latter may participate in a slight degree in its production. That a longer systole

of the heart is not inconsistent with its quickened pulsation, is evident; for we have only to suppose the period of repose shorter, and we may have the systole much lengthened, whilst the number of the contractions of the organ in a given time are increased.

The Salts of Zinc.—These salts are very closely allied to those of magnesia in their physiological action. When injected into the veins, in fact, the only difference that exists in the action of these two classes of salts is, that those of zinc produce the same effect as those of magnesia, even when exhibited in much smaller doses; the former also acting more powerfully on the nervous system. A solution, containing three grains of sulphate of zinc, when injected into the jugular vein, gave rise to a quickened pulsation of the heart ten seconds after its injection; the oscillations in the column of mercury were also diminished, and the pressure in the arterial system decreased; two minutes after the injection, it being less by a column of mercury equal to two inches; the action of the heart was also slower. Six grains, when injected into the veins, exerted a powerful influence on the heart; the pressure in the arterial system became diminished until it was only equal to a column of mercury of two inches. A solution, containing thirty grains of the salt, arrested the action of the heart eight seconds after it had been introduced into veins. On opening the thorax, the irritability of the heart was found to be more completely destroyed than in the experiments with magnesia.

On injecting a solution of this substance into the arteries, it is found not to exert any influence on the passage of the blood through the systematic capillaries. Sixteen grains introduced through the axillary artery destroyed life by the action it exerted on the nervous system; the heart continued beating for some time after respiratory movements had ceased. In the complete prostration of strength which follows its introduction into the veins, this salt agrees perfectly with the salt of magnesia. This prostration has been noticed by Orfila in his "*Traité des Poisons*," T. i. p. 576, where he states, that, after the injection of a quantity of this salt into the veins, the animal fell "*comme une masse inerte*." The sensibility does not seem affected, nor are there any convulsions, the animal lying, according to the expression of the French toxicologist, "*dans un état de calme parfait*." The injection of a solution containing six grains of sulphate of zinc into the veins of a dog, weighing about eight pounds, was followed by complete prostration of strength. A solution containing three grains more arrested the action of the heart a few seconds after its introduction into the veins. On opening the thorax, slight movements were still taking place in the ventricles; the blood in the left side of the heart was scarlet; it did not coagulate.* The

* This salt appears to possess the property of dissolving, or at least of preventing

various experiments I have made with this substance would lead me to conclude that it exerts a more marked influence in destroying the irritability of the auricles than of the ventricles. In one case, the ventricles were found contracting rythmically, when the auricles were quite still.

Sulphate of Copper.—This substance, when injected into the veins, exerts a more decided action on the heart, than does the sulphate of zinc. The injection of a solution, containing six grains of the sulphate of copper, gave rise to a quickened and fluttering action of the heart ten seconds after its introduction. This was followed in forty-five seconds by the pulsations of the heart becoming much slower, and a great diminution in the column of mercury, indicating the pressure in the arterial system, which fell full two inches. A solution, containing fifteen grains of the salt, arrested the action of the heart twelve seconds after its introduction into the veins. On opening the thorax, immediately after death, the heart was found motionless, with the exception of slight contractions of the interventricular septum; no movements of the auricles were observed. The right side of the heart contained a considerable quantity of dark blood; the left cavities contained a smaller quantity, which was rather scarlet, but not of so bright a tint as arterial blood; for it appears the sulphate of copper interferes with the change of colour which the blood undergoes in its passage through the lungs, a point in which it differs from the salts of zinc. Injected into the arteries, this salt does not exert any influence on the passage of the blood through the systemic capillaries. Ten grains injected into the axillary artery, produced death by its action on the nervous system, without causing an immediate alteration in the pressure in the arteries.

The general symptoms which follow the introduction of the sulphate of copper into the veins differs from those produced by the salts of magnesia and zinc, by the absence of that complete prostration of muscular power which is so strikingly characteristic of the action of these latter salts. A solution containing three grains of the salt, when injected into the veins, gave rise to efforts to vomit two minutes after its introduction. On injecting a solution containing four grains, into the veins of the same animal, there was dyspnœa produced in fifteen seconds. The animal fell immediately, but continued moving its legs, which were rather

the coagulation of the fibrine of the blood. When blood taken from animals in whose veins this substance has been injected is allowed to stand, a perfectly clear and limpid solution of the fibrine is obtained, the colouring matter being precipitated apparently unaltered. In fact, the sulphate of zinc appears to exert but little influence on the colouring matter of the blood, for when injected into the veins in a sufficiently concentrated state, to completely change the physical character of this fluid, and even of the tissue of the lungs; yet the blood in the left side of the heart is found of a bright scarlet colour.

stiff, although there were no marked spasms or convulsions. After the animal had been lying on its side four minutes, it again rose on its legs, during violent efforts to vomit, but it soon fell down again. The action of the heart was arrested on introducing six grains more into the veins; respiration was suspended in forty-five seconds; the conjunctiva was sensible one minute after the suspension of the respiratory movements. After every respiratory movement had been suspended during two minutes, respiration again commenced, and continued uninterruptedly during a minute and a-half. The animal did not in any degree recover its sensibility, or the power of voluntary motion, during this return of the respiratory movements. In one experiment with this substance, a full respiratory movement took place, on making an incision into the thorax, four minutes after the animal had manifested any sign of sensation, and five minutes after the action of the heart had ceased: this movement was probably reflex; it was not accompanied by any marked irritability of the muscles, which, in fact, did not offer those movements which are generally produced in the muscular fibre, by the contact of air, and the irritation of the scalpel, immediately after death. This absence of irritability in the muscles after death, produced by the injection of sulphate of copper into the veins, as also the absence of irritability in the heart, has been noticed by Orfila. I have also observed that the same effect has been produced on the muscular fibre by the injection of the salts of zinc and magnesia.

The Salts of Lime.—I am not aware that any observations have been published on the action of these salts when introduced into the veins. In their action on the heart they resemble very closely those substances which have been already noticed. When injected into the veins in sufficient quantities, they arrest the action of the heart in from eight to twelve seconds. To produce this effect, a drachm and a-half or two drachms of the chloride of calcium or nitrate of lime are required. When injected in small quantities, the pulsations of the heart are rendered quicker, and the oscillations in the column of mercury are diminished. In their subsequent action on the heart, the salts of lime differ from the salts of magnesia, copper, and zinc, by their tendency to make the action of the heart permanently quicker, the rapidity of the pulsations being increased, until they are introduced in sufficient quantities to paralyse the heart. An examination of the thorax immediately after death discovers the heart more or less deprived of irritability. Sometimes the organ has been found perfectly motionless. The blood in the left cavities is always of a bright scarlet colour. When injected into the arterial system the salts of lime exert no action on the general capillary circulation, the pressure in the arteries not being affected.

On injecting a solution of chloride of calcium through the axillary artery, I have noticed a curious fact, that would tend to prove, that, if there be not regurgitation of blood into the left ventricle during its diastole, there at least cannot be any onward current in the ascending and transverse part of the aorta. The fact to which I allude is, the quickening of the action of the heart four seconds after the introduction of this salt into the axillary arteries, its pulsations being affected in the same manner as when the salt becomes applied to it, after being introduced into the veins; the short interval (only four seconds) elapsing between the injection of the substance into the arteries, and its action on the heart, precludes the idea, that the salt would have been applied to that organ, after traversing the general and pulmonary circulation. The only explanation of the facts that presents itself is, that the injection must have directly reached the origin of the coronary arteries, and was thus applied to the heart. The force with which the injection was pushed would undoubtedly favour this, as the tube inserted into the axillary artery was small; but this would have been readily overcome by any onward movement of the blood in the aorta. I have often observed the same phenomenon when injecting other substances into the axillary artery; but sometimes the heart remains unaffected, until sufficient time has elapsed for the blood containing the substance to go the round of the general and pulmonary circulation.

The general symptoms that follow the introduction of the salts of lime into the veins, differ from those produced by the salts of copper, by the prostration of strength not being so great, and also by the occurrence of spasmodic and convulsive movements. A solution containing six grains of nitrate of lime, merely produced an expression of pain when injected into the veins; the injection of fourteen grains was followed by a spasmodic affection of the diaphragm; this was manifested only during expiration, which was performed by two or three distinct efforts: the same phenomenon has been noticed threetimes: sometimes the animal appeared to cough two or three times during each expiration. Efforts to vomit are sometimes produced. When a considerable quantity of the salt (from thirty to forty grains) has been injected into the veins, the strength of the animal becomes affected; it either lies down, or walks about very unsteadily; the sensibility does not seem altered. When the salt is introduced into the veins in sufficient quantities to arrest the action of the heart, general spasm is produced in about fourteen seconds. This soon relaxes, and respiration goes on for forty or fifty seconds. The respiratory movements are then suspended, but recommence in about a minute, when eight or ten full inspirations generally take place. A few seconds before respiration ceases, the whole of the muscles of the body are observed to be in motion. These motions appear to be owing to irregular contractions

of the fasciculi of the muscles, not affecting the whole of the muscle at once, for they do not give rise to any general movement. It resembles in fact, a sort of thrill running through the muscles, which is most perceptible in the muscles of the trunk, where it continues some few seconds after respiration has ceased. It will be seen that this phenomenon is much more strongly marked after the injection of the salts of strontia into the veins. In all the experiments I have made with the salts of lime, the lungs have always been found of a bright rosy tint ;—in other respects their tissue has appeared unaltered. The colour of the blood in the left cavities of the heart, has always been of a bright scarlet, and very firmly coagulated, the salts of lime evidently favouring the coagulation of the blood.

Strontia.—The salts of strontia, when injected into the veins, destroy the irritability of the heart more readily than the salts of lime. In smaller quantities they render the action of the heart slower, and produce a diminution of pressure in the arterial system, immediately after their injection. After a few seconds, the pressure in the arteries again becomes augmented, and the oscillations in the column of mercury are increased. Ten seconds after the introduction into the veins of a solution containing ten grains of chloride of strontium, the column of mercury in the hæmadynamometer fell one inch and a-half, and appeared to remain stationary for two or three seconds ; it then again rose, and fifteen seconds after the injection, it was two inches higher than before the introduction of the salt into the veins. The oscillations in the column were also much increased : before the injection, the systole of the heart only produced an oscillation of an inch in the column of mercury ; it now oscillated three inches at each pulsation ; the action of the heart was also slower. The introduction of forty grains into the jugular vein, arrested the action of the heart in fifteen seconds. On opening the thorax, both auricles were pulsating regularly ; the ventricles were motionless ; both the right and left cavities of the heart were full of blood ; that in the left side was of a bright scarlet colour. On introducing a solution of chloride of strontium into the arteries, the pressure in the arterial system becomes augmented about four seconds after the introduction of the first portion of the injection. The column of mercury in the hæmadynamometer is rendered about two inches higher, which is sufficient to show, that the passage of the blood through the systemic capillaries is affected. A solution containing thirty-six grains of the salt, arrested the action of the heart twenty seconds after its introduction into the artery. Thirty grains had been previously injected into the arteries of the same animal.

The salts of strontia, when introduced into the blood, appear to act but slightly on the nervous tissue, the action of the heart

being arrested previously to any general symptoms manifesting themselves. Even when injected into the arteries, these salts arrive at the heart in a sufficiently concentrated state to arrest its pulsations, without having given rise to any marked symptoms of their action on the nervous centres. In the experiment above alluded to, in which the heart was arrested by the injection of thirty-six grains into the arteries, the respiration continued uninterruptedly for three minutes after the action of the heart had ceased, and the conjunctiva was still sensible, two minutes after the arrest of the circulation: the animal seemed to suffer a great deal of pain, thirty seconds after the introduction of the salt into the arteries. Ten grains of nitrate of strontia injected into the veins, the animal being at liberty to move about, gave rise to no marked symptom. Sixteen grains of the salt arrested the action of the heart; the animal fell down ten seconds after the introduction of the salt into the veins; there were no convulsions, and the animal remained quite sensible for some time after the heart had stopped. There was expression of pain twenty-five seconds after the injection; the respiration continued for a minute and a-quarter; it then became suspended for a minute, and again commenced, six full inspirations taking place.

But the most interesting phenomenon which is produced by the injection of this salt into the veins, is the continuance of muscular movements after death; these movements last for a quarter of an hour or twenty minutes after death, giving rise to motions of the trunk and limbs. I have observed them to continue for forty minutes in the muscle that had been exposed on opening the thorax, and this, too, with sufficient vigour to move the ribs, although the temperature of the atmosphere was but 42° Fahrenheit. I propose to give a more complete analysis of this phenomenon, when treating of the action of the salts of baryta, which, together with the salts of lead, exert the same action on the muscular fibre.

Baryta.—The salts of baryta are very analogous in their action to those of strontia; in fact, the resemblance that exists between the action of these two classes of salts is quite as marked, if not more so, than that which has been noticed between the salts of magnesia and of zinc. The only difference appears to be, that the salts of baryta are much more poisonous than those of strontia; they are, in fact, the most deleterious substances, as far as regards their action on the heart, that I have experimented with. A solution containing a quarter of a grain of chloride of barium, affected the heart twelve seconds after its introduction into the jugular. The effect on the heart, and on the pressure in the arteries, was precisely the same as that produced by the injection of ten grains of chloride of strontium. On injecting two grains into the veins, the action of the heart was arrested in twelve seconds. On open-

ing the thorax, the state of the heart was the same as in the animal which had been killed by the injection of the chloride of strontium: the auricles were contracting; the ventricles were motionless.

When injected into the arteries, the same effects are produced as when a salt of strontia is used. I shall merely, therefore, refer to the description of the action of this latter salt, without going into any details on the injection of the salts of baryta into the arteries. Their action on the nervous system is so slight, as not to give rise to any marked symptoms. When the heart is paralysed, the animal falls—general spasm comes on, which lasts for a few seconds—respiration is not generally arrested until two minutes after the action of the heart has ceased—the conjunctiva generally retains its sensibility about a minute and a-half after paralysis of the heart—the lungs are found of a bright scarlet colour, but their texture appears unaltered. Neither the salts of strontia or baryta, appear to exert any influence on the coagulation of the blood.

I shall now offer a few observations on one of the most striking phenomenon which follow the introduction of the salts of baryta into the veins. I allude to the continuation of muscular movements after death. I have already noticed this point, when speaking of the action of the salts of strontia; but as this interesting fact has not attracted the attention of previous observers, I have thought it better to give a few details relating to it. In every case in which I have injected the salts of baryta into the veins or arteries, (in all eleven times,) these muscular contractions have been noticed after death. They are never observed to take place with equal force over the whole of the body at the same time, but are always most marked in some particular set of muscles. I have observed them most frequently in the muscles of the thorax and anterior extremity; but I know of no set of muscles in which these movements have not been seen. They do not always take place immediately after death, the animal sometimes remaining quite still for some minutes, after which the muscles may commence contracting. These muscular contractions are sometimes observed at various parts of the body at the same time, or they may be confined almost wholly to one part, or they may affect different parts at different periods. In one instance I have observed them commence a few minutes after death in the muscles attached to the angles of the mouth; from thence they appeared to be propagated to the muscles of the ear, the orbit, and of the side of the face; then to the muscles of the neck, and lastly to those of the thorax and the muscles moving the anterior extremity, in which they did not take place till ten minutes after death. The force with which the muscles contract is sometimes considerable. I have seen the posterior part of the trunk moved by the contractions of

the lumbar muscles, and have on two occasions observed the body pushed away, on furnishing a *point d'appui* to the sole of the foot. This last phenomenon took place ten minutes after the thorax had been opened. The longest period after death at which I have observed these movements take place spontaneously, has been thirty minutes, and this was in the muscles moving the vibræ or whiskers; but in those muscles which had been exposed on opening the thorax, I have seen vivid contractions forty minutes after death, and this when the temperature of the room in which the animal lay was at 42° Fahrenheit. As to the cause of these movements, whether they be owing to an effect produced by the substance on the nervous system, or to its direct contact with the muscles, it would be premature of me to give an opinion. There is one fact, however, which I have observed, which would tend to favour this latter opinion—it is, that when these substances had been injected into the arteries, the contractions are generally most marked in the limb, through the artery of which the injection has been introduced. No other substances that I am aware of, besides the salts of strontia, baryta, and lead, give rise to these movements.*

Lead.—The salts of this metal, in their physiological action, form a link between the salts of strontia and baryta, and those of silver and soda; they agree with the former, in their action on the heart and muscular tissue, and with the latter, by the effects they give rise to in the lungs. When a solution containing three grains of acetate of lead, was injected into the veins, the pressure in the arterial system became diminished in seven seconds; four seconds after this, the column of mercury in the hæmadynamometer regained its former height, and even indicated an increased pressure in the arteries; the action of the heart was rather quickened, and its oscillations less. On injecting a solution containing fourteen grains of the salt, the pressure in the arteries was first diminished, and then became much increased, the column of mercury being from three to four inches higher, about eighteen seconds after the injection; the action of the heart became slower, about two minutes after the injection, and the oscillations at each pulsation of the heart became increased. On injecting a solution containing

* The action of the acetate of lead on the muscles, appears not to have escaped the careful observation of Haller, for I find a passage in Scheele's work—(Ueber die transfusion des Blutes, B. 2, S. 253,) in which he states, when alluding to the experiments performed by Sproegel with this substance, "that the irritability of the other muscles did not appear to have been so completely destroyed as that of the heart, on the irritability of which organ after death, Haller was unable to institute any researches." Unfortunately, I have not been able to find in any of Haller's works the original passage to which Scheele refers. The only other allusion I find to this subject, is in Orfila's work on poisons, where he states, that on opening an animal which had been killed by an injection of chloride of barium into the veins—"les chairs étoient palpitantes."

Experiments that have been performed on birds since this memoir was written, prove that these salts exert the same action on the muscular fibre of this class of animals as they do on that of mammalia.—J. B. April 23.

a drachm of the salt, into the veins, the action of the heart appeared arrested in ten seconds, and the mercury in the instrument fell rapidly. The animal, apparently dead, lay perfectly still a minute and a-half after the injection, and remained so for two minutes; the heart again commenced pulsating, and the pressure in the arterial system was in a few seconds equal to a column of mercury of eight inches; the respiration was also renewed. On injecting another drachm of the salt, (making in the whole three drachms that had been introduced into the veins,) the action of the heart was arrested in ten seconds, but the pulsation of the auricles appeared to have been renewed after about a minute, producing no effect, however, on the pressure in the arterial system, which remained at zero. On opening the thorax, the right auricle was found contracting feebly, but no movements were observed on any other part of the organ; both cavities of the heart contained a great deal of blood; that in the left side was more scarlet than that in the right; but not of that bright colour which arterial blood ought to be. When introduced into the arteries, the acetate of lead is found to influence the passage of the blood through the general capillaries. A solution, containing six grains of the salt, when injected into the axillary artery, produced a sudden increase of pressure in the arterial system; in four seconds, the column of mercury, which, before the injection, was equal to from five to six inches, began rapidly to rise; thirty seconds after the injection, it was up to twelve inches; the respiration was arrested a minute and a-quarter after the injection; yet the pressure in the arteries diminished but slowly. Two minutes after respiration had ceased, the column of mercury was still equal to ten inches; at three minutes, it had fallen to six inches, the contractions of the heart continuing regularly, and producing oscillations of half-an-inch; at five minutes after respiration had ceased, the mercury had sunk to three inches and a-half; and at seven minutes, the pressure in the arteries was only equal to a column of mercury of two inches, the action of the heart being very weak. After having reached this point, it again became increased to six inches, on the animal taking a deep inspiration; the pulsations of the heart were also rendered quicker by this renewed arterialization of the blood. There was but one respiratory movement; and after this, the heart gradually became arrested as in asphyxia.

The general symptoms that this substance produces when introduced into the blood, are owing to two causes, either to its action on the lungs and heart, or on the nervous system; when injected into the veins, it kills by its action on the lungs, or by destroying the irritability of the heart; but when introduced into the arteries, it evidently proves fatal, by its action on the nervous system, with which it is brought into direct contact. Its action on the lungs is shown a few minutes after the introduction of the

salts of lead into the veins, by the escape of a great deal of froth from the mouth ; this froth, having its origin in the air cells, speedily prevents the aeration of the blood, the bronchi and trachea being filled with it, and the animal thus perishes by asphyxia. After death, the lungs present the appearances observed in the first and second stages of pneumonia, their tissue being engorged and hepatized ; livid spots were observed on the surface, and the air passages were either impermeable, or filled with a frothy fluid of a whitish colour, but slightly tinged with the colouring matter of the blood.* This secretion is formed in considerable quantities ; for although I have never been able to collect the whole that has been produced, yet I have certainly seen as much as four or five ounces escape at the trachea. In one experiment an animal was suffocated by this secretion, after only three grains of acetate of lead had been introduced into the veins.

The action of this salt on the nervous system is most strikingly shown when it is introduced into the arteries. In an experiment to which I have already alluded, respiration became suspended a minute and a-quarter after the introduction of six grains of acetate of lead into the arteries, (six grains had been previously introduced in divided doses), the animal lay apparently dead, with the exception of occasional movements of the tail during the space of seven minutes ; these movements of the tail were very violent, but were not accompanied by any movements in other parts of the body ; at the end of seven minutes, however, the animal gave a deep inspiration, and then lay perfectly lifeless. I have often observed this renewed respiration follow the injection of the acetate of lead into the veins ; but in these cases it took place about two minutes and a-half after respiratory movements had been suspended. When introduced into the veins in sufficient quantity, the acetate of lead arrests the action of the heart, by its direct influence on that organ. The quantity required to produce this effect seems to vary in different cases ; in the experiment above alluded to, three drachms were injected into the veins, before the action of the heart was stopped, whilst I have in another instance seen the heart lose its irritability, by the introduction of only twelve grains of the salt into the venous system. I would observe, that, in the former experiment, the animal was larger, weighing about twenty pounds ; and also that the salt was dissolved in spring-water, acidulated with a small quantity of acetic acid, distilled water being used in the other experiments. As I have before stated, this salt gives rise to contractions in the muscular fibre after death ; but not in so striking a manner, as do the salts of strontia and baryta. The acetate of lead alters the physical character of the blood, no

* A superficial analysis of this fluid shows that it contains a considerable quantity of albumen, and by incineration lead can be readily detected in it.

perfect coagulation taking place, when it has been injected into the veins.

Silver.—The salts of silver agree with those of lead in their action on the systemic capillaries. They exert a more marked action on the lungs; but do not appear to be so deleterious to the heart. The injection of a solution, containing three grains of nitrate of silver, into the veins, causes at first a diminution, and then an increase in the pressure in the arterial system. After a short time, the pressure in the arteries was again diminished, the blood being no longer aerated, owing to the production of a great quantity of frothy fluid in the air passages. The animal died from asphyxia, about twelve minutes after the introduction of the salt into the veins. On opening the thorax immediately after respiration had ceased, the heart was found pulsating rhythmically, and continued contracting for ten minutes. The right auricle was irritable half-an-hour after the thorax had been opened. If the salt is introduced into the veins in larger quantities, the passage of the blood through the lungs is suddenly arrested. On injecting into the veins a solution, containing a drachm of the salt, the column of mercury in the hæmadynamometer began to sink ten seconds after the injection. At thirteen seconds, it was equal to only two inches; nor were any oscillations produced in it, although the heart continued beating strongly. The pressure in the arteries never again rose, no blood being sent into the arterial system, although the heart could be seen to be beating violently, by the movements it produced in the parietes of the thorax. The heart was exposed three minutes after the injection; the left ventricle was still contracting feebly. The right side of the heart was so distended with blood, as to render contraction impossible. The left cavities appeared to have contracted on their contents, as much as the distended state of the right side would allow. There was a small quantity of dark blood in the left ventricle, which was coagulated, though not very firmly, having been evidently slightly affected by the salt, a small portion of which must have found its way to the left side of the heart; on opening the right ventricle, more than seven ounces of dark grumous blood escaped from it, and from the vessels leading to it. I have, on one occasion, seen the action of the heart arrested after this substance had been introduced into the veins. It was, when a solution containing twelve grains of the salt had been used, the jugular veins having been divided immediately after the injection, in order to prevent the brain from being affected, by the great venous congestion that supervenes on the arrest of the passage of the blood through the lungs. By this means, respiratory movements were not immediately arrested, and the blood containing the salt passed into the left side of the heart, and became circulated over its parietes,

giving rise to an arrest of its contractions ; but, even in this case, the ventricles manifested signs of irritability ten minutes after death.

When introduced into the arteries, the salts of silver evidently give rise to a great obstruction in the passage of the blood through the systemic capillaries. A solution containing two grains of the acetate of silver, when introduced into the arterial system, gave rise to an increase of pressure equal to a column of mercury of six inches, two grains more injected into the arteries were sufficient to produce death, giving rise to a pressure equal to fifteen inches of mercury. The heart was stopped by asphyxia about five minutes after the respiration had ceased.

When the salts of silver are injected into the veins, the nervous system appears to be affected solely by the action of these salts on the lungs ; for I have never noticed a symptom which I could attribute to their direct action on the nervous tissue. When the quantity injected is small, the circulation through the lungs is not arrested, but a secretion takes place, which, by obstructing the air-passages, prevents the aeration of the blood, thus indirectly affecting the nervous system. This secretion, however, is not always produced, and yet even then the animal seems to die of asphyxia, owing to extensive changes produced in the tissue of the lungs. When the quantity of the salt is sufficient to arrest the pulmonary circulation, the nervous system becomes violently affected, owing to the congestion which immediately takes place in the veins. In a few seconds, opisthotonos comes on, and the spasm relaxes in a short time by the death of the animal. The effects that follow the introduction of the salts of silver into the arteries closely resemble those produced by the acetate of lead, when injected into these vessels. Respiration was suddenly suspended by the injection of two grains of nitrate of silver into the arteries. This, however, was the second injection ; the respiratory movements were renewed three minutes after they had been suspended, five or six full inspirations taking place. I am inclined to think the symptoms which manifested themselves in this case, as well as those produced by the salts of lead when injected into the arteries, are owing to the great increase of pressure in the arterial system. The influence which the salts of silver exert over the passage of the blood through the lungs, is evidently much greater than that of the salts of lead, for I have never seen the circulation of the blood through these organs totally arrested by any dose of the latter salts.

Soda.—The salts of soda bear a close resemblance in their physiological action to those of silver. In the memoir I read at the Academie des Sciences at Paris, to which I have before alluded, I have related experiments showing the action of these salts when injected into the veins and arteries. As might be expected, the

quantities required to produce the same effects are much greater when the salts of soda are used, than when those of silver are injected. I have nothing to add to what has already been published on this subject, except the following experiment, which I performed at the suggestion of my esteemed friend, Dr Sharpey. It furnishes an additional proof of the correctness of the opinion that I had advanced, as to the cause of the violent nervous symptoms that manifest themselves when the pulmonary circulation is arrested; and it also shows how slight the action of the salts of soda must be on the nervous tissue. The animal was prepared by exposing the two jugular veins. A solution containing three drachms of nitrate of soda was injected into the venous system,—a quantity more than sufficient to arrest the passage of the blood through the lungs. A few seconds after the injection, violent opisthotonos came on. This was instantly relieved by dividing both the jugular veins,—congestion in the brain being thus prevented. The animal lost a great deal of blood, but, notwithstanding this, it survived the experiment two hours. During this time, no nervous symptoms were observed; the respiration continued regular, there were no convulsions, and sensibility was unimpaired; in fact, the animal seemed only to be suffering from loss of blood, the presence of so large a quantity of soda in the blood being apparently in no way injurious, except by the action it exerted on the lungs. At the end of two hours, the animal died asphyxiated, owing to the air-passages becoming filled with a frothy secretion.* Neither the salts of soda or silver exert any marked action on the general muscular tissue. The irritability of the muscles after death appears to be rather increased.

It now remains for me to notice the action of the salts of ammonia and potassa, when injected into the vascular system. The physiological action of these substances, does not allow of their taking a place in that natural arrangement we have hitherto pursued, and by which a substance has always stood in close physiological relation to that which preceded it, and that which followed it. There is undoubtedly a great difference between the action of the salts of magnesia and those of soda, yet have we observed the phenomena which characterize the action of these substances, gradually manifesting themselves, or disappearing; becoming more strongly marked or less perceptible; so that although we find no one phenomenon common to the substances which occupy the extremes of the series, yet we are unable to se-

* It would appear that Orfila has noticed the production of this frothy secretion, both when the salts of lead, and those of silver had been injected into the veins. He, however, seems to have attributed its origin to the stomach, as, he says, the animal vomited some whitish substance, together with a great quantity of frothy fluid.

lect any two substances placed together in the intermediate part of the scale, which, in their physiological action, do not present many striking points of resemblance.

Ammonia.—The salts of ammonia, like most of those substances we have hitherto examined, exert, when injected into the veins, a powerful action on the heart: they differ, however, from the other substances, with which they may agree in this respect, both by their influence on the systemic capillary circulation, and by the general symptoms they give rise to. When a weak solution of the salts of ammonia is injected into the veins, the action of the heart becomes quickened, and the pulse smaller, a few seconds after the injection; the pressure in the arteries soon becomes increased, owing to the passage of the blood through the systemic capillaries being impeded. A solution containing six grains of nitrate of ammonia, caused an increased pressure in the arterial system fifteen seconds after its injection, and at thirty-five seconds, the column of mercury, which, before the injection was from four to six inches, rose to eleven inches:—this pressure soon became reduced, and the action of the heart irregular, great oscillations taking place in the mercury. A solution containing forty grains of the salt, when injected into the jugular vein, caused an augmentation in the pressure in the arterial system, the column of mercury rising as high as fourteen inches a few seconds after the injection. The action of the heart was arrested ten seconds after the introduction into the veins, of a solution, containing two drachms of the salt. On opening the thorax, the auricles were contracting rhythmically, the ventricles appeared to have totally lost their irritability; both cavities of the heart were full of blood, that in the left side was scarlet, and firmly coagulated, that in the right was of the ordinary colour of venous blood. The alkali itself in much smaller quantity produces the same effects. A solution containing twenty drops of a saturated solution of ammonia, arrested the action of the heart.

When injected into the arteries, the salts of ammonia caused a great augmentation in the pressure in the arterial system. A solution containing two drachms of the nitrate of ammonia, arrested the action of the heart a minute after it had been injected into the axillary artery, respiratory movements continuing forty-five seconds after the heart had stopped.

The salts of ammonia appear to exert a powerful action on the nervous system, agreeing in this respect much more closely with poisons derived from the organic kingdom, than with any inorganic substances whose action I have yet investigated. Fourteen seconds after the injection into the veins, of a solution containing ten grains of nitrate of ammonia, the animal fell on its side, affected with slight convulsions. After lying on the ground for half-a

minute, it rose, and did not appear to suffer the least from the injection; the introduction of ten grains more into the veins, gave rise to similar symptoms; the animal fell, but was again on its legs thirty seconds after the injection, the action of the salt being apparently due to its contact with the nervous tissue, and passing off as soon as blood which does not contain the salt, becomes again circulated over the nervous centres. Some minutes after the injection, the hind legs appeared to be partially paralysed, and after this had passed off, first one fore leg, and then the other, appeared to suffer in the same manner. Introduced into the veins in larger quantities, these salts appear to deprive the animal of the power of controlling its own movements; the whole body becomes violently convulsed; these convulsions very much resemble those produced by strychnine, and like these, can be called out by the slightest jar or noise; the respiratory movements remain unaffected, and the sensibility appears unimpaired, even in the midst of the strongest convulsions. In two instances, after death produced by the salts of ammonia, I have observed a sort of convulsive movement of the left auricle, which was in violent motion, without, however, causing any regular contraction, the movement seeming to part from many distinct points.*

Potass.—The accounts I have already published of the action of the salts of potass, (see *Arch. Gen. de Med.* Nov. 1839), will enable me to pass them over, with but a slight notice of their action on the vascular system. In this respect they exactly resemble the salts of ammonia, producing the same action on the general capillaries, and on the heart; both these substances appear also not to exert the least action on the lungs. In their action on the nervous system, there is a marked difference between the salts of potass and those of ammonia,—the violent convulsions which follow the introduction of the latter class of salts into the veins, never being observed when the salts of potass are injected. After the injection of six grains of the bicarbonate of potass into the veins, I have seen the animal fall on its side, the posterior extremities slightly convulsed, and after a few seconds again rise. On introducing an additional quantity of the salt, the heart was arrested, the salts of potass, after those of baryta, being the most fatal to the action of the heart.

Having now related the more important facts that have been observed in my experiments, I shall conclude by offering one or two observations which they appear to suggest.

The most curious fact which a consideration of the above experiments presents to us, is the close connection which appears to exist between the physiological action of these substances and their

* Convulsive movements caused by the injection of hydrochlorate of ammonia into the veins have been noticed by previous observers.

chemical properties. The property to which I more particularly allude is that of isomorphism. The analogy in the physiological action of isomorphous compounds, was first forced on my notice, by the perfect resemblance that exists between the salts of silver and of soda in their action on the animal economy when injected into the veins; differing as these salts do in their sensible chemical properties, but closely connected as being isomorphous. Further investigation soon proved to me that this fact was of far more importance than I had anticipated, by showing that most striking points of resemblance exist generally between isomorphous compounds in their action on the animal tissues, when introduced directly into the blood.* In the present state of chemistry, it would be useless to attempt to derive from this fact any farther generalization, for the doctrine of isomorphism itself is still in a very imperfect state.

Before quitting this point, I would recall an observation I made in a memoir read at the Academie des Sciences in June 1839, and which follows as a corollary from the fact of the analogous action of isomorphous compounds, although I had no idea at that time that such was the case. In speaking of the connection between the chemical composition of substances, and their physiological action, I observed—"Ce rapport n'est pas il est vrai tel que l'état actuel de la science des affinités chimiques nous l'eût fait prévoir, car il est évident, que les décompositions effectuées dans le sang, par ces sels, décompositions sur lesquelles roule toute leur action physiologique, il est évident dis-je, que ces décompositions dependent de la base du sel, et ne sont que peu modifiées par la combinaison de cette base avec les acides, même les plus puissants." Should a more extended series of researches prove, that the identity which I have noticed in the physiological action of isomorphous compounds does not in all cases hold good, yet I think that the facts above related are sufficient to show that there exists some intimate connection between the chemical properties of substances, and their physiological action, the investigation of which promises to furnish a rich field for physiological researches.

* There is apparently an exception to this rule in the salts of lead, which, although agreeing with those of baryta and strontia, in most of the phenomena they present, yet exert an analogous action on the pulmonary tissue with the salts of silver, with which they are not isomorphous; but it has been remarked, that there exists a striking analogy in the combinations which the oxides of these two bases form with organic compounds; an analogy which does not exist between the oxide of silver and any other base. There is also a difference between the action of the salts of potass, and those of ammonia on the nervous tissue; for although these substances exert the same action on the heart and the systemic capillaries, yet the violent nervous symptoms that follow the introduction of the salts of ammonia into the veins are never produced by the salts of potass. The peculiar action of the salts of ammonia on the nervous tissue,—an action which connects it in its physiological relations with organic poisons,—is interesting when we consider the chemical composition of ammonia, standing as it does between organic and inorganic compounds.