12. Katira BH, Giesinger RE, Engelberts D, Zabini D, Kornecki A, Otulakowski G, Yoshida T, Kuebler WM, McNamara PJ, Connelly KA, et al. Adverse heart-lung interactions in ventilator-induced lung injury. *Am J Respir Crit Care Med* 2017;196:1411–1421.
13. Vieillard-Baron A, Loubieres Y, Schmitt JM, Page B, Dubourg O, Jardin F. Cyclic changes in right ventricular output impedance during mechanical ventilation. *J Appl Physiol* (1985) 1999;87:1644–1650.
14. Mekontso Dessap A, Boissier F, Charron C, Bégot E, Repessé X, Legras A, Brun-Buisson C, Vignon P, Vieillard-Baron A. Acute cor pulmonale during protective ventilation for acute respiratory distress syndrome: prevalence, predictors, and clinical impact. *Intensive Care Med* 2016;42:862–870.
15. Taylor RR, Covell JW, Sonnenblick EH, Ross J Jr. Dependence of ventricular distensibility on filling of the opposite ventricle. *Am J Physiol* 1967;213:711–718.
16. Scharf SM, Brown R, Saunders N, Green LH, Ingram RH Jr. Changes in canine left ventricular size and configuration with positive end-expiratory pressure. *Circ Res* 1979;44:672–678.
17. Whittenberger JL, McGregor M, Berglund E, Borst HG. Influence of state of inflation of the lung on pulmonary vascular resistance. *J Appl Physiol* 1960;15:878–882.
18. Vieillard-Baron A, Matthey M, Teboul JL, Bein T, Schultz M, Magder S, Marini JJ. Experts' opinion on management of hemodynamics in ARDS patients: focus on the effects of mechanical ventilation. *Intensive Care Med* 2016;42:739–749.
19. West JB, Tsukimoto K, Mathieu-Costello O, Prediletto R. Stress failure in pulmonary capillaries. *J Appl Physiol* 1985;1991:1731–1742.

Copyright © 2017 by the American Thoracic Society

Toward Early Identification of Clinically Relevant Interstitial Lung Disease

With advances in the treatment of pulmonary fibrosis (1, 2) comes growing awareness of the potential usefulness of early identification of at-risk individuals (or those affected but asymptomatic). The optimal approach to identification of at-risk individuals is undetermined, but the use of computed tomography (CT) scans hold promise.

Numerous studies have documented the relevance of visually identified, variously defined “interstitial lung abnormalities” (ILA), which are seen in approximately 7% of the general population (3). The presence of ILA is associated with increased risk of death in multiple international cohorts (4). ILA appear to be persistent in roughly 60% of reimaged subjects, with progression seen in 20–43% on interval follow-up scans obtained 2–6 years later (5, 6).

Use of subjective (visual) markers of a preclinical interstitial lung disease (ILD) phenotype is problematic because it is time-consuming and subjective, with high interobserver variability, and reveals indeterminate findings on up to 40% of scans (3). Many of these barriers may be surmountable by the development and validation of automated or semiautomated methods of CT analysis. High attenuation areas (HAA) are an automated, quantitative measure of lung density obtained via analysis of inspiratory CT scans. The presence of higher HAA on CT scanning done for nonpulmonary indications (i.e., cardiovascular disease screening) is associated with greater cumulative cigarette smoke exposure and spirometric restriction (reduced FVC) (7), elevated serum matrix metalloproteinase-7 and IL-6, lower 6-minute walk distance and cardiopulmonary exercise testing breathing reserve, increased odds of identification of visual ILA on subsequent whole-lung CT, and increased hazard of all-cause mortality (8). Visually identified ILA and automated measurement of HAA appear to capture somewhat different pathoanatomic and phenotypic characteristics. For example, the positive predictive value of high HAA (i.e., >10% HAA) for ILA is just 0.19, with very poor agreement between

the two measures ($\kappa = 0.13$) (9). In addition, HAA are not associated with *MUC5B* polymorphisms, whereas ILA are (9).

In this issue of the *Journal*, Podolanczuk and colleagues (pp. 1434–1442) report an important advancement in the effort to validate HAA as an index for preclinical ILD (10). The authors examined 6,808 community-dwelling adults enrolled in the MESA (Multi-Ethnic Study of Atherosclerosis) study, an NIH/NHLBI-funded observational study designed to investigate subclinical cardiovascular disease. All subjects lacked a history of cardiovascular disease or other life-limiting comorbidities, and underwent cardiac CT scanning on study enrollment, with HAA measured on these scans (defined as the percentage of imaged lungs having CT attenuation values between –600 and –250 Hounsfield units). During a median follow-up period of 12.2 years, 52 participants were identified to have ILD at the time of a hospitalization or death. After accounting for potential confounding factors, the authors found that higher HAA was associated with increased odds of having an adjudicated ILD diagnosis, an increased rate of respiratory and ILD hospitalizations, an increased rate of ILD death, and increased odds of subsequent self-reported dyspnea.

This is the first report linking a quantitative CT finding to serious ILD events, including ILD hospitalizations and deaths. The weak relationship between HAA and visible ILA suggests that HAA may be a surrogate for an interstitial abnormality that is not yet visible. It could also be an index of poor lung inflation due to increased lung stiffness and/or decreased muscle strength.

The long follow-up interval, and the robust event adjudication method, are clear strengths of this study. The authors also gave careful attention to sensitivity analyses, providing reassurance that these associations were not driven by a small group of subjects in the top quartile of ILA who may have had undiagnosed ILD. An important limitation is a lack of data on less severe ILD manifestations. Patients with ILD may have substantial morbidity despite lack of death or hospitalization for ILD during the study period, and would potentially benefit from early identification. The relationship of HAA with other ILD events is not captured here. Validation of the relationship of HAA with ILD events by whole-lung CT imaging, in addition to cardiac CT (which images roughly 70% of the lung), would also be useful. Technical issues that

should be explored include determination of the optimal upper and lower Hounsfield unit thresholds for HAA in risk prediction of ILD, and the value of adjustment for inspired lung volume, interscanner variability, and other sources of variation. A puzzling feature of this study is the fact that high HAA was associated with a higher proportion of females and never-smokers, and a lower proportion of white subjects, all features that should have been associated with decreased risk of death from ILD.

Various questions in this area remain unanswered. First, is the HAA index robust enough to be applied to modern-day CT scans, which use substantially lower CT dose and different reconstruction techniques than those used in the MESA study? Second, are the sensitivity and specificity of the technique adequate for screening populations with known (or suspected) risk factors such as smoking history or occupational exposure? Third, the confounding effect of smoking status is unclear. Current smoking is clearly associated with increased lung attenuation (11, 12). Smoking status was also a strong effect modifier of the association of HAA with matrix metalloproteinase-7, IL-6, and FVC, such that the association of these markers with HAA was lost in never-smokers and substantially stronger in ever-smokers, suggesting smoking may drive these relationships (8). Despite this, effect modification by smoking status was not identified for all-cause mortality (8), or for any of the ILD outcomes in the current study (10). Finally, is a particular automated method more useful than another in general, or when applied to a specific CT scan type or subpopulation of interest? Several alternative automated, quantitative CT analysis tools have been shown to identify patients with idiopathic pulmonary fibrosis at risk of disease progression or death (13–15), and smokers at increased risk of death (16). Eventual direct comparison of the performance of these technologies may be useful.

Although unanswered questions remain, this contribution from Podolanczuk and colleagues has increased our knowledge of the relevance of quantitatively measured lung density changes, moving us a step closer to understanding how to identify and manage individuals at risk for pulmonary fibrosis and other interstitial lung diseases. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Margaret L. Salisbury, M.D., M.S.
Department of Medicine
University of Michigan
Ann Arbor, Michigan

David A. Lynch, M.B.
Department of Radiology
National Jewish Health
Denver, Colorado

ORCID ID: 0000-0001-8217-6955 (M.L.S.).

References

1. King TE Jr, Bradford WZ, Castro-Bernardini S, Fagan EA, Glaspole I, Glassberg MK, Gorina E, Hopkins PM, Kardatzke D, Lancaster L, et al.;

- ASCEND Study Group. A phase 3 trial of pirfenidone in patients with idiopathic pulmonary fibrosis. *N Engl J Med* 2014;370:2083–2092.
2. Richeldi L, du Bois RM, Raghu G, Azuma A, Brown KK, Costabel U, Cottin V, Flaherty KR, Hansell DM, Inoue Y, et al.; INPULSIS Trial Investigators. Efficacy and safety of nintedanib in idiopathic pulmonary fibrosis. *N Engl J Med* 2014;370:2071–2082.
3. Hunninghake GM, Hatabu H, Okajima Y, Gao W, Dupuis J, Latourelle JC, Nishino M, Araki T, Zazueta OE, Kurugol S, et al. *MUC5B* promoter polymorphism and interstitial lung abnormalities. *N Engl J Med* 2013;368:2192–2200.
4. Putman RK, Hatabu H, Araki T, Gudmundsson G, Gao W, Nishino M, Okajima Y, Dupuis J, Latourelle JC, Cho MH, et al.; Evaluation of COPD Longitudinally to Identify Predictive Surrogate Endpoints (ECLIPSE) Investigators; COPDGene Investigators. Association between interstitial lung abnormalities and all-cause mortality. *JAMA* 2016;315:672–681.
5. Araki T, Putman RK, Hatabu H, Gao W, Dupuis J, Latourelle JC, Nishino M, Zazueta OE, Kurugol S, Ross JC, et al. Development and progression of interstitial lung abnormalities in the Framingham Heart Study. *Am J Respir Crit Care Med* 2016;194:1514–1522.
6. Jin GY, Lynch D, Chawla A, Garg K, Tammemagi MC, Sahin H, Misumi S, Kwon KS. Interstitial lung abnormalities in a CT lung cancer screening population: prevalence and progression rate. *Radiology* 2013;268:563–571.
7. Lederer DJ, Enright PL, Kawut SM, Hoffman EA, Hunninghake G, van Beek EJ, Austin JH, Jiang R, Lovasi GS, Barr RG. Cigarette smoking is associated with subclinical parenchymal lung disease: the Multi-Ethnic Study of Atherosclerosis (MESA)–lung study. *Am J Respir Crit Care Med* 2009;180:407–414.
8. Podolanczuk AJ, Oelsner EC, Barr RG, Hoffman EA, Armstrong HF, Austin JH, Basner RC, Bartels MN, Christie JD, Enright PL, et al. High attenuation areas on chest computed tomography in community-dwelling adults: the MESA study. *Eur Respir J* 2016;48:1442–1452.
9. Kliment CR, Araki T, Doyle TJ, Gao W, Dupuis J, Latourelle JC, Zazueta OE, Fernandez IE, Nishino M, Okajima Y, et al. A comparison of visual and quantitative methods to identify interstitial lung abnormalities. *BMC Pulm Med* 2015;15:134.
10. Podolanczuk AJ, Oelsner EC, Barr RG, Bernstein EJ, Hoffman EA, Easthausen IJ, Hinckley Stukovsky K, RoyChoudhury A, Michos ED, Raghu G, et al. High-attenuation areas on chest computed tomography and clinical respiratory outcomes in community-dwelling adults. *Am J Respir Crit Care Med* 2017;196:1434–1442.
11. Shaker SB, Dirksen A, Lo P, Skovgaard LT, de Bruijne M, Pedersen JH. Factors influencing the decline in lung density in a Danish lung cancer screening cohort. *Eur Respir J* 2012;40:1142–1148.
12. Zach JA, Williams A, Jou SS, Yagihashi K, Everett D, Hokanson JE, Stinson D, Lynch DA; COPDGene Investigators. Current smoking status is associated with lower quantitative CT measures of emphysema and gas trapping. *J Thorac Imaging* 2016;31:29–36.
13. Humphries SM, Yagihashi K, Huckleberry J, Rho BH, Schroeder JD, Strand M, Schwarz MI, Flaherty KR, Kazerooni EA, van Beek EJR, et al. Idiopathic pulmonary fibrosis: data-driven textural analysis of extent of fibrosis at baseline and 15-month follow-up. *Radiology* 2017;285:270–278.
14. Salisbury ML, Lynch DA, van Beek EJ, Kazerooni EA, Guo J, Xia M, Murray S, Anstrom KJ, Yow E, Martinez FJ, et al.; IPFnet Investigators. Idiopathic pulmonary fibrosis: the association between the adaptive multiple features method and fibrosis outcomes. *Am J Respir Crit Care Med* 2017;195:921–929.
15. Maldonado F, Moua T, Rajagopalan S, Karwoski RA, Raghunath S, Decker PA, Hartman TE, Bartholmai BJ, Robb RA, Ryu JH. Automated quantification of radiological patterns predicts survival in idiopathic pulmonary fibrosis. *Eur Respir J* 2014;43:204–212.
16. Ash SY, Harmouche R, Ross JC, Diaz AA, Hunninghake GM, Putman RK, Onieva J, Martinez FJ, Choi AM, Lynch DA, et al. The objective identification and quantification of interstitial lung abnormalities in smokers. *Acad Radiol* 2017;24:941–946.

Copyright © 2017 by the American Thoracic Society