but some centers must have used it substantially more often, considering that only half of the centers reported any NIV use on Days 1 and 2. Also, given that NIV use in patients with severe ARDS was as prevalent as in those with mild ARDS, and that more recent NIV guidelines have made no recommendation for its use in acute lung injury or pneumonia (10) or have suggested its use only for mild ARDS (11), it appears that some centers in the LUNG SAFE study were overzealous in their application of NIV to patients with ARDS.

Even more disturbing is the finding of the propensity-matched analysis showing increased ICU mortality associated with NIV over invasive ventilation in patients with ARDS with $\text{PaO}_2/\text{FiO}_2$ less than 150. In the context of the previous studies showing high NIV failure rates in patients with hypoxemic respiratory failure, as well as the more recent study by Frat and colleagues (6) showing advantages of HFNO over NIV, this observation raises the question of whether NIV should be used at all in sicker patients with ARDS. However, it seems premature to make such a recommendation, especially with the recent RCT (12) showing that in patients treated with NIV for ARDS, intubation (18 vs. 62%) and mortality rates were much lower during use of the helmet interface as compared with a full face mask. This suggests that the technique of administering NIV is an important determinant of success. We have learned not to rely too much on single RCTs to guide ICU management, so further studies are urgently needed to sort out the relative merits of NIV delivered via a face mask, helmet, or HFNO versus prompt intubation in the therapy of ARDS. In the meantime, though, NIV should be used with great caution, if at all, in patients with ARDS with more severe oxygenation defects. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Nicholas S. Hill, M.D.
Erik Garpestad, M.D.
Tufts University
Boston, Massachusetts

References

1. Nava S, Hill N. Non-invasive ventilation in acute respiratory failure. *Lancet* 2009;374:250–259.
2. Garpestad E, Schumaker G, Hill NS. Noninvasive ventilation for acute respiratory distress syndrome: breaking down the final frontier? *Crit Care Med* 2007;35:288–290.
3. Antonelli M, Conti G, Moro ML, Esquinas A, Gonzalez-Diaz G, Confalonieri M, Pelaia P, Principi T, Gregoretti C, Beltrame F, *et al.* Predictors of failure of noninvasive positive pressure ventilation in patients with acute hypoxemic respiratory failure: a multi-center study. *Intensive Care Med* 2001;27:1718–1728.
4. Antonelli M, Conti G, Esquinas A, Montini L, Maggiore SM, Bello G, Rocco M, Maviglia R, Pennisi MA, Gonzalez-Diaz G, *et al.* A multiple-center survey on the use in clinical practice of noninvasive ventilation as a first-line intervention for acute respiratory distress syndrome. *Crit Care Med* 2007;35:18–25.
5. Zhan Q, Sun B, Liang L, Yan X, Zhang L, Yang J, Wang L, Ma Z, Shi L, Wei L, *et al.* Early use of noninvasive positive pressure ventilation for acute lung injury: a multicenter randomized controlled trial. *Crit Care Med* 2012;40:455–460.
6. Frat JP, Thille AW, Mercat A, Girault C, Ragot S, Perbet S, Prat G, Boulain T, Morawiec E, Cottereau A, *et al.*; FLORALI Study Group; REVA Network. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015;372:2185–2196.
7. Bellani G, Laffey JG, Pham T, Madotto F, Fan E, Brochard L, Esteban A, Gattinoni L, Bumbasirevic V, Piquilloud L, *et al.* Noninvasive ventilation of patients with acute respiratory distress syndrome: insights from the LUNG SAFE study. *Am J Respir Crit Care Med* 2017;195:67–77.
8. Bellani G, Laffey JG, Pham T, Fan E, Brochard L, Esteban A, Gattinoni L, van Haren F, Larsson A, McAuley DF, *et al.*; LUNG SAFE Investigators; ESICM Trials Group. Epidemiology, patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA* 2016;315:788–800.
9. Ozsancak Ugurlu A, Sidhom SS, Khodabandeh A, leong M, Mohr C, Lin DY, Buchwald I, Bahhady I, Wengryn J, Maheshwari V, *et al.* Use and outcomes of noninvasive positive pressure ventilation in acute care hospitals in Massachusetts. *Chest* 2014;145:964–971.
10. Keenan SP, Sinuff T, Burns KE, Muscedere J, Kutsogiannis J, Mehta S, Cook DJ, Ayas N, Adhikari NK, Hand L, *et al.*; Canadian Critical Care Trials Group/Canadian Critical Care Society Noninvasive Ventilation Guidelines Group. Clinical practice guidelines for the use of noninvasive positive-pressure ventilation and noninvasive continuous positive airway pressure in the acute care setting. *CMAJ* 2011;183:E195–E214.
11. Ferguson ND, Fan E, Camporota L, Antonelli M, Anzueto A, Beale R, Brochard L, Brower R, Esteban A, Gattinoni L, *et al.* The Berlin definition of ARDS: an expanded rationale, justification, and supplementary material. *Intensive Care Med* 2012;38:1573–1582.
12. Patel BK, Wolfe KS, Pohlman AS, Hall JB, Kress JP. Effect of noninvasive ventilation delivered by helmet vs face mask on the rate of endotracheal intubation in patients with acute respiratory distress syndrome: a randomized clinical trial. *JAMA* 2016;315:2435–2441.

Copyright © 2017 by the American Thoracic Society

What Is in a Pattern? That Which We Call Idiopathic Pulmonary Fibrosis by Any Other Pattern Would Behave Alike!

The interstitial lung diseases (ILDs), and idiopathic pulmonary fibrosis (IPF) in particular, represent a heterogeneous group of lung disorders that are linked to genetic susceptibility, environmental exposures (including cigarette smoke), and aging (1). The diagnostic

approach to ILD and IPF has evolved significantly. Evidence demonstrating the presence of pleural-based and basilar honeycomb changes on a high-resolution computed tomographic (HRCT) scan of the chest is strongly correlated with the presence of usual interstitial pneumonia (UIP) on surgical lung biopsy (2, 3); this had led to HRCT becoming the primary modality for the evaluation and diagnosis of patients with ILD and IPF. In fact, the 2011 American Thoracic Society/European Respiratory

J.A.d.A. and T.L. are supported by National Institutes of Health grant P01HL114470.

Society/Japanese Respiratory Society/Latin American Thoracic Association guidelines stated that, in the right clinical setting, HRCT demonstrating a “definite UIP” pattern would suffice, whereas HRCT with a “possible UIP” pattern would have to be paired with a surgical lung biopsy to establish the diagnosis of IPF. Definite UIP criteria included subpleural, basal predominance of reticular abnormality and honeycombing without inconsistent features such as extensive ground-glass opacities or air trapping; whereas HRCT scans with extensive subpleural and basilar predominant reticular abnormalities but no honeycombing were classified as having a possible UIP pattern (4).

Those recommendations were intended to inform both clinical practice and the design of clinical trials. Although the new HRCT-based diagnostic criteria improved specificity, it did so at the expense of sensitivity. As we apply the recommended diagnostic algorithm to clinical practice and to the design of clinical trials, some of its practical shortcomings become apparent (5). For example, what else could an older white man, without a history of environmental, occupational, or pharmacological exposures or evidence of an underlying connective tissue disorder, presenting with an HRCT scan demonstrating a possible UIP pattern, have? Studies suggest that, within the context of a clinical trial, and considering advanced age, an HRCT scan with a possible UIP pattern could reliably diagnose IPF (6, 7). Recognizing the heterogeneity of presentations in patients with IPF, it is not surprising that attempts to classify, either by “lumping” or “splitting” the disease into defined categories, is challenging.

An important issue that has arisen from the recent diagnostic guidelines is whether patients with IPF with a possible UIP pattern represent a distinct phenotype, with distinct prognosis and clinical behavior compared with those presenting with definite UIP. Furthermore, if this were the case, would they respond to antifibrotic drugs in the same manner?

In this issue of the *Journal* (pp. 78–85), Raghu and colleagues seek to begin to address these questions (8). They conducted a *post hoc* subgroup analysis using pooled data from the INPULSIS trials of nintedanib. Their primary objective was to ascertain differences in disease behavior and progression and the response to treatment between patients with honeycombing on HRCT and/or UIP confirmed by surgical lung biopsy (i.e., diagnosed with IPF according to the 2011 guidelines) and those with “possible UIP.” The possible UIP pattern was defined by the presence of reticular opacities and traction bronchiectasis, which is a slight modification of the 2011 guidelines criteria. It is important to underscore that all HRCT scans and lung biopsies were centrally reviewed before subject enrollment. The study population consisted of 723 patients who underwent HRCT scanning and/or biopsy confirming honeycombing and 338 who had features of possible UIP with traction bronchiectasis. The two groups were well matched for baseline demographic and physiological characteristics, although fewer patients without honeycombing had centrilobular emphysema. The main findings of this study were that both groups had similar rates of change in lung function, incidence of acute exacerbations, change in St. George’s Respiratory Questionnaire total score, and response to nintedanib, comparing baseline values with Week 52. The incidence of adverse events was also not different between groups.

The authors conducted a robust analysis of a well-characterized cohort of patients with IPF, suggesting that we need

to consider expanding our current definition of IPF to include those patients presenting with a possible UIP pattern and traction bronchiectasis, within the appropriate clinical context. Interestingly, both groups responded similarly to the antifibrotic agent nintedanib, suggesting that even more “advanced” disease (with honeycomb change) may be amenable to treatment. Indeed, the concept that fibrosis is an end-stage, irreversible process is challenged by findings in age-relevant animal models (9). The study by Raghu and colleagues is consistent with data from other groups indicating that patients with a definite UIP pattern, and those with a possible UIP pattern, have similar disease courses (10).

Although the study by Raghu and colleagues would support more “lumping” than “splitting” when it comes to HRCT patterns in IPF, it is also important to recognize that distinction between the findings of traction bronchiectasis and honeycombing can be subtle, both biologically and physiologically (11). Some studies suggest that the presence of traction bronchiectasis may be an independent predictor of mortality in IPF (12). It is possible that the modification in the definition applied here might have led to the selection of a subgroup with significant fibrosis, and thus not too dissimilar from those with honeycombing. One could argue that the similar baseline physiology favors that possibility. Another issue that always comes up as one appraises studies dealing with the diagnostic process in IPF is external validity. Case definition in clinical trials is often dependent on a centralized and highly specialized process that is difficult to reproduce in clinical practice.

Notwithstanding those limitations, the findings in this report advance our understanding of HRCT patterns and their relationship to IPF disease course, physiology, and even biology (based on equivalent responses to nintedanib). It also highlights the limitations of our current tools for the diagnosis and management of IPF. It defends the use of nintedanib in a broader group of patients with a clinical diagnosis of IPF, who do not stringently meet the 2011 American Thoracic Society/European Respiratory Society/Japanese Respiratory Society/Latin American Thoracic Association definition. Finally, this study reinforces the need for broader education so that the currently accepted practice in academic medical centers, using a multidisciplinary approach to the diagnosis of IPF, can be implemented in the community setting. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Joao A. de Andrade, M.D.
Department of Medicine
University of Alabama at Birmingham
Birmingham, Alabama
and
Birmingham Veterans Affairs Medical Center
Birmingham, Alabama

Tracy Luckhardt, M.D.
Department of Medicine
University of Alabama at Birmingham
Birmingham, Alabama

ORCID IDs: 0000-0002-3743-4561 (J.A.d.A.); 0000-0002-4063-5989 (T.L.).

References

1. Thannickal VJ, Zhou Y, Gaggar A, Duncan SR. Fibrosis: ultimate and proximate causes. *J Clin Invest* 2014;124:4673–4677.
2. Hunninghake GW, Zimmerman MB, Schwartz DA, King TE Jr, Lynch J, Hegele R, Waldron J, Colby T, Müller N, Lynch D, *et al.* Utility of a lung biopsy for the diagnosis of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2001;164:193–196.
3. Flaherty KR, Thwaite EL, Kazerooni EA, Gross BH, Toews GB, Colby TV, Travis WD, Mumford JA, Murray S, Flint A, *et al.* Radiological versus histological diagnosis in UIP and NSIP: survival implications. *Thorax* 2003;58:143–148.
4. Raghu G, Collard HR, Egan JJ, Martinez FJ, Behr J, Brown KK, Colby TV, Cordier JF, Flaherty KR, Lasky JA, *et al.*; ATS/ERS/JRS/ALAT Committee on Idiopathic Pulmonary Fibrosis. An official ATS/ERS/JRS/ALAT statement: idiopathic pulmonary fibrosis: evidence-based guidelines for diagnosis and management. *Am J Respir Crit Care Med* 2011;183:788–824.
5. Wells AU. “Any fool can make a rule and any fool will mind it.” *BMC Med* 2016;14:23.
6. Raghu G, Lynch D, Godwin JD, Webb R, Colby TV, Leslie KO, Behr J, Brown KK, Egan JJ, Flaherty KR, *et al.* Diagnosis of idiopathic pulmonary fibrosis with high-resolution CT in patients with little or no radiological evidence of honeycombing: secondary analysis of a randomised, controlled trial. *Lancet Respir Med* 2014; 2:277–284.
7. Fell CD, Martinez FJ, Liu LX, Murray S, Han MK, Kazerooni EA, Gross BH, Myers J, Travis WD, Colby TV, *et al.* Clinical predictors of a diagnosis of idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2010;181:832–837.
8. Raghu G, Wells AU, Nicholson AG, Richeldi L, Flaherty KR, Le Mauff F, Stowasser S, Schlenker-Herceg R, Hansell DM. Effect of nintedanib in subgroups of idiopathic pulmonary fibrosis by diagnostic criteria. *Am J Respir Crit Care Med* 2017;195:78–85.
9. Hecker L, Logsdon NJ, Kurundkar D, Kurundkar A, Bernard K, Hock T, Meldrum E, Sanders YY, Thannickal VJ. Reversal of persistent fibrosis in aging by targeting Nox4–Nrf2 redox imbalance. *Sci Transl Med* 2014;6:231ra47.
10. Lee JW, Shehu E, Gjonbrataj J, Bahn YE, Rho BH, Lee MY, Choi WI. Clinical findings and outcomes in patients with possible usual interstitial pneumonia. *Respir Med* 2015;109:510–516.
11. Piciocchi S, Tomassetti S, Ravaglia C, Gurioli C, Gurioli C, Dubini A, Carloni A, Chilosi M, Colby TV, Poletti V. From “traction bronchiectasis” to honeycombing in idiopathic pulmonary fibrosis: a spectrum of bronchiolar remodeling also in radiology? *BMC Pulm Med* 2016;16:87.
12. Sumikawa H, Johkoh T, Colby TV, Ichikado K, Suga M, Taniguchi H, Kondoh Y, Ogura T, Arakawa H, Fujimoto K, *et al.* Computed tomography findings in pathological usual interstitial pneumonia: relationship to survival. *Am J Respir Crit Care Med* 2008;177:433–439.

Copyright © 2017 by the American Thoracic Society

Respiratory Muscle Strength: A Reliable Index for Predicting Survival in Amyotrophic Lateral Sclerosis?

Amyotrophic lateral sclerosis (ALS) is a progressive motor neuron disease that leads to respiratory and bulbar insufficiency. The eventual outcome is hypercapnic respiratory failure or pneumonia. Part of the clinical assessment includes evaluation of respiratory muscle force generation. The most readily available test is the FVC. Unfortunately, FVC may not decrease until there is severe muscle weakness. Hypercapnia occurs over a wide range of VC and maximal inspiratory and expiratory mouth pressures (MIP and MEP, respectively) (1, 2). Conversely, hypercapnia exhibits a wide range even in patients with close to normal VCs (3).

A decrease in FVC on assuming the supine position has been shown to provide a more sensitive indicator of diaphragmatic weakness (4, 5). MIP and MEP provide useful information related to survival, but are a challenge to perform in the presence of bulbar weakness or submaximal effort. Another approach is the maximal sniff, generated by rapid contraction of the diaphragm and scalene muscles (6). This maneuver has been used to assess diaphragmatic and global inspiratory muscle strength by measuring sniff transdiaphragmatic pressure (Sn Pdi) and sniff esophageal pressure (Sn Pes). A variation on this approach has been the sniff nasal inspiratory pressure (SNIP) (7, 8), which avoids use of a mouthpiece and esophageal balloons, but again is problematic in patients with bulbar impairment (9). SNIP declines linearly with progression of disease (8), and is better at predicting hypercapnia than MIP or VC (10); the best SNIP is generally 6–10% higher in the last 10 of 20 sniffs than in the first 10 (11). In general, sniff tests have greater sensitivity and specificity than VC for predicting survival (2, 3, 12). In a follow-up to a comprehensive study of tests of respiratory muscle strength in 81 patients with ALS with and without bulbar

dysfunction (2), in this issue of the *Journal*, Polkey and colleagues (pp. 86–95) report measures of respiratory muscle strength as they related to mortality and use of noninvasive ventilation (NIV) 15 years after the initial study was published (13). Complete data were available in 78 patients (22% women). Three patients declined testing because of distance from the medical center and discomfort from their condition. Sadly, all patients had expired by completion of the current study, although allowing assessment of the predictive power of several tests for death and NIV at intervals of up to 3 years and for relevant cutoff points to be obtained. In stepwise regression analysis for ventilation-free survival, all respiratory muscle strength assessments achieved statistical significance, whereas sex or age did not. The highest individual test scores were for Sn Pdi and twitch Pdi (Tw Pdi), using cervical magnetic stimulation. In multivariable analysis, age, Tw Pdi, Sn Pdi, Sn Pes, and MEP retained significance, whereas age and sex did not. SNIP also had strong predictive power.

A striking finding was that when receiver operator character analysis was used to determine the cutoff point for absolute survival and ventilation-free survival for each variable, VC was consistently >80% predicted at all cutoff points throughout the study beyond 3 months; even at 3 months before death, its cutoff point was still 78% predicted. Although SNIP, MEP, Tw Pdi, and Sn Pdi exhibited a general linear decline over the course of 5 years before death (in 25 patients for whom sequential measures were available), predicted VC values remained close to normal until about 1 year before death or the need for NIV, at which time it declined rapidly.

What is the explanation for the difference in decline between the VC and measures of respiratory muscle force generation? The