J Am Chem Soc. 2017 November 22; 139(46): 16666–16676. doi:10.1021/jacs.7b08089.

Department of Chemistry, and Center for Theoretical Biological Physics, Rice University, Houston, Texas 77005, United States

Using a predictive coarse-grained protein force field, we compute and compare the free energy landscapes and relative stabilities of amyloid- β protein (1-42) and amyloid- β protein (1-40) in their monomeric and oligomeric forms up to the octamer. At the same concentration, the aggregation free energy profile of $A\beta_{42}$ is more downhill, with a computed solubility that is about ten times smaller than that of $A\beta_{40}$. At a concentration of 40 μM , the clear free energy barrier between the prefibrillar tetramer form and the fibrillar pentamer in the $A\beta_{40}$ aggregation landscape disappears for $A\beta_{42}$, suggesting that the $A\beta_{42}$ tetramer has a more diverse structural range. To further compare the landscapes, we develop a cluster analysis based on the structural similarity between configurations and use it to construct an oligomerization map that captures the paths of easy interconversion between different but structurally similar states of oligomers for both species. A taxonomy of the oligomer species based on β -sheet stacking topologies is proposed. The comparison of the two oligomerization maps highlights several key differences in the landscapes that can be attributed to the two additional C-terminal residues that $A\beta_{40}$ lacks. In general, the two terminal residues strongly stabilize the oligomeric structures for $A\beta_{42}$ relative to $A\beta_{40}$, and greatly facilitate the conversion from prefibrillar trimers to fibrillar tetramers.

Figure 1: Kinetic model of A β 40 aggregation. The graph plots Force (F) against the number of monomers (n). A solid blue curve shows the force profile, with a peak at n=4 labeled "A β 42 fiber". A dashed green curve shows the force profile for "A β 40 backtracking", with a peak at n=4 labeled "Prefibrillar oligomers". The concentration is [A β] = 40 μ M.

During normal brain metabolism, the cleavage of the amyloid precursor protein generates several versions of the Amyloid- β peptide ($A\beta$).^{1,2} These rather insoluble pathological

products are major components of the senile plaques found in the brains of patients with Alzheimer's Disease (AD).^{3,4} Cleavage generally gives either a peptide of length 40 or one of length 42. The longer of the two predominant peptide species $A\beta_{42}$ has two additional hydrophobic residues (Ile41 and Ala42) at the C-terminus. $A\beta_{42}$ is the minor component in human cerebrospinal fluid and plasma, being found in approximately a 1:10 ratio to the $A\beta_{40}$ form.⁵ Nevertheless, $A\beta_{42}$ is found three times more often in the senile plaques characteristic of the Alzheimer's disease cortex according to studies that use end-specific monoclonal antibodies to differentiate the species.³ The shorter and more prevalent $A\beta_{40}$ form, on the other hand, seems to play a more important role in the later stages of plaque formation as evidenced by a strong correlation between detection of $A\beta_{40}$ and the maturity of the plaques.³ *In vitro*, both peptides aggregate into fibrils, but $A\beta_{42}$ aggregates more quickly at comparable concentrations.⁶ Systematic mutational studies reveal a strong correlation between the aggregation propensity and the hydrophobicity and the β -sheet propensities of the residues at the terminal position 41 and 42.⁷ In this paper, we use computation to compare the aggregation energy landscapes and structural ensembles for the two related proteins and explore the effects of these additional two residues on the early stages of aggregation.

In a micelle or in an apolar solvent environment, the monomers of both $A\beta_{40}$ and $A\beta_{42}$ are largely disordered but adopt α -helical structures in the central hydrophobic core region and in the C-terminal region.^{8,9} The NMR data in these environments reveal very little in the way of structural difference suggesting that the greater aggregation propensity of $A\beta_{42}$ is not primarily related to this monomeric helix-rich membrane-bound structure.⁸ In water at pH 7.3, at room temperature and in the concentration range between 5 and 10 mM, however, the $A\beta_{42}$ peptide appears to take on somewhat more β -sheet structure (79% β -sheet) compared with only 57% for $A\beta_{40}$. Under those more physiological conditions little α -helical structure appears to be present.¹⁰ We note however some recent studies using circular dichroism do report a dominant presence of coiled structure in the aqueous form with varying amount of α -helix and β -strand, depending on sample preparation.¹¹ Extensive All-atom simulations using both implicit and explicit solvent models have explored the monomeric configurations of $A\beta$ peptides. Much as in the laboratory, the results of these computational studies are highly divergent.^{12–16} The β content of the monomer is crucial for further aggregation but is sensitive to environmental conditions.

One explanation for the different aggregation propensities of the two peptides lies in the fibril structure. By monitoring hydrogen/deuterium exchange in the fibrils, Sanchez et al. measured the rate coefficient for a monomer to dissociate from its fibril. This rate is 60 times larger for $A\beta_{40}$ than it is for $A\beta_{42}$.¹⁷ The reported fibril structures for $A\beta_{40}$ and $A\beta_{42}$ are different. Different high resolution fibril structures of $A\beta_{40}$ obtained through several solid state NMR studies generally display the same U-shaped β -turn- β motif. The different polymorphs however do differ from each other mainly in the detailed way in which protofilaments stack; the turn region occurring between Asp23 and Gly29 is often stabilized by a salt bridge formed between Asp23 and Lys28.^{18,19} Determining the fibril structure of $A\beta_{42}$ has been more difficult owing to its much higher aggregation propensity in comparison with $A\beta_{40}$. In 2015 and 2016, three groups nevertheless successfully elucidated the $A\beta_{42}$ fibril structure arriving at consistent results reflecting most likely the

thermodynamically most stable form.^{20–22} This fibril structure can be recognized by the fibril-specific antibodies that bind to senile plaques in AD patients' brain, but is not recognized by other antibodies that do not bind to such plaques.²¹ The $A\beta_{42}$ fibril displays a triple parallel- β -sheet structure in an "S" shape but manifests several detailed differences from the above-mentioned " β arch"²³ structure for the $A\beta_{40}$ fibril. The two additional C-terminal residues that are not present in $A\beta_{40}$ seem to provide additional stabilization for the $A\beta_{42}$ fibril. In this fibre form the C-terminal carboxyl group of Ala42 forms a salt bridge with Lys28, and Ile41 is in close contact with Gly29.²² Without these interactions there would be space for water molecules to intercalate into the turn region and destabilize the fibril structure. This structural difference may explain why seeding with $A\beta_{42}$ fibril does not speed up the aggregation of $A\beta_{40}$ as much as seeding with $A\beta_{40}$ fibrils does.²⁰ Colvin et al. also suggest that the hydrophobic strip formed by residues V40-A42 that is located at the outer surface of the protofilament could enhance secondary nucleation, a possible mechanism for faster aggregation of $A\beta_{42}$.²² The newly discovered $A\beta_{42}$ fibril structure is clearly an important piece of the puzzle that motivates our present computational work which aims to explore the similarities and differences in aggregation mechanism for the two peptides.

It has become widely accepted that the soluble oligomeric forms of $A\beta$ can be potent neurotoxins and may be closely linked to the neuronal injury and death in AD patients.²⁴ Bitan et al. isolated low-molecular-weight $A\beta$ oligomers and used photo-induced cross-linking techniques to show that $A\beta_{40}$ and $A\beta_{42}$ oligomeric ensembles have different distributions of structures. Their studies indicated that $A\beta_{40}$ existed as monomers, dimers, trimers and tetramers in rapid equilibrium; while in contrast, $A\beta_{42}$ formed mainly pentamer and hexamer units which further assembled to protofibril-like structures.²⁵ Understanding the concentration dependence of these oligomer ensembles and their structural details, especially the effects of residues 41 and 42 on the stability of configurational ensembles calls for computational investigation. For all-atom simulations, however, simulating even just a dozen full-length $A\beta$ peptides in a box is a dauntingly computationally intense task because of the long time scales for association at laboratory concentrations.²⁶ This argues for using coarse-grained models, but while computationally efficient, many coarse-grained transferable force field models suffer from a lack of proven track record in successful structure prediction in folding and binding. This makes it hard to assess their predictive power for aggregation. Here, we employ an efficient force field, AWSEM, which in contrast to other coarse-grained models, has been shown to be effective at predicting folding structure²⁷ and thermodynamics.²⁸ Even with the coarse-grained model, we must use high concentrations to overcome the association time scale problem. Statistical mechanical analysis using physical cluster theory, however, allows us to extrapolate reliably simulation results to realistic laboratory concentrations. Using these tools, in this study we can compare the aggregation energy landscapes and aggregation mechanisms of $A\beta_{40}$ and $A\beta_{42}$ over a wide range of concentrations, side by side, illustrating their differences in solubility as well as thermodynamic stability at various oligomer sizes. Our previous work on $A\beta_{40}$ revealed an interconversion between prefibrillar and fibrillar forms where local stabilizing contacts in the prefibrillar form must be broken in order to form a new set of stabilizing contacts in the fibrillar form—the reaction bottleneck. By requiring the breakup of stable structures, this

backtracking slows primary nucleation significantly for $A\beta_{40}$. This “backtracking” event however is not prominent in the simulations for $A\beta_{42}$ for any oligomer sizes, perhaps providing an explanation for its outsize role in early pathology. The presence of the C-terminal residues 41I-42A in $A\beta_{42}$ greatly stabilizes a tetrameric fibrillar cluster which is the bottleneck for fibrillization in $A\beta_{40}$. We highlight differences in the oligomerization pathways of the two forms by constructing a detailed map with nodes that represent clusters of structures based on similarities in contact patterns and connect the nodes using the similarity between structures in different clusters to determine the strength of the connection. This local similarity based map of the landscape should resemble the map of the most kinetically relevant connections between structural assemblies since barriers generally grow with the amount of structural rearrangement²⁹ needed to interconvert the species.

Methods

Simulation model

In this study, we use AWSEM to compute and compare the aggregation landscapes of both $A\beta_{40}$ and $A\beta_{42}$. AWSEM (Associative memory, Water mediated, Structure and Energy Model) is a predictive and transferable coarse-grained protein folding force field. In the AWSEM coarse-grained scheme, each amino acid residue in a protein is described by three beads, which represent the C_{α} , C_{β} and carbonyl O atoms of the residue’s chemical structure with an ideal peptide bond geometry. In addition to maintaining protein backbone geometries, the energy function of AWSEM includes physically motivated potentials that embody the chemical and physical properties of residues in the context of secondary and tertiary interactions. These energy terms are transferable between different proteins. The tertiary interactions use parameters that are learned using the energy landscape theory.³⁰ Additionally, the formation of local sequence structure is biased through an associative fragment memory term, which selects input based on similarity in local sequence to peptide fragments where structures are available in the Protein Data Bank (PDB). The associative memory term encodes local structural tendencies.²⁷ Detailed descriptions of the setup in AWSEM, tertiary interaction, and fragment memory usage can be found in our previous work³¹ as well as the original AWSEM paper.²⁷

To study the concentration dependence of the aggregation process of $A\beta_{42}$, we first simulate 12 $A\beta_{42}$ monomers in a box of volume $272 \text{ \AA} \times 272 \text{ \AA} \times 272 \text{ \AA}$ with periodic boundary condition. This simulation thus has nominal monomer concentration of 1mM. Initially, the free monomers are distributed evenly in the box. The system is first equilibrated for a simulation time, equivalent of 1 μs in real time scale. Simulation snapshots are then collected for structural sampling and for calculating free energy profiles in this fixed ensemble. We use Reiss’s nucleation theory³² as a theoretical framework for this ensemble to compute the population of individual oligomers/clusters and convert their population into corresponding free energy. Suppose we have N protein monomers in a macroscopic box of volume V at temperature T . According to this theory (using a grand canonical approach), the probability of finding a single n -mer cluster and $N - n$ free monomers in the box is

$$P(n, N-n) = q_n(V, T) \cdot e^{n\mu_0/kT}, \quad (1)$$

where $q_n(V, T)$ is the partition function for the n -mer cluster and μ_0 is the chemical potential of the free monomers. See the Methods section in our previous work³¹ for details. The corresponding free energy is given by

$$F(n, N-n) = -k_B T \ln P(n, N-n) = F(n) - n\mu_0, \quad (2)$$

where $F(n) = -k_B T \ln q_n$ is the free energy of the cluster in the fixed ensemble. Eq. (1) and (2) are valid for a macroscopic system where the total number of free monomers is way larger than the size of the cluster, i.e., $N \gg n$ (μ_0 is therefore invariant). Owing to the theoretical treatment used, all simulation snapshots in principle should be properly filtered such that only some selected snapshots with the presence of exact n -mer cluster and $N-n$ free monomers in the box would be screened out for performing free energy calculation.

Because of the finite number of protein monomers in the simulation box, the effective monomer concentration changes as the cluster continues to grow. To account for this finite-size effect one adds a free energy correction as shown below:

$$F(n, N-n) = F^{sim}(n, N-n) + nk_B T \ln \frac{N-n}{N}, \quad (3)$$

where $F^{sim}(n, N-n)$ is the free energy obtained from simulation and the last term refers to the free energy correction term. In other words, the free energy of a macroscopic system is approximated using the free energy calculated from simulation with the finite-size correction. Once the finite-size issue has been taken into account, one can estimate the free energy $F(n, N-n)$ as a function of n at any particular initial concentration c_0 .

Since the unbound monomers are dilute, the free energy calculated at a particular c_0 can be extrapolated to another concentration of interest c_1

$$F(n, N-n, c_1) = F(n, N-n, c_0) - nk_B T \ln \frac{c_1}{c_0}. \quad (4)$$

All the simulations were carried out using the Nose-Hoover thermostat (for NVT ensemble) as implemented in the LAMMPS molecular dynamics simulation package.³³ To study the free energy profile along the fiber structural coordinate Q_{fibril} , we used umbrella sampling techniques as described in our previous work.³¹ The Q_{fibril} is a collective coordinate that calculates the fraction of contacts formed in an oligomer structure with respect to a reference fibrillar structure. In this study, a relaxed A β 42 structure is used as a reference. The reference A β 42 fibrillar structure we use was generated by relaxing a solid-state NMR

structure²² (PDB ID: 5KK3) with the AWSEM force field. We also studied a relaxed structure that was obtained using another recently solved NMR structure²¹ (PDB ID: 2NAO) as the starting point. The two initial structures used were both solved at neutral pH (pH 8 and 7.4, respectively) and are essentially identical in both the ordered region (15–42) and disordered N terminal tail (1–14). Notice also that the final relaxed $A\beta$ 42 fibrillar structure is consistent with the solid-state NMR structure of $A\beta$ 40¹⁸ used in our previous work.³¹ Recently after our paper was submitted, we learned of another recently solved cryo-EM structure that shows some polymorphic characteristics for the $A\beta$ 42 peptide in the fibril, e.g., the EM density map of the entire N terminus was now well resolved in contrast to the solid-state NMR structures we used as starting points.³⁴ We note that this cryo-EM structure was obtained under relatively acidic conditions. The effects of pH are quite interesting but in our view call for considerably more simulation work specifically focused on them.

Clustering and generating the oligomerization map

In order to quantify the differences and similarities of the structures of $A\beta$ 40 and $A\beta$ 42 oligomers, we have applied clustering analyses to subsets of structures from both the unbiased and the biased simulations. We use $1 - Q_{relative}$ as a distance measure between two snapshots, where $Q_{relative}$ is the fraction of the contacts that are the same for the two structures. The C-terminal residues 41I and 42A of $A\beta$ 42 are excluded when calculating $Q_{relative}$ so $A\beta$ 40 and $A\beta$ 42 structures can be compared to each other. To generate the set of structures for clustering, we extract all the trimer, tetramer and pentamer structures from the $A\beta$ 42 and $A\beta$ 40 unbiased simulations in which there are twelve $A\beta$ proteins. Then we randomly select 500 structures from the $A\beta$ 40 simulations and 500 structures from the $A\beta$ 42 simulations for the trimer, tetramer and pentamer. For the tetramer and the pentamer, we also select 250 structures from biased simulations of $A\beta$ 40 and 250 structures from biased simulations of $A\beta$ 42, so that the selected structures represent a more thorough sampling of the energy landscape. We select trimer, tetramer and pentamer structures for clustering analyses because this is the size range for the transition to the fiber for $A\beta$ 40 at the concentration of $40\mu M$ as shown in Fig. 2. This structural change is crucial for the understanding of the early stages of aggregation. We separate the selected structures into 5 non-overlapping groups: prefibrillar trimer (p-III), prefibrillar tetramer (p-IV), prefibrillar pentamer (p-V), fibrillar tetramer (f-IV) and fibrillar pentamer (f-V), regardless of whether the configuration represents an $A\beta$ 40 or $A\beta$ 42 structure. We then performed hierarchical clustering for each group individually and identify centroid structures of each cluster. To characterize the structural transitions between groups, we calculate the $Q_{relative}$ between all the cluster centroid structures for the following pairs of groups: p-III & p-IV, p-IV & p-V, p-III & f-IV, p-IV & f-V, f-IV & f-V. The fraction of contacts compared with the β -arched $A\beta$ 40 fibril structure, Q_{fibril} , for all the centroids is also calculated. This coordinate is also indicated on the oligomerization map. We want to point out several aspects of the calculation of the fraction of shared contacts between two structures. First, the last two residues of each chain of $A\beta$ 42 are always excluded from the calculation for consistency. Since the specific chain order for two structures of the same size (e.g., both are trimers) may affect the Q value, all permutations of the chain order for each structure are used for the calculation and the largest Q value of these permutations is assigned as the final result. For two structures of different size (e.g., a trimer and a tetramer), all possible subsets of chains of the larger

complex and their permutations are considered and the largest Q of the subset permutation is again chosen as the final result. Dealing with these combinatorial issues required deciding how many structures to use for the calculation, because the combinatorics of this Q calculation results in a rapid increase of the computational cost (with increasing cluster sizes).

The oligomerization map is essentially a graphical representation of the free energy landscape with nodes and edges. The nodes represent the clusters. We use the Q_{fibril} of the centroids from all the clusters as the x -coordinate of the map and the size of the oligomer as the y -coordinate for the center of each node representing its own cluster; the size of the disc representing the node is proportional to its weighted population calculated from the two-dimensional aggregation free energy at $40\mu M$ concentration for $A\beta_{40}$ and $A\beta_{42}$; the width of the edges is proportional to the $Q_{relative}$ between the two centroids. A large relative Q should indicate a facile transformation. We generated oligomerization maps both for $A\beta_{40}$ and $A\beta_{42}$ separately to visualize the difference between them.

Results

Comparing the aggregation free energy landscapes of $A\beta_{40}$ and $A\beta_{42}$

While secondary nucleation is important in many experiments³⁵ as well as in vivo, at some point, the formation of $A\beta$ amyloids must begin with the primary nucleation of fibers from peptide monomers. From a thermodynamic perspective, the progress of the primary nucleation of $A\beta$ can be described using a grand canonical free energy potential as a function of the size of the formed oligomer. In this grand canonical formulation, the thermodynamic potential of an $A\beta$ oligomer with a particular size is obtained by subtracting from the standard free energy of the n -mer, the summed chemical potential of the monomeric proteins which make it up (see Methods for details). In the grand canonical ensemble, the chemical potential is largely set by the concentration of free monomers. This effective thermodynamic potential for aggregation determines the extent to which $A\beta$ oligomerizes and whether an oligomer once formed will continue to grow by adding new monomer units from solution. Macroscopic fibrils will form when this thermodynamic potential decreases monotonically in a downhill manner after a critical nucleation size has been reached. The aggregation nucleus can be defined as that particular oligomer size at which the free energy peaks, assuming there are no additional barriers to structural rearrangement which, as we will see, is indeed an issue for amyloid β . The size of the nucleus depends on the protein concentration. Fig. 1 compares the aggregation free energy profiles for $A\beta_{40}$ and $A\beta_{42}$ at different monomeric protein concentrations. Each individual free energy curve is plotted as a function of the oligomer size. To obtain the aggregation free energy we start by first simulating twelve monomers in a simulation box of fixed size having periodic boundary condition, which yields a nominal initial protein concentration of 1mM . This free energy for a finite system having a fixed finite number of monomers is then corrected by using physical cluster theory to remove the finite-size effects so as to describe nucleation in a thermodynamically large system with a macroscopically defined monomer concentration. The resulting free energy (at 1mM) is then used to extrapolate to lower concentrations, again employing physical cluster theory. We show results for $40\mu M$, $4\mu M$,

0.4 μM , and 0.04 μM . The experimental value of the solubility for $A\beta 42$ is 0.04 μM ,³⁶ nearly the value at which we find that the corrected free energy curves flatten out for large oligomer size. In our previous paper,³¹ we used the same simulation protocol to obtain the aggregation free energy for $A\beta 40$ at 1 mM, and extrapolated the free energy curves to find the protein's solubility 0.4 μM ³⁷ which also compares favorably to experiment. We see the grand canonical free energy analysis for $A\beta 40$ and $A\beta 42$ at the macroscopic level successfully builds a thermodynamic correspondence between simulation and experiment.

The known fiber structures of both $A\beta 40$ and $A\beta 42$ differ considerably from the prefibrillar forms as witnessed by there being antibodies that bind specifically to the fiber and not to prefibrillar forms and vice versa. The structural difference between fiber and prefibrillar forms for $A\beta 40$ was confirmed in our earlier simulations of $A\beta 40$. As in that study in order to understand the mechanism of fibrillization, we must define a collective coordinate Q_{fibril} , which is the fraction of contacts formed within an oligomer structure with respect to a reference fibrillar structure. In this case we use a relaxed $A\beta 42$ structure for the hexamer as the reference state. Free energy profiles projected onto Q_{fibril} along with the oligomer size then monitor the overall progress of $A\beta 42$ aggregation using two reaction coordinates just as in our earlier $A\beta 40$ study. Fig. 2 compares the two-dimensional grand canonical free energy surfaces for $A\beta 40$ and $A\beta 42$ in each case using Q_{fibril} measured with respect to their respective reported fibril structure along with the oligomer size n as coordinates. The critical nucleus for $A\beta 42$ is smaller than it is for $A\beta 40$ at the same concentration of 40 μM at 300K. At this concentration for $A\beta 40$, there is a clear separate step for structural conversion when a monomer is added to the tetramer to form a pentamer. At this point the simulations show a free energy barrier that separates the prefibrillar and fibrillar basins. This reaction path feature, in contrast, is not prominent for $A\beta 42$. This difference in topography reflects the result that the individual oligomers for $A\beta 42$ with $n \geq 3$ display a shallower free energy basin along Q_{fibril} than does $A\beta 40$. This feature implies $A\beta 42$ has a higher kinetic propensity for fibrillization.

Constructing and comparing the oligomerization maps of $A\beta 40$ and $A\beta 42$ oligomers

To quantify and visualize the differences between the dominant oligomerization pathways for $A\beta 40$ and $A\beta 42$ species, we construct an oligomerization map by clustering the structures obtained from our simulations. This clustering uses a local measure of similarity based on the number of common contacts in any two structures. In random heteropolymers, the paths of escape from a trap can be characterized with a local reaction coordinate that is correlated to an overlap measure Q_{ij} , the fraction of common contacts.²⁹ Thus, we expect clustering based on Q_{ij} to give a first approximation to the kinetically significant basins of attraction on the rough landscape of a typical polymer. Using this notion, we carry out the clustering analysis for the trimers, tetramers and pentamers in order to cover thoroughly the prefibrillar oligomeric region of the aggregation landscape. We employ a scheme of clustering common to both the $A\beta 40$ and $A\beta 42$ species. To do this we clustered at the same time a large group of structures, 50% of the structures being selected from the $A\beta 40$ simulations and 50% being selected from the $A\beta 42$ simulations. By mixing the structures of the two species to set up the clustering one develops a common set of local basins of structures that can be used to visualize both the similarities of and the differences between

the landscapes for the two species. As shown in Fig. 3, after first identifying all the clusters using a mixture of $A\beta 40$ and $A\beta 42$ structures, we can then isolate the $A\beta 40$ structures and recalculate the population of each cluster without counting the $A\beta 42$ structures and vice versa. We then visualize the cluster populations and their stronger kinetic connections based on contact similarities. In the diagram the area of the discs in the figure scales with the equilibrium population of the clusters. The clusters of structures from the unbiased simulations are colored in blue. We have also included structures clustered from simulation runs that are biased along the Q_{fibril} reaction coordinate. These are indicated in light blue. Their populations reweighted to take into account the biasing are again indicated by the areas of the corresponding discs.

Comparing the map in Fig. 3 for $A\beta 40$ and the map in Fig. 4 for $A\beta 42$, we can see that the main differences in the oligomerization pathway for the two species can be traced to the two tetramer clusters A and B that are located in the region where $0.4 < Q_{fibril} < 0.6$. Cluster B is structurally quite similar to the pentamer fibril cluster C and can grow into it by the simple addition of a monomer. In contrast, some backtracking that would require breaking the hydrogen bonds of the β -hairpins is required for an oligomer in cluster A to grow into one in a fiber forming cluster C, hence there should only be a weak kinetic connection between these two clusters. In the case of $A\beta 40$, cluster A is well connected to prefibrillar trimers ($Q_{fibril} < 0.4$) whose kinetic linkage extends over prefibrillar tetramers and pentamers. These prefibrillar species exhibit a range of morphologies, as shown in Fig. 3 (see section below for taxonomy of these prefibrillar species). The structural transformations among these prefibrillar species represent a dynamical search in which individual $A\beta$ monomers explore their binding configurations on pre-existing molecular constructs having partial sheet structure. This search continues until the prefibrillar species undergo a proper structural rearrangement into the prefibrillar cluster A. As a result, backtracking is required for species in cluster A to turn into fibrillar species i.e., cluster B and finally goes through structures in cluster C. For $A\beta 42$ there are few connections from prefibrillar trimers to cluster A; however, these prefibrillar trimer clusters with cluster B remain connected, as shown in Fig. 4. These residual links to cluster B ensure that facile formation of fibril cluster B almost always occurs through the trimer species via direct monomer addition, not from structures in cluster A which must backtrack.

Cluster A is well-populated in the map of $A\beta 40$ (Fig. 3) while cluster B is dominant for $A\beta 42$, as shown in Fig. 4. This difference arises from the presence of C-terminal residues 41I-42A because these terminal residues stabilize the fibrillar oligomer structure. This is evident in Fig. 5 where for the $A\beta 40$ species which lacks the C-terminal residues, the contact map (shown on the left) indicates the β -arch structure becomes loosely packed. There are only scattered contacts in the map. We also see the arches break down and turn into β -hairpins at both ends. For $A\beta 42$ on the other hand, the β -arch structure is quite well formed along the fibril, the interchain contacts are densely formed and the C-terminal residues are deeply involved in the formation of β -hydrogen bond contacts. To further localize the regions where these intra- and inter- contacts are reorganized when converting into a fibrillar structure, we compute contact probability maps for the ensembles corresponding to clusters A and B (see Fig. S1 in supporting information). By subtracting the contact probability map of the cluster A ensemble from the map for the cluster B

ensemble, we obtain the difference map shown in the middle part of the figure. This difference map highlights the regions where changes occur. The locations where existing intra-monomer contacts must break are shown in blue while the newly formed inter-monomer contacts are shown in red. These regions agree with the loosely packed C-terminal regions found in the $A\beta$ 40 contact map (left). The difference map clearly demonstrates the backtracking in molecular detail where intra- and inter-monomer contacts are broken and formed, respectively, at both ends when converting from a prefibrillar to fibrillar structure.

Taxonomy of oligomer clusters

Fig. 6, Fig. 7, and Fig. 8 show structural ensembles for the clusters of the prefibrillar oligomers (trimer, tetramer, and pentamer, respectively) as obtained by our clustering analysis, using relative Q. These figures show all the clusters with more than one member. The centroid of each cluster is shown as a single dark blue Richardson trace while other members of the ensemble are shown in transparent gray to give an idea of the breadth of each ensemble. One also finds some largely unstructured clusters for the trimer with only single representative seen in each cluster. The oligomer structures exhibit a range of configurations with a mixture of secondary structure elements (α -helix, β -hairpin, and coil). The individual monomers sometimes align into a sheet structure but sometimes the secondary structure elements are randomly oriented into an overall disordered aggregate. The structure ensembles of prefibrillar trimer, tetramer, and pentamer share many common structural features. For each oligomer group, the most populated cluster always corresponds to a sheet-based structure in which the individual monomers by themselves are well formed into a β -hairpin structure and are then properly aligned and stacked on each other. The most populated clusters for the trimer, cluster 1 (31 %) and cluster 4 (26.7 %), both show a β -sheet structure formed by aligning two β -hairpins in the same plane but with the third β -hairpin loosely attached to the sheet structure (see Fig. 6). This additional β -hairpin in the context of the tetramer can either initiate the development of a second layer of the β -sheet shown in cluster 1 (25.3 %) or participate in elongating the pre-existing sheet structure shown in cluster 5 (33.9 %) (see Fig. 7). Similarly, for the pentamer, cluster 2 (29.6 %) exhibits a longer β -sheet structure formed by aligning four β -hairpins (see Fig. 8). Interestingly, owing to the chiral nature of β -sheets, the single sheet structure becomes twisted as the oligomer size grows. As a result, the cluster begins to resemble a cylindrin, as can be seen in the pentamer cluster 2 (see Fig. 8). In many cases, the sheet structure exhibits variations from perfection. For instance, a sheet structure is sometimes augmented by adding only a single β -strand which involves only partial sequence length of a single $A\beta$ monomers (see cluster 5 (2.3 %) and cluster 24 (16.3 %) in the trimer). Moreover, in some cases two β -strands from the same $A\beta$ monomer are shared by two different layers of β -sheets, respectively. This β -strand sharing behavior is mostly seen in larger oligomers, for example, cluster 3 (5.6 %) and cluster 4 (16.8 %) in the pentamer. This sharing phenomenon is not as common in the trimer clusters simply because there are fewer β -strands present. To clarify the structural heterogeneity outlined above, we have developed a notation specifically to describe the topologies of the sheet-based configurations. These classifications for the clusters of prefibrillar structures are shown in Fig. 6. Last but not least, we see that helical structure is not very prevalent but does occur in the smaller oligomers. In other words, there is a significant trend for α to β conversion as the oligomer size increases from trimer to

pentamer. Indeed, for the trimer there is a category of amorphous aggregates with such a broad range that each cluster is rarely populated being represented as singletons in our study. Most of the configurations of these amorphous aggregates show significant content of helical structures as shown in Fig. 9. As we mentioned earlier under appropriate solvent conditions, a single $A\beta$ monomer undergoes a dynamic conversion between α -helix and β -hairpin (see Fig. 10). This same structural conversion occurs then as oligomerization proceeds in aqueous solvent. Explicit all-atom simulation also supports this dynamic conversion of the monomer in the early stage of the oligomerization process.¹⁶ When the oligomer grows into a key prefibrillar tetramer species ($Q_{fibril} > 0.4$), the prefibrillar species may undergo a prefibrillar-to-fibrillar structural conversion, as summarized by the major oligomer clusters shown in Fig. 11. In this figure, the oligomer species from left to right correspond to the prefibrillar cluster A, fibrillar clusters B and C, respectively, as indicated in the maps shown in Fig. 3 and Fig. 4.

The taxonomy of oligomer clusters presented in the work should allow a comprehensive structural analysis of the oligomer landscape which may be important in computationally targeting specific oligomer structures which may be especially toxic in neurodegenerative diseases. The library of backbone structures can form the basis for a library of fully atomistic oligomer structures that could be used for drug design.

Discussion

Nucleation mechanism for fibrillization: the importance of backtracking

The comparison between the $A\beta_{40}$ and $A\beta_{42}$ free energy surfaces already highlights the importance of the two additional hydrophobic residues (Ile41 and Ala42) at the C-terminus of $A\beta_{42}$. The two residues should affect the kinetics of primary nucleation, as the nucleus size decreases from 4 (for $A\beta_{40}$) to 3 (for $A\beta_{42}$) at the concentration of 40 μM . The corresponding free energy profile for $A\beta_{42}$, exhibits a lower barrier of nucleation prior to forming fibrillar structures. This low nucleation barrier allows a more rapid fiber formation.

The oligomerization landscape maps for $A\beta_{40}$ and $A\beta_{42}$ can give us some ideas about which of the two species is better able to trigger a cascade of amyloid formation. From our calculations, the predicted solubility of $A\beta_{42}$ is one order of magnitude lower than that of $A\beta_{40}$. At the same time, the concentration of $A\beta_{42}$ measured in the spinal fluid of healthy controls is also about 10 times lower. Apparently, then, the two species are present at similar levels of supersaturation. From the perspective of the thermodynamics of primary nucleation alone, then, it would be unclear which species would first aggregate if the local concentrations of both species rise in parallel. On the other hand, the comparison of the two oligomerization landscape maps suggests there should be some significant kinetic differences arising from the mechanistic details of the primary nucleation process for the two species: $A\beta_{40}$ requires backtracking while $A\beta_{42}$ does not. Likewise, if oligomers are the pathological culprits rather than the fiber forms, backtracking may cause an accumulation of prefibrillar $A\beta_{40}$ in contrast to $A\beta_{42}$ which more readily goes on to the fiber form. We expect aggregation without backtracking to be more facile than when backtracking is required. Indeed, kinetic measurements of aggregation of $A\beta_{42}$ in general show a shorter kinetic lag time/half time than that do measurements on $A\beta_{40}$ at the same

initial monomer concentration.^{38,39} The two species also differ in their secondary nucleation behavior according to experiments.⁴⁰ We remark that some of the mis-stacked prefibrillar oligomer clusters resemble the structures we might imagine to occur in secondary nucleation. Further computational investigation will be needed however to elucidate the mechanism of secondary nucleation in molecular detail.

Another interesting issue is whether the aggregates of one form of $A\beta$ can act as seeds for the other. There have been mixed conclusions from experimental studies. If cross seeding occurs, then the initially formed $A\beta_{42}$ aggregate could further trigger the full amyloid cascade of the $A\beta_{40}$ via either further growth of one species or seeding for the other species.

Towards building a kinetic model for the amyloid funnel

We constructed the oligomerization map using the structural similarity between clusters as an indication for how the clusters would be dynamically connected. To a first approximation, the clusters should reflect the kinetically significant basins of attraction on the oligomerization landscape and their high contact overlap connections should highlight their kinetically dominant paths of interconnections. Noting that the thicker edges in the maps refer to kinetically more relevant connections between the two clusters, we see that $A\beta_{42}$ should have a more facile path to fibrillation than does $A\beta_{40}$ as we can clearly see a thicker edge between fibrillar cluster B and cluster C from the $A\beta_{42}$ oligomerization map. This rapid fibrillation pathway implies that the addition of single $A\beta_{42}$ monomers is in fact a kinetically favorable process that can go on with little rearrangement. Once nuclei are formed in a supersaturated system, a cascade of monomer additions becomes thermodynamically favorable, eventually yielding a macroscopic fiber. The oligomerization map thus offers additional structural insight into the “amyloid funnel” picture that we discussed in our previous paper.³¹ The amyloid funnel of $A\beta_{42}$ becomes more downhill due to the two additional C-terminal residues (41I and 42A). According to our aggregation landscape analysis, these two residues play an essential role in both thermodynamics and kinetics of aggregation. We note that these aggregation landscapes indicate that the early stages of aggregation are kinetically much more complex than has generally been considered in the kinetic descriptions used for quantitatively fitting experimental data for the different amyloid proteins that usually assume simple mass action kinetics for monomer addition at the primary stage.^{41–43} More appropriate quantitative kinetic analyses will require turning the oligomerization maps developed in the present work based on structural similarities into kinetic Markov state models having quantitatively correct rate coefficients.^{44,45} A deeper comparison of the structural differences between the clustered ensembles elucidated here should yield approximate reaction coordinates which can then be used to calculate intercluster conversion rate coefficients, a task we leave for the future.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by Grant R01 GM44557 from the National Institute of General Medical Sciences. Additional support was also provided by the D.R. Bullard-Welch Chair at Rice University, Grant C-0016.

References

1. Takami M, Nagashima Y, Sano Y, Ishihara S, Morishima-Kawashima M, Funamoto S, Ihara Y. *J Neurosci.* 2009; 29:13042–13052. [PubMed: 19828817]
2. Kukar TL, et al. *J Biol Chem.* 2011; 286:39804–39812. [PubMed: 21868378]
3. Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. *Neuron.* 1994; 13:45–53. [PubMed: 8043280]
4. McLaurin J, Yang DS, Yip C, Fraser P. *J Struct Biol.* 2000; 130:259–270. [PubMed: 10940230]
5. SAG, Ho L, Eckman C, Long K, Sovos L Jr, Younkin L, Suzuki N, Younkin S. *J Biol Chem.* 1995; 270:7013–7016. [PubMed: 7706234]
6. Jarrett JT, Lansbury PT. *Cell.* 1993; 73:1055–1058. [PubMed: 8513491]
7. Kim W, Hecht MH. *J Biol Chem.* 2005; 280:35069–35076. [PubMed: 16079141]
8. Shao H, Jao S-c, Ma K, Zagorski MG. *J Mol Biol.* 1999; 285:755–773. [PubMed: 9878442]
9. Crescenzi O, Tomaselli S, Guerrini R, Salvadori S, D'Ursi AM, Temussi PA, Picone D. *Eur J Biochem.* 2002; 269:5642–5648. [PubMed: 12423364]
10. Barrow CJ, Yasuda A, Kenny PT, Zagorski MG. *J Mol Biol.* 1992; 225:1075–1093. [PubMed: 1613791]
11. Nasica-Labouze J, et al. *Chem Rev.* 2015; 115:3518–3563. [PubMed: 25789869]
12. Sgourakis NG, Yan Y, McCallum SA, Wang C, Garcia AE. *J Mol Biol.* 2007; 368:1448–1457. [PubMed: 17397862]
13. Côté S, Derreumaux P, Mousseau N. *J Chem Theory Comput.* 2011; 7:2584–2592. [PubMed: 26606631]
14. Ball KA, Phillips AH, Wemmer DE, Head-Gordon T. *Biophys J.* 2013; 104:2714–2724. [PubMed: 23790380]
15. Zhu M, De Simone A, Schenk D, Toth G, Dobson CM, Vendruscolo M. *J Chem Phys.* 2013; 139:035101. [PubMed: 23883055]
16. Tarus B, Straub JE, Thirumalai D. *J Am Chem Soc.* 2006; 128:16159–16168. [PubMed: 17165769]
17. Sánchez L, Madurga S, Pukala T, Vilaseca M, López-Iglesias C, Robinson CV, Giralt E, Carulla N. *J Am Chem Soc.* 2011; 133:6505–6508. [PubMed: 21486030]
18. Petkova AT, Yau WM, Tycko R. *Biochemistry.* 2006; 45:498–512. [PubMed: 16401079]
19. Bertini I, Gonnelli L, Luchinat C, Mao J, Nesi A. *J Am Chem Soc.* 2011; 133:16013–16022. [PubMed: 21882806]
20. Xiao Y, Ma B, McElheny D, Parthasarathy S, Long F, Hoshi M, Nussinov R, Ishii Y. *Nat Struct Mol Biol.* 2015; 22:499–505. [PubMed: 25938662]
21. Wälti MA, Ravotti F, Arai H, Glabe CG, Wall JS, Böckmann A, Güntert P, Meier BH, Riek R. *Proc Natl Acad Sci USA.* 2016; 201600749.
22. Colvin MT, Silvers R, Ni QZ, Can TV, Sergeyev I, Rosay M, Donovan KJ, Michael B, Wall J, Linse S, Griffin RG. *J Am Chem Soc.* 2016; 138:9663–9674.
23. Kajava AV, Baxa U, Steven AC. *FASEB J.* 2010; 24:1311–1319. [PubMed: 20032312]
24. Haass C, Selkoe DJ. *Nat Rev Mol Cell Biol.* 2007; 8:101–112. [PubMed: 17245412]
25. Bitan G, Kirkitadze MD, Lomakin A, Vollers SS, Benedek GB, Teplow DB. *Proc Natl Acad Sci USA.* 2003; 100:330–335. [PubMed: 12506200]
26. Barz B, Olubiyi OO, Strodel B. *Chem Commun.* 2014; 50:5373–5375.
27. Davtyan A, Schafer NP, Zheng W, Clementi C, Wolynes PG, Papoian GA. *J Phys Chem B.* 2012; 116:8494–8503. [PubMed: 22545654]
28. Sirovets BJ, Schafer NP, Wolynes PG. *J Phys Chem B.* 2015; 119:11416–11427. [PubMed: 26102155]

29. Plotkin SS, Wang J, Wolynes PG. *J Chem Phys.* 1997; 106:2932–2948.
30. Schafer NP, Kim BL, Zheng W, Wolynes PG. *Isr J Chem.* 2014; 54:1311–1337. [PubMed: 25308991]
31. Zheng W, Tsai MY, Chen M, Wolynes PG. *Proc Natl Acad Sci USA.* 2016; 113:11835–11840. [PubMed: 27698130]
32. Reiss H, Bowles RK. *J Chem Phys.* 1999; 111:7501–7504.
33. Plimpton S. *J Comput Phys.* 1995; 117:1–19.
34. Gremer L, Schölzel D, Schenk C, Reinartz E, Labahn J, Ravelli RBG, Tusche M, Lopez-Iglesias C, Hoyer W, Heise H, Willbold D, Schröder GF. *Science.* 2017; 358:116–119. [PubMed: 28882996]
35. Cohen SIA, Linse S, Luheshi LM, Hellstrand E, White DA, Rajah L, Otzen DE, Vendruscolo M, Dobson CM, Knowles TPJ. *Proc Natl Acad Sci USA.* 2013; 110:9758–9763. [PubMed: 23703910]
36. Tycko R. *Neuron.* 2015; 86:632–645. [PubMed: 25950632]
37. Qiang W, Kelley K, Tycko R. *J Am Chem Soc.* 2013; 135:6860–6871. [PubMed: 23627695]
38. Vandersteen A, Hubin E, Sarroukh R, De Baets G, Schymkowitz J, Rousseau F, Subramaniam V, Raussens V, Wenschuh H, Wildemann D, Broersen K. *FEBS Lett.* 2012; 586:4088–4093. [PubMed: 23103738]
39. Cukalevski R, Yang X, Meisl G, Weininger U, Bernfur K, Frohm B, Knowles TPJ, Linse S. *Chem Sci.* 2015; 6:4215–4233. [PubMed: 29218188]
40. Meisl G, Yang X, Hellstrand E, Frohm B, Kirkegaard JB, Cohen SIA, Dobson CM, Linse S, Knowles TPJ. *Proc Natl Acad Sci USA.* 2014; 111:9384–9389. [PubMed: 24938782]
41. Xue WF, Homans SW, Radford SE. *Proc Natl Acad Sci USA.* 2008; 105:8926–8931. [PubMed: 18579777]
42. Schreck JS, Yuan JM. *J Phys Chem B.* 2013; 117:6574–6583. [PubMed: 23638987]
43. Tsai MY, Yuan JM, Lin SH. *J Chin Chem Soc.* 2015; 62:21–25.
44. Schafer NP, Hoffman RMB, Burger A, Craig PO, Komives EA, Wolynes PG. *PLOS One.* 2012; 7:1–8.
45. Noe F, Clementi C. *Curr Opin Struct Biol.* 2017; 43:141–147. [PubMed: 28327454]

Figure 1.

We plot the appropriate thermodynamic potential $F - n\mu$ in units of kcal/mol for an open grand canonical system. The aggregation free energy for $A\beta_{42}$ is first obtained by simulating twelve monomers in a fixed size box with 1mM concentration at 300K. The solid line with spheres represents the corrected free energy at the simulated concentration of 1mM after removing the finite-size effect and accounting for the poor sampling of fibril structures in the raw simulation data which need to be supplemented by umbrella sampling as described previously for $A\beta_{40}$.³¹ The free energy for $A\beta_{40}$ at the same condition is in the dashed line with open spheres. n is the size of the oligomer. The extrapolated free energy profiles are also shown at 40 μ M (square symbol), 4 μ M (asterisk), and 0.4 μ M (diamond). We also show the profile at the predicted solubility 0.04 μ M for $A\beta_{42}$ (triangles) and 0.4 μ M for $A\beta_{40}$ (dashed line with diamonds).

Figure 2.

The grand canonical free energy surface in units of kcal/mol at the concentration of $40\mu M$ at 300K is plotted using an added dimension of Q_{fibril} that reveals more structural details of the oligomer. Q_{fibril} is the fraction of contacts formed in a particular structure that match with those formed in the fibrillar counterpart which contains only parallel hydrogen bonds. The closer the value of Q_{fibril} of a structure is to 1, the more similar the structure is to a fibril form. For $A\beta_{40}$, there is a clear barrier between the tetramer and the pentamer, suggesting monomer, dimer, trimer and tetramer can be kinetically stable. On the other hand, for $A\beta_{42}$, the barrier is not apparent, and the trimer and tetramer appear to be quite polymorphic with different Q_{fibril} , once it reaches the pentamer and the hexamer, the free energy is downhill. This comparison is consistent with Bitan et al.'s cross-linking study on $A\beta$ oligomers.²⁵

Figure 3.

The oligomerization pathway map for $A\beta_{40}$ at the concentration of $40\ \mu M$. The nodes in the graph represent the clusters of oligomeric structures and two nodes are connected by an edge if two clusters are similar to each other. The area of each circle is proportional to the population of the corresponding cluster and the thickness of the line scales with the magnitude of $Q_{relative}$ between the centroid structures of two connected clusters (we use a cutoff value of 0.2 to filter insignificant connections). The navy blue points are formed using the unbiased simulations while the light blue points come from the biased simulations. Together they provide a rather complete coverage of the oligomerization map. The y-axis indicates the size of the oligomer, and the x-axis is the Q_{fibril} for the centroid structure of each cluster. Some of the centroid structures are illustrated using a cartoon representation showing the directions of alignment of the β -strands in the map: the N-terminal is colored in deep blue and the C-terminal is colored green or light blue depending on whether it is formed in a β -hairpin structure or in a β -arch structure, respectively. It's reassuring to see the topological similarities in the structural representations between connected nodes.

Figure 4.

The oligomerization pathway map for $A\beta_{42}$ at the concentration of $40\ \mu M$. The red circles represent clusters formed using the unbiased simulations and the pink circles indicate clusters formed for the biased simulations. The clusters B and C are much more populated in this map than they were in the map of $A\beta_{40}$ but cluster A is less well-populated. The more populated fibrillar clusters and thick edges between the tetramer and pentamer suggest a more facile fibrillization pathway for $A\beta_{42}$ species that does not require backtracking.

Figure 5.

(left and right) The contact maps of the two tetramer clusters with $Q_{fibril} > 0.4$. The contact cutoff is set at 6.5 between C_{β} atoms. The left contact map is calculated from the centroid of the cluster A that is highly populated in $A\beta_{40}$ species, while the right contact map represents the cluster B that is highly populated in $A\beta_{42}$ species. The presence of the two C-terminal residues (41I-42A) increases the stability of the fibril structure significantly compared with other oligomeric tetramer structures for $A\beta_{42}$. The representative structures of the clusters along with their cluster ensemble shown as transparent shadows are shown below. The difference map is created by subtracting the contact probability map of cluster A from the map of cluster B (see Fig. S1 in supporting information). The color indicates formation of new contacts formed (red) or breakage of existing contacts (blue). The dotted red and blue circles highlight the regions of the intermolecular contacts that are formed and the intramolecular contacts that are broken, respectively, in the process of converting to a fibrillar structure.

Trimer

Figure 6.

Structural ensemble for the clusters of the prefibrillar trimer are shown. These clusters all have more than one member. The centroid structure of each cluster is shown as a dark blue trace along with other members of the ensemble shown in transparent gray. The number in the parentheses represents the fractional population of the cluster in the prefibrillar oligomer. The notation in the square brackets is used to describe the topologies of the sheet-based configurations. Each digital number specifies a single layer of β -structure. The β -structure here refers to three different topologies: 1. single hairpin (use 1). 2. partial hairpin (use 1'), and 3. sheet structure (use the number of monomers >1). Depending on the size of the resulting sheet, the sheet structure consists of a different number of β -strands. Since each β -hairpin contributes two β -strands, the sheet structures formed by three, four, five, and six β -strands are represented by 1.5, 2, 2.5, and 3, respectively. Different layers of structural motifs are assigned another digital number, separated using a symbol "+". In some cases, we use subscripts to show special variations for sheet topologies. For example, 1_(0.5+0.5) refers to a sheet structure formed by two β -strands that come from different $A\beta$ monomers; the former 0.5 is underscored in order to distinguish it from one having another β -strand from a different $A\beta$ monomer. A similar notation also applies to compound cases, where the two β -strands from a single β -hairpin contribute to two different layers of sheet structures, respectively, as exemplified by 1.5_(0.5) + 1.5_(0.5); the former 0.5 and the latter 0.5 are both underscored in order to show that these two β -strands in fact do come from the same $A\beta$

monomer. Helical structure is denoted by “h”. Some $A\beta$ monomers are highlighted in cyan. Please see the main text for the explanation of this feature. Note that configurations belonging to the amorphous aggregates are shown in Fig. 9.

Tetramer

Figure 7.

Structural ensembles for the clusters of prefibrillar tetramers are shown. Cluster 1 shows double layers of the β -sheet structures (the second layer colored in cyan) while cluster 5 exhibits an elongation for monomer addition (in cyan). See Fig. 6 for a more complete definition of the notation scheme.

Pentamer

Figure 8.

Structural ensembles for the clusters of the prefibrillar pentamer are shown. Cluster 3 and 4 exhibit two β -strands from the same $A\beta$ monomers (in cyan) are shared by two different layers of β -sheets. Yellow moieties are shown as a separated layer of the sheet in the cluster in order to reduce visual confusion. See Fig. 6 for a more complete definition for the notation scheme.

Trimer (amorphous, 1.8%)

Figure 9.

Structural ensembles for the clusters of the amorphous aggregates of the prefibrillar trimer are shown. These clusters are rarely populated (each contributes only 0.09%) being represented as singletons in our clustering analysis. Most of the configurations of these amorphous aggregates show significant content of helical structures. The overall fractional population of these amorphous aggregates contributes a total of 1.8% of all the clusters of the prefibrillar trimer.

Figure 10.

A single $A\beta$ monomer in solution undergoes a dynamic conversion between α -helix and β -hairpin. Figure reproduced from our previous paper.³¹

Figure 11.

Structural ensembles for the major clusters that participate in prefibrillar-to-fibrillar structural conversion are shown. From left to right, the clusters correspond to the prefibrillar cluster A, fibrillar cluster B, and fibrillar cluster C, respectively, as illustrated in Fig. 3 and Fig. 4. See Fig. 6 for a more complete definition of the notation scheme.