ART. VI.—*On the Impregnation of the Ovum of the Amphibia.*
Section First. By GEORGE NEWPORT, F.R.S., &c. (From the Philosophical Transactions of the Royal Society for 1851. Part I, p. 169.) Received 20th June.—Read 20th June 1850.

THE Amphibia, of all the vertebrated animals, afford to us the readiest means of investigating the difficult subject of Impregnation by actual experiment; and it is only, perhaps, by combining experiment with careful observations on the physical conditions that affect the development of the germ, and comparing these with the facts of the natural history and instincts of the species, that we may hope, ultimately, to obtain some further insight into this one of Nature's most hidden secrets.

I shall endeavour, therefore, in this communication, to show the condition of the ovum in the Amphibia through its earliest changes, and also before and immediately after impregnation, and to detail experiments made with a view to learn by what means its fecundation is effected;—and in a future communication I propose to trace the development of the embryo from the time of fecundation to that of its liberation from the ovum, in the two chief divisions of the class,—the tailless and the tailed Amphibia. The subjects thus naturally form two series—Impregnation and Development.

IMPREGNATION OF THE OVUM.

The history of what we can now prove to be the agent of impregnation, the spermatozoon, deserves to be especially noticed. Although great attention has been paid by physiologists during the last thirty years to almost every point of inquiry connected with the production and physical composition of the seminal fluid of animals, and its relation to the fecundation of the ovum, we have remained to the present time without any acknowledged proof either of the part which the different constituents of this fluid take in impregnation, or of the mode in which it effects impregnation. This perhaps is little to be wondered at when we remember how many years elapsed before the great discovery of Ham and Leewenhoek of the existence of moving bodies in the fluid, as part of its normal composition, was admitted. It is now one hundred and eighty-three years since Leewenhoek communicated the important discovery of these bodies to the Royal Society,* and regarded them as in some way essential to the fecundation of the ova; although his conjecture that they be-

* Philosophical Transactions, 1667, vol. xii. p. 1040.

came the future embryos was erroneous. Eighty-five years afterwards, Dr Parsons, Foreign Secretary of the Royal Society, believed it to be "extreme nonsense to imagine that the insignificant animals called *spermatic animals* can contribute anything towards propagation," &c.,* and it was not until the publication of the observations of Ledermüller,† a few years after that, that the production of spermatozoa, as part of the fluid, began to be admitted. Dr Hill, in the notes to his English translation of Swammerdam's *Biblia Naturæ*,‡ two years later, mentioned them as abundant in the frog at the season of pairing, but that it was then the fashion to doubt even their existence. Yet Needham, ten years after this, while acknowledging that these bodies are found in the fluid of all animals, adopting the views of Buffon and Daubenton, stated that they do not exist until after the fluid is removed from the vessels, and decomposition has commenced.§ And later still, even in our own time, their existence has been denied in the most positive manner by Sir Everard Home.|| Spallanzani, however, was so well acquainted with them, as found in the frog and toad, that he has recorded his great surprise at not observing them in the latter on two occasions.¶ Bonnet** and Gleichen,†† also, well knew them to abound in the males of animals of distinct species at the season of impregnation, but discovered that they are usually absent in hybrids, a fact that has since been confirmed by Prevost and Dumas.‡‡ These two observers, regarding the spermatic bodies, with Leeuwenhoek, as essential elements of the semen, believed that they actually penetrate bodily into the ovum, and become by metamorphosis part of the future embryo. Still more recently it has been stated by Dr Barry§§ to this Society that he has actually seen the spermatozoon within the ovum, a statement which my own observations do not enable me to confirm.

Before any satisfactory conclusion could be arrived at respecting the importance of the spermatic bodies in impregnation,

* Philosophical Observations on the Analogy between the Propagation of Animals and that of Vegetables. 8vo. 1752 (note), p. 44.

† Physikalische Beobachtungen der Samenthierchen. Nuremb. 1756. And also, "Beyträge zu denen Beobachtungen deerer Saamenthiergen und Kleiste Aale gehörig. 12mo. Frankfurt und Leipzig, 1759."

‡ Book of Nature, folio, part 2 (note), p. 105, 1758.

§ Notes des Nouvelles Recherches sur les Découvertes Microscopiques de l'Abbé SPALLANZANI par M. NEEDHAM. Lond. 1769, tom. i. p. 196.

|| Lectures on Comparative Anatomy, 4to, vol. v. pp. 332 and 337. 1828.

¶ Dissertations relative to the Natural History of Animals and Vegetables. Lond. 1789, p. 151.

** "Contemplations de la Nature;" and "Œuvres d'Histoire Naturelle," 4to, tom. iii. p. 454, &c. 1779.

†† Abhandlung über der Samen-und Infusions thierchen. Nuremb. 1788.

‡‡ Annales des Sciences Naturelles (Prem. Série), tom. i. p. 182, 1824.

§§ Proceedings of the Royal Society, vol. iv. p. 432. Phil. Trans. part 1, 1843, p. 33.

it was necessary to ascertain their nature, to trace their mode of development and production, to establish the periods of their occurrence in different classes of animals, and to learn something of their chemical composition, as well also as of the fluid portion of the semen in which they move. Most of these inquiries have been well followed out by Wagner, Siebold, Muller, and more especially Klliker, and more recently by Wagner and Leuckardt, from whose labours we have now some positive information which enables us to deduce a fair conclusion respecting their function, although a direct proof of its correctness is still to be supplied. Most observers now believe with Klliker, that the spermatozoa (still so called) are not independent living organisms, but are merely elementary constituent parts of the male body, an opinion in which my own investigations lead me fully to coincide. This opinion, indeed, is not entirely new, as a like view was held by some observers at the beginning of the last century, when it was still questioned whether the spermatozoa are normal constituents of the semen. Dr Drake,* in his "New System of Anatomy," while acknowledging that he had seen the seminal animalcules, and combating on the one hand the theory of Leewenhoek respecting them, and on the other the view that had previously been held with regard to the ovum, doubted their separate organization, and suggested that they "may be nothing more than some large particles of mixed fluid, whose motions and different figure the microscope discovers to our eyes," &c. G. TREVIRANUS† more recently held a similar opinion, that they are not independent animals, but are analogous in their structure and properties to particles in the pollen of plants, and that their motion is of the kind discovered by ROBERT BROWN in vegetables. KLLIKER,‡ however, first distinctly referred them to a class of known organic constituents of the living body, the vibratile cilia, a view which had previously been discussed and inclined to by MULLER.§

But however much our knowledge has become settled in regard to the nature of the spermatc bodies themselves, and their mode of development, their relation to the fluid portion of the semen in which they are contained is still a matter of doubt. H. Goodsir|| regards certain albuminous flakes in the fluid portions of the semen of Crustacea as the debris of dissolved cells,

* New System of Anatomy, by JAMES DRAKE, M.D., F.R.S., vol. i. p. 352, 1707.

† TIEDEMANN, Zeitschrift, vol. v. part 2, 1835.

‡ Beitrge zur Kenntniss der Geschlechtsverhltnisse und der Samen flussigkeit wirbelloser Thiere. Berlin, 1841.

§ Elements of Physiology (Eng. ed.) part 6, 1841, p. 1478.

|| Anatomical and Pathological Researches, 1844, p. 40.

and as the source of nourishment and development of the spermatozoa; while a more recent observer, Dr Kirkes,* regards the spermatozoa as the elaborators of the fluid, and the conveyers of it to the ovum at the time of impregnation. This latter supposition was originally advocated by Wagner, Valentine and Bischoff. But two of these observers have recently changed their views,† and now regard the fluid portion as only of secondary importance in impregnation, and the spermatogenic bodies as of essential. This view, as Wagner states,‡ is founded chiefly on the fact, that in some of the invertebrata the whole mass of the semen appears to be constituted almost or entirely of spermatozoa, while scarcely any *liquor seminis* can be detected;—and further, on the great improbability, perhaps impossibility, of the liquor seminis of those animals which expel their ova into water before impregnation being brought into contact with the ovum. But the same author justly remarks, that “even up to the present day this hypothesis of the influence of the liquor seminis has not met with any direct refutation.” To this I may add, that however strong the presumption may be in favour of the agency of the spermatozoa in those instances in which the liquor seminis has not been observed it affords no sufficient reason for disbelieving that the spermatozoa are not resolved into fluid at the moment of fecundation; or that in those animals in which the liquor seminis occurs in abundance it is not that which impregnates the ovum.

The question then, so far as *proof* is concerned, both of the direct agency of the spermatozoa, and of the non-efficiency of the liquor seminis in impregnation, remains open, as well also as that which involves the knowledge as to how impregnation is effected.

It is to these questions that this communication which I have now the honour of laying before the Royal Society is chiefly directed. I propose *first* to show the time and mode of disappearance of the germinal vesicle, and the condition of the ovum in the Frog and Newt, immediately before and after impregnation, and to endeavour to supply proof from actual experiments that the spermatozoa alone, in all cases of communion of the sexes, are the sole agents in impregnating the ovum; and further, that impregnation *cannot* be effected by the *liquor seminis*; and next to examine in what way the agency of the spermatozoa is influenced, impeded, or exerted.

* Handbook of Physiology, 1848, p. 610.

† BISCHOFF in MULLER's Archiv, 1847. WAGNER in Article “Semen,” Cyclopædia of Anatomy and Physiology, part xxxvi., January 1849.

‡ Loc. cit. p. 507.

1. CHANGES IN THE OVUM WITHIN THE BODY.

The ovum of the Amphibia has so frequently been the subject of examination by the best observers that a further detailed account of its development may at first appear to be useless, after what we already know of its changes through the labours of SWAMMERDAM, LEEWENHOEK, ROESEL, SPALLANZANI, PREVOST and DUMAS, RUSCONI, BAER, REICHERT, VOGT, BELL and others. But apart from the fact already mentioned, that the ovum of the Amphibia affords us the best means of actual experiment on impregnation, there are questions which relate to its earlier conditions on which the observers named are not agreed, but which are of importance with regard to the physiology of reproduction in the whole of the vertebrata.

I shall state, therefore, what I have myself observed with regard to these questions from the time when the ovarian ovum is approaching to maturity to that of its expulsion from the body, before entering on the subject of its impregnation.

As our means of comparing and testing the accuracy of all observations in natural history, and of experimental results in physiology, depend mainly on the correct identification of the objects examined, I may here state at once that the objects of the following details have been the Frog, *Rana temporaria*, and the Toad, *Bufo vulgaris*, among the ANOURA; and the *Triton palustris*, *Lissotriton punctatus* and *L. palmipes*, among the Water Newts, the URODELA. Neither SPALLANZANI, PREVOST, and DUMAS, nor RUSCONI, to whose observations I shall have frequent occasion to refer, mention the species they have examined. SPALLANZANI has given only the popular names of his animals, but RUSCONI has given a figure of his, which appears to have been *Rana esculenta*, and as the description he has given of the ovum of this species agrees with the description given by PREVOST and DUMAS, it is probable that the species which they employed was the same.

The Ovarium Ovum.—When the frog, *Rana temporaria*, is examined in the autumn, after it has ceased to feed, and is preparing to retire to its winter quarters, the ova within it have already attained to more than two-thirds of their ultimate dimensions, before leaving the ovaries, and have begun to distend the abdomen. They remain in this state through the greater part of the winter, while the frog is hybernating, as Professor BELL remarks,* in the mud at the bottom of ponds and stagnant waters. But as the spring approaches, and the animal is aroused from its lethargy by increased temperature, the ova then rapidly acquire their full development. A female frog taken at the

* History of British Reptiles, 1832, p. 89.

end of September, with her body enlarged with ova, was confined in water in a cold room, undisturbed through the winter excepting only at intervals of examination. The weather being mild, and the temperature of the room during October and November, being sometimes at or but little below 50° FAHR., the frog remained active, and came frequently to the surface to respire. In December the temperature sunk to below 40° FAHR., when the frog became lethargic, and scarcely changed its place at the bottom of the water during nearly a fortnight, while the temperature was almost stationary, and ranged only from 35° FAHR. to 37° FAHR. On the 10th of January, when it had again risen to 40° FAHR., and that of the water to 37.5° FAHR., the animal was still submerged and motionless; but on sudden exposure to the light of a candle it crept languidly, and almost imperceptibly, to a distance of about two inches, and again became quiet. On the 26th of January the temperature of the room had risen gradually and continuously to 52° FAHR., and that of the water to 48° FAHR., but the frog remained submerged and perfectly quiet with its eyes partially closed. Although it was not cognizant of any object by sight, the irritability of its body was now much increased, as it moved instantly when touched ever so lightly, but quickly relapsed into its previous state of rest. On the morning of the 1st of February, the temperature being then 48° FAHR., it was still submerged and motionless, but in the evening, when the warmth was increased to 52° FAHR., I found it with its nostrils only above the surface of the water, and evidently beginning to respire freely, but its eyes were completely closed. When the light of a candle was suddenly cast upon it, the eyes were slowly opened, but its body remained immovable. On the following day, February 2, it was evident that its hybernation had been brought to a close, as it was then active, with its head out of the water, and its eyes widely open, and perfectly alive to external objects, as when the hand was slowly approached it withdrew beneath the water. The temperature at this time was 53° FAHR. During this period of hybernation the frog became slightly emaciated, and its abdomen, instead of being more enlarged, was somewhat diminished in bulk,—a good proof that during the inactivity of the respiratory and circulatory functions, the secretory also are lessened, and the development of the ova is arrested. But although no food was supplied to the frog at this time, and none probably is taken by the creature in its natural haunts, as at the time it comes forth but few of the objects on which it feeds are abroad, its body soon became enlarged, showing that the ova were then rapidly attaining their full development. Between the 2nd and 22nd of February the temperature of the

room was occasionally as low as 42° FAHR., yet the creature remained active beneath the water, without relapsing into its previous state of hybernation. It only continued longer beneath the surface without rising to respire. At this period, having found that some frogs in their natural haunts had already come forth, I removed the subject of these observations also from the water to a damp locality, and on the following day found it greatly changed in appearance. While confined in the water it was of a dull dirty brown colour, but some hours after its removal it cast its tegument, and changed to a bright yellow, with the usual brown markings, and had increased in size, both in its body and limbs.

The conclusion to which these circumstances seemed to lead was, that quickly after the frog leaves its hybernaculum, it casts its tegument as the insect escapes from its puparium, and acquires new vigour, while the ova are attaining their full growth. The Toad undergoes a similar change. About a fortnight later in the season than the frog, I have seen many toads in a shallow ditch of slow moving water in the act of casting their dark brown tegument, and acquiring one of a greenish yellow.

On examining several frogs taken from their natural haunts, I found them in, as nearly as possible, the same state of development with regard to the ova, judging from external appearance, as the specimen I had watched through the winter. A few had just paired, but the majority were still single. On opening the bodies of the latter, I found the ovaries greatly enlarged, and the ova, apparently ripe, but still contained in the ovisacs.

It is at this period, therefore, immediately after hybernation, and before the ova have left the ovaries, that the condition of the ovum is a matter of great interest with reference to the structure and contents of the germinal vesicle, the period at which the vesicle is changed or disappears, and the circumstances under which the ovum escapes from the ovary, and is received into the oviduct.

The Germinal Vesicle.—The fate of the germinal vesicle in the matured ovum is still a matter of doubt. Previous to the embryological researches of Dr MARTIN BARRY, it was usually believed that the vesicle entirely disappears before or at the time of fecundation of the ovum. But this view was combated by the author named, who, quoting the opinions of previous inquirers, contended that the germinal vesicle in Mammalia does not disappear, as believed by PURKINJE,* in Birds and Amphibia, by bursting during the generative act, and pouring its contents into the germinal layer of the fecundated ovum.

* Symbolæ ad Ovi Avium Historiam ante Incubationem, 1825; and Article "Ei," Encyclop. Wörterbuch, Band x. p. 112, 1834.

Neither, as supposed by BAER,* by being urged forwards and burst between the vitellus and its membrane before fecundation. Nor, as stated by RATHKE,† with reference to the Crustacea, by disappearing while still within the ovary. Or, as supposed by WHARTON JONES,‡ in Birds and Amphibia, by approaching the surface of the yelk, and by the giving way of the coats of the vesicle, and the effusion of its contents on the surrounding surface of the yelk. Or, further, as believed by BISCHOFF,§ by disappearing at the time of exit of the ovum from the ovary. On the contrary, Dr BARRY has stated that the germinal vesicle returns to the centre of the ovum, and the germinal spot to the centre of the vesicle, before the ovum leaves the ovary, and that these do not become dissolved, but only are changed in character by a process of cell development within them, which ends in the production of two cells in the centre of the yelk, which are the foundation of the body of the future embryo.

Perhaps what I am about to mention may assist us to reconcile or correct the views of this embryologist, as well as those of the authorities he has quoted. It is well known, that if the ovum of the frog is examined before the yelk has attained to one-half of its ultimate dimensions, the germinal vesicle is distinctly visible as a large circular body near the centre of the yelk, apparently granular in its interior, and more opaque than the yelk itself. In the centre of the vesicle its nucleus, the germinal spot, is then equally distinct. If the ovum is examined when it has nearly acquired its full dimensions, the vesicle is still found to exist, but, as compared with the size of the entire ovum, is relatively smaller than at earlier periods, and is recognized with more difficulty, owing chiefly to the yelk cells having both increased in number around it, and also acquired a darker colour. It is equally well known that all appearance of the vesicle is lost in the amphibia, as in birds, before the ovum is prepared for fecundation; but as to the way in which it disappears, or even as to the time, observers are not agreed.

Structure of the Germinal Vesicle.—Having collected a number of frogs that had recently left their hybernacula, and had not yet paired, I placed some in spirit for examination. On dissecting them afterwards, I found that in some the ova had nearly reached maturity, but had not left the ovary. The yelks being rendered firm by the spirit, I was able, by gentle pressure, to break open some of these beneath the microscope, without

* "Lettre sur la Formation de l'Œuf," in Breschet, *Repertorium*, 1829.

† *Untersuchungen ueber die Bildung und Entwickelung des Fluss Krebses*, fol. Leipzig, 1829.

‡ *Philosophical Transactions*, 1837, part 2.

§ R. WAGNER's *Lehrbuch*, &c., 1839.

diffuence of the contents. The aggregated yelk cells were then seen to consist of two kinds; the one dark-coloured, which form the upper portion of the yelk, and which diminish in intensity of colour from brown or black at the circumference to a leaden or grey in the centre; the other, which forms the inferior portion, consisting of cells that become lighter-coloured as they approach the surface. The whole form a mass, of nucleated cells of nearly uniform size, closely aggregated together. In the interior of the yelk, amidst the dark-coloured cells, and *much more near the surface of the dark* than that of the white portion of the ovum, I was surprised to find the germinal vesicle still entire, and of a somewhat oval, lenticular form, although irregularly compressed by the contraction of the whole egg in the spirit. It was of a dense white colour, and opaque, from the action of the spirit, and was in striking contrast to the dark cells of the yelk, which adhered to its surface, and amidst which it lay imbedded like the kernel in a peach or apricot. On the surface of the black portion of the yelk was a minute orifice, already noticed by PREVOST and DUMAS,* the outlet of a canal that passes through this portion of the yelk to the germinal vesicle in its interior. This I believe to be the result of the yelk cells having only imperfectly closed around the germinal vesicle. It is however of some consequence in the future development of the ovum, as it is in this canal, that the cleavage of the yelk is commenced. The vitelline membrane was already formed, but I could not discover any orifice or perforation in it, either corresponding to the canal in the yelk, or to any other part of the surface. It is not possible to mistake the germinal vesicle for any portion of the lighter substance of the yelk,—first, from the fact that the vesicle at this period is of an intense opaque, white colour, very different from that of the yelk substance,—next, from its being completely isolated from the lighter and imbedded in the dark substance,—and lastly, from its being still invested with a distinct envelope. On removing the vesicle, and examining it separately, first, as taken from the yelk, without crushing it, and next by gentle compression, and with the highest powers of the microscope, the interior was seen to be filled with secondary cells. Each of these, formed by a distinct envelope, *appeared* to contain other, or tertiary cells, and strongly reminded me of the developmental cells of the spermatozoa in the male organs, since these again seemed to contain granules, or quaternary cells. In the midst of the secondary cells I was able to distinguish, in the centre of the germinal vesicle, in some specimens, one or two cells of larger size than the rest, and which I regarded as the remains of the

* *Loct.* tom. ii. 104.

germinal spot, or central nucleus. In those ova which, from their size and general appearance, seemed to be the most mature, the peripheral series of cells within the germinal vesicle were of smaller diameter than those nearer to the centre, as if the earlier developed secondary cells had disappeared and liberated their contents. The cells of the dark portion of the yelk that adhered to the germinal vesicle were also nucleated, but were of much smaller size than the peripheral cells of this body. I must remark, however, that saving the fact noticed, of having seen in some vesicles one or two cells near the centre of larger size than the rest, I have not been able to distinguish any separate nucleus, or germinal spot, which seems, at this time, to have disappeared as a distinct body. The germinal vesicle thus near its maturity is a mother-cell that contains a multitude of daughter or secondary cells, each, apparently, including its own progeny.

It is thus certain that the germinal vesicle exists in the ovum of the frog until after the period of hybernation. But in individuals that have been several days abroad, and in which the ova are so far matured as greatly to distend the body, and be nearly ready to leave the ovisacs, I have not always been able to detect the vesicle. The yelk has then a greater proportion of white substance in its interior, intermingled with the dark, and this, I suspect, is the result of the disappearance of the vesicle.

In the ovum of the *Toad*, when as nearly mature as that of the Frog before leaving the ovisacs, I have been unable to detect any trace of the germinal vesicle. The dark portion of the yelk is then quite distinct from the light, and forms a cortical stratum of intensely black pigment, scarcely more than than one-sixth of the diameter of the yelk in thickness. The light portion is of a yellowish white, and from which the germinal vesicle, when present, is not readily distinguished. But at a little earlier period, although only a few days before the pairing of the Toad, I have found the vesicle fully as large and as distinct as before its disappearance in the egg of the Frog, but situated more distant from the centre, and nearer to the black, the future dorsal surface of the egg.

In the great Water Newt, *Triton palustris*, the yelk of the matured ovarian ovum is of a dull pea-green colour, which in spirit is changed to a greenish-yellow. On breaking open the yelk of this species, after it has been hardened in spirit, I have always found the germinal vesicle of the same opaque, intense white colour, of precisely the same structure, and situated in the same part of the yelk as in the Frog. It has also a canal

passing from the middle of the surface of the darker coloured portion of the yelk through its substance to the vesicle, as in the Frog.

In the lesser Newts, *Lissotriton punctatus* and *L. palmipes*, the yelk is of a brown or liver colour on the future dorsal surface, with a white central spot, and of a white or pale straw colour on the ventral. In ripe ovarian ova of these species also I have found the germinal vesicle of the same white colour, and having the same structure as in the larger Triton and the Frog.

In each of these instances, not merely when the ovum is immature, but even when nearly ready to leave the ovisac, the germinal vesicle is situated in the interior of the yelk, and not immediately at the surface, and thus far the fact is in accordance with the observations of Dr BARRY, that in the Mammalia the germinal vesicle is in the interior of the ovarian ovum, and does not disappear on the surface. But, nevertheless, it is not in the *centre* of the yelk; its place, as I have stated, is *excentric*.

Time of disappearance of the vesicle.—It is well known that in frogs and toads the ovum is never impregnated until after it has left the body, and consequently not until long after it has left the ovary, and the germinal vesicle has entirely disappeared. This disappearance does not, I think, take place at the instant the ovum is about to leave the ovisac, but a short time before, as I have found the ova of the Frog in one instance more matured than in that already described, still contained in the ovisacs, and yet most of them without any remains of the vesicle which I could identify as such, and with the yelks containing a larger proportion of white substance, as in the Toad. In a very few, however, the vesicle was still present, and exhibited the structure I have described. Certainly, then, the vesicle is not burst at the moment the egg escapes from the ovisac. There seems to be a short period of time between the disappearance of the vesicle and the full maturity of the ovum, during which the yelk itself undergoes some further change, and acquires the appearance noticed in the matured eggs of the Toad, and in the most advanced of those of the Frog. No trace of the germinal vesicle can be detected in any ova that have left the ovary and are contained in the cavity of the abdomen, before entering the oviducts. I have examined many ova, both of the Frog and Newt, from the abdominal cavity, but in every instance the vesicle has entirely disappeared. In some specimens, which seemed to have been in the act of escaping from the ovary when the animal was killed by immersion in spirit, I have found in the place occupied by the vesicle an

aggregation of white nucleated cells, which, examined by the microscope, exhibited a close resemblance to those seen in the interior of the vesicle. In the midst of these there has occasionally been one or two of larger size than the rest, and which I have imagined to be the remains of the germinal spot, and possibly the origin of the future embryo vesicle of the impregnated ovum, an opinion, however, which I have not had the means of verifying, and I must further state that I have failed to recognise these larger cells in ova that were free in the cavity of the abdomen. Each of the three species of Newt, as well as the Frog, have presented similar appearances in the germinal vesicle and ovum under similar circumstances.

Thus it is quite certain that the germinal vesicle disappears in the Amphibia before the ovum enters the oviduct. I believe it does so in the interior of the yelk, not in the centre, but nearer to the dorsal than to the future ventral or white surface; and not, as has been supposed, on the dorsal or dark surface, between the vitellus and the vitelline membrane. This view is supported by the fact, that that portion of the yelk which incloses the vesicle in an advanced stage of the ovum in the Frog is of a more or less intense black colour, while the vesicle is perfectly white; and that at a further advanced stage, after the vesicle has disappeared, and its place is occupied by a collection of white cells, the dark portion of the yelk still preserves its intense black colour, except at the point that corresponds to the central canal, which then has a leaden hue. PREVOST and DUMAS,* and also RUSCONI,† have mentioned that there is a yellow spot at a corresponding part of the dark surface of the egg of the species they have examined after impregnation, *Rana esculenta*?, but these appearances must not be mistaken for the general vesicle arrived at the surface. There is a similar spot, and that too of an elevated form, on the egg of each of the *Lissotritons*. But independent of the fact that the germinal vesicle has entirely disappeared from the interior of the egg before it escapes from the ovary, this spot is shown not to be the vesicle, both in the fact that in the egg of *Rana temporaria* the dark portion of the yelk is unchanged, while in each case the spot is perforated, and leads into the canal that passed originally to the vesicle. I regard the spot as simply a protrusion outwards of the edges of the canal while closing, after the vesicle has disappeared. I shall presently show that a similar white spot is formed on the under surface of the egg of the Frog soon after deposition, and which might equally well be mistaken for the germinal vesicle.

* Annales des Sciences Naturelles, tom ii. p. 104.

† Developpement de la Grenouille Commune, 4to. Milan, 1826, p. 9.

Mode of disappearance of the Vesicle.—The mode in which the vesicle disappears may be inferred from the facts of its structure. Being filled with a progeny of cells which we may regard as of different periods of growth, and these again containing others, it is fair to conclude that this process of cell-formation is that by which the parent vesicle is ultimately destroyed. At the time when the germinal vesicle has nearly attained its full size, the peripheral cells are smaller than those nearer to its centre, while the yelk cells that surround the vesicle are still smaller than either, and are of a dark colour. When, therefore, the vesicle has acquired its full size, by the simple vegetative endogenous growth of the contained cells, we may fairly presume that the death of the parent mother-cell, or germinal vesicle, takes place as the result of their enlargement, by the diffuence of its investing membrane; and the enclosed daughter cells, thus gradually set free in the midst of those of the yelk, as in the ideal, form one mass with the latter, and the moment of the actual disappearance of the vesicle thus escapes direct observation, its previous existence being indicated only by the unbroken outline of the investing membrane.

These views lead me to agree with WAGNER and BARRY in regard to the structure and mode of growth of the germinal vesicle, but not as to that by which it disappears. Dr BARRY indeed believes that the vesicle only becomes changed by its mode of development, and does not cease to exist. But most certainly it does disappear in the Amphibia, and, as I believe, through the growth of the young cells in its interior. I cannot therefore agree with Dr BARRY, that the changes in the vesicle end in the production of two cells in the centre of the yelk, that give immediate origin to the embryo; but rather believe that it is from one of the central cells of the germinal vesicle that the future *embryo vesicle* takes its origin, while the remainder of the liberated cells are distributed with this through the substance of the yelk, when the segmentation of this body takes place. This opinion is in accordance with that of VOGT,* who found that in the ova of *Alytes obstetricans* the germinal spots increase in number, and that a few hours after fecundation small vesicles, similar to these spots, are scattered through the yelk. I have myself found similar vesicles in the fecundated egg of the Frog about three hours after impregnation, but have not traced them to their origin. With regard to the disappearance of the spot in the germinal vesicle, the facts observed in the ova of the Frog agree with those noticed by KOLLIKER† in the ova

* Untersuchungen über die Entwicklungsgeschichte der Geburtshelfer-Kraute (*Alytes obstetricans*), 4to, 1842.

† Müller's Archiv, 1843.

of intestinal worms, that all appearance of the spot is lost before that of the vesicle. This circumstance, however, may be owing either to the spot having given origin to cells in the vesicle which quickly attain to similar dimensions, and from which it is not otherwise distinguished, or to its becoming entirely obscured by their multiplication. Although no observations have been made on the origin of the embryo vesicle that appears in the yolk after the disappearance of the germinal, I am still inclined to regard this as being in some way derived from the lost germinal spot, notwithstanding that KOLLIKER found a certain period of time elapse between the disappearance of the vesicle and that at which he was able to recognize this body.

Transit of the Ovum.—Thus, then, when the ovum escapes from the ovisac and ovary into the cavity of the abdomen, the germinal vesicle and spot have entirely disappeared, and it consists only of the yolk enclosed in an exceedingly delicate, structureless, vitelline membrane. This is its condition in the Frog, Toad, and Newts. It is then extremely delicate and easily lacerated. The mode in which the ovum passes into the oviduct has been the subject of much inquiry. I am quite certain, as SWAMMERDAM long ago showed, that the ova, when mature, pass from the ovaries into the cavity of the abdomen, and from thence into the oviducts, in the Frogs, Toads, and Newts, quite independent of any intercourse with the male, as I have myself recently had an opportunity of proving. The frog I have already mentioned as having watched through its season of hibernation, was kept apart from all others until the 2d of April, at which time she had not deposited any ova. But from the altered form of her body it was evident that the ova had passed, or were at that time in the act of passing, into the oviducts. I then placed her in a vessel with others, some of which were males, but not one of these joined with her. Nevertheless, on the 6th of April she cast her ova, without having paired, and died on the following day. On examination after death, I found that the whole of the ova had left the ovaries, all of which, excepting only two still free in the cavity of the abdomen, had passed through the oviducts without any intercourse with the male. These ova of course were sterile; but it was worthy of note that their envelopes did not expand to so great an extent as those of the eggs of paired individuals.

Passage into the Oviduct.—In what way the ova pass from the cavity of the abdomen into the mouth of the duct has never been satisfactorily explained. SWAMMERDAM examined the question with much care,* but was unable to form any decided opinion respecting it, as he correctly states that the mouths of

* Book of Nature, Part ii., pp. 108, 109.

the ducts are at a distance from the ovaries, and are not free to grasp the ova like the fimbriated extremities of the Fallopian tubes in Mammalia, but are confined in the peritonæum, which is continuous with that which passes over the pericardium and heart. PREVOST and DUMAS have since stated* that the ova after leaving the ovaries are seized by the tubes ("sont saisis par des trompes;") but they do not show in what way this seizure is effected. They have omitted to describe the structure of the parts concerned in the act, and have not mentioned the way in which the ova are conveyed to the tubes from the ovaries, or whether the mouths of the tubes approach the ovaries, as in Mammalia,

The entrance to the tubes in the Frog, as SWAMMERDAM has correctly shown, is in the peritonæum at each side of the heart, and I have found it in nearly the same place in the Newts. The apex of the pericardium in the Frog is attached to the cartilage of the sternum by two layers of peritonæum, which together form the mediastinum, and enclose between them the trunk of the median abdominal vein, a branch of the Vena Cava. Tracing one of these layers of peritonæum upwards and over the pericardium, we find in it an orifice, at the part where it is reflected on itself to form the lateral portion of the suspensory ligament of the liver. This orifice is elongated, oval and funnel-shaped, and, when dilated, forms a kind of pouch at the anterior boundary of the space or cavity between the liver and the heart, and laterally it is in free communication beneath the suspensory ligament with the common cavity of the abdomen. The oviduct commences in this dilated orifice, as a narrow tube with thick muscular parietes and with a thick mucous lining. It passes at first upwards and forwards, confined to the peritonæum, and then outwards above the base of the lung, gradually increasing in its dimensions. Immediately after it has passed the lung, it becomes more enlarged, and as it passes backwards to the side of the spine forms many convolutions, which end in the dilated oviduct or common receptacle for the eggs ready to be deposited.

The commencement of the oviduct in the Newts is very similar to that of the Frog. In the *Triton palustris* it differs only in the entrance being larger, and situated more to the side of the body, and above the lung, to the base of which, as well as to the peritonæal investment of the heart, it is confined, as in the Frog; but it has a more free communication laterally with the common cavity of the abdomen than in that animal.

The ova escape from the ovaries into the cavity of the abdomen among the viscera both in the Frogs and Newts, and

* Loc. cit. tom. ii. p. 105.

are carried forwards to the spaces between the liver and heart on each side to the dilated mouth of the oviduct. They certainly are not seized by the tubes as they escape from the ovaries, as they are constantly found free in the abdominal cavity, while the mouths of the tubes being confined in the peritonæum and having no appendages, cannot be extended to reach them. Their transfer seems to be effected in the Frog in part by the action of the abdominal muscles forcing them onwards in the spaces between the viscera, aided perhaps by the peristaltic action of the stomach and intestines; and their entrance into the tubes, when arrived in the vicinity, seems to be induced by an ingurgitory or suction action at the mouth, occasioned by the alternating and pulsatory motion of the heart, with which the tube is connected by means of the peritonæum. The tube itself is formed of strong longitudinal and transverse fibres, which are continued into the peritonæum, and the former especially into the suspensory ligament, the free external margin of which bounds the outer side of the orifice. The transverse fibres are strongly marked at the commencement of the orifice, where there is a slight pouch; so that when the eggs are entering, these fibres doubtless prevent their return and transfer them onwards.

This, I believe, is the way, in which the eggs enter the oviducts. It is quite certain from the anatomy of the parts, that they cannot be grasped by the oviducts until they are conveyed to them. I have not actually witnessed the passing of the eggs from the abdomen into the ducts in the Frog, but I have seen the eggs moved onwards in the smaller Newt, *Lissotriton palmipes*. Having deprived a female of this species of sensation and power of motion, by division of the spinal cord through the medulla oblongata, I proceeded to open the abdomen to obtain ova from the oviducts for experiments on artificial impregnation. I then found, that a number of ova were free in the abdominal cavity, and that some had very recently entered the ducts, while others were in the immediate vicinity of the mouths. The heart was still pulsating vigorously and with great regularity; and I then saw that at each pulsatory action the ova passed slowly forward between the liver and lung, towards the mouth of the oviduct, which still contained two or three ova that appeared to have entered at the moment of the operation. I did not witness the actual entrance of an ovum, but saw that the action of the heart certainly had the effect of inducing the advance of it to the mouth of the tube, and quite sufficient to lead me to regard this as one of the chief means of its entrance into the duct.

It is not until the ovum has become clothed in the oviduct with its gelatinous envelope that it is susceptible of impregna-

tion. This remark applies equally to the Frogs, Toads, and Newts. The ova of the Frog and Newt at large in the abdominal cavity are always entirely without this envelope, and consist simply of the yelk mass enclosed in an extremely delicate vitelline membrane. They are so easily injured that it is only with great difficulty that they can be removed from the abdomen for examination unbroken. Those of the Newt, when taken up ever so carefully by means of a hair pencil, often burst the membrane simply by their own weight. But immediately after they have entered the oviduct and begin to acquire their envelopes, the yelk appears to undergo some change, as it becomes much firmer and is less easily injured. Shortly after the egg has entered the duct, it gains the first layer of an investment, which, from the great similarity it bears to the gelatinous layer gained by the ovum of the Rabbit in the Fallopian tube, and regarded by its discoverer in that animal, Mr WHARTON JONES,* as the origin of the chorion, I am disposed, with him, to look upon as the analogue of that layer. This covering adheres very closely to the vitelline membrane, and is scarcely to be distinguished from it, except at certain periods of change. It is acquired before the egg has arrived at the first convolutions of the oviduct. During the remainder of its passage the egg gains two other distinct layers of similar investment, which, together, we afterwards recognise as the gelatinous envelope of the Frog and Toad, and the capsule of the Newts.

These envelopes are not merely simple means of protection to the egg during the production of the embryo, as has been supposed, but, as I shall presently show, are essential to it at the period of fecundation, and without which the egg is not susceptible of impregnation. The layer of envelope which I regard as the foundation of the *chorion*, is a dense, but very transparent thin covering, in immediate contact with the vitelline membrane, and is formed of cells so closely aggregated together as to have coalesced into a fibrous structure. The two layers external to this are also formed of cells, which, with their nuclei, are distinctly visible in the envelope of *Triton palustris*, in which they alternate in regular series. Although these layers, which constitute the jelly in the egg of the Frog, become detached in that of the Newts quickly after oviposition, and, expanding as in the Frog, they leave the egg at liberty in a chamber in their interior, they are nevertheless essential to the impregnation of the ovum, which takes place before or at the time of leaving the body, as in Frogs and Toads. RUSCONI† removed the envelopes of the egg of frogs, and found

* Phil. Trans. part 2, 1837.

† Développement de la Grenouille commune. Milan, 1826.

that the embryo still became developed, and thence concluded that these coverings serve only mechanical purposes during the changes: but it will presently be seen that they have a more important function at a much earlier period. During the time they are in course of formation around the egg the yelk undergoes some further change. The light portion becomes of a whiter, and the dark portion of a deeper colour. Internally the cells vary more in size, the lighter-coloured being the largest. I have not succeeded in recognizing any embryo or central vesicle up to this period.

2. CHANGES AFTER SPAWNING AND IMPREGNATION.

First Period of Development.—It has been long known that a division or cleavage of the yelk of the Frog's egg is one of the earliest and apparently invariable results of fecundation. The primary division was first seen by SWAMMERDAM, and was figured* and mentioned, but was not understood by him. SPALLANZANI long afterwards recognized it in the egg of the Toad (*Alytes obstetricans*†), which he says becomes about a day after fecundation marked “with two furrows which meet to form an angle,”—that the furrows afterwards become deeper, and that “two small tumours arise on each side of the furrows,”—changes which have since been more accurately and completely described by VOGT‡. SPALLANZANI also says that the egg of the common Toad is “marked with four furrows which intersect each other at right angles nearly like the husk of a chestnut half opened,” but he seems to have thought this was the usual condition of the ovum. To PREVOST and DUMAS,§ however, we owe the important discovery of the cleavage of the yelk as a process of the fecundated egg; to RUSCONI,|| BAER¶ and others, its full exemplification in the Amphibia; and to BARRY** and BISCHOFF†† its detection and elucidation in the Mammalia.

The agent immediately concerned in these changes is believed to be the embryo vesicle and its progeny, produced after the disappearance of the germinal vesicle. But it is yet uncertain what is the origin of the embryo vesicle, or whether it exists in the unfecundated ovum. As cleavage of the yelk certainly is not the result of the disappearance of the germinal vesicle, which disappears from all ova of the Amphibia, whether

* *Loc. cit.* tab. xlvi. figs. 5, 8.

† *Loc. cit.* vol. ii. p. 159.

‡ Untersuchungen über die Entwicklungsgeschichte der Geburtshelfer-Kräte (*Alytes obstetricans*), 1842.

§ Annales des Sc. Nat. tom. ii. p. 110, 1824.

|| *Loc. cit.*

¶ MÜLLER's Archiv, 1834.

** Philosophical Transactions, 1839, 1840.

†† Entwicklungsgeschichte des Kaninchen-eies, 1842.

they are afterwards impregnated or not, I was desirous, at the commencement of my experiments on impregnation, to learn *from direct observation* whether the unfecundated ovum ever passes through any stage of cleavage; since the ascertainment of the fact in the negative would be an important test in the experiments I was about to make. For this purpose it was necessary to collect many pairs of frogs at the proper season, and when, from symptoms which are soon recognized, it was found they were about to cast their ova, to wait patiently, perhaps for many hours, for the result, in order that the exact condition of the ovum, impregnated naturally, should be first ascertained. SWAMMERDAM long ago remarked that the spawning of the frog takes place very rapidly "by a single effort."* It is often completed, as I have found in the English species, in a few seconds, and usually in less than a minute, during which the male impregnates them, so that if the animals are not closely watched, the opportunity of observing the earliest appearances of the ovum is lost. Having noted the condition of the impregnated ova of several pairs of frogs within the first few minutes after spawning, I found those of different individuals vary much with respect to the white or inferior surface, and exhibit appearances that may readily be mistaken for the breaking up of a vesicle on the surface. This appearance is due to a more or less complete state of maturity of the eggs of different broods, and according as their spawning has been retarded or hastened. The peculiarities are the most marked in the least matured, the white surface of the egg being the last completed part, and forming the base of the egg in the ovisac. Having noticed the appearances of the eggs when impregnated naturally, I was enabled to compare them with others impregnated artificially, and these with some of the same brood not impregnated.

Immediately after the frog has spawned, the ova form a close rounded mass, which at first is scarcely so large as a walnut. They then seem to consist almost entirely of dark-coloured yolks with thin gelatinous envelopes. The form of the egg is then somewhat oval, with the white portion a little more conical than the dark, and differing slightly in different ova. In some there is a dark spot in the centre of the white, that looks like a depression or cavity, or perhaps a vesicle. I am not certain that this appearance is in reality a vesicle, and therefore am content to describe it as a spot, although it conveys the idea of being a vesicle. In some ova there are two, four, or six of these spots imbedded each in a small portion of white substance.

* *Loc. cit.* part 2, p. 111.

When only a single spot exists, the white surface of the egg for some space around it is more defined than afterwards, and exhibits faint indications of a crucial division of the yelk on this surface immediately around the spot. This is the condition of a few ova immediately after spawning; but the majority have then advanced farther in their changes, and show four or six rounded dark-coloured spots at a little distance in the place of the single central one. When four spots occur they are usually arranged in a quadrangle, and are less than their own diameter apart. They convey the idea of being derived from the central one; but I have never seen any division of this, and if such division takes place, I think it must occur before or at the very moment the ova are expelled. In a further advanced stage of the ovum the four dark spots have become larger, and are each imbedded in a distinct portion of the white surface.

One minute after deposition the spots are more widely separated, and are then each encircled by a separate patch of white substance. *Two minutes* after spawning six dark spots have made their appearance, one of which is situated nearly in the centre, and the remaining five are so arranged around this that the white patches in which they were imbedded seem to have coalesced. In *three minutes* the spots are further enlarged, and appear joined by a dark line of colour extending from each, so that the whole form, as it were, a knotted ring that includes a patch of the white surface of the yelk with one of the dark spots near the centre. In *four minutes* the ring around the included white substance is more distinct, and the white surface of the egg has increased in extent. In a further advanced stage at this period the white portion included in the ring exhibits the appearance of a white, very opake patch, the dark spot in the centre having disappeared. Around this opake white patch is the dark-coloured knotted ring, now become more uniform, and resembling a ragged chink or slight circular furrow or division in the white surface. The centre of this hemisphere of the egg thus comes to be occupied by a white patch instead of the dark spot. At *five minutes* this central white patch,—which, as before stated, and from what afterwards occurs, may readily be mistaken, on casual inspection, for the germinal vesicle, altered in its appearance and arrived at the surface,—were it not that we now know that this has long before entirely disappeared,—becomes more defined, and the dark circle around it is more uniform and distinct. At *ten minutes* the central patch is a little reduced in size, and the circle that encloses it begins to take the appearance of a diffused halo. At *fifteen minutes* the central white patch is more reduced, and the halo is spread wider, while the whole of this

hemisphere of the egg has acquired a whiter appearance, and become more distinct from the dark colour of the sides and future dorsal hemisphere. At *twenty minutes* the central patch has become still smaller and rounder, and the dark halo much broader.

At this period an interesting circumstance occurs, which may hereafter be found to have some reference to changes in the interior of the yelk, possibly to some rapid evolution of the so-called central or embryo vesicle in the locality originally occupied by the germinal vesicle and spot, which, as we have seen, is nearest to the dark surface. This is the partial rotation of the entire yelk. Up to about this period the ova remain undisturbed in the water in a mass as they are expelled, and lie indiscriminately, some with the dark and some with the white portion of the yelk uppermost, or horizontal. But during the time that has passed since the ova have been in contact with water, the envelopes have imbibed fluid and expanded until these investments of the yelk have acquired a thickness equal to about two-thirds of the diameter of the yelk itself. The yelks, that have remained to this time with their white surface uppermost, now change their position spontaneously by a partial rotation of the whole mass of each on its axis, within the vitelline membrane, until the dark surface of the whole is placed uppermost. Whether this change of position is merely the result of an expansion of the vitelline membrane at this period, when the ovum is rapidly ceasing to be susceptible of impregnation, as I shall presently show is the case after this lapse of time in the water, or whether it be also connected, as we may fairly believe, with changes going on in the interior of the yelk, I am not prepared to decide. It is important, however, to note, that the change takes place at about the time, at which I have found a great abundance of bright clear rounded vesicles distributed throughout the yelk, but chiefly in the place originally occupied by the germinal vesicle. In some of these vesicles, which I regard as the progeny of the germinal vesicle, I have seen irregular-shaped nuclei that appeared to be formed of a multitude of nucleoli. These vesicles convey to me the same idea as that conveyed by the vesicles seen by BISCHOFF in mammalian ova, excepting only that in the egg of the Frog they contain compound nuclei.

At *thirty minutes* the central patch on the white surface of the egg has almost disappeared, and the halo around it is still more diffused. At *forty-five minutes* it has entirely disappeared in most specimens, and its place is occupied by a broad dark area which includes the boundary of the previous halo, and which appears to be occasioned by a slight depression in the

centre of this surface of the yelk. *One hour* after spawning this depression is somewhat deeper. The white surface has become still more defined, and the dark has acquired a more intensely black colour. The egg remains in this state without further perceptible change during the succeeding *second* and *third* hour, excepting only that the depression in the white surface becomes a little deeper, but it has almost disappeared at the end of the *fourth* hour, when *segmentation* or cleavage of the yelk is about to take place. But this is not invariably the case. When it does remain it is always of an oval form, and the primary cleavage of the yelk, as it proceeds on either side, from above downwards, meets in its centre and invariably passes through it transversely to its long diameter. These are the first perceptible changes in ova that are impregnated by the natural union of the sexes, and when spawning has not been retarded. But in some broods of eggs that have been retained longer than usual in the oviducts, the whole of these changes have already taken place, in so far as regards those of the yelk, the white surface of which then exhibits an uniform appearance.

Changes immediately before segmentation or cleavage of the yelk.—Segmentation usually commences in from four to five hours. At about *one hour and a half* after spawning, the peripheral layer of cells on the middle of the dark or uppermost portion of the yelk of the impregnated ovum becomes separated from the inner surface of the vitelline membrane, and this separation goes on until a broad free space is left between this envelope and the superior layer of yelk-cells. This space, which we may designate the *respiratory chamber*, is at first but a small area above the middle of the dark surface of the yelk, and is commenced above the central canal. It seems to be occasioned by a recedence towards the interior, or a shrinking, at this period, of the yelk cells of the dark hemisphere of the egg, commencing in the centre of this part and extending gradually, but in a less degree, to the circumference. This recedence goes on until the space left between the vitelline membrane and the yelk is equal to about one-sixth of the diameter of the whole mass, when the space appears to be occupied by a very transparent fluid, interposed between the now depressed surface of the yelk and the vitelline membrane. In the centre of the black surface is the minute orifice noticed by PREVOST and DUMAS,* and BAER,† which leads into the central canal that communicated with the germinal vesicle in the ovarian ovum. It is in the margins of this canal that segmentation is commenced. While the space or chamber between the black

* *Loc. cit.* Tom. ii. p. 104, 1824.

† MULLER's Archiv, 1834.

portion of the yelk and the vitelline membrane is being formed, and from fifteen to thirty minutes before there is any sign of cleavage, the yelk becomes extended horizontally in a direction transverse to that in which the first cleft afterwards takes place, and assumes a transitory obtuse oval form, which it retains until the yelk begins to divide. The division, as correctly shown by BAER,* commences in the extension in opposite directions of at first a faint indentation in the margin of the central canal, which quickly becomes deeper, and is carried across the surface of the yelk, and gradually more and more deepening and widening as it proceeds, is carried round the sides, and meeting in the middle of the depression on the under surface, or of the remains of the white patch when this has not already disappeared, is completed by passing through the middle of the yelk ; which is thus divided into two portions. This first division occupies from twenty to thirty minutes before it is finished, and it is not until then that a second fissure is commenced.

I have not had any opportunity of proving whether this division is the direct result of subdivision of the central vesicle, and the attraction of the yelk-cells in equal proportions around each division of that body, as believed by KOLLIKER,† and, as it seems fair to infer, is the case ; but in addition to the observation by Professor SHARPEY,‡ that the contraction of the entire yelk at the commencement of these changes, and the movements he has observed among its granules as they proceed, are in favour of this opinion. - I may remark that the extension of the yelk of the Frog's egg in a direction transverse to the first cleavage, may also be advanced as conformable to the same view. It is further supported by the fact which I have seen in the egg both of the Frog and Newts, that before the second or crucial cleavage is commenced the yelk becomes contracted, so that the first cleft for a time is almost imperceptible while it is extended in the transverse direction, or line of axis of the first division, after which the second or crucial cleft is commenced. To this I may add another fact which appears to be equally significant, and which occurs in the unimpregnated eggs both of the Frog and Newts, but more especially in the latter. Although no recedence of the yelk from the vitelline membrane takes place in the unimpregnated egg of the Frog, it becomes, nevertheless, slightly oval after the first few hours, but returns to its original shape some time afterwards. But the unimpregnated egg§ of the Newts is not only separated from the vitelline membrane, but also is depressed, and has a

* *Loc. cit.*

‡ QUAIN'S *Anat.* fifth edition, 1848.

† MULLER'S *Archiv*, 1843.

§ Or, possibly, partially impregnated.

distinct pit in the centre of its upper surface, and also assumes an obtuse oval form, both which it retains, when the egg is preserved in water, until decomposition has commenced.

Changes in the impregnated and unimpregnated Ovum compared.—Having traced the egg impregnated by natural union of the sexes through its first period of development, I was able to compare the phases it exhibits with those of the artificially impregnated, and these with the appearances in unimpregnated eggs of the same brood, placed under precisely similar circumstances with reference to light, heat, air, water, and locality. The ova experimented on were all procured from the same female, and the seminal fluid from the male with which she was paired, and at the time the female was about to spawn.

SPALLANZANI obtained unimpregnated eggs of the Frog for his experiments by opening the body of the female and removing them from the distended oviducts. This mode is exposed to the objection, that in the removal of the ova they are liable to be brought into contact with the blood of the animal from the cut vessels, and that the ova thus obtained may not be the most mature, and fitted for experiment. It seemed desirable, therefore, to obtain them by another mode,—the total withdrawal of the influence of sensation, and power of tension in the muscles by division of the spinal cord through the medulla oblongata. The attempt was made with a female frog that had been paired for several days, and, from appearances, would have deposited her ova naturally in the course of a few hours. The spinal cord was divided as quickly as possible with a strong pair of scissors immediately behind the brain, and this organ was also destroyed, so that all consciousness was annihilated. The attempt was successful. The muscles deprived of voluntary power instantly became relaxed, and allowed of the ova being passed, by gentle compression of the body, through the natural passage without contact with the blood of the animal, in greater or smaller number at pleasure, and thus afforded easy means of experiment.

I may once for all state that it was in this way that the ova were always obtained in the following investigations.

Recourse was had to a similar expedient to procure the seminal fluid from the male. SPALLANZANI had obtained the fluid, by vivisection, from the seminal vesicles themselves. But there seemed more objection to the adoption of this mode with the male than the female, the certainty of much of the fluid being lost, independent of the severity of the operation. Indeed SPALLANZANI states, that he was never able to procure more than from two to three grains from a single individual. I therefore

availed myself of a habit in the male frog, and which SPALLANZANI had previously noticed and taken similar advantage of in the newt, to obtain the fluid in greater quantities than by the mode constantly adopted by that physiologist. When the male frog, like the newt, is taken in the hand, or slightly compressed, at the season of pairing, a quantity of fluid is immediately passed. This consists chiefly of seminal fluid mixed with water expelled from the effect of the compression, or during the efforts to escape, as water is passed by other animals at the moment of capture. It abounds with spermatozoa in their most active state, and thus is fitted for experiment. It required therefore only to secure the limbs of the animal and compress it slightly, to obtain the fluid without severe injury. This ready mode was adopted on all occasions when the fluid was required, and the precaution taken always to examine a portion with the microscope, to be assured of its nature before employing it. Spermatozoa have never, during the season of pairing, been absent from it. At the end of the season they have been less abundant, and spermatozoal cells in greater proportion than at an early period. But in these cases I had reason to think that the chief part of the fluid consisted of water. It is probable that this was the case in the two instances of apparent absence of spermatozoa in the Toad, mentioned by SPALLANZANI,* and that the fluid did really contain spermatozoa, although few in number, and consequently easily overlooked, and that the ova were impregnated by these, and not by the fluid portion of the semen, as he appears to have supposed.

On comparing the white surface of the yelk of the unimpregnated with that of the impregnated egg, whether the egg had been fecundated naturally or artificially, I was not able to detect any difference during the first twelve minutes. The changes went on in both, and appeared to be almost identical in each. But after the time specified no further progress was perceptible in the unimpregnated ovum, which continued to exhibit the same appearance for several hours. But the white surface of the impregnated egg became more and more changed, up to the time of cleavage of the yelk, when it was almost an uniform surface.

These observations were afterwards repeated with similar results, and the conclusion to which they led was, that changes take place in the yelk from the period when the germinal vesicle disappears and the ovum leaves the ovary to the moment of its expulsion from the body, and which changes may proceed for some time afterwards quite independent of impregnation; and

* Dissertations relative to the Natural History of Animals and Vegetables, 1789, vol. ii., p. 151.

that these have some reference to the evolution of the central or embryo vesicle: possibly also that they do not cease immediately, but subside gradually when the stimulus imparted by impregnation is not supplied.

The experiments made to ascertain whether the unimpregnated ovum passes through any stage of cleavage, consisted of four sets, placed in four vessels of equal size, containing each about two ounces of water. Ova were passed from the same female as quickly as possible at the same time into each of these vessels. To one of these marked A, a considerable quantity of a mixture of seminal fluid, one part to three parts of water, was immediately added; to a second, B, only a single drop of this mixture; while the third and fourth, C and D, contained only water with the unimpregnated ova, and the four vessels were then placed in every other respect under precisely similar conditions.

A few minutes after the impregnating fluid had been added to A, I examined some of the ova beneath the microscope, and found a vast abundance of spermatozoa adhering to every part of the surface of their gelatinous envelopes. On other ova from the set B, there were also many spermatozoa attached, but in much smaller number than on the ova of set A.

In *five hours and fifteen minutes*, the temperature of the room during the interval having ranged only from 53° FAHR. to 54° FAHR., segmentation had commenced vigorously, and was strongly marked in the whole of set A. But it had not commenced in set B. It did not occur in these until *five hours and twenty-two minutes*, when it began in these also. Thus there were *seven minutes'* difference in the commencement of the changes in these two sets of ova, a circumstance which led to the belief that this difference might have some reference to the relative quantities of the impregnating fluid employed,—an opinion which I had long before been led to by observations on the impregnation of the common Earwig, *Forficula*, in which it had appeared to me that deficiency in the quantity of the impregnating fluid is unfavourable to fecundation.

No segmentation or cleavage of the yolk took place in the sets of ova marked C and D, which, except in becoming a little oval, as already mentioned of unimpregnated eggs, remained, in so far as the appearance of the yolk surface was concerned, in the same state as at a few minutes after spawning, and they continued in exactly the same condition at the end of twenty-two hours. This I have since found constantly to be the case with unimpregnated ova, whether they happen to be exposed to a high or low temperature of the surrounding medium. The yolk of the impregnated egg gradually acquires a more intense

black colour, which strikingly contrasts with the dull colour of the unimpregnated.

At the end of *six days* the majority of the ova in A and B were producing embryos, while those in C and D were fast decomposing.

These trials afforded the positive test I required from direct observation, as a fixed point in the experiments about to be commenced,—that segmentation certainly does not take place in the unimpregnated ovum.

3. SUSCEPTIBILITY OF THE OVUM.

I was not aware, at the time of commencing my experiments in March 1849, nor indeed until very recently, of the extent to which the original investigations of SPALLANZANI, and of PREVOST and DUMAS had long ago been carried,* and it has only been since my experiments were completed, and during the preparation of this paper for presentation to the Royal Society, I have learned by careful reference to their first memoirs, that they have anticipated me in part of this inquiry; that of endeavouring to separate the spermatozoa by filtration from the more fluid portion of semen, and testing the effect of these two constituents in artificial impregnation. To them therefore be all honour for the result; although even they, as they honourably mention, had themselves been anticipated in this by SPALLANZANI, and that too with similar success. The extraordinary results obtained by SPALLANZANI† in artificial impregnation, and the imperfect knowledge which we possess of the nature of the means by which it is effected, has induced me to endeavour to repeat and vary his experiment, and to conceive others, which, so far as I am aware, have not yet been attempted. I have been the more urged to this from the circumstance mentioned by SPALLANZANI, and already alluded to (p. 373), the occasional supposed absence of spermatozoa from fluid that is capable of fecundating; and also from a belief formed long ago with regard to the Articulata, that the spermatozoa, nevertheless, certainly are the efficient agents in impregnation, although full proof of the fact has been wanted. I have been desirous therefore of learning how far this belief can bear the test of direct experiment, or the fact be capable of demonstration by artificial means in the Amphibia. As, however, the experiments I have myself made vary from those of the authors mentioned,—have not been influenced by the result they had previously arrived at,—have been somewhat more extended, and, as I believe, will now tend

* Annales des Sciences Naturelles, Tom. i. et ii. 1824.

† Dissertations, &c. vol. ii.

to place the fact of the direct agency of the spermatozoa in impregnating the ovum beyond doubt,—it has seemed desirable still to give them in detail, as assisting to establish an important point of knowledge by facilitating a comparison of the results of independent investigations.

Duration of susceptibility.—The length of time during which the ovum, after it has been passed, remains susceptible of fecundation, is affected by several circumstances. I had reason to believe at the commencement of my experiment that this time is very short. SPALLANZANI found that when the egg of the toad was expelled into water it was not susceptible of fecundation after a lapse of fifteen minutes.* This was at a raised temperature of the atmosphere, 81°·5 FAHR. On the other hand, he also found that, at this temperature, ova retained fourteen hours within the body of the female after death, and of course not in contact with air or water, might still be fecundated; and that when preserved in an ice-house fecundation might be effected at two days after the death of the parent.† But PREVOST and DUMAS arrived at the conclusion‡ that the time is much more extended in the Frog;—that fecundation may take place when the temperature ranges from 53° FAHR. to 59° FAHR. at the expiration of one hour after immersion of the eggs;—that some eggs are fecundated at two, and a very few even at the end of three hours, after which no fecundation takes place. The results obtained by myself both on the Frog and Toad have been most in accordance with those by SPALLANZANI. This is the more worthy of notice from the circumstance that a slight difference between his and mine is readily accounted for by a corresponding slight difference of temperature, which SPALLANZANI, and PREVOST and DUMAS, have remarked, and since them also Mr BELL,§ has great influence on the changes of the eggs and young. The temperature at which my test experiments were made, was a little lower even than that at which PREVOST and DUMAS made theirs; and yet I was not able to find any ova susceptible of fecundation after they had remained from thirty to forty minutes in water.

On careful examination of PREVOST and DUMAS' experiments, I think the difference may perhaps be due to a circumstance which seems equally to affect some of SPALLANZANI's results, namely, the mode in which the impregnating fluid employed was obtained. These authors state that the fluid they employed was expressed from the testicles of the frogs, so that from what we now know of the mode of origin of the spermatozoa, this fluid in all pro-

* *Loc. cit.* vol. ii. p. 178.

† *An. des Sc. Nat.* Vol. ii. p. 135.

‡ *Ibid.* pp. 176 and 177.

§ *British Reptiles*, p. 92.

bility contained a large proportion of developmental cells that included spermatozoa not fully matured, but which might become liberated in the water at a longer or shorter period. Or, possibly, the fluid added to ova that had been long in the water, had been very recently obtained; in which case the vigorous spermatozoa might effect the impregnation of ova that had become almost insusceptible through the imbibition of water by their envelopes. I am led to this view by the fact that the jelly-like envelope of the Frog's egg begins to imbibe and expand the instant it is brought into contact with fluid; and from having ascertained that there is a close relation between the degree of expansion and imbibition of this envelope and the susceptibility of the ovum to become impregnated, and that these conditions are also greatly affected by temperature. The act of expansion of the envelope is an act of endosmose, and possibly this is one of the means by which the impregnating agent is made to exert its influence on the yelk. The yelk is not a passive recipient during the endosmic action of its coverings, but seems to participate in that action, as I have seen portions of its surface heave and contract within the vitelline membrane during the first hour the egg has remained in water. It may thence be inferred, that if the impregnating stimulus be not supplied quickly, the fitness of the ovum to become impregnated is diminished in proportion as its envelopes are expanded. If then it be proved that the spermatozoon is the agent in impregnation, but, so far as can be discovered, does not penetrate bodily into the ovum or its envelopes, and yet, as may be shown, must always come into contact with their surface, the more rapidly and to the greater extent this expansion takes place, and removes the efficient body from that which it is in some way destined to affect, the less will be the chance of its impregnating the ovum, and the less will the ovum become susceptible of impregnation even by the most healthy and vibratile spermatozoa.

The extent and rate of expansion of the envelope of the frog's egg, during the first half hour it remains in water, very nearly coincide with the diminution of the fitness of the ovum to become fecundated. This is shown by observing the rate of expansion of the envelope during the first fifteen minutes of submersion, and then testing the fitness of the ovum, by experiment, during a similar period.

At the *moment* when the ovum is expelled from the body, the envelope is merely a thin gelatinous layer, its entire diameter being equal only to about one sixth of the diameter of the yelk. After it has been *one minute* in water, and begun to imbibe and

expand, it is then equal to about one-fourth of the diameter of the yelk. At the end of *two minutes* it is enlarged to one-third, and in *three minutes* to one-half the diameter of this body. In *four minutes* it exceeds three fifths, and in *six minutes* two-thirds, and it continues to imbibe fluid and expand at the same rate, until, at from *ten to fifteen minutes*, it very nearly equals in thickness the whole diameter of the yelk; and at *half an hour* it is one-fourth greater than this. PREVOST and DUMAS* noticed the expansion of the envelope during the first *six hours*, but entirely overlooked the rate of expansion during the most important period, the first hour, and noticed only the general fact that the diameter of the envelope, and the end of the first hour and a half, was as 5 to 2·5 at the time of spawning, and that it had nearly acquired its full size at the end of three hours. My own observations agree with this latter statement. The expansion of the envelope is greatly retarded at the end of the third or fourth hour, until after cleavage of the yelk has taken place, when it again proceeds, but much more slowly than at first. If then we bear in mind the rate of expansion of the envelope during the first half-hour, the following experiments will give some idea of the degree of susceptibility of the ovum to become impregnated during that period.

Mr Newport subsequently, on the 6th of April 1850, performed a series of six experiments, in which, after immersing in water, *ova* varying in number from eighty-three (83) to two hundred and five (205), and added impregnating seminal fluid, so as completely to cover the ova; and two experiments in which eighty-one ova, and one hundred and thirty nine ova (139), were exposed to dry air, and then subjected to the operation of seminal fluid applied in an effectual manner.

Of this series of experiments the following tabular summary indicates the results:—

• TABLE I.—Set F.

Experiment.	Ova.	Time.	Medium.	Fluid obtained.	Segmentation.	Embryos.	Per centage.
		"		"	" "		
No. 1	83	1	Water.	43	3 55	32	·38
No. 2	92	2	Water.	43	3 55	45	·49
No. 3	127	3	Water.	17	4 3	33	·26
No. 4	81	3	Air.	8	4 5	53	·65·5
No. 5	136	5	Water.	20	4 8	10	·07
No. 6	139	5	Air.	12	4 11	37	·26
No. 7	205	15	Water.	26	4 14	45	·22
No. 8	100	30	Water.	44		2	·02

Thus, then, at a temperature of 60° Fahr., the susceptibility of the ovum to become impregnated is greatest at the time it is passed into water, and for two or three minutes afterwards, and segmentation then takes place more quickly, even when the seminal fluid has been for nearly three quarters of an hour mixed with water, than after longer immersion. The fitness of the ovum to become impregnated is gradually diminished, and segmentation takes place more tardily, according to the length of time which the ovum has remained in water, as is seen by comparing the results of Nos. 1 and 2 with 7 and 8. On the other hand, while the desiccating effect of exposure to air more arrests the fecundation of the ovum and the occurrence of segmentation of the yelk than a continuance for a corresponding length of time in water, it seems to be less prejudicial to the fecundity of the ovum than immersion in that fluid, as appears to be shown by comparison of Nos. 4 and 6 with 3 and 5, the difference in the number of ova produced being too great to lead us to attribute this to difference in the length of time the impregnating fluid had been obtained.

In the foregoing set of experiments, the quantity of impregnating fluid supplied to the ova was but little attended to, it being added very freely in each case. In the following set I was desirous of knowing what difference would result from the fluid being applied more sparingly, or but for a very short space of time. SPALLANZANI had made experiments with a similar view, but his appeared to be open to some objections, as he had not noted some important circumstances which greatly affect the result, as the temperature of the medium, the length of time the fluid employed had been obtained, &c. In the experiments now made, these circumstances were attended to, and I noticed a curious fact which I first remarked in experiments in 1849. It is what I may designate *partial impregnation*, and is indicated by a portion only of the yelk becoming segmented. This frequently happens with ova that have been brought into contact with only very small quantities of seminal fluid, and but for short spaces of time, as in some of the following experiments. These ova, so far as I have observed, never produce embryos. Segmentation is arrested in some at the very commencement; in others it goes on to the second or crucial fissure; and in a very few cases may proceed somewhat further; but it is never completed to granulation of the yelk. This, I think, is a fact which deserves some consideration with reference to the formation of the embryo.

[Another set of six experiments (F) had been previously instituted on the 22d of March 1850, in which ova, varying in

VOL. L XXVII. NO. 191. E e

number from forty-two to seventy-seven, were passed into water at 46°. 5, F., in one exposed merely to air, and in two experiments passed into a solution of carmine for five minutes and fifteen minutes respectively, and then impregnating seminal fluid, was applied to them in various modes.]

From these experiments, which were performed, in order to ascertain the influence of temperature, it seemed evident that the susceptibility of the ovum to become impregnated is diminished in proportion to the degree of expansion of its envelope and its imbibition of fluid, conditions which are greatly affected by the temperature of the medium in which the ovum is placed during the first hour; and there seems reason to suppose, that this diminution may be due to the extent to which the envelope becomes influenced by temperature, rather than to any insusceptibility at that time in the yolk itself.

[From these experiments, and other three performed on a female frog in which the oporal chord had been divided four hours and a half previously, the following inferences are deduced]

These facts proved that the ova employed were still fitted to become impregnated when the fluid was supplied to them in sufficient abundance, and for a sufficient length of time, and within the period during which the envelope continues to expand and imbibe most rapidly. This condition is always promoted by an early removal from a low to a comparatively high temperature during the period of expansion, as in *Set G*, in which segmentation took place in from *five hours and a half to six hours*, and when the removal from low to high temperature was within the first half hour; while it did not occur, in the only experiment in which it happened in *Set F. No. 1*, until the end of *six hours and a-half*, when the removal from a similar low to a like high temperature was not made until one hour and a-half after impregnation.

The influence of temperature is thus as marked on the impregnation of the ovum as it can be proved to be on the future development of the embryo. Impregnation is accelerated, and its occurrence also is more certain in a high than in a low temperature. In the latter it becomes retarded and is less determined. This applies equally to the susceptibility of the ovum, and to the fitness of the impregnating fluid to effect impregnation. But in proportion as this fitness is exalted by increase of temperature, so is the duration of the capability to receive in the one, and the efficiency to communicate in the other diminished. SPALLANZANI found that the ova of toads placed in an ice-house could be impregnated at the end of forty-one hours.* PREVOST and DUMAS† also mention that they were

* Dissertations, &c., Vol. ii. p. 177.

† *Loc. cit.* Vol. ii. p. 140.

successful, and that too to a great extent, with ova that had been twenty-four hours *in water*, the temperature during the period ranging from 18° Cent. (64° 4 FAHR.) to 22° Cent. (71° 6 FAHR.), and with some eggs that had not been immersed even at thirty-six hours, * the temperature being then from 12° Cent. to 15° Cent. (53° 6 to 59 FAHR.). The results obtained by myself have been much less successful. Out of one hundred and forty ova obtained from a female frog, killed twenty-four hours before and preserved at or below the temperature of 55° 5 FAHR., at which the experiment was made, only a very few became partially segmented, but not one produced an embryo; although an abundance of impregnating fluid, abounding with spermatozoa, and obtained only a few minutes before it was employed, had been supplied to them. It is evident, therefore, that this failure was due chiefly to the ova, and not to inefficiency of the impregnating fluid. On the other hand, I have been equally unsuccessful with ova from a frog that had been killed only two hours and a-half, when the impregnating fluid employed had been more than four hours and a-half mixed with water. In this case the failure appeared to have been due chiefly to the spermatozoa, nearly the whole of which, on inspection by the microscope, were found to be motionless and appeared to have lost their vitality. At the same time it must be mentioned, that the female from which the ova employed in No. 3 of the last set of experiments were obtained, still existed, in so far as the vitality of the muscular system was concerned, and therefore can hardly be mentioned in comparison with MM. PREVOST and DUMAS's observation.

But while the numerical results obtained by myself have been less favourable than those of SPALLANZANI or the physiologists now mentioned, the general facts, so far as they are open to comparison, are in full accordance with them. The difference in the details of our respective observations appears to have been due in chief part to the influence of temperature at the time of the impregnation of the ova, or within the first two or three hours after the impregnating fluid has been supplied. Thus, if the temperature has been gradually rising at the time of impregnation, the fecundation of the ovum, as I have stated, has more certainly taken place than when the temperature was subsiding, the condition of the ova and of the impregnating fluid employed being equally fit in each case. SPALLANZANI has shown that in his experiments ova did not become impregnated after they had remained fifteen minutes in water. In the experiments by myself, I could rarely obtain fecundation after thirty minutes' immersion. The difference of time between these results may

* *Loc cit.* p. 134.

fairly be attributed to difference in the temperature at which the experiments were made, and in great measure to the influence of this on the endosmosis and expansion of the envelopes.

But it was possible that some other agent might be concerned in these results, and that light, as well as heat and immersion in water, might greatly influence them. To put this to the test, and to learn whether the difference depends entirely or chiefly on the amount of temperature, I have made two sets of experiments at precisely the same time, performed in the same way, with ova from the same female, and impregnating fluid from the same male, the only difference being that within a very few minutes after the impregnating fluid was supplied, one set was removed to a higher and slightly rising temperature, from which all light was excluded, while the other was allowed to remain freely exposed to light, but in a room of ten or twelve degrees lower temperature, and which was becoming still further reduced.

The influence of light and heat on the development of the embryo has already been referred to by SPALLANZANI, PREVOST and DUMAS, RUSCONI, Dr W. EDWARDS, and Mr BELL. RUSCONI expressly states that light has no influence on the development of the germ,* but his observations, as well as those before made by SPALLANZANI, show that heat has a very marked influence, and this has been fully confirmed by Dr W. EDWARDS, and Professor BELL. Very recently also the subject has been referred to by Mr HIGGINBOTTOM,† and I have great pleasure in stating that my own observations on the influence of heat, and the little effect of light on the development of the tadpole, are in accordance with the observations made by him. But the object I have had most in view has been, as above stated, to mark the effect of heat, without light, on the changes of the ovum, more especially during the period of fecundation, the first three or four hours after the egg is laid, and onwards to the termination of what I shall hereafter propose to consider when describing the development of the embryo,—as the end of the *third period*—the closure of the *laminæ dorsales* and the establishment of ciliary aëration on the surface of the body.

[From two sets of six experiments of the same nature performed on the 20th March 1850, the conclusions deduced are the following.]

Thus while segmentation took place in No. 1 H, in three hours and fifty-six minutes, when the temperature was rising

* *Loc. cit.* p. 20.

† Proceedings of the Royal Society, May 16, 1850, and Phil. Trans., Part ii., 1850. See also Ed. Med. Journ., lxxvi., 493, Oct. 1851.

from 59° FAHR. to 64° FAHR., it did not occur in No. 1 I, until seven hours and forty-five minutes, when the temperature during the interval was sinking from 48° FAHR. to 47° FAHR. This sufficiently marks the great influence of temperature during the earliest periods of change in the ovum; and this injurious effect of reduction of temperature at that period is further shown in the relative number of embryos in these comparative experiments. That the injurious effect of reduced temperature at the time of impregnation is mainly the cause of this result, and not the diminution of temperature after the period of impregnation, seems to be shown in the circumstance, that while at the end of the eighteenth hour the ova in the set H had already passed through all the stages of segmentation, and the surface of the yelk had become granulated, and the blastoderma had begun to be formed even although the temperature in that case subsided a little after segmentation had commenced,—from 59° to 57°,—the corresponding set of ova, No. 1 I, had advanced only to the octuple division of the yelk. A similar difference in the rate of development we have seen takes place in the growth of the embryo. At the end of three days the embryos of set H were advanced to the stage at which the laminae dorsales are proceeding rapidly to meet, and form the median dorsal sulcus of the growing body. But the corresponding ova in set I, had not been carried further than to the earliest perceptible indications of the *area germinativa*.

These facts sufficiently prove the great influence of temperature on the development of the embryo in its earliest stages, as the comparative numerical results do also its effects on the impregnation of the ovum. Out of eighteen ova placed in the higher and increasing temperature, thirteen produced embryos at nearly similar stages of growth; while of nineteen ova maintained in a low and diminishing temperature only eight became segmented, and but three of them arrived at the tadpole state.

A somewhat similar but more marked result took place with the ova of No. 2 in the two sets of experiments. The impregnating means employed in these trials had already been forty minutes mixed with water on the filter. Out of forty-eight ova employed in set H, twenty-five became segmented, and ten of these produced young. But in set I fifty-one ova gave birth to only two embryos.

In the third experiment of each set the difference is as strongly marked. As the filtered fluid employed in both was the same, and the very few spermatozoa contained in it were by the same means brought into contact with the ova in each, it might have been expected that each would have produced embryos. But while the production of a single tadpole in the

one case, at a high temperature, may be looked upon as leading to the inference that these bodies are the efficient agents in impregnation, the entire absence of all appearance of impregnation in the ova of the other set, to which the same fluid had been equally applied, seems to point to the cause of failure in this case as depending on the prejudicial effect of a low temperature of the surrounding medium on their agency.

The temperature of the surrounding medium ought, therefore, always to be borne in mind when we are attempting to deduce conclusions from experiment on impregnation and development. The presence of light appears to be only of secondary consideration as compared with heat; since in set H, from which light was carefully excluded, not only did impregnation take place more certainly and rapidly than in set I, which were exposed to light, but the embryos also were produced in greater number, and acquired maturity in less than one-half the space of time than in the latter; the only difference of circumstance between the two sets being degree of temperature.

Influence of Aëration.—Next in importance to heat is a free aëration of the ovum. This is of less consequence with reference to impregnation than to the subsequent production of the embryo. In every set of experiments there are always some ova more advanced than others. These are ova which have been nearest to the surface of the water, and which, consequently, have been more completely aërated as well as exposed to a slightly higher temperature than others at a greater depth. It is from this cause chiefly that the results of experiments on artificial impregnation, and even of observations on naturally impregnated ova, are always less complete and successful than what takes place with regard to the ova in the natural haunts of the species. The ova in a state of nature are usually deposited in well-aërated places, clear, slow-moving water, or shallow and but slightly turbid water. It is almost impossible to afford to ova that are the subjects of experiment, either in broad flat dishes, or in glass vessels in one's study, the amount of aëration required to ensure complete success. By too frequently changing the water in the vessels the embryos often become injured; while if the water be not changed, development is arrested, and decomposition commences, and the experiment entirely fails. Even when these difficulties are obviated by a gentle withdrawal of the water, and a renewal of it with equal care, the perfect stillness of the fluid in the interval of our observations does not allow of that extent of aëration to the embryo which it gains in a perfectly natural state, either in slow-moving waters, where I have usually found the eggs deposited, or in pools of still

water, the surface of which is agitated by currents of air, and affected by diurnal changes of temperature.

Thus, then, we may conclude that the procreative force of the germ, and of the impregnating fluid, is augmented by increase of heat, but the duration of the force is lessened. It becomes less and less energetic in proportion as the temperature is diminished, but the period during which it is capable of being exerted is extended. In each of these conditions aëration is of essential consequence, and becomes more and more necessary in proportion to the increase of heat.

(To be concluded in next Number.)

ART. VII.—*Account of a Case in which the Cæsarean Section was performed; with Remarks on the Peculiar Sources of Danger attendant on the Operation.* By CHARLES WEST, M.D., Physician Accoucheur to St Bartholomew's Hospital, and Lecturer on Midwifery in the Medical College.—(Received December 20th, 1850.—Read January 28th, 1851.) (From the Med. Chir. Trans., Vol. xxxiv. 1851.)

ELIZABETH WILLIAMS was one of eight children, of whom seven reached adult age. Her father, who had been insane for some years previously, died at the age of 73, her mother at that of 68. Two of her brothers and one sister died of phthisis, and this sister was insane for some years before her death. Two brothers and a sister survive in good bodily health, but the sister is of weak intellect: and Elizabeth herself had, on more than one occasion, shown indications of insanity.

As a child, E. Williams' health had been good, but it became much impaired about the age of 15, when she began to menstruate; and ever since then she had frequently, indeed almost constantly, suffered from pain in the back, referred especially to the pelvis. About the age of 23 the pain became much severer than it had ever been before; but, in spite of this, and of the advice which she received from some medical man not to marry, she married, at the age of 25, the husband of her deceased sister. She did not become pregnant until fifteen months after marriage; but conception was followed by a very considerable increase of pain, associated, during the latter half of her pregnancy, with a great and rapidly increasing difficulty in walking. For some weeks before her confinement she did not move further than from her bedroom on the second floor to her sitting-room on the first floor, where she generally spent the whole day lying or sitting upon her sofa. Sometimes, however, she moved about her room, and on the day before her confinement, was occupied in putting up curtains to the windows; her