ART. I.—*The Blood Corpuscle considered in its different Phases of development in the Animal Series. Memoir I. Vertebratu.*

Received 27th February, read 19th June 1845. *Memoir II.*

Invertebrata. Received 8th May, read 19th June 1845. *Memoir III.*

Comparison between the Blood Corpuscle of the Vertebrata and that of the Invertebrata. Received 7th May,

read 19th June 1845. By T. WHARTON JONES, F. R. S., Lecturer on Anatomy, Physiology, and Pathology at the Charing-Cross Hospital, &c. Philosophical Transactions of the Royal Society of London for the year 1846, Part II. London, 1846. 4to.

IN these papers the author examines very carefully the blood corpuscle in individuals of the different families of the Vertebrated and the Aspondylous animals.

Beginning with FISHES he takes the skate as the representative. The blood of this animal presents two kinds of *corpuscula*, which at first view under the microscope particularly attract the attention of the observer. The first kind is without colour, but appears dark in consequence of refracting the light strongly, is roundish in form, and about $\frac{1}{300}$ th part of one inch in diameter, or even larger. The other kinds, which are more numerous, are of a red colour and oval form, and measure about $\frac{1}{600}$ th part of one inch long, by $\frac{1}{700}$ th part of one inch broad, (Figs. 1, 2, 3, 7.)

The first-mentioned kind of corpuscle is composed of an agglomeration of granules surrounded by a cell-membrane. The granules, which are clear and strongly refract the light, measure about $\frac{1}{2000}$ th part of one inch in diameter; but they may appear at one time separate, and at another more fused together, like granules of quicksilver. This corpuscle Mr Jones designates by the name of *granule blood-cell*. Varieties of these from different animals are represented in the first compartment of the first column of Plate III.

On one side of this granule blood-cell may be seen a clear spot indicating the place of a nucleus. (Fig. 1.)

By the action of water the cell-wall becomes distended, so that it is enlarged in diameter. At the same time the aggregation of the granules becomes less close, and the appearance of fusion is diminished. Mr Jones sometimes saw the granules suddenly collapse, and flatten under the action of water, as if they were small vesicles bursting. (Fig. 2.)

By the action of diluted acetic acid the granulous mass is dis-

solved, leaving the cell-wall, in the interior of which is then distinctly seen a cellæform nucleus, about $\frac{1}{450}$ th part of one inch in diameter, (Fig. 2b.) He observed individual cells under the action of acid, and saw granule disappearing after granule, until the nucleus came into view.

The roundish or spherical shape of the granule blood cell belongs to it after the death of the animal. In some examples of the blood examined not long after death, the granules cells presented, at first, remarkable changes in shape, and other phenomena immediately to be noticed, which after a time were no longer visible, the cells settling into the round form. These changes and other phenomena, he thinks, will be found in the blood of the living skate.

"My attention," says Mr Jones, "was first attracted to the phenomena, by observing a granule cell with the granule apparently escaping from it, as if burst. (Fig. 3.) But the cell soon appearing again with all the granules collected, I was led to watch, and soon perceived that the appearance of granules escaping, as if from a burst cell, was owing to the transparent and colourless cell-wall bulging out on one side, leaving the granules agglomerated and adhering, though only for a short time; for some single granules were seen to separate and burst from the rest, and to enter the hitherto empty compartment produced by the bulging of the cell-wall. This took place in a manner very regular. I have seen the granules enter the compartment by one side, and circulate along the bulging cell-wall to the other side, until the whole compartment became filled with granules. This being finished, the bulging subsided, but was succeeded by bulging of another part of the cell-wall, into which a similar flow of granules took place; and so on all round the cell."

The red oval corpuscle is an oval shaped cell with a *nucleus* round or oval, and with red coloured contents. Hence it resembles the red corpuscle of the blood of the frog in structure as well as in appearance; the only difference being that its *nucleus* is cellæform, whereas that of the red corpuscle of the frog is rather solid-looking. The red oval corpuscle of the blood of the skate Mr Jones names the *nucleated blood-cell*, in contradistinction to the granule cell, which, though also nucleated, is not in general discovered to be so, until after solution of the granules by acetic acid, and which, in consequence of being filled with granules, is well designated granule blood-cell.* (Figs. 4, 5, 6, 7.)

Besides the two kinds of cells now described in the blood of the skate, there are others which, on account of their paleness, do not at first attract attention, but which it is of great importance to notice.

* This term the author adopts from Vogel of Göttingen, who employed it to designate a form of cell developed in inflammatory exudation.

One kind (Fig. 4.) have the characters of the granule-cell, excepting that the granular contents are very fine, and do not so strongly reflect the light, rendering them pale. This pale granule-cell the author considers an advanced stage of the dark-looking granule-cell; the latter he calls the *coarsely granular* stage, the former the *fine granular* stage of granule-cell.

Another of the corpuscles referred to have essentially the same structure as the red nucleated blood-cell above described, being a cell containing in its interior a cellæform nucleus, (Fig. 5.) The only difference is, that both cell and nucleus are circular, (the former about $\frac{1}{80}$ th of one inch in diameter, the latter about $\frac{1}{100}$ th,) and that the cell is either colourless or slightly red.

That this circular nucleated cell is an early stage of the oval red nucleated cell is shown by the circumstance, that besides it, there are cells having the same structure, but more or less oval shaped, and either altogether colourless, or slightly tinged red, (Fig. 6,) which evidently form a transition to the oval red nucleated blood-cell.

Two stages of the nucleated blood-cell may therefore be recognized, an *uncoloured* and a *coloured* stage.

The description here given of the nucleated blood-cell of the skate in the uncoloured stage, and comparison of the figures 5 *a*, 2 *b*, show that it resembles the granule blood-cell, after its granules have been dissolved by the action of acetic acid; the only difference being, that in the former the cellæform nucleus is larger and more developed.

This resemblance, and the circumstance that corpuscles occur in which the cellæform nucleus is large, though surrounded by a few granules, (fig. 4 *b*, and fig. 2, lamprey,) and which may also be viewed as a transition from the granule to the nucleated cell;—prove that the nucleated blood-cells are formed from the granule blood-cell, and are, in truth, the granule-blood-cells with the cellæform nucleus, developed at the expense of the granules which have disappeared.

From the facts now stated the author concludes that the relationship between the granule blood-cell and the nucleated blood-cell is, that they are to be viewed as two different *phases* of development of the same body; the granule-cell being the *first phase of development*, and the nucleated cell the *second phase of development*. Each phase of development further comprehends two stages;—the phase of granule-cell, a coarse and a fine granular stage; the phase of nucleated cell an uncoloured and a coloured stage.

In the osseous fishes examined by Mr Jones, the blood-corpuscle, in its phases and stages of development, is essentially the same as in the skate. In the lamprey, a cartilaginous fish, lower

in the series than the skate, the blood-corpuscle, in its phases and stages of development, is the same, with the exception that the nucleated cell in its coloured stage is circular, as well as that in the uncoloured stage (Fig. 4 a, 3).

In another genus, the lowest of the lowest division, *Branchiostoma lubricum* of Costa, (*Amphioloxus lanceolatus*, Yarrell,) in which the blood is quite colourless, the author thinks, that though the blood-corpuscles of this animal have not been examined, yet it will turn out, that some of them are the same as the granule-blood-cells of the skate and lamprey, and others the same as the nucleated blood-cells in their colourless stage. If this be the case, the blood of the *Branchiostoma lubricum* differs from that of other fishes, merely in the absence of the nucleated cells in the coloured stage. He allows, nevertheless, that it is possible that nucleated cells in the commencement of the coloured stage may exist.

REPTILES.—The *Frog*. In the blood of the frog, as in that of the skate, the following kinds of *corpuscula* are to be recognized; granule-cells and nucleated cells, the former in coarse and fine granular stages, the latter in colourless and coloured stages.

The objects usually spoken of as lymph and colourless corpuscles are the granule-cell in both its stages, and the nucleated cell in its colourless stage. They exist in the blood, and must be considered as blood-corpuscles.

The corpuscle in the blood of the frog, identical with the granule blood-cell of the skate, in the coarsely granular stage, is round, and contains clear granules strongly refracting the light, like those of the same cell in the blood of the skate, but smaller; (Fig. 1). These granules are usually collected in greatest number at one side. The cell itself Mr Jones has seen exhibiting changes in shape with movements of the granules, such as were described in regard to the granule-cell of the skate.

By adding water this granule blood-cell becomes distended; and if diluted acetic acid, or rather water acidulated with acetic acid, be added, the granules are dissolved, and a cellæform nucleus about $\frac{1}{3500}$ th of one inch in diameter comes into view (Fig. 1.) If stronger acetic acid be added, instead of one nucleus, the appearance of two or more smaller *nuclei*, lying close together, may be presented.

The granular blood-cell in the finely-granular stage of the frog, when distended by the action of water, presents a regular circular form, with a diameter usually about $\frac{1}{1200}$ th or $\frac{1}{1500}$ th of one inch, sometimes more, sometimes less (Fig. 2 b.) It contains an indistinctly granular matter, not much refracting the light, and therefore pale.

Before being acted on by water, the granule blood-cell usually presents an irregular shape, with processes shooting from it (Fig.

2 a.) After the blood is drawn it speedily undergoes the change to this shape, which is analogous to the change of shape which the coarsely-granular blood-cell presents. After distension by the action of water, a cellæform nucleus is sometimes already visible in the interior, through the finely-granulated contents (Fig. 2 b.); but by the action of acetic acid, one is always distinctly brought into view. The diameter of this cellæform nucleus, in the state of distension in which it is seen, may be stated at $\frac{1}{2200}$ th part of one inch.

Instead of one cellæform nucleus of this size, an appearance of two, three, or even four smaller ones may be brought out (Fig. 2 c.); and this even more readily than in the case of the coarsely-granular cell. This, however, the author regards as an appearance produced by shrivelling of the walls of the single cellæform nucleus, by the action of the acetic acid.

In short, when the blood is treated with water first, so as to distend fully the granule-cells, and then very dilute acetic acid added, never more than one single cellæform nucleus comes into view. If the acid be added a little less diluted, and before the distending action of the water exerts its full effect on the cell, generally an appearance of two or more small *cellæform nuclei* come into view. Though these nuclei appear sometimes detached, they are usually in contact.

Nucleated blood-cells; uncoloured stage. In the blood of the frog, besides granule-cells, are seen uncoloured but distinct and well-defined *corpuscula*, round or oval, finely-granulated, refracting the light, and about $\frac{1}{3200}$ th part of one inch in diameter.

These corpuscles are apt to be confounded with the granule-cells, and appear sometimes to have been taken for free *nuclei*. They are *nuclei*, but not free, being surrounded by a cell-wall, which, without a good microscope and good light, is readily overlooked, on account of its paleness. This pale cell, with distinct nucleus, which is the nucleated blood-cell in the uncoloured stage, may be circular, (Fig. 4, or oval, Fig. 5,) and of a size approaching that of the red corpuscle or nucleated cell in the coloured stage.

The pale cell-wall may embrace the nucleus so closely as not to be distinguishable at all, until distended by the addition of water, when it appears as represented in Fig. 3.

Nucleated blood-cell; coloured stage. This is the red, oval corpuscle of the blood of the frog (Fig. 6.) It is the counterpart of the red oval corpuscle or nucleated blood-cell in the coloured stage of the skate. The principal difference between the cells in question in the two genera consists in the character and appearance of the nucleus. In the red cell of the skate, the nucleus, though more or less oval, retains its cellæform appearance. In that of the frog, it is still more solid-looking and condensed than

it was in the nucleated cell in the uncoloured stage, and at last acquires the appearance as if consolidated by fusion into a small oval-shaped homogeneous mass.

The development of the red oval corpuscle, or nucleated cell in the coloured stage, from the nucleated cell in the uncoloured stage, may be readily understood to take place by the latter secreting into its interior colouring matter, becoming at the same time enlarged, in the manner already explained as to the red nucleated cell of the skate.

The development of the nucleated cell in the uncoloured stage from the granule-cell appears to take place in the following manner. The cellæform nucleus of the granule-cell becomes developed into the solid-looking nucleus at the expense of the granules which disappear. As this takes place, the cell-wall contracts around the nucleus, becoming thicker and stronger, and only becomes distended again by the accumulation of contents, as already described in the instance of the skate.

MAMMIFERA.—MAN AND MAMMIFEROUS ANIMALS.—In man and the mammalia, though the process is more complex, it follows the same general type of formation.

Mr Jones examines first the corpuscles of the early embryo of a mammiferous animal, comparing them with those of the oviparous Vertebrata; and then the corpuscles of the fully-formed blood of man and the mammalia, comparing them with those of the embryonic blood of the same class, and with those of the blood of the oviparous Vertebrata.

Blood corpuscles of the embryo ox about $1\frac{1}{4}$ of one inch long, and comparison of them with those of the oviparous vertebrate animals.

“44. The blood of this embryo presented the following kinds of corpuscles :—

“1. Granule-cells in coarsely granular and finely granular stages.

“2. Nucleated cells in uncoloured and coloured stages.

“3. Small red corpuscles similar to those of the fully-formed blood of the animal.

“45. *Granule-cells*.—Some of these contained pretty large and well-defined granules and refracted the light strongly,—coarsely granular stage (fig. 1.) ; others presented less defined granules, and refracting the light less, appeared very delicate and pale,—finely granular stage. Fig. 3 represents one of these cells with the cellæform nucleus already visible through the finely granular contents ; this may be viewed as a transition to the phase of nucleated cell, whilst fig. 2 represents a granule-cell intermediate between the coarsely and finely granular stages.

“46. The size of these granule-cells was from about $\frac{1}{1460}$ th to about $\frac{1}{1823}$ th of an inch in diameter.

"47.—*Nucleated cells—Uncoloured stage.*—In these, as in the corresponding cells of the blood of the frog, the nucleus, quite cellæform, however, was very distinct, presenting a contrast with the scarcely visible cell-wall (fig. 4.). In some, as above described in the frog (par. 36.), the cell-wall closely embraced the cellæform nucleus, so that this at first appeared free, until by the addition of water the extremely pale cell-wall was brought out. Often the distension of the cell-wall took place on one side only, presenting the appearance delineated in figure 5. In some again the cell-wall was not only distended but pretty distinct, and even slightly reddish (fig. 6).

"48. The size of the cellæform nucleus was from $\frac{1}{4860}$ th to $\frac{1}{3650}$ th of an inch in diameter, and the cell itself in those in which it was distended from about $\frac{1}{4480}$ th to $\frac{1}{1460}$ th of an inch in diameter.

"49. *Nucleated cells—Coloured stage.*—Most of these were circular (fig. 7 a), but some were oval (fig. 8). The circular measured about $\frac{1}{1950}$ th of an inch in diameter. The oval measured about $\frac{1}{1460}$ th in the long diameter, and about $\frac{1}{2125}$ th of an inch in the short diameter. The cellæform nucleus, when not at first evident, was brought distinctly into view by the addition of very dilute acetic acid, or even of water merely. It was circular, and measured from about $\frac{1}{4170}$ th to about $\frac{1}{3650}$ th of an inch in diameter (fig. 7 b).

"50. As to the *small red corpuscles similar to those of the fully-formed blood* (fig. 9), they were rather larger than in the grown animal. They closely resembled the cellæform nuclei of the nucleated cells, only they were red, whilst the latter were uncoloured. There were some free corpuscles, however, uncoloured which resembled the cellæform nuclei of the nucleated cells, and also the small red corpuscles *minus* their colour.

"51. Not having yet had an opportunity of examining the blood on its first appearance in the mammiferous embryo, I do not know if small red corpuscles already exist then, but in embryos a little further advanced than that, the blood of which is at present under consideration, I found the number of small red corpuscles greater—that of the nucleated red corpuscles less; and in fœtuses still farther advanced, the well-marked nucleated red corpuscles had altogether given place to the small red corpuscles.

"52. Putting the small red corpuscles out of view, the blood of the early mammiferous embryo here examined resembles the blood of the oviparous vertebrata in presenting, besides granule-cells, nucleated cells both in uncoloured and coloured stages.

"53. In regard to the red nucleated cells it is to be remarked, that the circular ones were the most numerous, and as they are identical in appearance, size, and structure, with the red corpuscles of the blood of the lamprey, with the exception that their cellæform nuclei are larger, the blood of the mammiferous embryo might be compared to a mixture of the blood of the lamprey, with a small

quantity from a skate or frog to supply the few oval red nucleated cells, and a small quantity from a calf to supply the small red corpuscles."

"*Examination of the corpuscles of the fully-formed blood of man and the mammifera, and a comparison of them with those of the blood of the early mammiferous embryo and of oviparous vertebrate animals.*

"57. *Granule blood-cells.*—The corpuscles in the blood of man and the mammifera corresponding to the granule-cells of the blood of the animals above considered, are certain of those commonly spoken of under the name of 'lymph' or 'colourless' corpuscles—not all, as will be shown below, par. 66.

"58. Both coarsely and finely granular stages of the granule-cell may be recognised. In human blood they in general become speedily collapsed after the blood is drawn, and some, especially those in the finely granular stage, may be seen on careful inspection shooting out processes (fig. 2 *a*.) like the same cells in the blood of the frog (par. 27.) In this state they look as if they had burst, but such is not the case in general.

"59. By the addition of water, the collapsed granule-cells become distended and their appearance is thus distinctly brought out. Fig 1 *a* represents the human granule-cell in its coarsely granular stage thus distended; the contained granules, it will be observed, are minute, and on attentive observation they are seen to be in active molecular motion. Fig. 2 *b* represents the finely granular stage of granule-cell of human blood, distended by water.

"60. The granule-cell of human blood is sometimes seen to present, though not very strikingly, a clear spot indicating, as I think, the place of the nucleus, similar to that above represented in the case of the granule-cell of the skate.

"61. When distended by the action of water the human granule blood-cell is about $\frac{1}{2400}$ th of an inch in diameter.

"62. Fig. 3 *a* represents the granule-cell, coarsely granular stage, and fig. 4 the granule-cell in its finely granular stage of the blood of the horse. Figs. 5 *a* and 6 represent the coarsely and finely granular stages of the granule-cell of the blood of the elephant. It will be observed that the diameter of these cells is somewhat greater than that of the granule-cell of human blood, but the principal difference is the size of the contained granules in the coarsely granular stage. Whilst in the granule-cell, coarsely granular stage, of the horse, the granules may be estimated in round numbers at about $\frac{1}{13000}$ th of an inch in diameter, those in the same cell of the elephant are about $\frac{1}{20000}$ th, and those in the human granule blood-cell about $\frac{1}{25000}$ th.

"63. If, after the granule blood-cell has been distended by the action of water, very dilute acetic acid be added, the contained granules are dissolved, and a cellæform nucleus brought into view. Fig. 1 *b* represents a human granule-cell so acted on; fig. 3 *b* a

granule blood-cell of the horse similarly acted on; and fig. 5 b a granule blood-cell of the elephant in the process of being acted on by acetic acid.

“64. If, before the addition of the acid, the granule-cell has not been distended by the action of water, and if the acid has not been much diluted, instead of one cellæform nucleus of the size represented, and which in the human granule-cell may be about $\frac{1}{360}$ th of an inch in diameter, but larger in the elephant and smaller in the horse, an appearance of several smaller ones variously aggregated may be brought out. This, however, in opposition to what is generally believed, and in opposition to what I myself once believed, I can affirm most positively is merely an appearance artificially produced by the corrugating action of the acid on the walls of the single cellæform nucleus, altogether in the manner above shown to be the case with the cellæform nucleus in the corresponding cells of the blood of the frog.

“65. *Nucleated blood-cells—Coloured stage.*—In human blood to which water has been added, besides the distended granule-cells, cells of the same size may be seen, not containing granules, but presenting most distinctly a single cellæform nucleus (fig. 7.) Such cells resemble the granule-cell deprived of its granules by the action of acetic acid. Moreover, it will be observed that they are essentially similar to the nucleated blood-cells in the uncoloured stage of the blood of the mammiferous embryo and of the oviparous vertebrate animals. In their mode of development from granule-cells, as indicated by the occurrence of cells in transition stages (fig. 10) as well as in their structure, they are in fact clearly identical. I recognise them, therefore, as the corpuscles in human blood in the *phase of nucleated cell, uncoloured stage.*

“66. In consequence of the collapsed state into which they fall, immediately on the blood being drawn (fig. 8), these uncoloured nucleated cells of human blood are scarcely to be distinguished from the collapsed granule-cells (fig. 2 a) until after distention by water. They are usually confounded with them under the same name of ‘lymph,’ or ‘colourless’ corpuscle.

“67. In the blood of the mammiferous animals which I have yet examined in reference to the point under consideration, viz. the horse, elephant, paco, sheep, goat, and rabbit, uncoloured nucleated cells likewise exist, those of the blood of one animal resembling those of another, and all resembling those of human blood in all essential particulars of structure. What difference there is consists in the size of the cell, but especially in the size of the cellæform nucleus, as may be seen by comparing the figures 9 and 11 of the nucleated blood-cells of the elephant and goat for example; the cellæform nucleus of the former being about $\frac{1}{360}$ th, that of the latter only about $\frac{1}{600}$ th of an inch in diameter.

“68. *Nucleated blood-cells—Coloured stage.*—In the blood of man unacted on by re-agents I have not yet observed such corpuscles, but in the blood of other mammifera which I have examined, especially in the blood of the horse and elephant, I have observed them.

Fig. 13 represents one of several such corpuscles observed in the blood of the horse which had been drawn from the animal for a day or two, but to which no water or other reagent had been added either before or during the examination. The transparent and colourless circumferential ring, the optical expression of the transparent and colourless cell-wall doubled on itself, formed a striking contrast with the red-coloured interior of the cell, as did also the uncoloured circumference of the cellæform nucleus. This nucleated cell differed from the nucleated cell in the uncoloured stage in not being collapsed. The one delineated was not plumply distended, but others were. Fig. 14 represents a coloured nucleated cell from the blood of the elephant more fully distended.

“69. Fig. 15 *a* represents a nucleated cell slightly tinged red, which was observed in human blood, but only after the addition of water; fig. 15 *b* represents a still more slightly tinged nucleated cell, observed likewise after the addition of water, in the blood of the horse, and figs. 16 and 17 represent similar ones from the blood of the sheep and paco. These may be looked on as transitions from the uncoloured to the coloured stage of nucleated cell, such as have been above referred to in the blood of the early mammiferous embryo and in that of the oviparous vertebrata.”

Mr Jones next examines and describes the characters of the red corpuscle of the fully formed blood of man and the mammifera.

“In consequence of similarity of colour, it has been generally taken for granted that this is the exact analogue of the ‘red corpuscle’ or coloured nucleated blood-cell of the oviparous vertebrata. Physiologists have accordingly supposed that it should contain a nucleus; but though unsuccessful in the attempt to demonstrate one, they have not altogether ceased to believe in the existence of an exact analogy between it and the red ‘corpuscle’ or coloured nucleated blood-cell of the oviparous vertebrata; they have rather had recourse to conjecture to account for the absence of a nucleus.

“72. In reference to this point, the red nucleated cell of the blood of the early mammiferous embryo has been especially appealed to; it being considered by Mr Gulliver, and I think correctly, as the exact analogue of the ‘red corpuscle’ or coloured nucleated blood-cell of the oviparous vertebrata. But to take it for granted that the ‘red corpuscle’ of the fully-formed blood of the mammifera is the exact analogue of the red nucleated cell of their early embryonic blood, is not more warranted than to take it for granted that the ‘red corpuscle’ of the fully-formed blood of the mammifera is the exact analogue of the red nucleated blood-cell of the oviparous vertebrata.

“73. Under the impression, however, that the ‘red corpuscle’ of the fully-formed blood of man and the mammifera is the exact analogue of the red nucleated cell of the blood of the early mammiferous embryo, Mr Gulliver endeavours to account for the absence of a nucleus in the former by supposing that it disappears at an early

period of intra-uterine life. This apology for the absence of a nucleus in the 'red corpuscle' of the fully-formed blood of man and the mammifera, would have had weight if the 'red corpuscle' had been an object persistent throughout life like a limb or an eye, but as it is an object constantly disappearing and being regenerated, we should expect, if it were really a nucleated cell originally, to meet with it in a stage when it does contain a nucleus.

“74. Professors H. Nasse, Schultz, and Henle, who appear to have observed the nucleated cell of the fully-formed mammiferous blood both in its uncoloured and coloured stages as above described, have supposed it to be the 'red corpuscle' in such a stage. But its larger size (compare figs. 13 and 20, 14 and 23, 11 and 24) seems to forbid such a supposition. Supposing, however, that this objection could be waived, still it is nothing but unsupported conjecture to say that the nucleated cell and the 'red corpuscle' are exactly the same body in different stages of development.

“75. On the other hand, there is a chain of evidence which appears to me to point forcibly to a different view of the nature of the 'red corpuscle' of the fully-formed blood of man and the mammifera, whilst by implication it as forcibly forbids the supposition that the nucleated cell and 'red corpuscle' are exactly the same body—the latter the same as the former, only *minus* its nucleus, and with an increase of colour.

“76. The view of the nature of the 'red corpuscle' of the fully-formed blood of man and the mammifera to which I refer is this: the 'red corpuscle' of the fully-formed blood of man and the mammifera is the cellæform nucleus of the nucleated cell set free by the bursting of this cell itself, and become filled and red by the secretion of globuline and colouring matter into its interior; and the chain of evidence which appears to me so forcibly to point to it is the following:

“77. 1st. *Correspondence in size between the cellæform nucleus of the nucleated cell and the 'red corpuscle.'*—Whilst, as we have seen, there is a marked difference in size between the nucleated cell and the 'red corpuscle' of the same blood, there is, on the contrary, a striking correspondence in size between the cellæform nucleus of the nucleated cell and the 'red corpuscle.' A glance at the figures will show this. In human blood, in which the 'red corpuscle' is large, the cellæform nucleus of the nucleated cell is large. In the blood of the horse, in which the 'red corpuscle' is less, the cellæform nucleus of the nucleated cell is also less. Again, in sheep's blood, in which the 'red corpuscle' is still less, the cellæform nucleus of the nucleated cell is still less.

“78. These facts, and the fact already pointed out by Mr Gulliver, that the nucleus of the red nucleated corpuscle of the blood of the early mammiferous embryo is about the same size as the 'red corpuscles' in the same blood, which are similar to those of the fully-formed blood of the animal, were sufficient to draw my attention to the correspondence in size between the cellæform nucleus of the

nucleated cell and the 'red corpuscle,' but not sufficient to satisfy me that it was more than a coincidence. I therefore became desirous of testing further the correspondence; and for this purpose it occurred to me that the blood of the elephant (in which the 'red corpuscles' are larger than those of any mammiferous animal, as Mr Gulliver has shown) and the blood of the napu musk deer (in which, according to the same authority, the red corpuscles are smaller than those of any other mammiferous animal) would be well-adapted.

"79. Accordingly I made application to the Council of the Zoological Society for leave to obtain a minute quantity of blood from the animals I have mentioned. With this request the Council of the Society readily complied, so far as regards the elephant, but could not as regards the musk deer, their being at present no specimen in their menagerie. As a substitute for the blood of the musk deer I have had recourse to that of the goat, as distinguished for very small 'red corpuscles.'

"80. The result of my examination of the blood of the elephant is, that the cellæform nucleus of the nucleated cell is larger than that of any other animal, the blood of which I have examined, and that it presents a remarkable and exact correspondence in size with the 'red corpuscle' of the same animal. (See figs. 9 and 23.)

"81. The result of my examination of the blood of the goat is, that the cellæform nucleus of the nucleated cell is smaller than that of any other animal the blood of which I have examined. As to its correspondence in size with the red corpuscles, this does not at first view appear so complete as in the preceding instances; the large majority of 'red corpuscles' being smaller than the cellæform nucleus of the nucleated cell. There are, however, some 'red corpuscles' which exactly correspond in size with the cellæform nucleus of the nucleated cell (see figs. 11 and 24), and these have all the characters of the early stage. (See below, par. 86.)

"82. Notwithstanding this slight discrepancy in the blood of the goat, which moreover is rather apparent than real, I believe that it will be admitted as a general proposition, that there is a close and constant correspondence in size between the cellæform nucleus of the nucleated cell and the 'red corpuscle' of the same blood.

"83. 2d. *Correspondence in form between the cellæform nucleus of the nucleated cell and the 'red corpuscle.'*—In the blood of the animals which I had hitherto examined, I had found the cellæform nucleus of the nucleated cell circular, and, therefore, so far corresponding in shape with the 'red corpuscle;' but not considering this conclusive as to the existence of an essential relation in respect of form, it occurred to me that the point might be tested by a reference to the blood of the camel tribe, in which the red corpuscles are of an oval shape.

"84. Accordingly, availing myself of the liberal permission granted me by the Council of the Zoological Society, I procured a small quantity of the blood of the paco, and subjected the cellæform nucleus of its nucleated cells to a comparison with its 'red corpuscles.' The result was that in the nucleated cells generally the cellæform nucleus

was circular (fig. 12.) In a few nucleated cells, indeed, the cellæform nucleus was of a somewhat oval shape (fig. 17), but this did not seem to be more than what I had occasionally observed in the blood of those mammifera, in which the 'red corpuscles' are circular.

"85. My reference to the blood of the paco thus appeared to decide against an essential correspondence in form between the cellæform nucleus of the nucleated cell and the 'red corpuscle.' A little further study of the blood of the paco, however, soon cleared away this apparent difficulty, and showed me a state of things more corroborative of my view regarding the nature of the 'red corpuscle,' and in much closer correspondence with analogy, perhaps, than would have been that which I supposed possible. I discovered, in fact, corpuscles in different stages of transition, from the circular form similar to that of the cellæform nucleus within the nucleated cell, to the oval form of the perfect 'red corpuscle;' and I also discovered that in these different stages of transition in respect of form there are different degrees of coloration (figs. 26, 27, 28, 29, 30).*

"86. I had already observed progressive degrees of coloration in the circular 'red corpuscles' of the mammifera generally. Thus, in the blood of the horse, for example, I had observed corpuscles (fig. 19) similar on the one hand to the 'red corpuscle' (fig. 20) but uncoloured, and on the other to the cellæform nucleus of the nucleated cell (fig. 15), and this in all respects. Such corpuscles I was disposed to regard as cellæform nuclei recently set free from the nucleated cells, but not yet reddened by the secretion of colouring matter into their interior. Again, I had observed corpuscles similar to those just mentioned, but presenting a red tinge in the middle, the circumference untinged appearing like a transparent and colourless ring (fig. 21); but the combination of progressive stages of coloration of the 'red corpuscle' with its progressive changes of form in the blood of the paco is peculiarly interesting.

"87. The transition from the circular to the oval shape presented by the 'red corpuscle' of the paco, is analogous to that which we have seen presented by the red nucleated cell of the oviparous vertebrata. It is also worthy of remark, that like the latter the oval red corpuscle of the paco is reduced to a circular form by the action of water (fig. 31.) The progressive degrees of coloration are also analogous to what are presented by the nucleated cells of the blood of the oviparous vertebrata. But it is curious to observe, that whilst among the mammifera the attainment of the oval form by the 'red corpuscle' is exceptional, among the oviparous vertebrata the attainment of the oval form by the red nucleated cell is the rule.

"88. But to return to the bearing of the facts I have now related on the question before us. They show that originally the 'red corpuscles' of the paco are of a circular form like the 'red corpuscles'

* These numbers, which refer to figures in the last compartment of second column, though not inserted, correspond to the figures as arranged in that compartment.

of other mammifera, but that whilst the latter remain circular, the former gradually acquire an oval shape. From this it follows that the circular form of the cellæform nucleus of the nucleated cell of the blood of the paco, so far from being an objection to the view of the nature of the 'red corpuscle' of the mammifera which I am advocating, is just what should be according to it. In short, I consider the 'red corpuscle' of the paco, in its circular form and as yet uncoloured state, to be, like the similar corpuscles referred to in par. 86, cellæform nuclei but recently set free from the nucleated cell.

"89. 3d. *Correspondence in structure between the cellæform nucleus of the nucleated cell and the 'red corpuscle.'*—The cellæform nucleus of the nucleated cell is a simple cell or vesicle without nucleus or nucleolus, and such as exactly resembles the 'red corpuscle' minus its red coloured contents. Indeed, when by the action of water the 'red corpuscle' is deprived of its red coloured contents without being rendered indistinct, the resemblance in respect of structure is very striking, as indicated by the lumen and transparent circumferential ring—the optical expression of the double contour of the wall of the collapsed cell or vesicle.

"90. As already said, 'red corpuscles' may occur uncoloured or with the lumen only red, the circumferential ring just referred to remaining transparent and colourless. This indicates that the colouring matter is contained in the interior of the cell or vesicle. In its perfect state, however, the 'red corpuscle' appears red at the circumference also, owing, apparently, to its wall being impregnated with colouring matter.

"91. The absence of anything like a nucleus in the 'red corpuscle' of the fully-formed blood of the mammifera is in correspondence with the absence of anything of the kind in the cellæform nucleus of the nucleated cell. The appearance of dots, as if of nucleoli, occasionally seen in the latter, I consider to be mere corrugations, similar to those which give rise to the mulberry appearance which the red corpuscle is so prone to assume.

"92. Before admitting the correspondence in size, form and structure, between the cellæform nucleus of the nucleated cell and the 'red corpuscle' of the blood of the mammifera, which I have now demonstrated, as evidence in favour of their identity, it will be proper to inquire if there be any points of difference between them sufficient to constitute evidence to the contrary effect.

"93. *Points of difference between the cellæform nucleus of the nucleated cell and the 'red corpuscle.'*—1st. The cellæform nucleus of the nucleated cell refracts the light more than the 'red corpuscle.' 2d. The cellæform nucleus is not affected by water or acetic acid like the 'red corpuscles'—the former being by the action of these reagents rendered more distinct, the latter rendered indistinct. These circumstances indicate a chemical difference between the cellæform nucleus and the 'red corpuscle,' and if a satisfactory explanation of them could not be offered they would constitute a powerful argument against my view. But I believe the following explanation will remove the difficulty.

" 94. *The objection against the identity of the cellæform nucleus and 'red corpuscle' which might be founded on the different modes in which they refract light, and are affected by chemical reagents, explained away.*—The differences mentioned are striking enough between the cellæform nucleus and the 'red corpuscle' in its perfect state; but if we take the 'red corpuscle' in the stage when, as I have already described, it is as yet uncoloured or presents colour only in its interior, then we find but little difference between it and the cellæform nucleus of the nucleated cell, either in the mode in which it refracts light, or in which it is affected by water or acetic acid. Even when fully red, the 'red corpuscles' are not all equally affected by the action of water or acetic acid; some being always seen which resist to a greater or less degree the action of these reagents. Such corpuscles have been considered to be advanced formations, they appear to me on the contrary to be recent.

" 95. From this it appears that the 'red corpuscle' in its advanced stage does not differ, in the respects under consideration, more from the cellæform nucleus of the nucleated cell than it does from what is undoubtedly itself in an early stage. The differences mentioned therefore argue nothing against the identity of the cellæform nucleus and the 'red corpuscle.'

" 96. There being thus no evidence to the contrary effect, the correspondence in size, form, and structure between the cellæform nucleus of the nucleated cell and the 'red corpuscle' of the blood of the mammifera, which I have now demonstrated, will I believe be admitted to constitute pretty forcible evidence as to their identity. Whether any additional evidence can be adduced will be seen in the sequel, par. 106.

" 97. If the view of the origin and nature of the 'red corpuscle' of the fully-formed blood of man and the mammifera which I have now given be correct, this 'red corpuscle' must be considered as a third phase of development of the blood corpuscle, and of which there occur uncoloured and coloured stages. This third phase I propose to call the phase of *free cellæform nucleus.*—Pp. 74-79.

The author then gives the following summary of his observations on these different phases.

98. " In all the vertebrate animals examined, oviparous and mammiferous, we find blood corpuscles in what I call the first phase of development, or phase of granule-cell, this presenting in a more or less marked manner two principal stages, a coarsely and a finely granular stage. In all the animals examined we likewise find blood-corpuscles in what I call the second phase of development or phase of nucleated cell; this again presenting two principal stages, an uncoloured and a coloured. In *Branchiostoma lubricum* alone of all the oviparous vertebrata, we have seen reason to conjecture (par. 20,) that the nucleated blood-cell occurs in the uncoloured stage only, or if the coloured stage exists, that the coloration is only in a very slight degree.

" 99. In regard to the nucleated blood-cell in the coloured stage,

it is to be observed that it occurs in its highest degree of development and in great number only in the oviparous vertebrata, in which it constitutes the 'red corpuscle,' and in the early mammiferous embryo. In the fully-formed blood of the mammifera it occurs in a comparatively low degree of development and in very small number.

"100. It is in the mammifera alone that we find the blood-corpuscle in what I call the third phase of development or phase of free cellæform nucleus. This exists in both uncoloured and coloured stages. In the former stage it is rare, in the latter it is the 'red corpuscle' of the fully-formed blood of man and the mammifera. In the early mammiferous embryos which I have examined there have always been, in addition to the numerous 'red corpuscles' similar to those of the oviparous vertebrata, *i. e.* red nucleated blood-cells, some 'red corpuscles' similar to those of the fully-formed mammiferous blood. Is there a period in the development of the mammiferous embryo when there exist in the blood no such 'red corpuscles,' viz. free cellæform nuclei in the coloured stage?"—Pp. 79–80.

The author next gives a description of the corpuscle of the lymph of vertebrate animals, comparing it with the corpuscle of the blood. The following results deserve particular attention.

"101. *The Skate and Mackerel*.—The lymph of the skate examined was obtained from the orbit in very small quantity, and could not be considered a good specimen. The corpuscles were besides a few granule-cells, nucleated cells in the uncoloured stage,—the cell-wall of many extremely pale,—and some nucleated cells in the earlier stage of coloration,—all identical with the corresponding corpuscles of the blood of the animal. (See figs. 1 to 6.)

"102. The lymph of the mackerel, taken as an example of an osseous fish, was likewise obtained from the orbit, and might be viewed as a good specimen. It contained granule-cells, nucleated cells in the uncoloured stage,—many with the cell-wall so pale as to be readily overlooked without attentive examination,—and nucleated cells in the earlier part of the stage of coloration,—all identical with the corresponding corpuscles of the blood of the animal."

"103. *Frog*.—The lymph subjected to examination was obtained from one of the posterior lymph hearts. The corpuscles were,—1st. Corpuscles at once recognisable as identical with the granule-cells of the blood of the animal, and these in both coarsely and finely granular stages. There were also cells in transition from granule to nucleated cells. 2d. Corpuscles which at first sight appeared to be free nuclei, but were discovered on closer examination to be inclosed within cells of a circular or oval form, and extremely pale. The identity of these cells with the nucleated cells, uncoloured stage, of the blood of the animal could not be overlooked. 3d. A considerable number of red nucleated cells, the same as the 'red corpuscles' of the blood. Many of these no doubt were derived from small blood-vessels which had been cut in opening the lymph heart, but some of them I am inclined to believe belonged to the contents

of the lymph heart, those especially which appeared to be in an early stage of coloration. (See the figures of the blood of the Frog.)

“104. *Common Fowl*.—The lymph subjected to examination was obtained from a lymphatic gland. It contained,—1st. Corpuscles about $\frac{1}{3000}$ th of an inch in diameter, less or more, identical with the granule blood-cells of the blood of the same animal; some being in coarsely and some in finely granular stages (figs. 1 to 2.) 2d. Corpuscles in much greater number than the preceding, of the form of circular discs, with a depressed point in the centre about $\frac{1}{7000}$ th of an inch in diameter, and though colourless very distinct. At first glance these corpuscles appear to be free, but subjected to more careful examination, they were observed to be contained within a cell of a diameter of about $\frac{1}{3000}$ th of an inch, the wall of which was extremely pale. Having once perceived it, however, there was no difficulty in recognising the whole as a nucleated cell identical with the nucleated cell of the blood of the animal, in its uncoloured stage (fig. 3.) There were some nucleated cells similar to those just noticed, but slightly tinged red in the interior; and besides such there were some oval nucleated cells, altogether similar to the oval red corpuscles of the blood of the animal, but coloured only, and that slightly in the interior, the circumferential doubling of the cell-wall appearing quite transparent and colourless.’

“105. *Man and the Mammifera*.—The description of the corpuscles of the lymph, here to be given, was drawn from the examination of those presented by the contents of the thoracic duct. Besides the contents of the thoracic duct of the human subject, I have also particularly examined the contents of the thoracic duct of the rabbit, and found their corpuscles to be in all essential respects identical.

“106. The corpuscles were,—1st. Granule cells in both coarsely and finely granular stages, altogether similar to those of the blood (figs. 1, 2.) 2d. Besides many cells in transition from granule to nucleated phase, nucleated cells in both uncoloured and coloured stages, quite the same as those of the blood (figs. 7 and 15 a.) The nucleated cells were fully distended independent of the addition of water, and were comparatively the most numerous of the corpuscles in the lymph; those in the coloured stage being more so than those in the uncoloured stage. Slight as the coloration of the coloured stage of the nucleated cell was generally, some few cells presented it perhaps in a more marked manner than is usually presented by the same cells as they occur in the fully-formed blood, in this respect approaching to the red nucleated cell of the blood of the early mammiferous embryo. Some even were of an oval shape. 3d. A considerable number of free cellæform nuclei, both uncoloured and in different degrees of progressive coloration. These free cellæform nuclei were not much affected by the action of acetic acid or water. Lastly, there were seen here and there among the other corpuscles the empty shell of a nucleated cell in the coloured stage, with a free cellæform nucleus beside it as if just extruded by the bursting of the cell-wall. This is a circumstance which may be viewed as additional evidence, and this of a direct kind, in support

of the view above given of the nature and origin of the 'red corpuscle' of man and the mammifera.

"107. From the account now given of the corpuscle of the lymph of vertebrate animals, it appears that it is identical with the corpuscle of their blood. In the oviparous vertebrata it occurs, like the corpuscle of their blood, in the two phases of granule and nucleated cell; whilst in man and the mammifera it occurs, like the corpuscle of their blood, in the three phases of granule-cell, nucleated cell, and free cellæform nucleus.

"108. The only difference that exists between the corpuscle in the lymph and the corpuscle in the blood, is, as regards the oviparous vertebrata, the little degree of coloration which the coloured stage of nucleated cell as yet presents, and as regards the mammifera, the small degree of coloration which the coloured stage of free cellæform nucleus has as yet attained.

"109. The comparatively small total amount of corpuscles in the lymph is well known. As regards the relative number in the different phases of the corpuscle, it is to be remarked that in the oviparous vertebrata, the nucleated cells are, as in the blood, more numerous than the granule-cells. The nucleated cells are also more numerous than the granule-cells in the lymph of man and the mammifera, which is different from what obtains in their blood. Taking into account this relatively great number of nucleated cells, and the existence, though in less numbers, of free cellæform nuclei, we might compare the contents of the thoracic duct of a mammiferous animal, in respect both of the kinds of corpuscles and of their relative but not absolute numbers, with the blood of its early embryo state. The circumstances above mentioned, that some of the nucleated cells of the contents of the thoracic duct present in the coloured stage coloration in a pretty marked degree, and that some are of an oval shape, give additional exactness to the comparison."—Pp. 80–83.

Mr Jones remarks, what must have occurred to every one who has observed the statements of different authorities on this subject, that in the description of the lymph corpuscles given by different observers, much confusion exists. The causes of this confusion he explains in the following manner.

"1st. Observations made on the corpuscles of the lymph of oviparous vertebrate animals have been adopted as applicable to the corpuscles of the lymph of man and the mammifera, whereas it will have been seen from what has been above said, that the corpuscles of the lymph of the oviparous vertebrata can no more be taken as an exemplification of the corpuscles of the lymph of man and the mammifera, than the blood-corpuscles of the former as an exemplification of those of the latter.

"2d. Hewson, it is known, gave the name of 'central particles' to the nuclei of the red corpuscles of the blood of the oviparous vertebrata. He at the same time supposed that he had seen similar particles in the interior of the red corpuscles of the blood of man and the mammifera.

“ Having observed in the juice of lymphatic glands, bodies which he supposed to be free, and which he recognised to be similar to the central particles of the red corpuscles of the blood, Hewson came to the conclusion that the central particles are at first free, and that they subsequently become surrounded with the red coloured cell-wall, which he called ‘vesicular portion.’

“ How Hewson should have come to this conclusion is explicable, as regards the juice of the lymphatic glands of birds, on the following supposition. In the juice of the lymphatic glands of birds, it has been above seen, that nucleated cells in the uncoloured stage exist in great numbers; but that whilst the nucleus is very distinct, the cell-wall is extremely pale. In consequence of this Hewson appears to have overlooked the cell-wall and distinguished the nucleus alone, but when the cell-wall had already become evident around the nucleus by having acquired colour, he naturally supposed that it was a new formation.

“ How Hewson should have come to the same conclusion as regards the lymph of man and the mammifera, however, is scarcely explicable on any supposition. For to say nothing of the circumstance that there is not a ‘central particle’ in the red corpuscle of the blood of the mammifera, Hewson’s description of ‘central particles’ in the juice of the lymphatic glands of the mammifera cannot apply to the cellæform nuclei of the uncoloured nucleated cells, seeing that the cellæform nucleus of the uncoloured nucleated cell is nearly, if not as large as the perfect red corpuscle itself, which he alleged to contain the ‘central particle.’ The difficulty is still the same, even supposing Hewson to refer to free cellæform nuclei in the uncoloured stage, and still greater supposing him to refer to granule-cells.

“ 3d. Since Hewson’s time the cell-wall of the uncoloured nucleated cell in the lymph, especially of the oviparous vertebrata, has continued to be generally overlooked, and the nucleus described as free. The nucleus, moreover, has been confounded with the granule-cell, which is what is commonly taken as the type of the ‘lymph-corpuscle.’ The same confusion prevails in regard to the lymph of man and the mammifera, with this additional complication, viz., that granule-cells have been confounded with free cellæform nuclei in the uncoloured stage.

“ Wagner participating in the error of confounding granule-cells with cellæform nuclei, but having seen and distinguished the cell-wall of nucleated cells, he has taken their cellæform nucleus for a granule-cell, which had acquired a cell wall around it.

“ Of all the authors I am acquainted with, Henle alone has accurately distinguished the different forms of the corpuscle of the lymph of man and the mammifera, without, however, pointing out their relationship; but in his account of the corpuscles of the lymph of the frog, he has fallen into inaccuracies similar to some of those which have been above pointed out.”—P. 83.

The present is the most complete and the most accurate ac-

count hitherto placed before the physiological world of the constitution and true nature of the blood corpuscles and granules, as ascertained by microscopical observation. These observations have been evidently made with the greatest care, and with every precaution to exclude errors, from every source. They deserve, and will doubtless obtain, a general reception among all anatomists and physiologists, who are desirous to know correct facts only. It is important to observe, that Mr Jones has not in these observations employed, a very high magnifying power, though greater than that used by several microscopical observers; and this circumstance renders his observations the more likely to be correct and faithful.

Our limits will not allow us to enter on the other division of the researches of Mr Jones, and while we defer them only for another number, we have been desirous to put our readers in possession of the most important facts ascertained in relation to the vertebrated animals, with the history of which they may be supposed to be most interested.

ART. III.—*On the Pathology and Treatment of Scrofula; being the Prize Essay for the year 1846.* By R. M. GLOVER, M. D., Corresponding Member of the Medical Society of London, Lecturer on Materia Medica in the Newcastle Medical School. London, Churchill, Soho. Pp. 315. Illustrated by Engravings.

WITHIN the short period of two years we have had occasion to direct attention to the merits of two new works on the subject of scrofula and scrofulous diseases; their causes; their characteristic effects; their prevention; and their treatment. Of the work of M. Lugol an account was given in our sixty-fourth volume;* and the elaborate performance of Mr Phillips we considered in our sixty-sixth volume at some length.† We should not again, after having occasion so amply to investigate this subject, have been disposed soon to recur to it. The appearance of a third work, however, by Dr Glover, of Newcastle-upon-Tyne, has in a great degree induced us to depart from this resolution, and to bring before the notice of our readers a work which is certainly entitled to attention.

Dr Glover is the more entitled to patient attention on this subject, that he has made the causes, characters, and treatment of scrofulous diseases the subject of particular study and investigation. In 1844, he, along with another professional gentleman,

* Edinburgh Medical and Surgical Journal, Vol. lxiv. p. 234. Edin. 1845.

† Ibid., Vol. lxvi. p. 117. 1846.

The Blood Corpuscle

1. Granule Cell.

2. Nucleated Cell.

Lamprey

Skate

Frog

Fowl

Mam. E.

Man, & Mam.

3. Free Cellæform Nucleus

Mam. Embryo

Man, & Mam.