8. That this power of altering the pelvis upon the spine, so as to apply to midwifery, exists only in the mobility of the three or four superior lumbar, and four or five inferior dorsal vertebræ.

9. That if any single position could be fixed upon for the parturient woman, that midway, between lying on the back and side, is least objectionable.

ART. VIII.—*Secretion of Carbon by Animals.* By ROBERT RIGG, F.R.S. ✓

To the Editor of the Edinburgh Medical and Surgical Journal.

SIR,—The scientific world is at present much occupied with the application of chemistry to animal and vegetable physiology, and it may be interesting to some of your readers to know, that, by a few very simple experiments, they may satisfy themselves upon that branch of the subject which relates to the secretion of carbon by animals.

Suppose an animal, which comprises in its whole system 50 parts by weight of carbon, to be kept for five days, during which it consumes other 50 parts, it is evident that if, during these five days, it gives off to the surrounding atmosphere 60 parts, and at the end of that time is found to have increased its weight of carbon 10 parts, there is a positive gain of carbon equivalent to 20 per cent.

The experiment may be easily made upon young small animals. Take two of these so nearly alike that there can be no material difference in the weight of carbon they comprise. Kill one of these, and expose it to a temperature not exceeding 220° for two or three days; it may then be powdered, and by subjecting an average sample to analysis with oxide of copper, the weight of carbon comprised in the entire animal may be determined with the greatest certainty. The other being supplied with food, the weight and chemical constitution of which is ascertained, should be kept in a limited atmosphere, which must be changed and tested every one, two, or three hours.* The proportion of carbonic acid in that atmosphere will show the quantity of carbon given off by the animal in the course of the experiment, and the increase or decrease of carbon in the animal itself may be ascertained in the manner above mentioned.

In this manner I have experimented upon many animals, and without taking account of the carbon which passes off otherwise than by respiration, the results have invariably been a great increase of carbon; an increase which cannot be accounted for unless we conclude that carbon is secreted by animals.

* The health of the animal appears to be affected by an atmosphere containing more than 5 per cent. of its volume of carbonic acid.

Amongst my best experiments are those made with young mice. A healthy young mouse, weighing 200 grains, comprises in its constitution from 25 to 30 grains of carbon. When fed daily with 60 grains of bread moistened with water containing about 16 grains of carbon, it increases in weight and imparts to the atmosphere from 20 to 26 grains of carbon, this quantity varying generally with the quietness or the activity of the habits of the animal. A kitten, from six to ten weeks old, when supplied daily with four fluid ounces of skim milk containing 66 grains of carbon, will increase in weight, and impart to the atmosphere from 80 to 100 grains of carbon.

Either of these animals may be kept without food until they give off by respiration a weight of carbon equal to 80 per cent., and retain from 60 to 70 per cent. of that which they comprised at the commencement, shewing that a weight of carbon equal to 40 per cent. has been secreted. A tomtit was placed under experiment without food : the bird was violent and restless during its imprisonment. In sixteen hours it imparted to the atmosphere 65 per cent. of carbon, when it appeared to die of exhaustion, and retained 77 per cent. of the weight of carbon it originally contained, shewing a secretion of carbon in sixteen hours, when under violent exertion, equal to 42 per cent.

On making the carbon in the food and that in the air respired by a full-grown person the basis of our calculation, we obtain results which favour the conclusion that carbon is likewise secreted by man. Physiologists estimate the weight of carbon in air respired by an adult at from 5000 to 6000 grains per diem. I have subjected to ultimate analysis many articles of food, and find this weight of carbon far exceeds that in the food consumed by most labouring men, who may be supposed to impart to the atmosphere the greatest weight of carbon. A person eating each day after the following rate will consume 6000 grains of carbon :

Rump steak,	1 pound,	containing	1050 grains of carbon,	
Bread, -	1½	-	2830	—
Potatoes, -	½	-	310	—
Porter, -	2 pints,	-	760	—
New milk,	2 fluid-ounces,		57	—
Butter, -	1½ ounce,		320	—
Cheese, -	1 do.	-	150	—
Sugar, -	2 do.	-	350	—
Coffee, -	1 do.	-	96	—
Tea, -	1 do.	-	80	—
<hr/>				
				6003
				—

This weight of carbon is not more than is consumed by some persons who are actively employed, but it very far exceeds that in the food of our labouring population ; and on comparing it with that allowed for each adult in the different workhouses, &c., we have, in the dietary of the

City of London Union,	-	75	per cent. of this 6000.
Brentford Union,	-	50	do.
Uxbridge,	-	55	do.
Alesford,	-	56	do.
Macclesfield,	-	44	do.
Westminster New Prison,	-	57	do.
Millbank Penitentiary,	-	80	do.
House of Correction, Clerkenwell,	-	53	do.
Hanwell Lunatic Asylum,	-	75	do.

And if we make the carbon in the food of some of our agricultural labourers the subject of comparison, we find the deficiency greater than in any of the above-mentioned dietaries. I would add to these many other experiments which furnish, in my opinion, irresistible evidence of the secretion of carbon by animals. If an animal be kept without change of circumstances, or of diet, except as to the quantity of food, it will be found that the weight of carbon in the air respired does not vary in proportion to that consumed in the shape of food; on the contrary, the deficiency of carbon supplied seems to be met by an extraordinary effort of the animal system, as appears from the following results of accurate observation.

	The food.	The air respired.
If an animal in the first 24 hours has a plentiful supply of food, there is in	80 grs. of carb.	100 grs. of carb.
In the next 24 hours a less quantity,	70 do.	94 do.
Do. - sparing do.,	60 do.	87 do.
Do. - small do.,	50 do.	78 do.
Do. - very small do.,	40 do.	65 do.

But if the animal, instead of having its quantity of food varied, should be sometimes left in a quiescent state, and sometimes excited to great activity, the weight of carbon given off will be found to vary in proportion, within certain limits, to the activity of the habits of the animal, and the exertion called forth.

If the carbon in the food be represented by	-	100
That given off by an animal of easy habits will be		110
Do. do. active do.	-	130
Do. do. when under exertion,		140
Do. do. great do.		150

If the animal be both stinted in food and excited to great activity, the difference between the carbon comprised in the food and that given off by the animal is as follows:—

	In the food.	Given off.	Difference.
Carbon, - -	100	120	20
Do. - -	80	105	25
Do. - -	60	90	30
Do. - -	50	85	35

Hence, food is a substitute for expenditure of animal strength. To this may be added, that when an animal is distressingly exercised, the weight of carbon in the carbonic acid given off by respiration is at first increased, afterwards gradually diminishes, and becomes much less when the animal is in a state of exhaustion.

Rest alone is not then sufficient, but rest and food soon restore the strength of the animal, and with its strength its power of secreting carbon,—a power which I conceive to be essential to animal life, and which will probably furnish a solution to some of the most difficult problems of animal physiology, including that of the generation of animal heat. I am, Sir, your most obedient servant,

ROBERT RIGG.

Greenford, Middlesex,
August 15, 1844.

ART. IX.—*On the Colour and Structure presented by Corpora Lutea in the early stage.* By ROBERT PATERSON, M.D. &c., Leith. See Plate VIII.

CASE I.—A respectable and very fine-looking young female, about 19 years of age, committed suicide under most painful circumstances.

It appeared from the account given by the relations, that the deceased had been very much attached to a young man who visited the house, and with whom they feared an improper intimacy existed. The deceased had menstruated about a month previous to her death, and precisely three weeks before that event she had remained out till very late in the company of the young man referred to, and for which his visits to the house were prohibited. It was distinctly stated that the deceased had never seen him from that night. The suicide was committed about 11 o'clock at night; and an opportunity was afforded of examining the body next morning. The uterus and its appendages were carefully removed for examination afterwards, and with which I was assisted by my friend, Professor Allen Thomson.

A general glance at the uterus and appendages at once suggested the remark that they were plump and rounded in form, more so, indeed, than I had ever previously witnessed them. The uterus was slightly larger, and certainly more rounded in its appearance than the natural unimpregnated organ. Its mouth was completely filled up with thick, tenacious, but clear mucus. Upon cutting into its cavity, through the anterior wall, it was found to be lined with a very delicate decidual membrane. This membrane lined the cavity of the uterus throughout, but did not descend either into the neck or orifices of the Fallopian tubes; it was of a delicate flesh colour, with slight irregularities on its internal surface, giving the appearance of gentle elevations and rugæ or furrows between them, in fact resembling much the furrows produced in the palm of the hand by gentle flexion of the fingers. It was adherent to the internal surface of the uterus, could with difficulty be separated from it, and was a little firmer in texture to the feel