Preventive Percutaneous Coronary Intervention in ST-elevation Myocardial Infarction – The Primacy of Randomised Trials

David S Wald¹ and Jonathan P Bestwick²

1. Professor of Cardiology; 2. Lecturer in Medical Statistics, Wolfson Institute of Preventive Medicine, London, UK

Abstract

Randomised trials show a benefit of preventive (non-infarct artery) percutaneous coronary intervention in patients with acute ST elevation myocardial infarction, but non-randomised studies do not. The evidence on each is quantified and assessed. The primacy of randomised trials reveals the danger of using non-randomised studies that can, as in this case, give the wrong answer.

Keywords

STEMI, preventive PCI, non-infarct artery

Disclosure: The authors have no conflicts of interest to declare.

Received: 2 December 2014 **Accepted:** 4 February 2015 **Citation:** *Interventional Cardiology Review*, 2015;10(1):32–4

Correspondence: David S Wald, Queen Mary University of London, Wolfson Institute of Preventive Medicine, Charterhouse Square, London EC1M 6BQ, UK.
E: d.s.wald@qmul.ac.uk

The management of patients with acute ST-elevation myocardial infarction (STEMI) relies on restoring blood flow to the occluded infarct artery. Prompt percutaneous coronary intervention (PCI) and stenting of the stenosis causing the occlusion reduces the risk of cardiac death and recurrent infarction.¹ In about half of patients,² stenoses are identified in non-infarct arteries at the time of the PCI, leading some operators to extend the procedure and carry out an immediate 'preventive PCI' in the non-infarct arteries, on the basis that this may prevent future serious cardiac events. Until recently, however, clinical guidelines recommended that PCI be limited to the infarct artery, because of concern that the benefit of preventive PCI may not outweigh the risks of the extended procedure.^{3,4}

This recommendation was based on non-randomised studies in which the outcome of patients with STEMI and multivessel disease who received immediate preventive PCI at the time of their infarct artery PCI was compared with the outcome of patients treated by infarct artery PCI alone. In these studies, doctors determined who received preventive PCI and who did not. *Figure 1* shows a meta-analysis plot of 16 such studies (42,817 patients, median follow-up 12 months).^{5–20} The summary estimate in the non-randomised studies is not statistically significant but suggests a possible increased risk of all-cause death or non-fatal MI in the preventive PCI group (odds ratio 1.24, 95 % confidence interval [CI] 0.97–1.60; $p=0.085$). The use of all-cause rather than cardiac death as an outcome is a limitation, because it includes non-cardiac causes that are not influenced by PCI (e.g. cancer) and dilutes the relevant cardiac outcomes, but only one study⁷ reported cause-specific death. A more serious limitation, in the non-randomised studies, is selection bias: the extent to which patients who received preventive PCI, for example, were sicker than those receiving infarct artery-only PCI and were therefore heading for a worse outcome regardless of the treatment strategy adopted. Selection bias is not avoided by increasing study size or by adjusting

for confounding, because not all confounding factors are measured or known, so even in large propensity score-matched studies^{11,14} it is not possible to be sure which treatment is better. Selection bias is avoided in a randomised trial, since the use of preventive PCI is determined by random allocation rather than physician choice. Four such trials (979 patients, median follow-up 18 months),^{21–24} in which cardiac deaths were reported in all but one²³, have been completed and a meta-analysis plot of these trials is also shown in *Figure 1*. The relative risk of cardiac death or non-fatal MI is 0.39 (95 % CI 0.23–0.69; $p<0.001$), showing that selection bias is an important source of error in the non-randomised studies and indicating that preventive PCI, performed as an immediate extension of the infarct artery PCI, reduces the risk of cardiac death and MI by about 60 %.

How can these findings be reconciled with the view that PCI improves symptoms but not prognosis?^{25–27} The value of angioplasty has been studied in different groups of patients with coronary artery disease. The evidence of benefit in reducing the risk of cardiac death and MI in patients with STEMI is known.¹ In patients with non-STEMI, there is a short-term hazard but a long-term benefit.²⁸ In patients with stable angina, prior studies have shown no evidence of prognostic benefit.^{26–27} This gradation of effect may, at least in part, be due to the fact that PCI causes adverse cardiac events as well as preventing them, and in a lower-risk group, for example, in patients with angina rather than a MI, the benefit may not outweigh the harm.

In patients with STEMI, coronary artery plaque rupture remote from the infarct artery has been demonstrated in autopsy^{29,30} and intravascular ultrasound studies,^{31–34} which suggest that plaque instability is not a localised vascular event but a generalised process throughout the coronary tree. In an angiographic study of 253 patients with STEMI, the finding of multiple complex coronary artery lesions (>50 % stenosis) remote from the infarct artery

was associated with a 10.6-fold excess risk of recurrent acute coronary syndrome within 1 year.³⁵ The simplest, albeit speculative, explanation for the preventive benefit of PCI in such patients is by stabilising plaques prone to rupture and spontaneous thrombosis,³⁶ so reducing subsequent infarction. There is uncertainty whether the benefit outweighs the risk of PCI in stenoses <50 %, since these were not included in the randomised trials. Preventive PCI may also reduce ischaemia by improving coronary flow in severe stenoses and by preventing progression over time, explaining the observed reduction in refractory angina. The concordance between the components of the primary outcomes in Preventive Angioplasty in Myocardial Infarction (PRAMI) (hazard ratios 0.34 [0.11–1.08], 0.32 [0.13–0.75] and 0.35 [0.18–0.69] for cardiac death, non-fatal myocardial infarction and refractory angina, respectively)²⁴ suggests that both mechanisms may be similarly important in the prevention of future cardiac events.

In the PRAMI trial the evidence of benefit emerged early on.²⁴ The curves in the Kaplan–Meier plot (see *Figure 2*) diverged within a few days and the maximum effect was evident within a few months, suggesting that the immediacy of preventive PCI is important and that staged preventive PCI (undertaken after a few weeks) may not be as effective. The same observations were apparent in the Complete Versus culprit-Lesion only PRimary PCI Trial (CVLPRIT) trial²² in which all but about a quarter of patients in the preventive PCI group had immediate preventive PCI, the remainder having a staged procedure within a few days. Neither of the two trials was designed to compare immediate versus staged preventive PCI. Such a trial, if judged necessary, would need to be large to demonstrate a difference in outcome on top of the 60 % reduction in cardiac death or MI from preventive PCI alone.

The results of PRAMI²⁴ and CVLPRIT²² have prompted a rethink in the way we manage non-infarct artery stenoses in STEMI. The European Revascularisation Guidelines were recently changed (September 2014) and now recommend that immediate preventive PCI be considered in selected patients with STEMI,³⁷ but do not indicate how this selection should be made. The use of a physiological measure of blood flow, such as fractional flow reserve (FFR), may be better than visual angiographic assessment in guiding preventive PCI,³⁸ but it may also worsen outcomes if non-flow limiting stenoses are left untreated and become the sites of future infarction. Three trials of preventive PCI in patients with STEMI are in progress that are using FFR to decide which non-infarct artery stenoses to treat.^{39–41} No trial is designed to directly compare FFR with angiography in STEMI to determine which, if either, is better in guiding preventive PCI. Further research would be needed to resolve this uncertainty.

A limitation of the trials of preventive PCI, as in all randomised trials, is that they only provide an average effect of treatment. Some patients will benefit more from preventive PCI than others, but in the absence of knowing who they are, no special selection can be recommended. The trials excluded patients with cardiogenic shock, previous coronary artery bypass graft (CABG), significant stenosis of the left main stem or in whom the only non-infarct artery disease was a chronic total occlusion. Therefore, while the benefits of preventive PCI may apply in these selected groups, there

Figure 1: Immediate Preventive Percutaneous Coronary Intervention versus Infarct Artery-only Percutaneous Coronary Intervention – Outcomes in Non-randomised Studies (All-cause Death or Myocardial Infarction) and in Randomised Trials (Cardiac Death or Myocardial Infarction)

*All-cause death or myocardial infarction (MI) (cardiac death reported in only one of the 16 studies);⁷ **Cardiac death or MI (cardiac death reported in all but one trial).²² CI = confidence interval; PCI = percutaneous coronary intervention.

Figure 2: Kaplan–Meier Curves for the Primary Outcome (Composite of Cardiac Death, Non-Fatal Myocardial Infarction or Refractory Angina) in the Preventive Angioplasty in Myocardial Infarction (PRAMI) Trial²⁴

is uncertainty. In others, for whom the non-infarct artery stenoses are judged treatable by PCI, the evidence from the trials completed so far, is clear – that immediate preventive PCI confers substantial clinical benefit. The primacy of randomised trials reveals the danger of using non-randomised studies, which can, as in this case, give the wrong answer. ■

1. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet*, 2003;361:13–20.
2. Park D-W, Calre RM, Schuler PJ, et al., Extent, location, and clinical significance of non-infarct-related coronary artery disease among patients with ST-elevation myocardial

- infarction. *JAMA*, 2014;312:2019–27.
3. The Task Force on the management of ST-segment elevation acute myocardial infarction of the European Society of Cardiology (ESC). ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J*, 2012;33:2569–619.
4. O'Gara PT, Kushner FG, Ascheim DD, et al., 2013 ACCF/

- AHA guideline for the management of ST-elevation myocardial infarction: executive summary: A report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*, 2013;127:529–55.
5. Toma M, Buller CE, Westerhout CM, et al., for the APEX-AMI Investigators, Non-culprit coronary artery percutaneous

- coronary intervention during acute ST-segment elevation myocardial infarction: insights from the APEX-AMI trial, *Eur Heart J*, 2010;31:1701–7.
6. Dziewierz A, Siudak Z, Rakowski T, et al., Impact of multivessel coronary artery disease and noninfarct-related artery revascularization on outcome of patients with ST-elevation myocardial infarction transferred for primary percutaneous coronary intervention (from the EUROTRANSFER registry), *Am J Cardiol*, 2010;106:342–7.
  7. Xu F, Chen YG, Li JF, et al., Multivessel Percutaneous Coronary Intervention in Chinese Patients with Acute Myocardial Infarction and Simple Nonculprit Arteries, *Am J Med Sci*, 2007;333:376–80.
  8. Varani E, Balducci M, Aquilina M, et al., Single or Multivessel Percutaneous Coronary Intervention in ST-Elevation Myocardial Infarction Patients, *Catheter Cardiovasc Interv*, 2008;72:927–33.
  9. Roe MT, Cura FA, Joski PS, et al., Initial experience with multivessel percutaneous coronary intervention during mechanical reperfusion for acute myocardial infarction, *Am J Cardiol*, 2001;88:170–3.
  10. Cavender MA, Milford-Beland S, Roe MT, et al., Prevalence, predictors, and in-hospital outcomes of non-infarct artery intervention during primary percutaneous coronary intervention for ST-segment elevation myocardial infarction (from the National Cardiovascular Data Registry), *Am J Cardiol*, 2009;104:507–13.
  11. Iqbal MB, Ilesley C, Kabir T, et al., Culprit vessel versus multivessel intervention at the time of primary percutaneous coronary intervention in patients with ST-segment-elevation myocardial infarction and multivessel disease: real-world analysis of 3984 Patients in London, *Circ Cardiovasc Qual Outcomes*, 2014;7:936–43.
  12. Jin Z, Rha SW, Chen KY, et al., Culprit-lesion revascularization versus complete revascularization in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention with drug-eluting stents. Poster session presented at: Angioplasty Summit-TCT Asia Pacific; 2007 Apr 25–27; Seoul, South Korea.
  13. Karawani D, Nahir M, Abboud M, et al., Culprit only versus complete coronary revascularization during primary PCI, *Int J Cardiol*, 2008;123:288–92.
  14. Hannan EL, Samadashvili Z, Walford G, et al., Culprit vessel percutaneous coronary intervention versus multivessel and staged percutaneous coronary intervention for ST-segment elevation myocardial infarction patients with multivessel disease, *J Am Coll Cardiol Interv*, 2010;3:22–31.
  15. Khattab AA, Abdel-Wahab M, Röther C, et al., Multi-vessel stenting during primary percutaneous coronary intervention for acute myocardial infarction. A single-center experience, *Clin Res Cardiol*, 2008;97:32–8.
  16. Rahman M, Nfor T, Allaqaband S, et al., Clinical and angiographic outcomes in patients with ST-segment elevation myocardial infarction undergoing single versus multiple vessel percutaneous coronary intervention, *JACC*, 2010;55(10A):A98.E921.
  17. Corpus RA, House JA, Marso SP, et al., Multivessel percutaneous coronary intervention in patients with multivessel disease and acute myocardial infarction, *Am Heart J*, 2004;148:493–500.
  18. Kong JA, Chou ET, Minutello RM, et al., Safety of single versus multi-vessel angioplasty for patients with acute myocardial infarction and multi-vessel coronary artery disease: report from the New York State Angioplasty Registry, *Coron Artery Dis*, 2006;17:71–5.
  19. Seo J-S, Park D-W, Kim SS, et al., Long-term outcomes of culprit only versus complete revascularization during primary percutaneous coronary intervention, *J Am Coll Cardiol*, 2009;53(10s1):A1–A99.
  20. Estevez-Loureiro R, Rodriguez-Vilela A, Salgado-Fernandez J, et al., Effect of multivessel revascularization during primary percutaneous coronary intervention on outcomes of patients with ST-segment elevation myocardial infarction. Oral abstract presented at: 2010 Angioplasty Summit Transcatheter Cardiovascular Therapeutics Asia Pacific; 2010 Apr 28–29; Seoul, South Korea.
  21. Di Mario C, Mara S, Flavio A, et al., Single vs multivessel treatment during primary angioplasty: results of the multicentre randomised HEPacoat for culprit or multivessel stenting for Acute Myocardial Infarction (HELP AMI) study, *Int J Cardiovasc Interv*, 2004;6:128–33.
  22. Gershlick AH, The Complete versus Lesion only Primary PCI Trial. Available at: <http://www.escardio.org/about/press/esc-congress-2014/press-conferences/Documents/gershlick.pdf> (accessed 1 February 2015).
  23. Politi L, Sgura F, Rossi R, et al., A randomised trial of target-vessel versus multi-vessel revascularisation in ST-elevation myocardial infarction: major adverse cardiac events during long-term follow-up, *Heart*, 2010;96:662–7.
  24. Wald DS, Morris JK, Wald NJ, et al., for the PRAMI investigators, Randomized trial of preventive angioplasty in myocardial infarction, *N Engl J Med*, 2013;369:1115–23.
  25. Hochman JS, Steg PG, Does preventive PCI work?, *N Engl J Med*, 2007;356:15.
  26. Parisi AF, Folland ED, Hartigan P, A comparison of angioplasty with medical therapy in the treatment of single-vessel coronary artery disease, *N Engl J Med*, 1992;326:10–6.
  27. Boden WE, O'Rourke RA, Teo KK, et al., Optimal medical therapy with or without PCI for stable coronary disease, *N Engl J Med*, 2007;356:1503–16.
  28. Mehta SR, Cannon CP, Fox KA, et al., Routine vs selective invasive strategies in patients with acute coronary syndromes: a collaborative meta-analysis of randomized trials, *JAMA*, 2005;293:2908–17.
  29. Falk E, Shah PK, Coronary plaque disruption, *Circulation*, 1995;92:657–71.
  30. Davies MJ, Thomas A, Thrombosis and acute coronary-artery lesions in sudden cardiac ischemic death, *N Engl J Med*, 1984;310:1137–40.
  31. Schoenhagen P, Stone GW, Nissen SE, et al., Coronary plaque morphology and frequency of ulceration distant from culprit lesions in patients with unstable and stable presentation, *Arterioscler Thromb Vasc Biol*, 2003;23:1895–900.
  32. Rioufol G, Finet G, Ginon I, et al., Multiple atherosclerotic plaque rupture in acute coronary syndrome: a three-vessel intravascular ultrasound study, *Circulation*, 2002;106:804–8.
  33. Hong MK, Mintz GS, Lee CW, et al., Comparison of coronary plaque rupture between stable angina and acute myocardial infarction: a three-vessel intravascular ultrasound study in 235 patients, *Circulation*, 2004;110:928–33.
  34. Rioufol G, Gilard M, Finet G, et al., Evolution of spontaneous atherosclerotic plaque rupture with medical therapy: long-term follow-up with intravascular ultrasound, *Circulation*, 2004;110:2875–80.
  35. Goldstein JA, Demetriou D, Grines CL, et al., Multiple complex coronary plaques in patients with acute myocardial infarction, *N Engl J Med*, 2000;343:915–22.
  36. Meier B, Plaque sealing by coronary angioplasty, *Heart*, 2004;90:1395–8.
  37. The Task Force on Myocardial Revascularization of the European Society of Cardiology (ESC) and the European Association for Cardio-Thoracic Surgery (EACTS). 2014 ESC/EACTS Guidelines on myocardial revascularization, *European Heart Journal*, 2014;35:2541–619.
  38. Tonino PAL, De Bruyne B, Pijls NHJ, et al., Fractional flow reserve versus angiography for guiding percutaneous coronary intervention, *N Engl J Med*, 2009;360:213–24.
  39. Complete vs Culprit-only Revascularization to Treat Multivessel Disease After Primary PCI for STEMI (COMPLETE). Available at: <https://clinicaltrials.gov/ct2/show/NCT01740479?term=COMPLETE+culprit&rank=2> (accessed 1 February 2015).
  40. Primary PCI in patients with ST-elevation myocardial infarction and multivessel disease: Treatment of Culprit Lesion Only or Complete Revascularization (PRIMULTI). Available at: <https://clinicaltrials.gov/ct2/show/NCT01960933?term=COMPLETE+culprit&rank=5> (accessed 1 February 2015).
  41. Comparison Between FFR guided revascularization versus conventional strategy in acute STEMI patients with MVD. (CompareAcute). <https://clinicaltrials.gov/ct2/show/NCT01399736?term=COMPARE-acute&rank=1> (accessed 1 February 2015).