Published in final edited form as:

Mol Phylogenet Evol. 2018 April ; 121: 23–34. doi:10.1016/j.ympev.2017.12.015.

A Phylogenomic Analysis of Lichen-Feeding Tiger Moths Uncovers Evolutionary Origins of Host Chemical Sequestration

Clare H. Scott Chialvo^{1,*}, Pablo Chialvo², Jeffrey D. Holland³, Timothy J. Anderson³, Jesse W. Breinholt⁴, Akito Y. Kawahara⁴, Xin Zhou^{5,6}, Shanlin Liu^{7,8}, and Jennifer M. Zaspel³

¹Department of Biological Sciences, PO Box 870344, University of Alabama, Tuscaloosa, AL 35487, USA

²Department of Biology, 320 Stanley Avenue, Lander University, Greenwood, SC 29649, USA

³Department of Entomology, 901 West State Street, Purdue University, West Lafayette, IN 47907, USA

⁴Florida Museum of Natural History, 3215 Hull Road, University of Florida, Gainesville, FL 32611, USA

⁵Beijing Advanced Innovation Center for Food Nutrition and Human Health, China Agricultural University, Beijing 100193, People's Republic of China

⁶Department of Entomology, China Agricultural University, Beijing 100193, People's Republic of China

⁷China National GeneBank, 8/F, Beishan Industrial Zone, Yantian District, BGI-Shenzhen, People's Republic of China

⁸BGI-Shenzhen, Beishan Industrial Zone, Yantian District, China

Abstract

Host species utilize a variety of defenses to deter feeding, including secondary chemicals. Some phytophagous insects have evolved tolerance to these chemical defenses, and can sequester secondary defense compounds for use against their own predators and parasitoids. While numerous studies have examined plant-insect interactions, little is known about lichen-insect interactions. Our study focused on reconstructing the evolution of lichen phenolic sequestration in the tiger moth tribe Lithosiini (Lepidoptera: Erebidæ: Arctiinae), the most diverse lineage of lichen-feeding moths, with 3,000 described species. We built an RNA-Seq dataset and examined the adult metabolome for the presence of phenolics. Using the transcriptomic dataset, we recover a well-resolved phylogeny of the Lithosiini, and determine that the metabolomes within species are more similar than those among species. Results from an initial ancestral state reconstruction suggest that the ability to sequester phenolics produced by a single chemical pathway preceded

*Correspondence: Clare H. Scott Chialvo, Department of Biological Sciences, PO Box 870344, University of Alabama, Tuscaloosa, AL 35487, Fax: (205) 348-1786, clare.scott@ua.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

generalist sequestration of phenolics produced by multiple chemical pathways. We conclude that phenolics are consistently and selectively sequestered within Lithosiini. Furthermore, sequestration of compounds from a single chemical pathway may represent a synapomorphy of the tribe, and the ability to sequester phenolics produced by multiple pathways arose later. These findings expand on our understanding of the interactions between Lepidoptera and their lichen hosts.

Graphical abstract

Keywords

Metabolomics; Transcriptomics; Lichenivory; Ancestral State Reconstruction; Lichen Moths; Allelochemicals

1. Introduction

To defend against enemies, such as phytophagous insects and pathogens, plants and fungi have evolved a wide variety of defenses including secondary metabolites (allelochemicals (Fraenkel 1959)). Secondary metabolites act by making the host distasteful or toxic to herbivores. However, the impact of these chemical defenses can vary depending on the feeding strategy of an organism (*e.g.*, specialist or generalist (Reudler *et al.* 2011)). Unique and highly toxic allelochemicals (*e.g.*, cardiac glycosides) typically act to exclude non-specialized feeders that are unable to detoxify or sequester the compound (Becerra 1997; Krieger *et al.* 1971; Whittaker and Feeny 1971). Theoretically, generalist feeders evolve broad-spectrum mechanisms of detoxification that can be applied to an array of compounds traditionally thought of as less acutely toxic or “digestibility-reducing” (Ali and Agrawal 2012), while organisms that specialize on hosts containing highly toxic compounds can lose the ability to feed on many host plants (Whittaker and Feeny 1971). In addition to evolving tolerance to their host(s) allelochemicals, some phytophagous insects sequester the secondary metabolites for use as a chemical defense against their own predators and parasitoids (Opitz and Müller 2009).

With more than 157,000 described species (van Nieukerken *et al.* 2011), butterflies and moths (Lepidoptera) are the largest radiation of phytophagous animals on the planet (Scoble 1992), specializing on photosynthetic (vascular, non-vascular plants) and saprophytic

(macrofungi, lichens) food sources alike. Within the order, the tiger moths (Arctiinae), particularly the tiger moth tribe (Arctiini), are emerging as an evolutionary model for understanding the interactions of larval host chemistry, recruitment of detoxification enzymes (Hartmann 2004; Langel and Ober 2011), and the effect of stored host-derived compounds on predators (*e.g.*, bats; (Barber and Conner 2007; Corcoran *et al.* 2009; Hristov and Conner 2005)) and parasitoids (Bernays 2001; Singer *et al.* 2009; Singer *et al.* 2004). Larvae of some Arctiini sequester an array of secondary metabolites from their host plants including pyrrolizidine alkaloids, cardiac glycosides, and iridoid glycosides (Bowers and Stamp 1997; Hartmann *et al.* 2005; Rothschild *et al.* 1977; Rothschild *et al.* 1973). These compounds may be used for self-medication, as a chemical defense, or in chemical courtship (Conner 2009; Weller *et al.* 1999). Chemically defended Arctiini species advertise their chemical defenses and distastefulness using warning signals (aposematism) that are visual, acoustic, or a combination of both (Conner 2009; Corcoran *et al.* 2009). Furthermore, the ability to retain sequestered pyrrolizidine alkaloids from the larva to adult is a synapomorphy of Arctiini (Weller *et al.* 1999; Zaspel *et al.* 2014). Although numerous studies have examined plant-insect interactions (*e.g.*, chemical defense and larval feeding behavior) in the subfamily Arctiinae (Bowers and Stamp 1997; Conner 2009; Hartmann *et al.* 2005; Rothschild *et al.* 1977; Rothschild *et al.* 1973; Weller *et al.* 1999), very little is known about the lichen-insect interactions in the lichen moths, Lithosiini.

Lithosiini are a cosmopolitan group of approximately 3,000 described species best known for their larval feeding behavior, lichenivory (Rawlins 1984), which is thought to be ubiquitous among the described species. While the tribe is considered to be the most biodiverse lineage of lichenivores, some species primarily graze on algae (McCabe 1981). Like the more derived Arctiini, Lithosiini (Fig. 1) utilize chemical defenses against predators. Palatability studies that included lichen moth species found that the adults were distasteful to both bats and birds (Acharya and Fenton 1992; Collins and Watson 1983; Dowdy and Conner 2016; Hristov and Conner 2005), but did not include analyses to identify the source of their chemical defenses. Furthermore, unpalatable adult lichen moths appear to exhibit visual and acoustic aposematism, homologous to strategies in Arctiini (Dowdy and Conner 2016; Simmons 2008). For example, the lichen moth *Hypoprepia fucosa* is a visual mimic of cyanogen-sequestering fireflies (Coleoptera: Lampyridae) (*e.g.*, the adult color patterns of the moth resemble those of the fireflies; Forbes 1960), and *H. fucosa* is chemically defended with compounds that are as unpalatable to bats as non-lichen feeding tiger moths defended with cardiac glycosides, the most potent secondary metabolite sequestered by Arctiini (Hristov and Conner 2005). In addition, species in the lithosiine genus *Cisthene* engage in acoustic aposematism and are as unpalatable as *Pygarctia roseicapitis* (Dowdy and Conner 2016), an unrelated, non-lichenivorous arctiine that sequesters cardiac glycosides. While the chemical basis of unpalatability in adult Lithosiini has not been identified, it is hypothesized that the source of defense is phenolic compounds sequestered from the lichen host by lichen moth larvae (Hesbacher *et al.* 1995; Wagner *et al.* 2008). Thus, examining the adult metabolome not only provides a mechanism for identifying lichen derived phenolics that potentially contribute to unpalatability, it also serves as a means of assessing larval feeding behavior in the adult stage.

In two previous metabolomic studies (Hesbacher *et al.* 1995; Scott *et al.* 2014) that collectively examined 35 species of adult Lithosiini representing four of the seven lichen moth subtribes, lichen phenolics were identified in the metabolite profiles of 32 species. Both studies identified multiple phenolics in adults and found intraspecific variation in the compounds present. These phenolics are products of lichens, symbiotic organisms comprised of photosynthetic and fungal components. The fungal component of lichens produces numerous unique secondary metabolites (>1,000 metabolites identified; Stocker-Wörgötter (2008)) through three, primary chemical pathways (Rankovi and Kosani (2015); Fig. 2): acetate–polymalonate, shikimic acid, and mevalonic acid pathways. The most common phenolics (*e.g.*, depsides, depsidones, and dibenzofurans) are products of the acetate–polymalonate pathway (Rankovi and Kosani 2015), within which the majority of phenolics arise from the orcinol series through the oxidative coupling of small phenolic units related to orcinol and β -orcinol. The products of the orcinol series can be further modified to form other classes of phenolics including diphenyl ethers (Fig. 2). The phenolics produced by the three pathways are distributed throughout the layers of the lichen. They act to defend the photosynthetic component from the impacts of herbivores and solar radiation (Solhaug and Gauslaa 2012) and possess antimicrobial, anticancer, antiviral, and antioxidant properties (Rankovi and Kosani 2015). Lichen metabolites have been found to deter herbivory in gastropods and other invertebrates (Asplund 2011; Asplund *et al.* 2009), yet the precise mechanisms and ecological functions of lichen-invertebrate defenses are virtually unknown to science.

While metabolomic studies (Hesbacher *et al.* 1995; Scott *et al.* 2014) and the behavior of lichen moth adults (Dowdy and Conner 2016) suggest that lichenivory and subsequent phenolic sequestration is widespread within the tribe, the origins of this association and precise mechanisms of sequestration are currently unknown. In order to recover broad patterns of this historical relationship, a more comprehensive investigation of the compounds sequestered and the reliability with which they can be recovered both within biological replicates and across major clades is needed. For example, can the sequestered lichen phenolics be recovered reliably using metabolomic analyses of adult specimens stored under a range of conditions? Are chemical profiles consistent within and among closely related species, or do the metabolomes exhibit significant intraspecific variation? Given that lichens can contain a variety of phenolics derived from different chemical pathways, is lichen metabolite sequestration selective or more generalized (*i.e.*, do tiger moth larvae sequester lichen compounds produced by one or more chemical pathways)?

Our ability to examine the evolution of the relationship between these moths and their lichen hosts is hampered by the lack of a robust phylogenetic framework. While six separate phylogenetic studies (Jacobson and Weller 2002; Scott and Branham 2012; Scott *et al.* 2014; Wink and von Nikisch-Rosenegk 1997; Zahiri *et al.* 2012; Zaspel *et al.* 2014) recovered the monophyly of the lichen moth tribe using molecular or morphological data, they did not resolve relationships among tribes and genera. In the most recent study of Lithosiini, Scott *et al.* (2014) assessed the evolutionary relationships among 65 lichen moth species representing four of the subtribes using four gene fragments (*cytochrome oxidase C subunit I*, *cytochrome oxidase B*, *ribosomal protein S5*, and *nuclear large subunit rRNA 28S D2 loop*). Although these gene fragments had been shown to be phylogenetically informative at the

higher level in Lepidoptera and Noctuoidea (Mutanen *et al.* 2010; Wahlberg and Wheat 2008; Zahiri *et al.* 2012), they did not recover strong branch support for the deeper relationships within Lithosiini. Furthermore, the monophyly of each of the three subtribes represented by multiple taxa was not supported. Thus, there is need to develop a novel dataset that can provide a reliable evolutionary framework for further testing of the phylogenetic relationships within Lithosiini.

One of the main goals of the present study is to utilize a massively large transcriptomic dataset for lichen moths in order to establish the phylogenetic architecture of the group and combine it with data from a metabolomic analysis of lichen host compounds. With these data, we inferred a phylogeny of the tribe. We used hierarchical clustering and a permutational multivariate analysis of variance to assess the similarity of the overall phenolic profiles among and within species included in our metabolomic dataset. To test whether lichen moth clades exhibit a pattern of specialized versus generalized phenolic sequestration, we conducted an initial ancestral reconstruction of lichen chemical pathways known to produce compounds sequestered in the moth species we sampled. Our results provide strong support for the monophyly of two subtribes and demonstrate that closely related species in our sample consistently share chemical profiles regardless of specimen age or origin. Our analysis of chemical pathways provides initial evidence of chemical specialization vs generalization in this hyperdiverse moth lineage.

2. Materials and methods

2.1 Phylogenomic taxon sampling

Thirty-seven species, 27 from Arctiinae (22 Lithosiini, one Syntomiini, and four Arctiini), nine from the erebid subfamilies Aganainae, Calpinae, and Lymantriinae were sampled for the phylogenetic analysis. The 1K Insect Transcriptome Evolution Bioproject (1KITE; http://www.1kite.org/1kite_species.php; <https://www.ncbi.nlm.nih.gov/bioproject/183205>) provided the transcriptomes of three of the species sampled (*e.g. Phragmatobia fuliginosa*, *Amata phegea*, and *Phyllodes eyndhovii*). We selected these 37 species to test the monophyly of the tribe Lithosiini and its subtribes, as proposed by Bendib and Minet (1999), and the *Eugoa* group of Holloway (2001). The *Eugoa* group represents a generic grouping that was established to place genera that were not treated by the work of Bendib and Minet (1999) and includes genera found in the Old World Tropics, which is an area of high diversity. The 22 Lithosiini represented 19 genera, four of the seven subtribes (Cisthenina, Endrosina, Lithosiina, and Nudariina), and the *Eugoa* group (Table S1, supporting information). The sampling included 15 Old World (OW) and seven New World (NW) species (Table S1, supporting information). Diverse genera whose wing color patterns have caused taxonomic confusion (*e.g., Lyclene*) or have been found to be polyphyletic in previous studies (*e.g., Eilema*; (Scott and Branham 2012; Scott *et al.* 2014)) were represented by multiple species. Phylogenetic analyses were rooted using *Bombyx mori*, whose position well outside the Arctiinae is supported by many studies (Kawahara and Breinholt 2014; Mutanen *et al.* 2010; Regier *et al.* 2013; Regier *et al.* 2009).

2.1.1 RNA-Seq—We generated 33 novel transcriptomes for this study. These transcriptomes were produced from total RNA extracted from whole body flash-frozen or RNAlater-preserved adult specimens using either an RNeasy MiniKit (Qiagen) or a Trizol-Chloroform extraction. For *Haematomis uniformis*, total RNA was extracted from a larval specimen. Wing vouchers and collection location data for each specimen were deposited into the Purdue Entomological Research Collection. Extractions obtained using the Trizol – Chloroform protocol were purified with a DNA-Free RNA Kit (Zymo Research). Transcriptome data (ca. 2.5 GB raw data per sample) available from the 1KITE initiative was used for species representing Arctiina (*Phragmatobia fuliginosa*: BioProjectID: 267893, TSA: GCJS01000000), Syntomiini (*Amata phegea*: BioProjectID: 267859, TSA: GCNU01000000), and Phyllodini (*Phyllodes eyndhovii*: BioProjectID: 267894, TSA: GCOO01000000). Sequence data was obtained from the NCBI Short Read Archive for *Lymantria dispar* (Bazinet *et al.* 2013). The most recent proteome of *Bombyx mori* was obtained from Ensembl Genomes (Kersey *et al.* 2014).

For the 33 novel transcriptomes generated in this study, RNA was quantified using a Bioanalyzer 2100 (Agilent Technologies) at the Purdue University (PU) Genomics Core. RNA-Seq library construction and paired end sequencing on an Illumina HiSeq 2000 was also completed at the PU Genomics Core. Library construction was performed using the Illumina TruSeq Stranded Total RNA with the Ribo-Zero™ Gold Kit (epicentre). The libraries were barcoded and sequenced on five lanes (six or eight pooled samples in a lane). Adapters were removed and poor quality bases (Phred < 20) trimmed from the 5' and 3' ends of reads using Trimmomatic (Bolger *et al.* 2014). Following this processing, any reads of less than 30 bases were discarded. *De novo* assembly of the transcriptomes from these quality and adapter clipped reads was completed using Trinity (Grabherr *et al.* 2011). Trans-chimeras were removed from each of the novel transcriptomes using the pickH-cap-blast method that produces a dataset with fewer redundancies (Yang and Smith 2013). Coding regions of 100 amino acids or longer in the sequences were then determined using the TransDecoder plugin within Trinity. The new sequence data is available from GenBank (NCBI BioProject PRJNA312728; BioSamples SAM04569175-04569206). Data provided by 1KITE is available under the Umbrella BioProject PRJNA183205 (for full description of protocol used to sequence, assemble, and subsequently check for cross contamination in 1KITE transcriptomes see Appendix S1, supporting information).

2.1.2 Phylogenetic methods—Single copy ortholog groups were identified in OrthoMCL (Fischer *et al.* 2011) using the Basic Protocol 2 outlined in Fischer *et al.* (2011). In total, 1,328 single copy orthologous loci groups were identified that were present in more than 75% (28) of the taxa in the study. The sequences within these groups were aligned using the L-INS-i algorithm in MAFFT v.7 (Katoh and Standley 2013). After trimming the alignments using the 'automated1' setting in trimAl v1.4 (Capella-Gutiérrez *et al.* 2009), a consensus sequence for representative sequences of each ortholog group was extracted with EMBOSS v6.5.7 (Rice *et al.* 2000). To identify any contamination in the ortholog groups, the consensus sequence of each group was blasted against the NCBI non-redundant protein database using BLASTp, and the five most likely hits were retained along with the 'qseqid, qlen, sallseqid, sscinames, salltitles, pident, length, mismatch, gapopen, evalue, and bitscore'

for each hit. Groups with consensus sequences that did not align with any lepidopteran sequences or best aligned with non-invertebrate sequences were examined further. Individual sequences from these groups identified in the previous step were BLASTp searched against the non-redundant protein database to assess the identity using the above settings. All sequences from each of seven ortholog groups identified as prokaryotic or mammalian sequences were excluded using this method. We concatenated the remaining 1,321 orthologs into a supermatrix using FASconCAT (330,542 amino acids; Appendix S2, supporting information; Kück and Meusemann 2010). See Appendix S3 (supporting information) for a complete list of ortholog groups and their putative annotation. Table S2 (supporting information) provides a list of the coverage of each ortholog group in the species sampled. Substitution models and data partitions were determined using the r-cluster search strategy (rcluster-percent = 10; rclustermax = 1,000) implemented in PartitionFinder v2 with RAxML (Lanfear *et al.* 2017; Lanfear *et al.* 2014; Stamatakis 2014). All 128 models of amino acid evolution were considered using the ‘-raxml’ option. Following the program recommendation for analyses that will use RAxML, two additional searches were completed in PartitionFinder to identify the optimal model of rate heterogeneity (+I+G or +G) for all partitions (*i.e.*, the model of rate heterogeneity that produces the lowest BIC score when applied to all partitions). These two runs were completed using the best partitioning scheme identified in the first PartitionFinder analysis. See Appendix S4 (supporting information) for the partitioning scheme used and the model of evolution applied to each partition. An initial ML analysis was conducted in RAxML – HPC2 (Stamatakis 2006) on the CIPRES Science Gateway (Miller *et al.* 2010) using the ‘-f a’ option with 1,000 bootstrap replicates. We ran an additional 100 searches for the best likelihood tree in RAxML using the ‘-f d’ option and mapped bootstrap support values onto the best tree using the ‘-f b’ option and the previously obtained bootstrap replicates. The tree was rooted using *Bombyx mori*.

2.2 Metabolomic taxon sampling

Fifteen species were sampled from OW and NW taxa to represent the four subtribes of Bendib and Minet (1999) (Cisthenina, Eudesmiina, Lithosiina, and Nudariina; Table S1, supporting information; Fig. 1). When possible, taxa that were sampled for the phylogenomic analyses were included. Eleven of the 22 Lithosiini included in the phylogenomic study were sampled. These species were also chosen to exemplify the variation of wing color patterns found in the tribe. Both brightly colored species and drably colored species were assessed. When material was available, three biological replicates were sampled for each species.

2.2.1 Metabolomic analysis—Fifteen species of adult Lithosiini were assessed for the presence of 129 lichen phenolics (Appendix S5, supporting information). Metabolites were obtained using the entire body of each specimen. Dorsal and ventral photographs of these specimens and their labels were taken as vouchers and deposited within the Purdue Entomological Research Collection. The specimens were homogenized using a Precellys® 24 tissue homogenizer. Metabolites were obtained from the homogenate using a methanol-water: chloroform extraction (1.5ml methanol; 0.5ml water; 1ml chloroform). The targeted phenolics are polar compounds; therefore, only the methanol-water phase of the extraction, which would contain polar compounds, was used in the final analysis. After the phases

separated, the methanol-water phase was removed with a pipette and dried by spinning at room temperature for 14-16 hours in a Savant SpeedVac Concentrator (Thermoscientific). The concentrated pellets of metabolites were resuspended in 50µl of a 95% water : 5% acetonitrile solution.

The LC-MS analysis of these metabolites was performed with an Agilent 1100 system (Palo Alto, CA) using a Waters T3 column (3 µm, 150 × 2.1 mm i.d) and an Agilent MSD-TOF spectrometer following the methodology of Scott *et al.* (2014). These analyses were conducted in the Metabolite Profiling Facility within the Bindley Bioscience Center at Purdue University. Reads were analyzed using Agilent MassHunter Qualitative Analysis software. For each sampled species, this software was used to compare the peaks that were identified to determine differences in quantity and types of compound present (Appendix S6, supporting information). All replicates from each species were then assessed for 129 lichen phenolics, which have previously been reported to exhibit pharmaceutical activity and have a known chemical formula. Any peak with a mass that differed by less than 10 ppm from that of a given phenolic was considered a presumptive positive. Commercially available, chemical standards (*e.g.*, pure compounds) for five of the candidate phenolics (resorcinol, β -thujone, divarin, didymic acid, and methyl β -orcinolcarboxylate) were also analyzed at the Metabolite Profiling Facility using the same conditions as the biological samples.

2.2.2 Clustering analysis of phenolic compounds and ancestral state reconstruction

—Among tiger moths, lichen phenolic sequestration is only known from species in Lithosiini. To date, sequestered lichen-derived metabolites have been documented in five of the seven subtribes (Hesbacher *et al.* 1995; Scott *et al.* 2014). Our metabolomic examination of 15 of 22 Lithosiini species (68% of ingroup sampling) included in the phylogenetic dataset identified presumptive positives for 33 unique lichen phenolics. To further examine this dataset, and test whether replicates within species shared similar patterns of chemical sequestration, we completed a hierarchical clustering analysis that grouped individuals based only on their phenolic profiles. The presence/absence of the compounds was used to calculate the Jaccard's dissimilarity (Legendre and Legendre 1998) between chemical profiles of individuals. This asymmetric association measure works well with presence/absence data when double negatives are not considered to represent similarity. We clustered the profiles using Ward's clustering algorithm (Murtah and Legendre 2014). To determine the pruning height, and thus cluster delineation, we subjected the same data to k-means clustering with k varying 1–20. The resulting curve of within-group error sum of squares versus k showed a weak elbow at k=8. We used this pruning height to assign individuals to eight chemical profile groups. We used a permutational multivariate analysis of variance (adonis) to test for a significant different among the chemical profiles. Cluster and adonis analyses were carried out with adonis using the 'vegan' package (Oksanen *et al.* 2015) in R (R Core Team 2014).

Due to the high diversity of compounds recovered in a relatively small number of taxa sampled for our phylogeny, coupled with the unavailability of chemical standards for a large majority of phenolic compounds (Betz *et al.* 2011), we took a more conservative and biologically meaningful approach, and coded terminal taxa according chemical pathways instead of individual compound. The chemical pathways are based the groupings identified

and described by lichen biologists (Culberson 1969; Rankovi and Kosani 2015). A denser taxon sampling and comprehensive testing against chemical standards will be important in future comparative work, but beyond the scope of this initial study. The chemical pathways are A-F: (A) “orcinol” (phenolics that are members of the orcinol pathway including precursors, monocyclic derivatives, *para*-depsides, depsidones, dibenzofurans, depsones, and *meta*-depsides); (B) “aliphatic acids”; (C) “anthraquinones”; (D) “chromones”; (E) “diphenyl ethers”; (F) “mono-terpenes” (Appendix S7, supporting information).

Published feeding records for tiger moths suggest patterns of chemical acquisition and sequestration are generally conserved at the genus level (Conner and Jordan 2009); our clustering analysis of individual phenolic compounds by taxon provide additional support for this claim. Therefore, in cases where a chemical profile for a given species were absent, available profiles for congeners were used (approach as in Hwang and Weirauch 2012; Zaspel *et al.* 2014). Chemical pathway profiles of ancestral nodes were reconstructed using a Bayesian approach. We reconstructed ancestral areas using the Bayesian Binary MCMC ancestral state reconstruction (BBM) method implemented in RASP (Yu *et al.* 2015). The analysis was run for 5,000,000 cycles using 4 chains, which were sampled every 100 generations; the first 1000 iterations were discarded as burn-in, and the Fixed Jukes-Cantor model was used with a null root distribution.

3. Results

3.1 Phylogenetic analysis

To examine the evolutionary relationships among lithosiine species, we constructed a dataset of single copy orthologs identified from the transcriptomic data of 22 lithosiine species, 14 non-lithosiine erebid outgroups, and rooted the tree with *Bombyx mori* (Bombycidae). The final dataset contained 1,321 single copy orthologs (330,542 amino acids). A maximum likelihood (ML) analysis of the partitioned, dataset recovered a well-supported tree with all but one node receiving non-parametric bootstrap (BS) support score of greater than 80% (Fig. 3; See Appendix S4 (supporting information) for partitioning scheme used and substitution model applied to each partition).

The outgroup taxon sampling included four erebid subfamilies (Arctiinae, Aganainae, Calpinae and Lymantriinae). The analysis recovered Lymantriinae as the most basal subfamily. Calpinae was reciprocally monophyletic with BS = 100 to Aganainae + Arctiinae (BS=100). Within Arctiinae, the ML analysis recovered the monophyletic Lithosiini (BS=100) sister to Arctiini + Syntomiini (BS=100).

Within the Lithosiini, we included species representing four of the seven subtribes and the *Eugoa* species group. The ML analysis recovered the subtribes Nudariina and Lithosiina as well-supported, monophyletic clades (both BS=100; Fig. 3). Cisthenina was polyphyletic: the OW Cisthenina species are recovered as a monophyletic clade (BS=100) at the base of Lithosiini (BS=100). The NW Cisthenina species were recovered in a more derived position (BS=100) as the sister group to the OW Nudariina clade (BS>87). The ML analysis found the single representative of the subtribe Endrosina (*Stigmatophora roseivena*) as the sister group to *Trischalis subaurana* (BS=100) a representative of the *Eugoa* group (BS=100),

which was represented by two species. The Lithosiina are sister to the *Eugoa* group/Endrosina clade (BS>81).

We also included multiple representatives of three diverse Lithosiini genera associated with the subtribes Lithosiina and Nudariina, which previous studies (Scott and Branham 2012; Scott *et al.* 2014; Zaspel *et al.* 2014) found not to be monophyletic: *Eilema*, *Lyclene*, and *Miltochrista*. Although representatives of the latter two genera were both confined as Nudariina in our ML analysis, only *Lyclene* was recovered as monophyletic (BS = 100). The ML analysis recovered a polyphyletic *Miltochrista* within Nudariina, and a polyphyletic *Eilema* occurring in Lithosiina and Nudariina.

3.2 Metabolomic analysis

The metabolomic analysis of 15 lichen moth species (Fig. 1) revealed presumptive positives for 33 compounds representing 10 classes of lichen phenolics derived from two of the major metabolic pathways that produce phenolics (Fig. 4; Table S3, supporting information). When possible, three biological replicates were included for each species to assess for intraspecific variation due to preservation techniques and possible host switching by larvae. We also analyzed the spectra of commercially available standards of five of the compounds: resorcinol, β -thujone, divarin, didymic acid, and methyl β -orcinolcarboxylate. The retention times of these standards were compared to those obtained for the presumptive positives; however, the retention times of the standards did not match those found in the moths. Thus, we limit the rest of our results from the metabolomic analysis to the remaining 28 phenolics.

Of the remaining 28 compounds identified in this study, 17 are precursors, products, or monocyclic derivatives (*i.e.*, breakdown products) of the orcinol series. In addition to being the most common type of phenolic class, the orcinol series is also the only class of phenolics present in all of the sampled species. With the exception of the aliphatic acids and anthraquinones, we did not identify multiple presumptive positives for any other phenolic compound class. The most commonly identified phenolics were anziac acid (orcinol series: *para*-depside) and micareic acid (diphenyl ether). We identified presumptive positives for each of these phenolics in the metabolomic profiles of six species. Eleven of the phenolic compounds identified were unique to a single species.

In addition to identifying a wide range of phenolic compounds, our metabolic analysis identified interspecific variation in the phenolics present in the sampled species (Table S3, supporting information). The species with the highest number of phenolics identified, *Cyana meyricki* (13; Fig. 1B) represents the subtribe Nudariina, which has the highest, per-species average number of phenolics ($M=8.23\pm3.77$). The species with the lowest number of phenolics is *Eilema bicolor* (2; Fig. 1N), a member of Lithosiina. We identified six or more phenolics in three Cisthenina species known to be unpalatable to bats and birds (*Cisthene martini*, *C. tenuifascia*, and *Hypoprepia fucosa*; Fig. 1E-G; (Dowdy and Conner 2016; Hristov and Conner 2005)). The metabolomic profile of the firefly mimic, *Rhabdatomis laudamia* contained presumptive positives of seven phenolics.

Based on prior studies (Hesbacher *et al.* 1995; Scott *et al.* 2014), we hypothesized that lichen moths selectively sequester phenolics from specific chemical classes, but the phenolics

sequestered can vary intraspecifically. To examine the pattern of overall phenolic sequestration among and within our species, and to test our hypothesis of specialized chemical sequestration, we used a hierarchical clustering analysis to group individuals based solely on their phenolic profiles. We found that the chemical profile clustering grouped individuals within ten species with conspecifics, while individuals of *Cisthene martini*, *Eilema plana*, *Euryptidia ira*, *Hypoprepia fucosa*, and *Nishada sambara* were spread among more than one group (Fig. 5A). In all clusters except Cluster 3, species from the same geographic region were grouped together. Furthermore, the phenolic profiles of the different clusters were significantly different (adonis: df=8,33, F=7.32, p-value=0.0001) as shown in the NMDS plot (Fig. 5B).

3.2.1 Evolution of Phenolic Sequestration—A representation of the most likely states from the BBM of chemical profiles for Lithosiini is presented in Fig. 6. Outgroup taxa included in our study are not lichenivores, and are not known to sequester lichen-derived compounds. Therefore, we only report ancestral reconstructions for clades within Lithosiini. The ancestral state for the Lithosiini node could not be reconstructed. This ambiguity is most likely due to a lack of metabolic profile data (e.g., phenolics sequestered) in the earlier lithosiine lineages. However, while the most probable ancestral state at the node uniting all Lithosiini except the OW Cisthenina could not be reconstructed (60.87%), sequestration of orcinol series-associated compounds was the next most probable state (25.28%). We also reconstructed a single origin of orcinol series-associated lichen compounds (40.74%) at the node uniting Nudariina and NW Cisthenina (P=0.20). Although the ancestral state reconstructed at this position represents an origin of phenolic sequestration for compounds in the orcinol series, the chemical profiles obtained for the subtribes (and species) therein contain a diverse array of phenolics. For example, 20 of the 33 lichen-derived compounds reported were recovered in the Nudariina, and 18 were detected in the NW Cisthenina. Within the NW Cisthenina, support at the node uniting *Hypoprepia* and *Ptychoglène* in this reconstruction was relatively strong (P=0.79). A stark transition from primarily orcinol associated compound sequestration (i.e., specialized) to a more “generalized” pattern of lichen derivative sequestration (i.e., sequestration of lichen products arising from more than one chemical pathway) was observed in the common ancestor of Nudariina. Although not strongly supported (P=0.35), the putative ancestor of Nudariina is inferred to have sequestered a broad range of lichen compounds. The novel character state reconstruction for the putative ancestor of Nudariina included orcinol compounds, diphenyl ethers, and mono-terpenes (69.68%); these lichen phenolics were particularly prevalent in the OW genera *Lyclene* (P=0.69) and *Miltochrista* (P=0.98). There was no definitive ancestral reconstruction for the *Eugoa* group or subtribe Lithosiina, again, perhaps due to missing chemical profile data in those terminal taxa. However, among the Lithosiina, *Crambidia* and *Eilema* species both tested positive for compounds associated with the orcinol series resulting in strong support (P=0.86) for reconstruction of this state at the node uniting the two genera. Our results yielded weak support (P=0.05) for the reconstruction of orcinol-associated compounds at the node joining genera *Macotasa*, *Nishada*, and *Teulisna*. This state was recovered in 21.61% of the topologies, and the next most probable state was sequestration of both orcinol-associated compounds and diphenyl ethers (18.44%).

Sequestration of diphenyl ether compounds (21.11%) was reconstructed with weak to moderate support ($P=0.21$) for the common ancestor of *Nishada* and *Teulisna*.

4. Discussion

Elucidating the deeper evolutionary relationships within Lithosiini has proven to be a difficult problem. This inhibits our ability to use this lineage as a model system for understanding the relationships between Lepidoptera and lichens. By analyzing a phylotranscriptomic dataset, we are able to recover strong support for the early relationships within the lichen moths. These findings provide the first evidence that the chemical profiles within species and among closely related species are more similar to each other than expected by chance, and species sequester specific classes of phenolics. Furthermore, our initial metabolomic examination and reconstruction of chemical pathways support the idea that lichen phenolics are consistently, and perhaps even selectively, sequestered in Lithosiini.

4.1 Evolution of lichen moths

Previous phylogenetic studies of the evolutionary relationships of Lithosiini used either adult morphological characters (Scott and Branham 2012) or molecular datasets (Scott *et al.* 2014) that could not resolve the deeper relationships of the lithosiine genera or subtribes. However, transcriptome sequencing provides a means to obtain large quantities of data even for non-model organisms (Metzker 2010). Transcriptome-based datasets have been found to be informative in resolving the deeper relationships at a range of taxonomic levels and a variety of organisms (Hittinger *et al.* 2010; Misof *et al.* 2014; Wen *et al.* 2013). Our study is the first to use a phylogenomic (*e.g.*, transcriptome-based) dataset to reconstruct the evolutionary history of lichenivory and sequestration of lichen based secondary metabolites. The ML analysis of Lithosiini transcriptomes resulted in 20 of the 21 internal relationships that are well supported, adding to a growing body of evidence that transcriptome based datasets are a powerful approach to resolve previously intractable phylogenetic questions, and confirming other studies that have used transcriptomes for lepidopteran phylogenetics (*e.g.*, Kawahara and Breinholt 2014, Breinholt and Kawahara 2013, and Bazinet *et al.* 2013).

Among the four subfamilies represented in our taxon sampling for the phylogenomic analysis, the most likely topology supported the monophyly of all subfamilies. Relationships recovered among these subfamilies are congruent with those recovered in recent phylogenetic analyses of Erebiidae (Zahiri *et al.* 2012) and Arctiinae (Zaspel *et al.* 2014). Within Arctiinae, the topology we recovered provides further support for the hypothesis that Lithosiini is basal to the Arctiini + Syntomiini clade (Zahiri *et al.* 2012; Zaspel *et al.* 2014).

Our study provides a framework phylogeny that supports some of the previously proposed subtribal classifications of Lithosiini (Bendib and Minet 1999; Holloway 2001) and also highlights groups that will need further investigation. Our results support the subtribal hypotheses of Bendib and Minet (1999) for two lineages (Lithosiina and Nudariina). However, we found the subtribe Cisthenina to be polyphyletic. As currently defined, Cisthenina contains OW and NW species (Bendib and Minet 1999; Holloway 2001), and our analysis included representatives of both geographic regions. In our result, the two, non-adjacent clades representing Cisthenina are composed exclusively of only OW or NW taxa.

Scott *et al.* (2014) also recovered a polyphyletic Cisthenina represented by two clades that contained only NW taxa. These findings suggest that as currently defined the taxa placed within Cisthenina could represent three independent lineages. Thus, an intensive study is needed to determine the composition of Cisthenina and what morphological characters may reliably diagnose taxa to this subtribe. The fourth subtribe, Endrosina, is found in the clade containing the two representatives of the *Eugoa* group of Holloway (2001).

We included multiple representatives of three diverse genera (*e.g.*, *Eilema*, *Lyclene*, and *Miltochrista*). Each of these genera is composed of a large number of species with similar wing color patterns (within each genus) that have caused confusion and led to previous studies identifying them as either paraphyletic (*e.g.*, *Miltochrista* (Scott and Branham 2012)) or polyphyletic (*e.g.*, *Eilema* (Scott and Branham 2012; Scott *et al.* 2014), *Lyclene* (Scott *et al.* 2014; Zaspel *et al.* 2014), *Miltochrista* (Zaspel *et al.* 2014)). Our analyses recovered only *Lyclene* as monophyletic. We found all representatives of *Miltochrista* placed in the Nudariina, and representatives of *Eilema* in both the Lithosiini and Nudariina clades. The species *Eilema plana*, which was found in the Nudariina clade, was also recovered in Nudariina by Scott *et al.* (2014). Our findings suggest that monophyletic groups might be recovered from these large genera; however, an in-depth analysis of these lineages will be needed to determine their exact composition.

4.2 Phenolic sequestration

Metabolomic analysis demonstrated that sequestration of lichen phenolics in the larval stage and successful retention of these compounds through diapause to the adult stage occurs in four of the lithosiine subtribes: Cisthenina, Eudesmiina, Lithosiina, and Nudariina. Hesbacher *et al.* (1995) identified lichen phenolics in the metabolome of species representing Endrosina. However, to our knowledge, no metabolomic studies examined the chemical profiles of the remaining two subtribes (Acsalina and Phryganopterygina). We identified presumptive positives for 33 unique phenolics in the Lithosiini sampled; pyrrolizidine alkaloids, cardiac glycosides, iridoid glycosides, or tetrahydrocannabinol were not found in the metabolome of any species included in our analysis. While Scott *et al.* (2014) identified physodic acid in the metabolic profile of all sampled species, we did not find any peaks whose mass was within 10 ppm to this phenolic. Thus, the more stringent protocol for identifying presumptive positives utilized in this study decreases the probability of false positive. Genera within Arctiini sequester these chemicals for defense against predators and parasitoids (Bowers and Stamp 1997; Hartmann *et al.* 2005; Rothschild *et al.* 1977; Rothschild *et al.* 1973). Untargeted metabolic analyses such as the ones completed for this study generate large numbers of unknown metabolites (Goodacre *et al.* 2004). Preliminary identifications of unknown peaks can be made based on the mass of compounds of interest, and then these identifications should be confirmed using chemical standards for each unknown compound. Our spectral analysis of chemical standards representing five of the compounds identified with presumptive positives did not confirm their identities with matching retention times. However, due to a limited number of chemical standards for botanical products (Bowen and Northern 2010) and a lack of available databases, the best option for establishing a preliminary identification of phenolics is to use their mass. Furthermore, retention times can vary between analyses due to experimental drift (Patti *et al.*

2012). The number of samples run in an analysis or running samples at different times on the same machinery can cause this variation. Thus, the difference in retention time between the peaks identified as presumptive positives and the chemical standards could be due in part to this phenomenon. In addition, the matrices in which botanical compounds, such as lichen phenolics, occur are by nature highly variable (Betz *et al.* 2011), which can also contribute to differences in retention times and make chemical confirmation of them using standards difficult. To confirm the presence of the remaining phenolics identified, analyses of additional standards will be needed. For many of these compounds, chemical standards are not commercially available. Thus, absolute certainty of the compound identities is limited and will require more studies and the use of additional techniques, such as H^+ NMR. Studies such as this add to a growing body of literature regarding the association between lithosiines and lichens and can be used to guide future studies examining questions related to phenolic sequestration.

Our examination of the metabolomes combined with those of previous studies (Hesbacher *et al.* 1995; Scott *et al.* 2014) found intraspecific variation in the phenolic profiles of adult lichen moths. However, our clustering analysis demonstrated that within the majority of species, profiles are more similar within than among species. Thus, our findings demonstrate proof of concept that the use of adult specimens preserved in a variety of conditions and LC-MS provides a potentially reliable means of confirming lichen compound sequestration in the absence of known or observed lichenivory. Given the consistency of the chemical adult profiles and the number of presumptive positives identified, we reconstructed the evolution of sequestration by pathway, not compound. This protocol allowed us to look for broader patterns of evolution within the context of chemical specialization vs generalization. Since lichen host chemistry can vary considerably within and among species depending on environmental conditions (Betz *et al.* 2011) but the chemical pathways do not, a reconstruction of chemical pathways is potentially more biologically meaningful than by individual compound.

Our initial reconstruction of the origins of lichen phenolic sequestration demonstrated that the ability to selectively sequester phenolics from the orcinol series is the ancestral state for the NW Cisthenina + Nudariina clade and some members of Lithosiina as currently defined (Fig. 6). While this analysis could not resolve whether sequestration of orcinol series phenolics occurs in the more basal OW Cisthenina clade or in the *Eugoa* group/Endrosina, these taxa have not been sampled in any metabolomic studies. Thus, the ability to selectively sequester phenolics from the orcinol series is a strong candidate for a metabolic synapomorphy of Lithosiini.

Regardless of whether this is the case, results from our analyses indicated generalized sequestration of phenolics produced by more than one chemical pathway represents the derived condition. Species of Nudariina sequester compounds from two of the three main metabolic pathways (Figs. 2, 4 and 6) that produce phenolics: acetate-palmitate and mevalonic acid pathway. It is possible that generalist sequestration of phenolics would allow the species in Nudariina to utilize a broader range of host lichens. A similar pattern of evolution of “specialist to generalist” feeding behaviors was also supported in the derived

tiger moth clades (Zaspel *et al.* 2014), yet whether the detoxification and sequestration systems of herbivorous vs. lichenivorous tiger moths are similar, remains unknown.

In all species included in our metabolomic analysis, we found presumptive positives for lichen phenolics derived from the orcinol series. This result holds both for brightly colored, presumptively aposematic lineages (*Cisthene* spp. and *Ptychoglene phrada*) and for firefly mimics (*e.g.*, *Rhabdatomis laudamia*). The orcinol series refers to lichen phenolics that are produced by the acetate-polymalonate pathway and occur as two types: orcinol and β -orcinol (Rankovi and Kosani 2015). Phenolics from the orcinol series have antibacterial properties (Cheng *et al.* 2013) and cause contact dermatitis in humans. Hesbacher *et al.* (1995) and Wagner *et al.* (2008) hypothesized that sequestered lichen phenolics may be used by Lithosiini as a defense against predators and/or pathogens. As such, phenolics from the orcinol series could provide the chemical basis for the observed unpalatability in adult lithosiines.

These results suggest that lichen feeding is common and lithosiine larvae may selectively forage on a broad range of lichens, potentially making use of a variety of chemical defenses. We recognize the relatively small sample size used in our reconstruction analysis, but we have selected exemplar taxa that represent the major lithosiine lineages across their geographic ranges. Our approach to reconstructing the states is relatively basic, yet it provides a way to interpret the data in a phylogenetic context. Additionally, the results of our clustering analyses (Fig. 5) provide strong evidence that our chemical data is consistent and reliable both within and among the related taxa, thus even with an increased sampling, we predict close relatives will share similar chemical profiles. While further studies will be needed to determine what compounds are sequestered among early lithosiines, and to confirm whether the phenolics identified effectively deter predators and parasitoids, these findings help to broadly expand upon the poorly understood interactions between Lepidoptera and lichens.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding and facilities for this study were provided by Purdue University, National Science Foundation (DEB-0919185 to JZ and Susan Weller (UN-Lincoln)), and National Institutes of Health (R01 GM098856 to Laura Reed (University of Alabama)). John Rawlins (Carnegie Museum of Natural History) provided the two *Euryptidia ira* specimens in the metabolomics study. Bruce Cooper (Bindley Science Center) and Arun Seetharam (PU Bioinformatics Core) provided training in LC-MS and bioinformatic analysis respectively. Ortholog group identification and partitioning analyses were completed on the Georgia Advanced Computing Resource Center, and phylogenetic analyses were conducted using the CIPRES science gateway. The 1KITE community provided assistance in obtaining 1KITE transcriptomes – we acknowledge in particular Bernhard Misof, Karl M. Kjer, Alexander Donath (sequence submission), Lars Podsiadlowski (removal of contaminants), Ralph S. Peters, Carola Greve, Sven Nekum (collecting) and Karen Meusemann (collecting and data transfer). Adam Bazinet (UMD), and Charles Mitter (UMD) provided additional transcriptomes to expand the taxon sampling in the final analysis. Susan Weller provided guidance on taxon sampling.

References

- Acharya L, Fenton MB. Echolocation behavior of vertebrate bats (*Lasiurus cinereus* and *Lasiurus borealis*) attacking airborne targets including arctiid moths. *Canadian Journal of Zoology*. 1992; 70:1292–1298.
- Ali JG, Agrawal AA. Specialist versus generalist insect herbivores and plant defense. *Trends in Plant Science*. 2012; 17:293–302. [PubMed: 22425020]
- Asplund J. Snails avoid the medulla of *Lobaria pulmonaria* and *L. scrobiculata* due to the presence of secondary compounds. *Fungal Ecology*. 2011; 4:356–358.
- Asplund J, Solhaug KA, Gauslaa Y. Fungal depsidones – an inducible of constitutive defence against herbivores in the lichen *Lobaria pulmonaria*? *Basic and Applied Ecology*. 2009; 10:273–278.
- Barber JR, Conner WE. Acoustic mimicry in a predator-prey interaction. *Proceedings of the National Academy of Science USA*. 2007; 104:9331–9334.
- Bazin AL, Cummings MP, Mitter KT, Mitter CW. Can RNA-seq resolve the rapid radiation of advanced moths and butterflies (Hexapoda: Lepidoptera: Apoditrysia)? An exploratory study. *PLoS ONE*. 2013; 8:e82615. [PubMed: 24324810]
- Becerra JX. Insects on plants: macroevolutionary chemical trends in host use. *Science*. 1997; 276:253–256. [PubMed: 9092474]
- Bendib A, Minet J. Lithosiine main lineages and their possible interrelationships. I. -Definition of new or resurrected tribes (Lepidoptera: Arctiidae). *Annales de la Société Entomologique de France*. 1999; 35:241–263.
- Bernays EA. Neural limitations in phytophagous insects: Implications for diet breadth and evolution of host affiliation. *Annual Review of Entomology*. 2001; 46:703–727.
- Betz JM, Brown PN, Roman MC. Accuracy, precision, and reliability of chemical measurements in natural products research. *Fitoterapia*. 2011; 82:44–52. [PubMed: 20884340]
- Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*. 2014; btu170
- Bowen BP, Northern TR. Dealing with the unknown: metabolomics and metabolite atlases. *Journal of the American Society for Mass Spectrometry*. 2010; 21:1471–1476. [PubMed: 20452782]
- Bowers MD, Stamp NE. Fate of host-plant iridoid glycosides in lepidopteran larvae of Nymphalidae and Arctiidae. *Journal of Chemical Ecology*. 1997; 23:2955–2965.
- Breinholt JW, Kawahara AY. Phylotranscriptomics: saturated third codon positions radically influence the estimation of trees based on next-gen data. *Genome Biology and Evolution*. 2013; 5:2082–2092. [PubMed: 24148944]
- Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. 2009; 25:1972–1973. [PubMed: 19505945]
- Cheng B, Cao S, Vasquez V, Annamalai T, Tamayo-Castillo G, Clardy J, Tse-Dinh Y. Identification of anziaic acid, a lichen depside from *Hypotrachyna* sp., as a new topoisomerase poison inhibitor. *PLoS ONE*. 2013; 8:e60770. [PubMed: 23593306]
- Collins CT, Watson A. Field observations of bird predation on neotropical moths. *Biotropica*. 1983; 15:53–60.
- Conner, WE. *Tiger Moths and Woolly Bears: Behavior, Ecology, and Evolution of the Arctiidae*. Oxford University Press; New York, NY: 2009. p. 303
- Conner, WE., Jordon, AT. From armaments to ornaments: the relationship between chemical defense and sex in tiger moths. In: Conner, WE., editor. *Tiger Moths and Woolly Bears: Behavior, Ecology, and Evolution of the Arctiidae*. Oxford University Press; New York, NY: 2009. p. 155-172.
- Corcoran AJ, Barber JR, Conner WE. Tiger moth jams bat sonar. *Science*. 2009; 325:325–327. [PubMed: 19608920]
- Culbertson, CF. *Chemical and botanical guide to lichen products*. The University of North Carolina Press; Chapel Hill, NC: 1969.
- Dowdy NJ, Conner WE. Acoustic Aposematism and Evasive Action in Select Chemically Defended Arctiine (Lepidoptera: Erebidae) Species: Nonchalant or Not? *PLoS ONE*. 2016; 11:e0152981. [PubMed: 27096408]

- Fischer S, Brunk BP, Chen F, Gao X, Harb OS, Iodice JB, Shanmugam D, Roos DS, Stoeckert CJ Jr. Using OrthoMCL to assign proteins to OrthoMCL-DB groups or to cluster proteomes into new ortholog groups. *Current Protocols in Bioinformatics*. 2011; 35:6.12.11–16.12.19.
- Forbes, WTM. The Lepidoptera of New York and neighboring states, part 4: Agaristidae through Nymphalidae including butterflies. Cornell University Agriculture Experiment Station Memoir 371, Cornell University; Ithaca, NY: 1960.
- Fraenkel GS. The raison d'être of secondary plant substances. *Science*. 1959; 129:1466–1470. [PubMed: 13658975]
- Goodacre R, Vaidyanathan S, Dunn WB, Harrigan GG, Kell DB. Metabolomics by numbers: acquiring and understanding global metabolite data. *Trends in Biotechnology*. 2004; 22:245–252. [PubMed: 15109811]
- Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, Adiconis X, Fan L, Raychowdhury R, Zeng Q, Chen Z, Mauceli E, Hacohen N, Gnirke A, Rhind N, di Palma F, Birren BW, Nusbaum C, Lindblad-Toh K, Friedman N, Regev A. Full-length transcriptome assembly from RNA-seq data without a reference genome. *Nature Biotechnology*. 2011; 29:644–652.
- Hartmann T. Plant-derived secondary metabolites as defensive chemicals in herbivorous insects: A case study in chemical ecology. *Planta*. 2004; 219:1–4. [PubMed: 15042370]
- Hartmann T, Theuring C, Beuerle T, Bernays EA, Singer MS. Acquisition, transformation and maintenance of plant pyrrolizidine alkaloids by the polyphagous arctiid *Grammia geneura*. *Insect Biochemistry and Molecular Biology*. 2005; 35:1083–1099. [PubMed: 16102415]
- Hesbacher S, Giez I, Embacher G, Fiedler K, Max W, Trawöger A, Türk R, Lange OT, Proksch P. Sequestration of lichen compounds by lichen-feeding members of the Arctiidae (Lepidoptera). *Journal of Chemical Ecology*. 1995; 21:2079–2089. [PubMed: 24233908]
- Hittinger CT, Johnston M, Tossberg JT, Rokas A. Leveraging skewed transcript abundance by RNA-Seq to increase the genomic depth of the tree of life. *Proceedings of the National Academy of Science USA*. 2010; 107:1476–1481.
- Holloway JD. The moths of Borneo: Family Arctiidae, subfamily Lithosiinae. *The Malayan Nature Journal*. 2001; 55:279–469.
- Hristov NI, Conner WE. Effectiveness of tiger moth (Lepidoptera, Arctiidae) chemical defense against an insectivorous bat (*Eptesicus fuscus*). *Chemoeology*. 2005; 15:105–113.
- Hwang WS, Weirauch C. Evolutionary history of assassin bugs (Insecta: Hemiptera: Reduviidae): insights from divergence dating and ancestral state reconstruction. *PLoS ONE*. 2012; 7:e45523. [PubMed: 23029072]
- Jacobson NL, Weller SJ. A cladistic study of the Arctiidae (Lepidoptera) by using characters of immatures and adults. *Thomas Say Publications in Entomology, Entomological Society of America*. 2002; 98
- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*. 2013; 30:772–780. [PubMed: 23329690]
- Kawahara AY, Breinholt JW. Phylogenomics provides strong evidence for relationships of butterflies and moths. *Proceedings of the Royal Society of London (B)*. 2014; 281:20140970.
- Kersey PJ, Allen JE, Christensen M, Davis P, Falin LJ, Grabmueller C, Hughes DST, Humphrey J, Kerhornou A, Khobova J, Langridge N, McDowall MD, Maheswari U, Maslen G, Nuhn M, Ong CK, Paulini M, Pedro H, Toneva I, Tuli MA, Walts B, Williams G, Wilson D, Youens-Clark K, Monaco MK, Stein J, Wei X, Ware D, Bolser DM, Howe KL, Kulesha E, Lawson D, Staines DM. Ensembl Genomes 2013: scaling up access to genome-wide data. *Nucleic Acids Research*. 2014; 42:D546–D552. [PubMed: 24163254]
- Krieger RI, Feeny PP, Wilinson CF. Detoxication enzymes in the guts of caterpillars: an evolutionary answer to plant defenses? *Science*. 1971; 172:579–581. [PubMed: 5555079]
- Kück, P., Meusemann, K. FASconCAT, Version 1.0. Zoological Research Museum Alexander Koenig; Germany: 2010.
- Lanfear R, Calcott B, Frandsen P. PartitionFinder 2: new methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analysis. *Molecular Biology and Evolution*. 2017; 34:772–773. [PubMed: 28013191]

- Lanfear R, Calcott B, Kainer D, Mayer C, Stamatakis A. Selecting optimal partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology*. 2014; 14:82. [PubMed: 24742000]
- Langel D, Ober D. Evolutionary recruitment of a flavin-dependent monooxygenase for stabilization of sequestered pyrrolizidine alkaloids in arctiids. *Phytochemistry*. 2011; 72:1576–1584. [PubMed: 21288541]
- Legendre, P., Legendre, L. *Numerical Ecology*, Second Ed Developments in Ecological Modeling. Elsevier Science; New York: 1998.
- McCabe TL. *Clemensia albata*, An algal feeding arctiid. *Journal of the Lepidopterists' Society*. 1981; 35:34–40.
- Metzker ML. Sequencing technologies - the next generation. *Nature Reviews Genetics*. 2010; 11:31–46.
- Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010:1–8.
- Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG, Niehuis O, Peterson M, Izquierdo-Carrasco F, Wappler T, Rust J, Aberer AJ, Aspöck U, Aspöck H, Bartel D, Blanke A, Berger S, Böhm A, Buckley TR, Calcott B, Chen J, Friedrich F, Fukui M, Fujita M, Greve C, Grobe P, Gu S, Huang Y, Jermin LS, Kawahara AY, Krogmann L, Kubiak M, Lanfear R, Letsch H, Li Y, Li Z, Li J, Lu H, Machida R, Mashimo Y, Kapli P, McKenna DD, Meng G, Nakagaki Y, Navarrete-Heredia JL, Ott M, Ou Y, Pass G, Podsiadlowski L, Pohl H, von Reumont BM, Schütte K, Sekiya K, Shimizu S, Slipinski A, Stamatakis A, Song W, Su X, Szuesich NU, Tan M, Tan X, Tang M, Tang J, Timelthaler G, Tomizuka S, Trautwein M, Tong X, Uchifune T, Walz MG, Wiegmann BM, Wilbrandt J, Wipfler B, Wong TKF, Wu Q, Wu G, Xie Y, Yang S, Yang Q, Yeates DK, Yoshizawa K, Zhang Q, Zhang R, Zhang W, Zhang Y, Zhao J, Zhou C, Zhou L, Ziesmann T, Zou S, Li Y, Xu X, Zhang Y, Yang H, Wang J, Wang J, Kjer KM, Zhou X. Phylogenomics resolves the timing and pattern of insect evolution. *Science*. 2014; 346:763–767. [PubMed: 25378627]
- Murtah F, Legendre P. Ward's hierarchical agglomerative clustering method: which algorithms implement Ward's criterion? *Journal of Classification*. 2014; 31:274–295.
- Mutanen M, Wahlberg N, Kaila L. Comprehensive gene and taxon coverage elucidates radiation patterns in moths and butterflies. *Proceedings of the Royal Society (B)*. 2010; 277:2839–2848. [PubMed: 20444718]
- Oksanen, J., Blanchet, FG., Kindt, R., Legendre, P., Minchin, PR., O'Hara, FB., Simpson, GL., Solymos, P., Stevens, HH., Wagner, H. *vegan: Community Ecology Package*. R package version 2.2-1. 2015. <http://CRAN.R-project.org/package=vegan>
- Opitz SW, Müller C. Plant chemistry and insect sequestration. *Chemoecology*. 2009; 19:117–154.
- Patti GJ, Tautenhahn R, Siuzdak G. Meta-analysis of untargeted metabolomic data from multiple profiling experiments. *Nature Protocols*. 2012; 7:508–516. [PubMed: 22343432]
- Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, Podsiadlowski L, Petersen M, Lanfear R, Diez PA, Heraty J, Kjer KM, Klopsfstein S, Meier R, Polidori C, Schmitt T, Liu S, Zhou X, Wappler T, Rust J, Misof B, Niehuis O. Evolutionary history of Hymenoptera. *Current Biology*. 2017; 27:1013–1018. [PubMed: 28343967]
- Powell, JA., Mitter, C., Farrell, BD. Evolution of larval feeding habits in Lepidoptera. In: Kristensen, NP., editor. *Lepidoptera, Moths and Butterflies, Vol 1: Systematics, Evolution, and Biogeography Handbook of Zoology*. De Gruyter; Berlin: 1998. p. 403-422.
- R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing; 2014. Vienna, Austria. URL <http://www.R-project.org>
- Rankovi , B., Kosani , M. Lichens as a potential source of bioactive secondary metabolites. In: Rankovi , B., editor. *Lichen Secondary Metabolites: Bioactive Properties and Pharmaceutical Potential*. Springer International Publishing; New York; 2015. p. 1-26.
- Rawlins, JE. Mycophagy in Lepidoptera. In: Wheeler, Q., Blackwell, M., editors. *Fungus-Insect Relationships*. Columbia University Press; New York: 1984. p. 382-423.
- Regier JC, Mitter C, Zwick A, Bazinet AL, Cummings MP, Kawahara AY, Sohn JC, Zwickl DJ, Cho S, Davis DR, Baixeras J, Brown J, Parr C, Weller S, Lees DC, Mitter KT. A large-scale, higher-level,

- molecular phylogenetic study of the insect order Lepidoptera (moths and butterflies). *PLoS ONE*. 2013; 8:e58568. [PubMed: 23554903]
- Regier JC, Zwick A, Cummings MP, Kawahara AY, Cho S, Weller S, Roe A, Baixeras J, Brown JW, Parr C, Davis DR, Epstein M, Hallwachs W, Hausmann A, Janzen DH, Kitching IJ, Solis MA, Yen SH, Bazinet AL, Mitter C. Toward reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia): an initial molecular study. *BMC Evolutionary Biology*. 2009; 9:280. [PubMed: 19954545]
- Reudler JH, Biere A, Harvey JA, van Nouhuys S. Differential performance of a specialist and two generalist herbivores and their parasitoids on *Plantago lanceolata*. *Journal of Chemical Ecology*. 2011; 37:765–778. [PubMed: 21691810]
- Rice P, Longden I, Bleasby A. EMBOS: The European Molecular Biology Open Software Suite. *Trends in Genetics*. 2000; 16:276–277. [PubMed: 10827456]
- Rothschild M, Rowan MG, Fairbair JW. Storage of cannabinoids by *Arctia caja* and *Zonocerus elegans* fed on chemically distinct strains of *Cannabis sativa*. *Nature*. 1977; 266:650–651. [PubMed: 859633]
- Rothschild M, von Euw J, Reichstein T. Cardiac glycosides (heart poisons) in the polka-dot moth *Syntomeida epilais* Walker (Ctenuchidae: Lep.) with some observations on the toxic qualities of *Amata* (= *Syntomis*) *phegea* (Linnaeus). *Proceedings of the Royal Society of London (B)*. 1973; 183:227–247.
- Scoble, MJ. *The Lepidoptera Form, Function and Diversity*. Oxford University Press; New York, NY: 1992.
- Scott CH, Branham MA. A preliminary phylogeny of the lichen moth tribe Lithosiini (Lepidoptera: Erebidae: Arctiinae) based on morphological characters. *Insect Systematics and Evolution*. 2012; 43:321–369.
- Scott CH, Zaspel JM, Chialvo P, Weller SJ. A preliminary molecular phylogenetic assessment of the lichen moths (Lepidoptera: Erebidae: Arctiinae: Lithosiini) with comments on palatability and chemical sequestration. *Systematic Entomology*. 2014; 39:286–303.
- Simmons, R. Adaptive coloration and mimicry. In: Conner, WE., editor. *Tiger Moths and Woolly Bears: Behavior, Ecology, and Evolution of the Arctiidae*. Oxford University Press; New York, NY: 2008. p. 115–126.
- Singer MS, Mace KC, Bernays EA. Self-medication as adaptive plasticity: Increase ingestion of plant toxins by parasitized caterpillars. *PLoS ONE*. 2009; 4:e4796. [PubMed: 19274098]
- Singer MS, Rodrigues D, Stireman JO, Carrière Y. Roles of food quality and enemy-free space in host use by a generalist insect herbivore. *Ecology*. 2004; 85:2747–2753.
- Solhaug, KA., Gauslaa, Y. Secondary lichen compounds as protection against excess solar radiation and herbivores. In: Lüttge, U. Beyschlag, W. Büdel, B., Francis, D., editors. *Progress in Botany*. 2012. p. 283–304.
- Stamatakis A. RAXML-VI-HPC: Maximum Likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*. 2006; 22:2688–2690. [PubMed: 16928733]
- Stamatakis A. RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014; 30:1312–1313. [PubMed: 24451623]
- Stocker-Wörgötter E. Metabolic diversity of lichen-forming ascomycetous fungi: culturing, polyketide and shikimate metabolite production, and PKS genes. *Natural Products Reports*. 2008; 25:188–200.
- van Nieukerken EJ, Kaila L, Kitching IJ, Kristensen NP, Lees DC, Minet J, Mitter C, Mutanen M, Regier JC, Simonsen TJ, Wahlberg N, Yen SH, Zahiri R, Adamski D, Baixeras J, Bartsch D, Bengtsson BÅ, Brown JW, Bucheli SR, Davis DR, De Prins J, De Prins W, Epstein ME, Gentili-Poole P, Gielis C, Hättenschwiler P, Hausmann A, Holloway JD, Kallies A, Karsholt O, Kawahara AY, Koster S, Kozlov MV, Lafontaine D, Lamas G, Landry JF, Lee S, Nuss M, Park KT, Penz C, Rota J, Schintlmeister A, Schmidt BC, Sohn JC, Solis MA, Tarmann GM, Warren AD, Weller S, Yakovlev RV, Zolotuhin VV, Zwick A. Order Lepidoptera Linnaeus, 1758. (ed. by Z.-Q. Zhang) *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*. *Zootaxa*. 2011; 3148:212–221.

- Wagner DL, Rota J, McCabe TL. Larva of *Abablemma* (Noctuidae) with notes on algivory and lichenivory in macrolepidoptera. *Annals of the Entomological Society of America*. 2008; 101:40–52.
- Wahlberg N, Wheat CW. Genomic outposts serve the phylogenomic pioneers: designing novel nuclear markers for genomic DNA extractions of Lepidoptera. *Systematic Biology*. 2008; 57:231–242. [PubMed: 18398768]
- Weller SJ, Jacobson NL, Conner WE. The evolution of chemical defences and mating systems in tiger moths (Lepidoptera: Arctiidae). *Biological Journal of the Linnean Society*. 1999; 68:557–578.
- Wen J, Xiong Z, Nie ZL, Mao L, Zhu Y, Kan XZ, Ickert-Bond SM, Gerrath J, Zimmer EA, Fang XD. Transcriptome sequences resolve deep relationships of the grape family. *PLoS ONE*. 2013; 8:e74394. [PubMed: 24069307]
- Whittaker RH, Feeny PP. Allelochemics: chemical interactions between species. *Science*. 1971; 171:757–770. [PubMed: 5541160]
- Wink M, von Nikisch-Rosenegk E. Sequence data of mitochondrial 16S rRNA of Arctiidae and Nymphalidae: evidence for a convergent evolution of pyrrolizidine alkaloid and cardiac glycoside sequestration. *Journal of Chemical Ecology*. 1997; 23:1549–1568.
- Yang Y, Smith SA. Optimizing de novo assembly of short-read RNA-seq data for phylogenomics. *BMC Genomics*. 2013; 14:328. [PubMed: 23672450]
- Yu Y, Harris AJ, Blair C, He XJ. RASP (Reconstruct Ancestral State in Phylogenetics): a tool for historical biogeography. *Molecular Phylogenetics and Evolution*. 2015; 87:46–49. [PubMed: 25819445]
- Zahiri R, Holloway JD, Kitching IJ, Lafontaine D, Mutanen M, Wahlberg N. Molecular phylogenetics of Erebiidae (Lepidoptera, Noctuoidea). *Systematic Entomology*. 2012; 37:102–124.
- Zaspel JM, Weller SJ, Wardwell CT, Zahiri R, Wahlberg N. Phylogeny and evolution of pharmacophagy in tiger moths (Lepidoptera: Erebiidae: Arctiinae). *PLoS ONE*. 2014; 9:e101975. [PubMed: 25036028]

Highlights

- Phylotranscriptomic analysis helps resolve early relationships in Lithosiini
- Phylogeny highlights subtribal and generic relationships in need of further study
- Phenolic sequestration patterns more similar within than among lichen moth species
- Sequestering phenolics produced by a single pathway may represent a synapomorphy
- Phenolics from orcinol series may form the basis of chemical defense in Lithosiini

Figure 1. Species of Lithosiini examined for phenolic compounds

A: *Miltochrista* sp.; **B:** *Cyana meyricki*; **C:** *Lyclene acetola*; **D:** *Miltochrista ziczac*; **E:** *Cisthene martini*; **F:** *Cisthene tenuifascia*; **G:** *Hypoprepia fucosa*; **H:** *Rhabdatomis laudamia*; **I:** *Ptychoglene phrada*; **J:** *Macotasa nubeculoides*; **K:** *Nishada sambara*; **L:** *Crambidia cephalica*; **M:** *Eilema plana*; **N:** *Eilema bicolor*; **O:** *Euryptidia ira*.

Figure 2. Three metabolic pathways that produce the majority of lichen phenolics
Modified from Rankovi and Kosani (2015).

Figure 3. Phylogeny of the lichen moth tribe

The maximum likelihood tree recovered from the RAXML analysis, rooted with *Bombyx mori* (lnL = -3478648.753226). Bootstrap support values 50 shown beside each node.

Figure 4. Phenolic classes identified in the metabolomic analysis

Examples of the chemical classes of phenolics identified in our analysis, the metabolic path they arise from, and the coding used for each in the RASP analysis. Acetate-Polymalonate Pathway: β -Orsellinic Acid (A), para-depside (A)-anziac acid, meta-depside (A)-homosekikaic acid, depstone (A)-picrolichenic acid, depsidone (A)-pannarin, dibenzofuran (A)-didymic acid, aliphatic acid (B)-caperatic acid, anthraquinone (C)-solorinic acid, chromone (D)-siphulin, diphenyl ether (E)-micareic acid. Mevalonic Acid Pathway: mono-terpene (F)- β -thujone.

Figure 5. Statistical analyses of the lichen phenolic profiles of all analyzed moths

A: Cluster analysis of individual chemical profiles of individual moths. Ward's clustering algorithm (Murtah and Legendre 2014) was used on the Jaccard's dissimilarity (Legendre and Legendre 1998), which measured the overall dissimilarity in the occurrence of the individual chemicals. To delineate groups of similar metabolic profiles, the pruning height was determined with a scree plot of the within-groups error sums of squares from k-means clustering. **B:** Nonmetric multidimensional scaling of individual moths colored according to chemical group. Individuals with identical metabolic profiles treated as a single point. Dissimilarity ranks were ordered according to Jaccard's dissimilarity of the chemical profiles. The chemical groups correspond to the species groups found in Fig. 5A.

Figure 6. Evolutionary hypothesis of lichen phenolic sequestration in the lichen moth tribe

The most likely ancestral lichen phenolic sequestration strategy reconstructions and alternative reconstructions with a likelihood >5% are indicated on each branch of the Lithosiini clade along with the probability of the most likely ancestral state. The probability (P) of the most likely ancestral state is provided at each node. The topology shows only the relationships among the lichen moths and is reformatted from the phylogeny in Fig. 3. The coding used for each species in the analysis is provided next to the species name. A = orcinol series compounds, B = aliphatic acids, C = anthraquinone, D = chromone, E = diphenyl ether, F = mono-terpene.