Published in final edited form as:

Heterocycles. 2011 ; 82(2): 1577–1584. doi:10.3987/COM-10-S(E)114.

DE NOVO ASYMMETRIC APPROACH TO THE DISACCHARIDE PORTION OF SCH-47554

Xiaomei Yu[†], Miaosheng Li[‡], and George A. O'Doherty[§]

[†]Department of Chemistry, West Virginia University, Morgantown, WV 26506, USA.

[‡]Protea Biosciences, 955 Hartman Run Road, Morgantown, WV 26507.

[§]Department of Chemistry and Chemical Biology, Northeastern University, Boston, MA 02115, USA

Abstract

A method for the asymmetric synthesis of the disaccharide portion of SCH-47554 has been developed in 6 steps. The route is shorter than the reported route to a related disaccharide. The route involves the use of the Noyori reduction to establish the asymmetry of the D- and L-sugar portion of the molecule. Diastereoselective Pd-glycosylation reaction and subsequent post-glycosylation transformation are used to establish the remaining stereocenter.

INTRODUCTION

The angucycline family of antibiotics have been of interest to both the synthetic and biological communities due to their unique structures and potent antifungal, antibacterial and antitumor activities.² The structural diversity of this class of natural products is nicely typified by the trisaccharide natural product: SCH-47554 (**1**).³ SCH-47554 (**1**) was isolated by Schering-Plough from the fermentation broths of a strain of *Streptomyces* sp. (SCC-2136) and gained particular interest because of its display of antifungal activity against a variety of yeasts and dermatophytes. Like all of the angucycline antibiotics, SCH-47554 shares a β -C-glycoside linked sugar *ortho* to the phenol on the anthroquinone portion of the aglycone. A less common feature of SCH-47554 as compared to the typical angucyclines, is that its C-glycoside disaccharide portion consists of the rare β -D-amictose and α -L-aculose sugars (Scheme 1).

As part of our ongoing interest in the synthesis and study of this class of natural products we became interested in the synthesis and study of the disaccharide portion of SCH-47554.⁴ Retrosynthetically, we envisioned that SCH-47554 could be prepared by a convergent annulation of arylbromide **2** with a suitably metalated fragment like **3**. In turn, we hoped bromide **2** could be prepared from the direct C-glycosylation of **4** with bromo-phenol **5**. In turn disaccharide **4** could be prepared from pseudo-enantiomeric pyranones **6** and **7**. It is worth noting that a route to a molecule similar to disaccharide **2** was reported by Barrett.⁵

Correspondence to: Miaosheng Li; George A. O'Doherty.

¹This paper is dedicated to Professor Albert Eschenmoser on the occasion of his 80th birthday.

We set out to try to prepare a similar disaccharide (**4** and/or **2**) via our de novo methodology in significantly fewer steps from 2-acylfuran. Herein we report our successful efforts at the de novo synthesis of disaccharide **4**, as well as our unsuccessful efforts to couple it with **5** to form *C*-aryl disaccharide **2**.

RESULTS AND DISCUSSION

Previously, we have described a stereospecific Pd-catalyzed glycosylation reaction, which uses glycosyl-donors α -L-pyranones **7** and α -D-pyranones (*ent*)-**7**.⁶ The enantiomeric donors are readily prepared in 3 steps from acyl furan **8** (Scheme 2). The stereodivergent route employs an enantioselective Noyori reduction (**8** to **9(R)**/**9(S)**),^{7,8} an Achmatowicz oxidation and diastereoselective *t*-butyl carbonate formation (Scheme 2).⁹ By this route, the α -L-pyranone **7** could be isolated in ~50% overall yield, when the Boc-protection was performed selectively prepared in 82% (α : β = 3:1) at low temperature (–78 °C). For our purposes, we needed a more acid stable anomeric leaving group than a *t*-butyl carbonate group for our desired *C*-glycosylation. To these ends we decided to prepare in an analogous fashion the α -D-pyranone **6** with an anomeric pivalate group in similar yields and slightly better selectivity (α : β = 10:1).

The synthesis of the desired α -D-amiceto-sugar **11** began with a Luche reduction (NaBH₄/CeCl₃) of pyranone **6**, which diastereoselectively gave allylic alcohol **10** (68% yield) with the requisite C-4 equatorial alcohol (Scheme 3). A diimide reduction (excess *o*-nitrobenzenesulfonylhydrazide/Et₃N) was then performed to remove the pyran double bond yielding the D-amiceto-sugar **11** (80%). Previously, we had found that a significant amount of hydrogenolysis of the allylic bonds can occur under typical hydrogenation conditions.¹⁰ Our investigation into a cobalt catalyzed one-pot concomitant 1,4-/1,2-reduction of the enone **6** to the equatorial alcohol **11** led only to decomposition.

We turned to the palladium glycosylation condition to install the L-sugar of the desired disaccharides **4** (Scheme 4). Exposure of **11** to a slight excess of α -L-Boc-pyranone **7** under our typical Pd-catalyzed glycosylation conditions (2.5% Pd₂(dba)₃•CHCl₃/10% PPh₃)¹¹ gave a good yield of disaccharide **4** (70% yield). Finally, another Luche reduction (NaBH₄/CeCl₃) of pyranone **4** followed by acylation (Ac₂O/DMAP) gave the backup disaccharide **12** in 70% for the two steps.

With the two D,L-disaccharides **4** and **12** in hand, we explored the use of various Lewis acids to promote its *C*-glycosylation coupling with phenol **5** (Scheme 5). Unfortunately, all the Lewis acids (TiCl₄, Ti(Oi-Pr)₄, SnCl₄, BF₃, and TMSOTf) used for the coupling of **5** and the D,L-disaccharides **4** led to decomposition of the disaccharide **4**. Similarly, when the D,L-disaccharides **12** was used, the only coupling products detected were structure consistent with C-O glycosidic bond cleavage.

In conclusion highly efficient and stereocontrolled synthesis of the amicetose/aculose disaccharide **12** was achieved in 6 steps from acylfuran. This route to the disaccharide portion of SCH47554 was significantly shorter than previous carbohydrate based approaches. Preliminary efforts to transfer the disaccharide to naphthol containing molecules

proved to be unsuccessful. Future works along these lines are ongoing and will be reported in due course.

EXPERIMENTAL

General Methods and Materials

^1H and ^{13}C NMR spectra were recorded on 270 MHz, and 600 MHz spectrometers. Chemical shifts were reported relative to internal tetramethylsilane (δ 0.00) or CDCl_3 (δ 7.26) for ^1H NMR and CDCl_3 (δ 77.0) for ^{13}C NMR. Infrared (IR) spectra were obtained on FT-IR spectrometer. Optical rotations were measured with a digital polarimeter in the solvent specified. Flash column chromatography was performed on 60–200 mesh silica gel. Analytical thin-layer chromatography was performed with precoated glass-backed plates (60 Å, F_{254}) and visualized by quenching of fluorescence and by charring after treatment with *p*-anisaldehyde or phosphomolybdic acid or potassium permanganate stain. R_f values are obtained by elution in the stated solvent ratios (v/v). Ether (Et_2O), THF, methylene chloride (CH_2Cl_2) and triethylamine (Et_3N) were dried by passing through activated alumina columns with argon gas pressure. Commercial reagents were used without purification unless otherwise noted. Air- and/or moisture-sensitive reactions were carried out under an atmosphere of argon/nitrogen using oven-dried glassware and standard syringe/septa techniques.

(6*R*)-6-pivaloyloxy-(2*R*)-2-methyl-2*H*-pyran-3(6*H*)-one (6)

(6*R*)-6-Hydroxy-(2*R*)-2-methyl-2*H*-pyran-3(6*H*)-one (2.22 g, 8.59 mmol), 30 mL of CH_2Cl_2 , and 1.5 mL of Et_3N were added to a round bottom flask and cooled to -78°C . A catalytic amount (53 mg, 0.43 mmol) of DMAP and pivaloyl chloride (1.06 mL, 8.59 mmol) were added and the solution was stirred for 3 h at -78°C . The reaction was quenched with 50 mL of sat. aq. NaHCO_3 , extracted (3×50 mL) with Et_2O , dried (Na_2SO_4), and concentrated under reduced pressure. The crude product was purified using silica gel flash chromatography eluting with 10 % EtOAc /hexanes to give 2.41 g (7.04 mmol, 82 %) of **6**. R_f (20 % EtOAc /hexanes) = 0.4; $[\alpha]_D^{21} -66$ (c 0.1, CH_2Cl_2); IR (thin film, cm^{-1}) 2935, 2910, 2855, 1738, 1692, 1461, 1399, 1279; ^1H NMR (600 MHz, CDCl_3) δ 6.88 (dd, J = 10.2, 3.6 Hz, 1H), 6.46 (d, J = 3.6 Hz, 1H), 6.19 (d, J = 10.2 Hz, 1H), 4.57 (q, J = 6.6 Hz, 1H), 1.40 (d, J = 6.6 Hz, 3H), 1.23 (s, 9H), ^{13}C NMR (150 MHz, CDCl_3) δ 195.9, 177.0, 141.8, 128.2, 86.9, 72.3, 39.2, 27.0, 15.4; CI HRMS calculated for $[\text{C}_{11}\text{H}_{16}\text{O}_4]^+$: 212.1049. Found 212.1045.

((2*R*,5*S*,6*R*)-5,6-dihydro-5-hydroxy-6-methyl-2*H*-pyran-2-yl pivalate (10)

A solution of pyranone **6** (580 mg, 2.74 mmol) in CH_2Cl_2 (1.5 mL) and 0.4 M $\text{CeCl}_3/\text{MeOH}$ (1.5 mL) was cooled to -78°C . NaBH_4 (58.4 mg, 1.54 mmol) was added and the reaction mixture was stirred for 4 h at -78°C . The resulting solution was diluted with ether (10 mL), quenched with 5 mL of saturated NaHCO_3 , extracted (3×5 mL) with Et_2O , dried (Na_2SO_4), and concentrated under reduced pressure. The crude product was purified using silica gel chromatography eluting with 40% EtOAc /hexane to give enol **10** (400 mg, 1.90 mmol, 68%) as a white solid, mp: 62.5–64.0 $^\circ\text{C}$; R_f = 0.30 (40% EtOAc /Hexane); $[\alpha]_D^{21} +21$ (c 0.55, CH_2Cl_2); IR (thin film, cm^{-1}) 3391, 2972, 2933, 2837, 1720, 1612, 1513, 1246, 1029, 999,

819; ^1H NMR (600 MHz, CDCl_3) δ 6.21 (dd, $J = 2.4, 1.8$ Hz, 1H), 6.04 (d, $J = 9.6$ Hz, 1H), 5.75 (ddd, $J = 10.2, 2.4, 2.4$ Hz, 1H), 3.90 (d, $J = 8.4$ Hz, 1H), 3.72 (m, 1H), 2.20 (s, 1H), 1.31 (d, $J = 6.6$ Hz, 3H), 1.20 (s, 9H); ^{13}C NMR (150 MHz, CDCl_3) δ 177.7, 134.7, 125.1, 88.4, 70.3, 69.2, 39.1, 27.1, 18.0; CIHRMS: Calculated for $[\text{C}_{11}\text{H}_{18}\text{O}_4\text{Na}^+]$: 237.1102. Found: 237.1099.

((2R,5S,6R)-3,4,5,6-tetrahydro-5-hydroxy-6-methyl-2H-pyran-2-yl pivalate (11))

A solution of allylic alcohol **10** (370 mg, 1.73 mmol) was dissolved in CH_2Cl_2 (11 mL) at room temperature. *Ortho*-nitrobenzenesulfonylhydrazide (1.33 g, 6.55 mmol) and Et_3N (0.88 g, 8.74 mmol) were added while stirring. The reaction was tracked by TLC. After 12 h, the reaction was quenched by adding 10 mL of satd. aq. NaHCO_3 , extracted (3×50 mL) with methylene chloride, dried over Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified using silica gel flash chromatography eluting with 30% EtOAc/hexanes to give **11** (420 mg, 1.09 mmol, 80%) as a yellow oil. R_f (50% EtOAc/hexanes) = 0.15; $[\alpha]_D^{21} = -105$ ($c = 1.0$, CH_2Cl_2); IR (thin film, cm^{-1}) 3390, 2933, 1724, 1443, 1368, 1246, 1166, 1114, 1066, 1011, 940, 818; ^1H NMR; ^1H NMR (600 MHz, CDCl_3) δ 6.01 (s, 1H), 6.04 (d, $J = 9.6$ Hz, 1H), 3.65 (dq, $J = 9, 6.6$, 1H), 3.30 (ddd, $J = 10.2, 5.4, 1.8$, 1H), 1.98-1.65 (m, 4H), 1.23 (d, $J = 6.6$ Hz, 3H), 1.20 (s, 9H); ^{13}C NMR (150 MHz, CDCl_3) δ 177.0, 90.7, 71.8, 71.5, 39.1, 28.5, 27.2, 27.0, 17.9; CIHRMS: Calculated for $[\text{C}_{11}\text{H}_{20}\text{O}_4\text{Na}^+]$: 239.1259. Found: 239.1264.

((2R,5S,6R)-4-((2'S,6'R)-5,6-dihydro-6-methyl-5-oxo-2H-pyran-2-yloxy)-3,4,5,6-tetrahydro-5-hydroxy-6-methyl-2H-pyran-2-yl pivalate (4))

A CH_2Cl_2 (1.0 mL) solution of Boc pyranone **7** (422 mg, 1.85 mmol) and alcohol **11** (200 mg, 0.93 mmol) was cooled to 0 °C. $\text{Pd}_2(\text{DBA})_3 \cdot \text{CHCl}_3$ (24 mg, 2.5 mol%) and PPh_3 (21 mg, 10 mol%) were added to the reaction mixture at 0 °C. After stirring at room temperature for 8 h, the reaction mixture was quenched with 10 mL of satd. aq. NaHCO_3 , extracted (3×10 mL) with Et_2O , dried over Na_2SO_4 and concentrated under reduced pressure. The crude product was purified using silica gel flash chromatography eluting with 20% EtOAc/hexanes to give pyranone **4** (210 mg, 0.65 mmol, 70%) as a yellow oil: R_f (30% EtOAc/hexanes) = 0.5; $[\alpha]_D^{21} = -105$ ($c = 1.0$, CH_2Cl_2); IR (thin film, cm^{-1}) 2933, 1731, 1701, 1443, 1368, 1246, 1166, 1114, 1066, 1011, 940, 818; ^1H NMR; ^1H NMR (600 MHz, CDCl_3) δ 6.75 (dd, $J = 10.2, 3.6$, 1H), 6.04 (d, $J = 10.2$, 1H), 5.98 (s, 1H), 5.25 (d, $J = 9.6$ Hz, 1H), 4.55 (q, $J = 6.6$, 1H), 3.73 (dq, $J = 9, 6.6$, 1H), 3.30 (m, 1H), 2.09-2.01 (m, 2H), 1.89-1.79 (m, 2H), 1.33 (d, $J = 6.6$ Hz, 3H), 1.19 (d, $J = 6.6$ Hz, 3H), 1.20 (s, 9H); ^{13}C NMR (150 MHz, CDCl_3) δ 196.4, 176.6, 142.7, 127.2, 94.7, 90.4, 80.2, 70.2, 69.8, 38.9, 28.2, 26.9, 25.8, 17.9, 15.0; CIHRMS: Calculated for $[\text{C}_{17}\text{H}_{26}\text{NaO}_6^+]$: 349.3745. Found: 349.3750.

((2R,5S,6R)-4-((2'S,6'R)-5,6-dihydro-5-(4-acetate)-6-methyl-2H-pyran-2-yloxy)-3,4,5,6-tetrahydro-5-hydroxy-6-methyl-2H-pyran-2-yl pivalate (12))

A solution of pyranone **4** (50 mg, 2.74 mmol) in CH_2Cl_2 (1.5 mL) and 0.4 M $\text{CeCl}_3/\text{MeOH}$ (1.5 mL) was cooled to -78 °C. NaBH_4 (58.4 mg, 1.54 mmol) was added and the reaction mixture was stirred for 4 h at -78 °C. The resulting solution was diluted with ether (10 mL), quenched with 5 mL of saturated NaHCO_3 , extracted (3×5 mL) with Et_2O , dried (Na_2SO_4),

and concentrated under reduced pressure. The crude product was dissolved in CH₂Cl₂ (1.5 mL) and DMAP (10 mg) and acetic anhydride (2 mL) was added. After 10 min the reaction was quenched by adding of Et₂O (5 mL) and 10 mL of satd. aq. NaHCO₃. Resulting mixture was extracted (3 × 10 mL) with Et₂O, dried over Na₂SO₄ and concentrated under reduced pressure. The crude product was purified using silica gel flash chromatography eluting with 20% EtOAc/hexanes to give allylic acetate **12** (65 mg, 0.65 mmol, 70%) as a yellow oil: *R*_f (30% EtOAc/hexanes) = 0.5; [α]_D²¹ −65 (*c* 1.0, CH₂Cl₂); IR (thin film, cm^{−1}) 2933, 1731, 1725, 1443, 1368, 1246, 1166, 1114, 1066, 1011, 940, 818; ¹H NMR; ¹H NMR (600 MHz, CDCl₃) δ 6.02 (s, 1H), 5.85 (d, *J* = 10.2, 1H), 5.77 (ddd, *J* = 10.2, 2.4, 2.4 1H), 5.05 (m, 1H), 5.03 (m, 1H), 3.99 (dq, *J* = 9.6, 6.6, 1H), 3.76 (dq, *J* = 9.6, 6.6, 1H), 3.30 (m, 1H), 2.06 (s, 3H), 1.89-1.79 (m, 2H), 1.20 (d, *J* = 6.6 Hz, 3H), , 1.21 (s, 9H) 1.19 (d, *J* = 6.6 Hz, 3H); ¹³C NMR (150 MHz, CDCl₃) δ 177.1, 170.7, 130.2, 127.6, 96.3, 91.0, 80.1, 71.0, 70.5, 65.0, 39.3, 28.7, 27.3, 26.6, 21.2, 18.3, 18.1; CIHRMS: Calculated for [C₁₉H₃₀NaO₇⁺]: 393.4271. Found: 393.4269.

ACKNOWLEDGEMENTS

We are grateful to Kung Wang for his supervision of this project. We are grateful to NIH (GM088839 & 090259) and the NSF (CHE-0749451) for their financial support of our research program.

REFERENCES AND NOTES

2. (a) Rohr J, Thiericke R. Nat. Prod. Rep. 1992; 9:103. [PubMed: 1620493] (b) Krohn K, Rohr J. Top. Curr. Chem. 1997; 188:127.(c) Andriy L, Andreas V, Andreas B. Mol. BioSyst. 2005; 1:117. [PubMed: 16880973]
3. For the isolation of SCH-47554, see: Chu M, Yarborough R, Schwartz J, Patel MG, Horan AC, Gullo VP, Das PR, Puar MS. J. Antibiot. 1993; 46:861. [PubMed: 8514642]
4. For our synthesis as biological study of related molecules see: Yu X, O'Doherty GA. Org. Lett. 2008; 10:4529. [PubMed: 18785743]
5. The Barrett route to a related disaccharide occurred in 21 total steps from D-glucose and L-rhamnose Morton GE, Barrett AGM. Org. Lett. 2006; 8:2859. [PubMed: 16774275]
6. (a) Shan M, Xing Y, O'Doherty GA. J. Org. Chem. 2009; 74:5961. [PubMed: 20560564] (b) Guo H, O'Doherty GA. J. Org. Chem. 2008; 73:5211. [PubMed: 18563936] (c) Zhou M, O'Doherty GA. Org. Lett. 2008; 10:2283. [PubMed: 18461951] (d) Zhou M, O'Doherty GA. J. Org. Chem. 2007; 72:2485. [PubMed: 17338573]
7. Fujii A, Hashiguchi S, Uematsu N, Ikariya T, Noyori R. J. Am. Chem. Soc. 1996; 118:2521.
8. (a) Li M, Scott JG, O'Doherty GA. Tetrahedron Lett. 2004; 45:1005.(b) Li M, O'Doherty GA. Tetrahedron Lett. 2004; 45:6407.
9. (a) Babu RS, O'Doherty GA. J. Carb. Chem. 2005; 24:169.(b) Guo H, O'Doherty GA. Org. Lett. 2005; 7:3921. [PubMed: 16119932]
10. Haukaas MH, O'Doherty GA. Org. Lett. 2002; 4:1771. [PubMed: 12000295]
11. (a) Babu RS, Zhou M, O'Doherty GA. J. Am. Chem. Soc. 2004; 126:3428. [PubMed: 15025462] (b) Babu RS, O'Doherty GA. J. Am. Chem. Soc. 2003; 125:12406. [PubMed: 14531673]

Scheme 1.
Retrosynthetic analysis of SCH-47554

Scheme 2. Synthesis of the key pyranone coupling partners **6** and **7**

Scheme 3.
Synthesis of the desired D-amiceto-sugar **11**

Scheme 4.
Synthesis of the desired D-amicetose/L-aculose disaccharide **4**

Scheme 5.
Attempted Lewis coupling of **4** and **12** with **5**.