EPA Public Access

Author manuscript

Atmos Environ (1994). Author manuscript; available in PMC 2018 November 21.

About author manuscripts

Submit a manuscript

Published in final edited form as:

Atmos Environ (1994). 2017 November 21; 174: 214–226. doi:10.1016/j.atmosenv.2017.11.026.

Factors associated with NO₂ and NO_x concentration gradients near a highway

J. Richmond-Bryant^{a,*}, M.G. Snyder^b, R.C. Owen^c, and S. Kimbrough^d

^aNational Center for Environmental Assessment, Office of Research and Development, United States Environmental Protection Agency, 109 TW Alexander Drive, B243-01, Research Triangle Park, NC, 27711, USA

^bInstitute for the Environment, University of North Carolina, 100 Europa Drive, Suite 490, Campus Box 1105, Chapel Hill, NC, 27599-1105, USA

^cOffice of Air Quality Planning and Standards, Office of Air and Radiation, United States Environmental Protection Agency, C439-01, Research Triangle Park, NC, 27711, USA

^dNational Risk Management Research Laboratory, Office of Research and Development, United States Environmental Protection Agency, 109 TW Alexander Drive, E343-02, Research Triangle Park, NC, 27711, USA

Abstract

The objective of this research is to learn how the near-road gradient, in which NO₂ and NO_x (NO + NO₂) concentrations are elevated, varies with changes in meteorological and traffic variables. Measurements of NO₂ and NO_x were obtained east of I-15 in Las Vegas and fit to functions whose slopes (dC_{NO_2}/dx and dC_{NO_x}/dx , respectively) characterize the size of the near-road zone where NO₂ and NO_x concentrations from mobile sources on the highway are elevated. These metrics were used to learn about the near-road gradient by modeling dC_{NO_2}/dx and dC_{NO_x}/dx as functions of meteorological variables (e.g., wind direction, wind speed), traffic (vehicle count), NO_x concentration upwind of the road, and O₃ concentration at two fixed-site ambient monitors. Generalized additive models (GAM) were used to model dC_{NO_2}/dx and dC_{NO_x}/dx versus the independent variables because they allowed for nonlinearity of the variables being compared. When data from all wind directions were included in the analysis, variability in O₃ concentration comprised the largest proportion of variability in dC_{NO_2}/dx , followed by variability in wind direction. In a second analysis constrained to winds from the west, variability in O₃ concentration remained the largest contributor to variability in dC_{NO_2}/dx , but the relative contribution of variability in wind speed to variability in dC_{NO_2}/dx increased relative to its contribution for the all-wind analysis. When data from all wind directions were analyzed, variability in wind direction was by far the largest contributor to variability in dC_{NO_x}/dx , with smaller contributions from hour

*Corresponding author. richmond-bryant.jennifer@epa.gov (J. Richmond-Bryant).

Disclaimer

The study was reviewed by the National Center for Environmental Assessment, U.S. EPA and approved for publication. Mention of trade names or commercial products does not constitute endorsement or recommendation for use. The views expressed in this article are those of the authors and do not necessarily reflect the views or policies of the U.S. Environmental Protection Agency.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.atmosenv.2017.11.026>.

of day and upwind NO_x concentration. When only winds from the west were analyzed, variability in upwind NO_x concentration, wind speed, hour of day, and traffic count all were associated with variability in dC_{NO_x}/dx . Increases in O_3 concentration were associated with increased magnitude near-road dC_{NO_2}/dx , possibly shrinking the zone of elevated concentrations occurring near roads. Wind direction parallel to the highway was also related to an increased magnitude of both dC_{NO_2}/dx and dC_{NO_x}/dx , again likely shrinking the zone of elevated concentrations occurring near roads. Wind direction perpendicular to the road decreased the magnitude of dC_{NO_2}/dx and dC_{NO_x}/dx and likely contributed to growth of the zone of elevated concentrations occurring near roads. Thus, variability in near-road concentrations is influenced by local meteorology and ambient O_3 concentration.

Keywords

Near-road; NO_2 ; Oxides of nitrogen; Nitrogen dioxide; Dispersion

1. Introduction

Human exposures to nitric oxide (NO), nitrogen dioxide (NO_2), and NO_x (i.e., $\text{NO} + \text{NO}_2$) have been documented to be higher near the road compared with locations away from the road (Janssen et al., 2001; Roorda-Knappe et al., 1998; Van Roosbroeck et al., 2006). Several studies have measured near-road gradients to characterize the change in concentration (C) over distance (x) away from the road, as summarized in a review by Karner et al. (2010): $C = f(x)$. For this purpose of considering near-road exposures, we define the *near-road zone* as the region proximal to the road where the concentration of traffic-related air pollution is elevated relative to concentrations further downwind (or upwind) of the road (*background* levels are considered to be *urban-scale*, or 4–50 km based on the United States Environmental Protection Agency (U.S. EPA) regulations for siting ambient air quality monitors described in 40 CFR Part 58 App. D). The gradient of concentration with distance from the road, dC/dx , provides an indicator of the size of the near-road zone where NO_2 and NO_x concentrations from mobile sources on the highway are elevated. A sharp gradient (i.e., large magnitude dC/dx) would suggest that the near-road zone is narrow because concentrations return from the elevated concentrations at the edge of the road to background levels over a shorter distance. Conversely, a small gradient (i.e., small magnitude dC/dx) would suggest that the near-road zone is wide because concentrations return from the elevated levels at the edge of the road to background levels over a larger distance. While studies have focused largely on the magnitude of change, none have explicitly identified the factors that influence dC/dx . This type of analysis is important in that it would allow exposure and risk assessors to account for variability in traffic-related pollution in the near-road environment and to better predict exposures occurring in these locations having varying meteorological, geological, or built features.

The NO, NO_2 , and NO_x gradients change dynamically in the near-road zone. NO is emitted from mobile sources and then converts to NO_2 over a time-scale of seconds to minutes, mainly via oxidation of NO to NO_2 by available ozone (O_3) (Atkinson, 2000; Seinfeld and Pandis, 2006). NO_2 rapidly converts back to NO through photolysis and forms other reactive

nitrogen species, such as nitric acid (HNO_3) or nitrate radical ($\text{NO}_3\cdot$) during oxidation. Evolution of the NO_2 and NO_x gradients are influenced by available sunlight, O_3 , and other photooxidants.

Mechanical mixing may play a role in dispersion of the near-road plume, so that prevailing winds, atmospheric turbulence, and traffic volumes may also have an influence on the NO_2 and NO_x gradients. Large-scale mixing due to thermal gradients in the atmosphere has long been recognized as a driver of atmospheric turbulence (Obukhov, 1962). This mixing can be enhanced due to heat flux in urban areas (Britter and Hanna, 2003). Gordon et al. (2012), Durant et al. (2010), and Hu et al. (2009) all observed elevated near-road NO and NO_2 concentrations before sunrise during high atmospheric stability conditions. Hu et al. (2009) observed that the near-road zone of elevated pollutant levels was extended during atmospheric stability conditions. When studying the role of meteorological variables on near-road NO concentrations, Venkatram et al. (2007) observed that vertical velocity fluctuations had the largest influence among the meteorological variables measured.

Several studies noted that vehicle motion can create localized turbulence near roadways to enhance dispersion of traffic-related air pollution (Baldauf et al., 2008; He and Dhaniyala, 2011; Kastner-Klein et al., 2001; Sedefian et al., 1981; Wang and Zhang, 2009). di Sabatino et al. (2003) observed a relationship between vehicle-induced turbulence and traffic levels. In light traffic, the wake behind a vehicle is singular. For increasing traffic, the wakes combine so that turbulence is related to the number of vehicles, vehicle length scale, and vehicle speeds. When traffic is moving but densely packed, vehicle-induced turbulence becomes independent of the number of vehicles. At the same time, drag coefficient declines with decreasing space between vehicles, so the traffic-related turbulence is expected to decline as well.

The objective of this research is to estimate how the near-road gradient varies with changes in meteorological, ambient, and traffic variables to better understand the factors that influence human exposure to traffic-related NO_2 and NO_x . The gradient measure, dC/dx , serves as a surrogate for the size of the near-road zone and is statistically modeled as a function of meteorological, ambient, and traffic variables. The work is driven by the hypothesis that the gradient is influenced by different meteorological, traffic, and ambient air pollutant concentration variables. A description of the results is followed by a discussion and conclusions. It is hoped that this exercise improves our understanding of the extent of the near-road environment and hence improves our understanding of human exposure to traffic-related air pollutants. This work is also expected to benefit dispersion model development because it studies factors that are associated with the spatial gradient of NO_2 and NO_x concentrations in the near-road environment.

2. Methods

Methodology for this study is an extension of that presented in Richmond-Bryant et al. (2017). The information from Richmond-Bryant et al. (2017) is briefly summarized here, and then the extended meteorological analysis is described in detail.

2.1. Study area

Near-road measurements of air quality, traffic, and meteorology were obtained at a study area in Las Vegas, NV (Fig. 1). The study area was adjacent to Interstate-15 (I-15) with Annual Average Daily Traffic of approximately 206,000 vehicles per day, 10% of which were heavy-duty diesel trucks (Nevada, 2008). I-15 runs north-south at this location with a below-grade highway having a 20° embankment. The downwind sampling sites sit on flat terrain 25 m, 115 m, and 300 m east of the highway edge. An upwind sampling site was also sited approximately 100 m west of the highway edge. The Las Vegas climate is arid with hot summers and plentiful sunshine year-round. Las Vegas is surrounded by mountains, a geographic feature that frequently fosters stable atmospheric conditions during the evening and nighttime hours. A detailed description of the study area and pollutant measurements can be found in Kimbrough et al. (2013).

2.2. Data collection

Continuous monitoring of NO and NO_x was performed using chemiluminescence with a trace NO_x analyzer (Ecotech, Model EC 9841 B, Knoxfield, VIC, Australia). NO₂ concentrations were estimated over each sampling time interval by taking the difference between NO_x and NO concentrations. Data were logged every five minutes and averaged every hour. Multipoint calibration was performed at the start of the study, and zero and span checks occurred nightly.

Surface meteorological observations were measured at McCarran International Airport (LAS), which was 1.5 km north of the study area. Surface measurements included wind speed and direction, air temperature, relative humidity, precipitation, and solar radiation. Upper air measurements were made at the Universal Rawinsonde Observation (RAOB) station in Mercury/Desert Rock, NV at an elevation of 1006 m. Upper air measurements included elevation, temperature, moisture, and wind components. Surface and upper air data were processed using the American Meteorological Society/Environmental Protection Agency Model (AERMOD) Meteorological Processor (AERMET) package (Cimorelli et al., 2005), the meteorological preprocessor for the U.S. EPA's regulatory dispersion model, AERMOD. AERMET computes hourly average data for ambient air temperature, relative humidity, sensible heat flux, convective and mechanical mixing heights, atmospheric pressure, roughness height, eddy length scale and volume, Monin-Obukhov length scale, and vertical and horizontal friction velocity. Surface wind speed and direction measurements were obtained each minute and averaged over an hour (using AERMINUTE, https://www3.epa.gov/ttn/scram/7thconf/aermod/aerminute_userguide.pdf); the hourly averaged data were used in the analysis. The ambient O₃ concentrations were obtained from Clark County Department of Air Quality monitoring locations 6.6 km north-northeast (Orr site) and 7.4 km west-northwest (Paul Meyer site) of the I-15 field measurement site. Ozone measurements from the Orr site were used when winds were from the 120-degree sector centered at east, the Paul Meyer site measurements were used when winds were from the 120° sector centered at west, and averaged between the two sites when winds were from the 60° sectors centered at north and south. Ozone concentrations from the two sites had a Pearson correlation of 0.91 and a 21.8% percent difference (calculated as the average of the absolute value of the difference between the O₃ concentration at the Paul Meyer site and the

O₃ concentration at the Orr site, all divided by the mean across both sites). Negative values were censored from the dataset, and all remaining O₃ measurements were shifted up by a value of 0.01 ppb to prevent errors related to handling zero-valued data without compromising correlations among the variables. Traffic data were obtained from Nevada (2008) and aggregated to hourly traffic volumes and speeds.

Data were collected from the near-road site between December 12, 2008 and January 21, 2010, and meteorological data processed by AERMET were merged to the dataset for that time period. 8466 (81.8%) full hours of data were complete (as determined by the presence of NO₂ concentration data at the three downwind sites). Data were then excluded if the NO_x concentration increased with distance from the road during any given hour. This exclusion was made to remove potential outside sources (i.e., a truck parked next to a downwind monitor), and it was also used by Kimbrough et al. (2017) to examine the NO to NO₂ conversion, leaving 6361 (61.5%) hours of NO₂ and NO_x data. Records where meteorological and other variables were missing were further removed, leaving 5525 (53.4%) complete hours of data. A second analysis focused only on winds from the west (210–330°) so that the monitors were downwind of I-15. There were 3426 (33.1%) hours of data where winds emanated from the west, and when records with missing data were removed, 2622 (25.3%) hours of data remained for this analysis.

2.3. On-road concentration estimation

The near-road gradient was estimated by fitting an ordinary least squares (OLS) regression to the NO₂ and NO_x concentrations reported at the three downwind monitors at each hour. Richmond-Bryant et al. (2017) found a *logit* model to produce the most sensible fit to the data, given that the *logit* model produces an S-shaped curve rather than producing an infinitely large concentration at the road. From Richmond-Bryant et al. (2017):

$$\text{logit}(C) = \beta_1 \ln(x) + \beta_0 \quad (1)$$

where C = concentration, x = distance from the road to the monitor, β_0 = y-intercept, and β_1 = gradient (dC/dx). The concentration at the 300 m downwind monitor was used as the reference value.

2.4. Modeling the near-road gradient with respect to meteorological variables

The gradient dC/dx was modeled as a function of several variables (Table 1) to estimate which variables influence the size of the near-road zone. The gradients for NO₂ and NO_x were tested here because NO and NO₂ are interconverted through photochemistry, but their sum, NO_x, is mostly conserved. NO₂ may be oxidized further, but at time scales exceeding those of dispersion in the near-road environment (Seinfeld and Pandis, 2006). Therefore, loss of total NO_x mostly occurs through dispersion rather than chemistry (Carslaw, 2005). It is also instructive to study the association of the NO₂ gradient with meteorological variables to learn how meteorological variables may influence NO₂ generation via photochemical reaction of NO with O₃.

A generalized additive model (GAM) was applied to represent the response of $\beta_1 = dC/dx$ to the meteorological variables studied (Hastie and Tibshirani, 1986, 1990; Simon, 2006). The GAM enabled examination of the relative contributions of each variable as well as the response of the gradient to each meteorological variable. The GAM was chosen over a linear model to accommodate the nonlinear relationship between the gradient and several of the meteorological variables. Each term was modeled as a smooth function, s , which can take either a parametric or non-parametric form. The model was of the form:

$$dC/dx = s(\varphi) + s(T) + s(L_{Mon}^{-1}) + s(W) + s(WD) + s(TR) + s(O_3) + s(NO_{X,up}) + s(H) \quad (2)$$

where φ = sensible heat flux (W/m^2), T = ambient temperature (K), L_{Mon} = Monin-Obukhov length scale (Foken, 2006; Monin and Obukhov, 1954) of atmospheric turbulence above the roughness sublayer (m), W = wind speed (m/s), WD = wind direction ($^\circ$), TR = hourly traffic volume (number), O_3 = ozone concentration (ppb), $NO_{X,up}$ = upwind NO_X concentration (ppb), and H = hour of day. The variables included in this model were tested for Pearson correlation to support the assumption of independence among the variables. The correlation matrix is provided in Table 2. Maximum of convective and mechanical mixing height, roughness height, relative humidity, atmospheric pressure, lateral turbulent velocity, eddy volume, and friction velocity were removed from the final GAM models because their correlation with other variables was judged to be sufficiently high ($r > 0.7$) to artificially inflate the model fit. High correlation likely could be attributed to the fact that these terms were derived variables from AERMET (Cimorelli et al., 2005). Stepwise addition and subtraction of these variables in the GAM allowed for R^2 to be kept reasonably high while minimizing the sum squared residuals (SSR) normalized by the sum of the effective degrees of freedom (edf). Additionally, we tested the impact of removing O_3 or variables that were moderately correlated with O_3 (such as temperature and hourly traffic) through stepwise addition and subtraction, but these changes caused lower R^2 and higher SSR without changing the statistical significance or influence of the other variables.

Proc *gamm* from the *mgcv* package v.1.8–3 (Wood, 2016) was run in R v.3.1.2 (RCore_Team, 2015) to implement the model. Proc *gamm* allows for several different types of basis functions to fit the smoothed functions included as independent variables. The models were tested with the different basis functions, and a cubic splines basis function was selected as that which produced the lowest SSR normalized by the effective degrees of freedom of the model (Simon, 2006). To understand the sensitivity of the model results to basis function type, the rank order F-statistics of each variable were compared across basis functions. All basis functions produced the same two variables with highest F-statistics for both the NO_2 and NO_X models, and there was reasonable agreement among the models' rank order F-statistics for the other variables for both the dC_{NO_2}/dx and dC_{NO_X}/dx models. These results are shown in Supplemental Table S1. The data were not adjusted for autocorrelation because they were not present as a complete time series, since data were removed based on the criteria for a decreasing gradient away from the road. As such, the data were treated as individual observations. The model of gradient vs. each variable, along with associated point data, was plotted using proc *scatter.smooth*.

3. Results and discussion

Ambient concentration, meteorological, and traffic data are provided in Table 1 and in Kimbrough et al. (2013). Declining NO₂ and NO_x concentrations with distance from the road were observed on average over the three monitors sited to the east of I-15, while the average upwind NO_x concentration was close to the average NO_x concentration at the 300 m site. Variability in those concentrations, as indicated by large standard deviations and interquartile ranges, likely reflects the fact that these data included all meteorological conditions. During stable conditions, typically at nighttime and in early morning hours, the majority of winds came from the southwest; while during unstable conditions, typically occurring in the middle of the day, most winds came from the northeast (Fig. 2). Most variables tested were not highly correlated with each other and so were included in the model (Table 2).

The GAM was useful in evaluating the overall influence of the meteorological, ambient, and traffic variables on dC_{NO₂}/dx and dC_{NO_x}/dx. For the NO₂ model, R² was 0.409 (SSR/edf = 0.108). R² was higher for the NO_x model at 0.509 (SSR/edf = 0.873) when all wind directions were included in the analysis (Table 3). Hence, the models predicted some but not all of the variability in dC_{NO₂}/dx and dC_{NO_x}/dx. These values for R² were relatively low and likely reflected high uncertainty in dC_{NO₂}/dx and dC_{NO_x}/dx around some of the smooth functions.

It would be expected that wind direction influences human exposure to near-road NO₂ and NO_x. Richmond-Bryant et al. (2017) presented gradients for winds perpendicular and parallel to the road. When winds were from the west (i.e., perpendicular to I-15), dC_{NO₂}/dx and dC_{NO_x}/dx represented a more gradual decrease in NO₂ or NO_x concentration with distance from the road that would imply elevated exposures over a larger distance from the road. When winds were perpendicular to I-15, dC_{NO₂}/dx and dC_{NO_x}/dx represented sharp decreases in NO₂ or NO_x concentration from the road, implying that concentrations quickly decreased to background levels. This prompted focus on winds from the west in the analyses presented in Table 4 and Figs. 4 and 6.

When winds were constrained to those from the west, R² was 0.506 (SSR/edf = 0.0711) for the NO₂ model and R² = 0.475 (SSR/edf = 0.773) for the NO_x model (Table 4). Prior to removal of correlated variables, R² was larger for both the dC_{NO₂}/dx and dC_{NO_x}/dx models (R² = 0.538 and R² = 0.661, respectively).

GAM modeling produced an analysis of variance (ANOVA) with a set of F-statistics for the predictor variables to describe how variability in the smoothed GAM model as a whole compared to the variability within each predictor variable (Hastie and Tibshirani, 1990; Simon, 2006). In the case of a large F-statistic, the model variability far exceeded the variability in the predictor variable. Variability in ambient O₃ concentration had the most influence on variability in dC_{NO₂}/dx (47.5%), followed by wind direction (29.7%) when all wind directions were included in the analysis (Table 3). Variability in all other variables had less than 5% influence on variability in dC_{NO₂}/dx. When winds were restricted to those from the west (Table 4), variability in ambient O₃ concentration still had the largest influence on

variability in dC_{NO_2}/dx (36.0%). In this case, variability in wind speed accounted for a larger proportion of the variability in dC_{NO_2}/dx (27.7%), and variability in heat flux, traffic count, inverse Monin-Obukhov length scale, hour of day, and upwind NO_X concentration, and temperature each accounted for less than 10% of the variability in dC_{NO_2}/dx . These relationships persisted even when the influence of wind direction was removed. However, the influence of wind speed became much more important when wind direction was removed.

Variability in dC_{NO_X}/dx was most influenced by variability in wind direction (61.7%) when all wind directions were included in the analysis (Table 3). Variability in hour of day and upwind NO_X concentration had smaller influence on dC_{NO_X}/dx (15.8% and 9.5%, respectively). Variability in all other variables, including ambient O_3 concentration, had less than 5% influence on variability in dC_{NO_X}/dx . Variability in wind direction had the most influence on variability in dC_{NO_X}/dx , possibly given that overall NO_X is mostly conserved so that mechanical dispersion is important to NO_X concentration but atmospheric chemistry is not (Seinfeld and Pandis, 2006). When winds were restricted to those from the west (Table 4), variability in upwind NO_X concentration (26.9%), wind speed (23.8%), hour of day (18.9%), and traffic (16.5%) each had moderate influence on variability in dC_{NO_X}/dx . Variability in other variables influenced approximately 5% or less of the variability in dC_{NO_X}/dx . The biggest difference between the analyses involving all winds vs. winds just from the west was that variability in wind speed comprised a much larger proportion of the variability in dC_{NO_X}/dx and dC_{NO_2}/dx when winds were restricted to those from the west.

Inspection of the relationships between dC/dx and each independent variable provided insight about the contribution of each independent variable to the near-road gradient. Because concentration declined with distance from the road, dC/dx was always negative. Decreasing dC/dx (i.e., dC/dx becoming more negative) with an increasing variable value meant that the magnitude of dC/dx was increasing as the variable value increased; dC/dx thus moved farther away from zero. Hence, this suggested that the near-road zone was shrinking in size because concentration varied from its peak near the road to background levels over a shorter distance. Conversely, increasing dC/dx (i.e., dC/dx becoming less negative) with an increasing variable value meant that the magnitude of dC/dx was declining as the variable value increased; dC/dx hence moved towards zero. This suggested that the near-road zone was growing in size because concentration varied from its near-road peak to background levels over a longer distance.

Fig. 3 illustrates how each variable was related to dC_{NO_2}/dx when all wind directions were included. dC_{NO_2}/dx decreased (i.e., became more negative) with increasing ambient O_3 concentration measured at a fixed-site monitor, which also had the largest influence in the ANOVA of the model (Table 3). As O_3 concentration increased, the near-road zone of elevated NO_2 shrank to ambient levels over a shorter distance. This was an unexpected result, given that O_3 tends to react with NO to form NO_2 (Seinfeld and Pandis, 2006). Rodes and Holland (1981) also measured NO , NO_2 , and O_3 downwind of a Los Angeles highway and found a larger decrease in NO_2 concentration downwind of the highway with higher ambient O_3 concentrations. Chaney et al. (2011) measured NO , NO_2 , and O_3 downwind of two suburban roadways north of Leeds, UK and observed a similar relationship between

NO₂ concentration and distance from the road. Wind direction had the next largest influence in the ANOVA, and dC_{NO_2}/dx decreased as wind direction changed from an eastern origin (90°) to a southern origin (180°) and also as wind direction changed from a western origin (270°) to a northern origin (0°), where the highway ran north-south. Hence, as the wind direction approached parallel to the road, the near-road zone would likely shrink. Conversely, as wind direction approached the normal to the road, the size of the near-road zone would likely grow, as given by regions of increasing dC_{NO_2}/dx . This finding was consistent with Venkatram et al. (2013), where the authors observed concentrations of NO normalized by traffic counts at 7 m and 17 m downwind of a Raleigh, NC highway under different wind directions. When winds were nearly parallel to the road, the difference between normalized concentrations measured at the two locations was greater than when wind was normal to the road. This suggests that the gradient would decrease (i.e., become more negative), and the near-road zone of elevated concentration would shrink, when the wind was parallel to rather than perpendicular to the road. This was consistent with our model of dC_{NO_2}/dx as a function of wind direction. Decreasing dC_{NO_2}/dx (i.e., becoming more negative with a shrinking near-road zone) was associated with increasing wind speed for wind speeds less than 4 m/s. A change in this relationship for wind speeds greater than 4 m/s might be related to greater uncertainty from less data at higher winds. Changes in available O₃ had the most influence on changes in dC_{NO_2}/dx , likely because reactivity of NO to form NO₂ was influenced by the available ambient O₃, as also observed in Chaney et al. (2011), Rodes and Holland (1981) and Sander et al. (2011).

Fig. 4 displays the relationship between dC_{NO_2}/dx and each variable when winds were constrained to those from the west (210–330°). The curves of dC_{NO_2}/dx with respect to each variable followed the same general trend as when winds from all directions were included. However, the confidence intervals were smaller when winds were constrained to one direction. The slope of the relationship between dC_{NO_2}/dx and ambient O₃ concentration was more negative in this case, implying that increasing O₃ was associated with shrinkage of the near-road zone of elevated NO₂. Increasing wind speed was associated with shrinkage of the near-road zone of elevated NO₂ for low speeds. Above a wind speed of 4 m/s, increasing wind was associated with dC_{NO_2}/dx decreasing in magnitude (i.e., becoming less negative), so that the near-road zone of elevated NO₂ would grow with increasing wind speed. These results imply that increasing wind speed above 4 m/s may act to transport the NO₂ farther away from the highway, but below 4 m/s the gradient increased in magnitude (i.e., became sharper). This stands in contrast to Baldwin et al. (2015), who observed that normalized concentration decreased with increasing wind speed, although error bounds were high for the high wind speed category. The effect persisted for the all wind direction and westerly wind analyses. It is possible that the bilinear curve of dC_{NO_2}/dx vs. wind speed is influenced by competing effects of inter-reactivity between NO and NO₂ (Seinfeld and Pandis, 2006), causing the addition of NO₂ to the plume downwind of the road, and transport and dispersion. Another possible cause of the bilinear relationship between dC_{NO_2}/dx and wind speed is that, when winds were of greater speed and from the west, flights taking off from McCarran International Airport would fly directly over the study site from the runway oriented east-west and located within 0.5 km from the study site. Airport emissions have

been shown to contain NO_2 and NO_x (Carslaw et al., 2006). However, airport impacts would only be anticipated when winds are from the east (Henry et al., 2011).

Fig. 5 displays the relationship between dC_{NO_x}/dx and each variable when all wind directions were included. Attention to the y-axis of dC_{NO_x}/dx vs. each variable compared with that for dC_{NO_2}/dx shows that the range in dC_{NO_x}/dx was more than two times higher than for the range in dC_{NO_2}/dx . This roughly mirrored the 67% difference in estimated dC_{NO_x}/dx and dC_{NO_2}/dx based on the average NO_x and NO_2 concentrations from Table 1. ANOVA results indicate that variation in wind direction had the largest influence on variability in dC_{NO_x}/dx (Table 3). The behavior of dC_{NO_x}/dx with respect to wind direction was similar to the behavior of dC_{NO_2}/dx . However, more scatter in the data was observed between wind directions of 180° – 270° , and the magnitude of dC_{NO_x}/dx was larger for winds from the southwest compared with those from the northeast. As a result, the smooth fit to the data show that dC_{NO_x}/dx became more negative as wind direction shifted from an eastern origin (90°) to a western origin (270°), and the near-road zone likely would shrink over this change in wind direction. The ANOVA analysis also showed that variability in hour of day (15.8%) and upwind NO_x concentration (9.5%) also had larger associations with variability in dC_{NO_x}/dx (Table 3). The magnitude of dC_{NO_x}/dx decreased during the hours of 8:00 and 13:00 (i.e., suggesting near-road zone growth) and then increased from 13:00 to 0:00 (i.e., suggesting near-road zone shrinkage). dC_{NO_x}/dx appeared relatively stable between 0:00 and 8:00. Hour of day was not highly correlated with sensible heat flux, Monin-Obukhov mixing height, or traffic count (Table 2). The period from 8:00–13:00 coincided with convective daytime mixing. Rush hour traffic was at a maximum from approximately 10:00 to 16:00 (Kimbrough et al., 2017), where heavy traffic has the potential to cause localized turbulence. During this time period, median northbound traffic speed ranged from 34 to 48 miles per hour with wide variability due to construction several miles north of the monitoring site. Southbound traffic speed was maintained between 60 and 70 miles per hour (Supplemental Fig. S5). Greater mixing on the road may have caused shrinkage of the near-road zone after 13:00, resulting in enhanced dispersion of NO_x and therefore increased magnitude of dC_{NO_x}/dx . On the whole, increasing upwind NO_x concentration appeared to be associated with increased magnitude dC_{NO_x}/dx and therefore with potential shrinkage of the near-road zone of elevated NO_x concentration. Most of the NO_x data were for concentrations less than 50 ppb, so that uncertainty in the function increased as concentrations increased beyond than 50 ppb.

Fig. 6 displays the relationship between dC_{NO_x}/dx and each variable when winds were constrained to those from the west (210 – 330°). Unlike for dC_{NO_2}/dx , many of the curves of dC_{NO_x}/dx with respect to each variable varied in shape from the case when winds from all directions were included, and the confidence intervals for the case of winds from the west were not always smaller than when all winds were included in the analysis. For all wind directions and where wind directions were constrained to those from the west, dC_{NO_x}/dx increased in magnitude with increasing upwind NO_x concentration with similar slopes, suggesting that higher levels of upwind NO_x are associated with a smaller near-road zone of elevated NO_x in both cases. One important difference is that dC_{NO_x}/dx did not vary substantially with increasing wind speed beyond a wind speed of 3 m/s when all winds were included (and for wind speeds lower than 3 m/s, the confidence intervals became enlarged).

For winds from the west, dC_{NO_X}/dx decreased in magnitude with increasing wind speed, suggesting that higher wind speeds caused growth of the near-road zone of elevated NO_X concentration when winds were perpendicular to the road. The relationship between dC_{NO_X}/dx and traffic did not vary substantially when limiting the analysis to winds from the west. The relationship between dC_{NO_X}/dx and hour of day reached a minimum around 7:00, in the time period when atmospheric conditions transitioned from stable to convective (Kimbrough et al., 2017).

The most notable difference between the dC_{NO_2}/dx and dC_{NO_X}/dx models was attributed to ambient O_3 concentration. The magnitude of dC_{NO_2}/dx increased for increasing ambient O_3 concentration. Ambient O_3 reacts with NO emissions from the road to produce NO_2 downwind, and the size of the zone over which this occurs would possibly increase with greater available O_3 . Similarly, Rodes and Holland (1981) observed the steepest dC_{NO_2}/dx for the upper tertile of measured O_3 concentration with the middle and lower tertiles having progressively smaller dC_{NO_2}/dx . They estimated that concentration of a non-reactive gas would decline exponentially with distance from the road, but NO_2 decreased approximately linearly with distance from the road. The slower decline observed in the Rodes and Holland (1981) paper was attributed to the contribution of O_3 consumption by NO to produce NO_2 . Our results add to the findings of Rodes and Holland (1981) by demonstrating O_3 to be the most important among the variables tested. The curve for dC_{NO_X}/dx as a function of ambient O_3 concentration had an opposite slope to dC_{NO_2}/dx , and variance in ambient O_3 concentration only accounted for 1.3% of variation in dC_{NO_X}/dx . This was probably related to the fact that overall NO_X concentration was conserved, so it would be anticipated that O_3 would have little impact on dC_{NO_X}/dx . Likewise, Rodes and Holland (1981) observed little difference for the change in NO_X concentration over distance from the road among high, medium, and low O_3 groups.

A number of studies have used GAMs (Carslaw et al., 2007; Carslaw and Carslaw, 2007; Pan et al., 2013; Richmond-Bryant et al., 2011; Westmoreland et al., 2007) or other statistical models, such as boosted regression trees (Carslaw and Taylor, 2009; Sayegh et al., 2016) or nonparametric regression (Bloomfield et al., 1996) to investigate the role of meteorological variables on dispersion of near-road pollutants. These studies were instructive about the influence of meteorology and mobile sources on ambient concentrations. To our knowledge, our study was the first to use GAMs to model the near-road gradient (rather than concentration) as a function of meteorological variables and mobile sources. Sayegh et al. (2016) measured NO_X concentration at three sites (urban, near-road, and tunnel) in Leeds, UK and examined four categories of traffic (light and free-flowing, heavy but moving at speed limit, congested, and severely congested) using boosted regression trees. The near-road site was most similar to the Las Vegas site, and Sayegh et al. (2016) found that variability in wind direction was responsible for 31% of predicted variability in NO_X at the near-road site, compared with 61.7% of the variability in dC_{NO_X}/dx . Traffic counts on the opposite side of the road, traffic counts near the road on the same side of the road, temperature, wind speed, relative humidity, and Julian day (as a measure of seasonality) each accounted for 8–11% of the predicted variability in NO_X , while upwind NO_X levels, hour, and day of the week accounted for 3–5% each for the Sayegh et al. (2016) study. In comparison with our study, variability in hour of day, upwind

NO_x concentration, and traffic were each associated with 15.8%, 9.5%, and 4.7% of the variability in dC_{NO_x}/dx , respectively, and variability in all other factors were responsible for less than 3% of the variability in dC_{NO_x}/dx . Differences in these contributions were likely partially attributable to the fact that NO_x gradient, rather than concentration, was studied in our paper.

GAM was used to model concentrations of NO₂, NO_x, CO, benzene, and 1,3-butadiene as smoothed functions of year, north-south and east-west components of wind, temperature, Julian day, light duty traffic, heavy duty traffic, and (for NO₂ only) ambient O₃ concentration with good fit ($R^2 = 0.81$) for prediction of trends in the concentrations over the years 1998–2005 (Carslaw and Carslaw, 2007). The authors did not provide information on the relative influence of each variable but did illustrate that omission of meteorological variables from the model resulted in a smaller decline in NO_x concentrations and a slightly larger increase in NO₂ concentrations over the study period. The latter finding adds support for our adoption of meteorological variables into the model. Furthermore, Pan et al. (2013) measured fine PM across five cities in Southern California and then modeled the local PM concentration as a function of background PM_{2.5} concentration, wind speed and direction, momentum flux, sensible heat flux, vertical velocity fluctuation, and traffic count. Variability in background PM_{2.5} dominated the fine PM models in four cities (89.2–93.5%). Omitting background PM_{2.5} from the analysis in those four cities, wind direction was the next most important variable in lower density cities (Anaheim, Pasadena, Long Beach), and vertical velocity fluctuation was most important in one (Los Angeles), probably due to recirculation in street canyons. In the town of Huntington Beach, background PM_{2.5} was only responsible for 34.8% of the variability in fine PM. Sensible heat flux (23.6%) and vertical wind speed (12.0%) were also important in the Pan et al. (2013) study.

Carslaw and Taylor (2009) used boosted regression trees to model NO_x concentration near Heathrow Airport, London, UK in terms of heavy and medium aircraft departures from two runways, wind speed, wind direction, temperature, hour, weekday, Julian day, and a temporal trend term. None of these variables individually dominated the model. Based on analysis of plots of NO_x concentration as a function of each individual factor, wind speed had the largest influence, which is similar to our findings for dC_{NO_x}/dx . Temperature, temporal trend, hour, wind direction, and Julian day were also influential components in Carslaw and Taylor's (2009) model. The aircraft departure terms each contributed less than 5% of the variability in the model. Similar to our findings for wind speed, those lower contributions may have been related to wide confidence intervals around the smoothed functions, especially as the departure rates increased.

Limitations of this study need to be considered when interpreting our model results. Several limitations were already articulated in Richmond-Bryant et al. (2017) and are restated here for the reader's convenience. First, inclusion of scenarios with winds from the east beyond meandering speeds hamper interpretation of the results, since the road was located to the west of the monitors and the objective of this study was to assess near-road gradients. Second, data were removed from the analysis when the 115 m or 300 m site had concentrations higher than those closer to the road, to ensure that the only source considered was Interstate-15. While this measure alleviates concern about the influence of other

sources, it may have also introduced some selection bias and suppression of uncertainties. Third, Interstate-15 was depressed below grade with a shallow embankment. The shallow embankment would have the effect of elevating concentrations at the roadway, but a near-road gradient would still be anticipated based on the wind patterns at and downwind of the road (Baldauf et al., 2009; Kimbrough et al., 2013). Because the depression was not accounted for in the gradient model, estimated on-road concentrations are likely underpredicted, contributing error to the gradient models. Fourth, it is noted in Figs. 3 and 5 that dC_{NO_2}/dx and dC_{NO_X}/dx were not equivalent at 0° and 359° . This discrepancy may derive from the relatively small proportion of winds from the north. A much larger proportion of the data have winds from the south to support any analysis related to winds parallel to the road. Last, this analysis was only performed for Las Vegas. The goal of parameterizing the near-road concentration gradients and exploring how they responded to changes in meteorological and other variables was to develop generalizable relationships. However, until this approach is tested for other cities, we cannot know whether our models are generalizable. Comparison to other models of the relationship between meteorology and concentrations, from the peer-reviewed literature, add some confidence to our analysis. However, to our knowledge, generalized additive modeling of the near-road concentration gradient with respect to meteorological and other variables is novel. Therefore, we cannot draw a direct comparison with other studies in the literature. Relatedly, the GAM assigned large uncertainties when a variable used in a smoothing function had limited data. We saw in Figs. 3 through 6 that there were regions of uncertainty, characterized by some inconsistency between the model and data, where data were more limited. Regions of uncertainty might have also introduced some inconsistencies between the magnitude of the smoothed functions and the F-test analysis. Therefore, it is important to review all of the data together to develop a picture of what variables influence the near-road concentration gradient.

4. Conclusions

Our study lends insight into the local processes that drive the NO_2 or NO_X gradient in the near-road environment. The results of this study suggested that the near-road concentration gradient was influenced by meteorology, traffic, ambient upwind NO_X concentrations, and ambient O_3 concentrations. For the NO_2 gradient, ambient O_3 concentrations and wind direction were the most influential variables. For the NO_X gradient, wind direction, hour of day, and upwind NO_X concentration were most influential. Our results imply that the magnitude of human exposure in the near-road environment would also vary with these meteorological variables. This work may influence future development of NO_2 and NO_X dispersion models. Moreover, the models developed herein provided a means to understand human exposure estimates in the near-road environment by describing those exposures with respect to measurable variables. Therefore, these principles can be applied to studying the effects of heavily trafficked roadways adjacent to side-walks and dwellings for urban design and for forecasting increased exposure conditions to minimize human exposure to air pollution from mobile sources.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank Dr. Stephen McDow, Dr. Stephen Graham, Dr. Steven Dutton, and Dr. John Vandenberg for their thoughtful reviews of this manuscript. The U.S. Environmental Protection Agency, through its Office of Research and Development, partially funded and collaborated in the research described here under Contract No. EP-D-12-044 work assignment 4-10 and 5-05 to the University of North Carolina at Chapel Hill.

References

- Atkinson R. Atmospheric chemistry of VOCs and NO_x. *Atmos. Environ.* 2000; 34:2063–2101.
- Baldauf R, Thoma E, Hays M, Shores R, Kinsey J, Gullett B, et al. Traffic and meteorological impacts on near-road air quality: summary of methods and trends from the Raleigh Near-road Study. *J. Air Waste Manag. Assoc.* 2008; 58:865–878. [PubMed: 18672711]
- Baldauf R, Watkins N, Heist D, Bailey C, Rowley P, Shores R. Near-road air quality monitoring: factors affecting network design and interpretation of data. *Air Qual. Atmos. Health.* 2009; 2:1–9.
- Baldwin N, Gilani O, Raja S, Batterman S, Ganguly R, Hopke P, et al. Factors affecting pollutant concentrations in the near-road environment. *Atmos. Environ.* 2015; 115:223–235.
- Bloomfield P, Royle JA, Steinberg LJ, Yang Q. Accounting for meteorological effects in measuring urban ozone levels and trends. *Atmos. Environ.* 1996; 30:3067–3077.
- Britter RE, Hanna SR. Flow and dispersion in urban areas. *Annu. Rev. Fluid Mech.* 2003; 35:469–496.
- Carslaw DC. Evidence of an increasing NO₂/NO_x emissions ratio from road traffic emissions. *Atmos. Environ.* 2005; 39:4793–4802.
- Carslaw DC, Beevers SD, Ropkins K, Bell MC. Detecting and quantifying aircraft and other on-airport contributions to ambient nitrogen oxides in the vicinity of a large international airport. *Atmos. Environ.* 2006; 40:5424–5434.
- Carslaw DC, Beevers SD, Tate JE. Modelling and assessing trends in traffic-related emissions using a generalised additive modelling approach. *Atmos. Environ.* 2007; 41:5289–5299.
- Carslaw DC, Carslaw N. Detecting and characterising small changes in urban nitrogen dioxide concentrations. *Atmos. Environ.* 2007; 41:4723–4733.
- Carslaw DC, Taylor PJ. Analysis of air pollution data at a mixed source location using boosted regression trees. *Atmos. Environ.* 2009; 43:3563–3570.
- Chaney AM, Cryer DJ, Nicholl EJ, Seakins PW. NO and NO(2) inter-conversion downwind of two different line sources in suburban environments. *Atmos. Environ.* 2011; 45:5863–5871.
- Cimorelli AJ, Perry SG, Venkatram A, Weil JC, Paine R, Wilson RB, et al. AERMOD: A dispersion model for industrial source applications. Part I: General model formulation and boundary layer characterization. *J. Appl. Meteorol.* 2005; 44:682–693.
- di Sabatino S, Kastner-Klein P, Berkowicz R, Britter RE, Fedorovich. The modelling of turbulence from traffic in urban dispersion models - Part I: Theoretical considerations. *Environ. Fluid Mech.* 2003; 3:129–143.
- Durant JL, Ash CA, Wood EC, Herndon SC, Jayne JT, Knighton WB, et al. Short-term variation in near-highway air pollutant gradients on a winter morning. *Atmos. Chem. Phys.* 2010; 10:8341–8352.
- Foken T. 50 years of the Monin-Obukhov similarity theory. *Bound. Layer Meteorol.* 2006; 119:431–447.
- Gordon M, Staebler RM, Liggio J, Li SM, Wentzell J, Lu G, et al. Measured and modeled variation in pollutant concentration near roadways. *Atmos. Environ.* 2012; 57:138–145.
- Hastie T, Tibshirani R. Generalized additive models. *Stat. Sci.* 1986; 1:297–318.
- Hastie T, Tibshirani, RJ. Generalized Additive Models. Chapman & Hall/CRC; London: 1990.
- He M, Dhaniyala S. A dispersion model for traffic produced turbulence in a two-way traffic scenario. *Environ. Fluid Mech.* 2011; 11:627–640.
- Henry R, Vette A, Norris G, Vedantham R, Kimbrough S, Shores RC. Separating the air quality impact of a major highway and nearby sources by non-parametric trajectory analysis. *Environ. Sci. Technol.* 2011; 45:10471–10476. [PubMed: 22044064]

- Hu S, Fruin S, Kozawa K, Mara S, Paulson SE, Winer AM. A wide area of air pollutant impact downwind of a freeway during pre-sunrise hours. *Atmos. Environ.* 2009; 43:2541–2549.
- Janssen NAH, Van Vliet PHN, Aarts F, Harssema H, Brunekreef B. Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmos. Environ.* 2001; 35:3875–3884.
- Karner AA, Eisinger DS, Niemeier DA. Near-roadway air quality: Synthesizing the findings from real-world data. *Environ. Sci. Technol.* 2010; 44:5334–5344. [PubMed: 20560612]
- Kastner-Klein P, Fedorovich E, Rotach MW. A wind tunnel study of organised and turbulent air motions in urban street canyons. *J. Wind Eng. Ind. Aerod.* 2001; 89:849–861.
- Kimbrough ES, Baldauf RW, Watkins N. Seasonal and diurnal analysis of NO₂ concentrations from a long-duration study conducted in Las Vegas, Nevada. *J. Air Waste Manag. Assoc.* 2013; 63:934–942. [PubMed: 24010374]
- Kimbrough S, Owen RC, Snyder M, Richmond-Bryant J. NO to NO₂ conversion rate analysis and implications for dispersion model chemistry methods using Las Vegas, Nevada near-road field measurements. *Atmos. Environ.* 2017; 165:23–34.
- Monin AS, Obukhov AV. Osnovnye zakonomernosti turbulentnogo peremeshivaniya v prizemnom sloye atmosfery (Basic laws of turbulent mixing in the atmosphere near the ground). *Tr Geofiz. Inst., Akad. Nauk. SSSR.* 1954; 24:163–187.
- Nevada, DOT. 2007 Annual Traffic Report. Carson City, NV: 2008.
- Obukhov AM. Some specific features of atmospheric turbulence. *J. Geophys. Res.* 1962; 67:3011–3014.
- Pan H, Bartolome C, Gutierrez E, Princevac M, Edwards R, Boarnet MG, et al. Investigation of roadside fine particulate matter concentration surrounding major arterials in five southern Californian cities. *J. Air Waste Manag. Assoc.* 2013; 63:482–498. [PubMed: 23687733]
- RCore_Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing; Vienna, Austria: 2015.
- Richmond-Bryant J, Bukiewicz L, Kalin R, Galarraga C, Mirer F. A multi-site analysis of the association between black carbon concentrations and vehicular idling, traffic, background pollution, and meteorology during school dismissals. *Sci. Total Environ.* 2011; 409:2085–2093. [PubMed: 21406309]
- Richmond-Bryant J, Owen RC, Graham S, Snyder M, McDow S, Oakes M, et al. Estimation of on-road NO₂ concentrations, NO₂/NO_x ratios, and related roadway gradients from near-road monitoring data. *Air Qual. Atmos. Health.* 2017; 10:611–625.
- Rodes CE, Holland DM. Variations of NO, NO₂ and O₃ concentrations down-wind of a Los Angeles freeway. *Atmos. Environ.* 1981; 15:243–250.
- Roorda-Knappe MC, Janssen NAH, De Hartog JJ, Van Vliet PHN, Harssema H, Brunekreef B. Air pollution from traffic in city districts near major motorways. *Atmos. Environ.* 1998; 32:1921–1930.
- Sander, SP., Abbatt, JPD., Barker, JR., Burkholder, JB., Friedl, RR., Golden, DM., et al. Section 1. Biomolecular Reactions. Jet Propulsion Laboratory; Pasadena, CA: 2011.
- Sayegh A, Tate JE, Ropkins K. Understanding how roadside concentrations of NO_x are influenced by the background levels, traffic density, and meteorological conditions using boosted regression trees. *Atmos. Environ.* 2016; 127:163–175.
- Sedefian L, Rao ST, Czapski U. Effects of traffic-generated turbulence on nearfield dispersion. *Atmos. Environ* (1967). 1981; 15:527–536.
- Seinfeld, JH., Pandis, SN. *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*. Second. John Wiley & Sons; Hoboken, New Jersey: 2006.
- Simon, NW. *Generalized Additive Models : An Introduction with R*. Chapman & Hall/CRC. Boca Raton, Florida: 2006.
- Van Roosbroeck S, Wichmann J, Janssen NAH, Hoek G, Van Wijnen JH, Lebret E, et al. Long-term personal exposure to traffic-related air pollution among school children, a validation study. *Sci. Total Environ.* 2006; 368:565–573. [PubMed: 16650461]
- Venkatram A, Isakov V, Thoma E, Baldauf R. Analysis of air quality data near roadways using a dispersion model. *Atmos. Environ.* 2007; 41:9481–9497.

- Venkatram A, Snyder M, Isakov V, Kimbrough, Su. Impact of wind direction on near-road pollutant concentrations. *Atmos. Environ.* 2013; 80:248–258.
- Wang YJ, Zhang KM. Modeling near-road air quality using a computational fluid dynamics model, CFD-VIT-RIT. *Environ. Sci. Technol.* 2009; 43:7778–7783. [PubMed: 19921893]
- Westmoreland EJ, Carslaw N, Carslaw DC, Gillah A, Bates E. Analysis of air quality within a street canyon using statistical and dispersion modelling techniques. *Atmos. Environ.* 2007; 41:9195–9205.
- Wood, S. Package ‘mgcv’. 2016. Available: <https://cran.r-project.org/web/packages/mgcv/mgcv.pdf>

Fig. 1.
Map of the Las Vegas near-road sampling site (Kimbrough et al., 2017).

Fig. 2. Concentration (in ppb) roses for NO_2 during stable conditions (top left) and unstable conditions (top right), and concentrations roses for NO_x during stable conditions (bottom left) and unstable conditions (bottom right). Concentrations presented in this figure were measured at the 25 m site to the east of I-15.

Fig. 3. GAM models of NO₂ gradient (dC_{NO_2}/dx) with respect to each smooth meteorological function. (first row l-r): Temperature, sensible heat flux, inverse Monin-Obukhov length scale; (second row l-r): wind speed, wind direction, traffic count; (third row, l-r): ambient O₃ concentration, upwind NO_x concentration, hour of day.

Fig. 4. GAM models of NO₂ gradient (dC_{NO_2}/dx) with respect to each smooth meteorological function when winds are constrained to come from the west (210–330°). (first row l-r): Temperature, Sensible heat flux, inverse Monin-Obukhov length scale; (second row l-r): wind speed, traffic count; (third row, l-r): ambient O₃ concentration, upwind NO_x concentration, hour of day.

Fig. 5. GAM models of NO_x gradient (dC_{NO_x}/dx) with respect to each smooth meteorological function. (first row l-r): Temperature, Sensible heat flux, inverse Monin-Obukhov length scale; (second row l-r): wind speed, wind direction, traffic count; (third row, l-r): ambient O₃ concentration, upwind NO_x concentration, hour of day.

Fig. 6. GAM models of NO_x gradient (dC_{NOX}/dx) with respect to each smooth meteorological function when winds are constrained to come from the west (210–330°). (first row l-r): Temperature, Sensible heat flux, inverse Monin-Obukhov length scale; (second row l-r): wind speed, traffic count; (third row, l-r): ambient O₃ concentration, upwind NO_x concentration, hour of day.

Table 1

Summary of the full set of data for meteorological, traffic, and concentration of ambient O₃, NO, and NO_x at the Las Vegas study site.

	Average (SD)	Median (IQR)
Maximum of convective and mechanical mixing height (z_i , m)	831.6 (1063)	301.0 (1233)
Sensible heat flux (ϕ , W/m ²)	67.61 (121.8)	-1.400 (137.8)
Ambient temperature (T, K)	293.5 (10.36)	292.7 (17.80)
Relative humidity (R, %)	25.29 (16.27)	21.00 (18.00)
Atmospheric pressure (P, mbar)	934.8 (6.057)	934.0 (9.000)
Monin-Obukhov length scale (L_{Mon} , m)	-89.79 (906.0)	4.700 (18.30)
Eddy volume at 25 m from road (Pvol, m ³)	26.71 (33.16)	13.02 (37.69)
Standard deviation of velocity (σ_v , m/s)	0.7315 (0.6732)	0.5028 (1.055)
Roughness height (z_0 , m)	326.9 (405.5)	161.2 (386.8)
Streamwise friction velocity (u^* , m/s)	0.2347 (0.1997)	0.1700 (0.2535)
Vertical friction velocity (w^* , m/s)	1.710 (0.9673)	1.741 (1.552)
Wind direction (WD, °) ^a	219.3	229.1
Wind speed (W, m/s)	2.655 (1.879)	1.920 (2.250)
Hourly traffic count (TR, number)	6757 (2633)	6996 (4473)
Ambient O ₃ concentration (O ₃ , ppb)	30.2 (19.4)	31.5 (33.0)
Upwind NO _x concentration (NO _{x,up} , ppb)	30.4 (28.8)	20.8 (35.0)
Hour	11.82 (6.85)	12 (12)
NO₂ Concentration		
25 m downwind (ppb)	25.9 (12.0)	25.2 (19.9)
115 m downwind (ppb)	22.4 (12.1)	21.0 (21.4)
300 m downwind (ppb)	19.5 (12.3)	17.1 (22.5)
NO_x Concentration		
25 m downwind (ppb)	51.0 (36.4)	39.8 (41.7)
115 m downwind (ppb)	37.5 (32.0)	26.2 (35.6)
300 m downwind (ppb)	31.3 (29.7)	20.0 (33.6)

^aCalculated as $\tan^{-1}(\Sigma \sin \Theta / \Sigma \cos \Theta)$ for the average and as $\tan^{-1}[\text{median}(\sin \Theta) / \text{median}(\cos \Theta)]$ for the median. Wind direction calculations were all made in radians and then converted to degrees for presentation in this table.

Table 2

Pearson correlations among meteorological variables. Bold text indicates correlation greater than 0.7 (with the exception of numbers on the diagonal), which prompted removal of some variables from the model to reduce collinearity. H = hour of day, z_i = maximum of convective and mechanical mixing height (m), ϕ = sensible heat flux (W/m^2), T = ambient temperature (K), P = atmospheric pressure (mbar), L_{Mon} = Monin-Obukhov length scale of atmospheric turbulence above the roughness sublayer (m), P_{vol} = eddy volume, σ_V = dispersion, z_0 = roughness height (m), u^* = streamwise friction velocity (m/s), w^* = vertical friction velocity (m/s), W = wind speed (m/s), WD = wind direction ($^\circ$), TR = hourly traffic count (number), O_3 = ambient ozone concentration (ppb), JD = Julian day (day), and $NO_{X,up}$ = upwind NO_X measurement.

	H	z_i	ϕ	T	R	P	L_{Mon}	P_{vol}	σ_V	z_0	u^*	w^*	WD	W	TR	O_3	$NO_{X,up}$
H	1.00																
z_i	0.11	1.00															
ϕ	-0.07	0.67	1.00														
T	0.15	0.51	0.37	1.00													
R	-0.18	-0.37	-0.30	-0.66	1.00												
P	-0.11	-0.34	-0.09	-0.49	0.24	1.00											
L_{Mon}	0.02	-0.02	0.02	0.07	-0.03	-0.06	1.00										
P_{vol}	-0.05	0.60	0.76	0.35	-0.26	-0.05	0.02	1.00									
σ_V	0.04	0.85	0.78	0.42	-0.35	-0.33	-0.02	0.65	1.00								
z_0	0.06	0.55	0.29	0.20	-0.20	-0.42	-0.06	0.10	0.72	1.00							
u^*	0.07	0.60	0.33	0.25	-0.24	-0.44	-0.07	0.17	0.76	0.99	1.00						
w^*	0.32	0.77	0.89	0.60	-0.52	-0.37	0.35	0.50	0.92	0.31	0.36	1.00					
WD	0.01	-0.21	-0.38	-0.13	0.09	-0.09	0.02	-0.40	-0.23	0.12	0.07	-0.09	1.00				
W	0.10	0.46	0.14	0.17	-0.19	-0.43	-0.07	-0.05	0.62	0.96	0.28	0.15	1.00				
TR	0.31	0.62	0.58	0.42	-0.35	-0.19	-0.08	0.54	0.61	0.33	0.39	0.49	-0.28	0.23	1.00		
O_3	0.18	0.72	0.49	0.61	-0.50	-0.50	0.00	0.44	0.72	0.55	0.62	0.74	-0.19	0.54	0.53	1.00	
$NO_{X,up}$	-0.14	-0.37	-0.13	-0.36	0.25	0.42	0.00	-0.04	-0.39	-0.47	-0.51	-0.46	-0.11	-0.53	-0.14	-0.64	1.00

Table 3

GAM results for the gradients dC_{NO_2}/dx and dC_{NO_X}/dx modeled as a linear combination of smooth functions of the meteorological variables. Variables are ordered as ranked for percent of variance.

	dC_{NO_2}/dx				dC_{NO_X}/dx			
	edf ^a	F-statistic	% of Variance	p-value	edf ^a	F-statistic	% of Variance	p-value
s(O ₃)	1.000	199.549	47.5%	< 2.00E-16	8.728	229.787	61.7%	< 2.00E-16
s(WD)	8.832	124.746	29.7%	< 2.00E-16	8.587	58.997	15.8%	< 2.00E-16
s(W)	4.194	19.543	4.7%	2.22E-16	6.979	35.371	9.5%	< 2.00E-16
s(NO _{X,up})	7.723	15.361	3.7%	< 2.00E-16	3.892	17.633	4.7%	7.59E-14
s(ϕ)	5.453	14.854	3.5%	2.95E-15	3.220	8.866	2.4%	4.78E-06
s(H)	7.823	13.829	3.3%	< 2.00E-16	5.685	7.134	1.9%	3.48E-07
s(T)	2.603	12.506	3.0%	4.48E-07	1.762	5.495	1.5%	0.006649
s(L _{Mon} ⁻¹)	6.466	11.714	2.8%	1.14E-13	4.199	4.769	1.3%	0.000648
s(TR)	5.177	8.026	1.9%	1.11E-07	3.159	4.429	1.2%	0.003591
R ² -adj	0.409				0.509			
SSR/edf	0.108				0.873			

^a edf = effective degrees of freedom.

Table 4

GAM results for the gradients dC_{NO_2}/dx and dC_{NO_x}/dx modeled as a linear combination of smooth functions of the meteorological variables when winds are constrained to come from the west (210–330°). Variables are ordered as ranked for percent of variance.

dC _{NO₂} /dx					dC _{NO_x} /dx				
	edf ^a	F-statistic	% of Variance	p-value		edf ^a	F-statistic	% of Variance	p-value
s(O ₃)	3.580	59.351	36.0%	< 2.00E-16	s(NO _{x,up})	6.589	43.807	26.9%	< 2.00E-16
s(W)	2.351	45.655	27.7%	< 2.00E-16	s(W)	1.464	38.678	23.8%	2.59E-12
s(φ)	4.455	13.876	8.4%	6.70E-12	s(H)	7.477	30.767	18.9%	< 2.00E-16
s(TR)	6.023	11.200	6.8%	2.09E-12	s(TR)	4.073	26.759	16.5%	< 2.00E-16
s(L _{Mon} ⁻¹)	6.002	11.175	6.8%	2.36E-12	s(L _{Mon} ⁻¹)	3.896	8.441	5.2%	1.44E-06
s(H)	6.753	9.546	5.8%	2.90E-11	s(O ₃)	4.499	6.147	3.8%	3.30E-05
s(NO _{x,up})	7.457	9.403	5.7%	5.40E-12	s(T)	4.253	4.514	2.8%	1.01E-03
s(T)	6.753	4.445	2.7%	1.07E-03	s(φ)	3.346	3.540	2.2%	1.15E-02
R ² -adj	0.506					0.509			
SSR/edf	0.0711					0.873			

^a edf = effective degrees of freedom.