Published in final edited form as:

Drug Alcohol Depend. 2017 October 01; 179: 13–17. doi:10.1016/j.drugalcdep.2017.06.013.

OPIOID MEDICATION MISUSE AMONG UNHEALTHY DRINKERS

Gerald Cochran^{a,b}, Rebecca McCarthy^c, Adam J. Gordon^{b,c,d,e}, and Ralph E. Tarter^f

^aUniversity of Pittsburgh, School of Social Work, 4200 Forbes Ave. #2006, Pittsburgh PA, 15260, USA, (412) 624-2325

^bUniversity of Pittsburgh, School of Medicine, M240 Scaife Hall, 3550 Terrace St, Pittsburgh, PA 15261, USA

^cVA Pittsburgh Healthcare System, University Drive (151C), Pittsburgh, PA 15224, USA

^fUniversity of Pittsburgh, School of Pharmacy, 3501 Terrace St, Pittsburgh, PA 15213, USA

Abstract

Background—Combining opioid medications and alcohol has serious implications for patient health, including overdose. Information regarding those who use/misuse opioid medications and engage in unhealthy alcohol use is limited to pharmacological and epidemiological descriptions. This study presents opioid medication misuse and behavioral, mental, and physical health characteristics of persons filling opioid medications that are engaged in unhealthy alcohol use.

Methods—We conducted a cross-sectional survey at 5 community pharmacies in Southwestern, Pennsylvania among patients filling opioid medications. Respondents completed validated opioid medication misuse, alcohol use, illicit drug use, depression, posttraumatic stress disorder (PTSD), and physical health functioning assessments. We present univariate and multivariate statistics describing opioid medication misuse and health risks among those positive for unhealthy alcohol use.

Results—A total of 344 patients completed the survey (75.8% response). A total of 15.9% of respondents screened positive for opioid medication misuse, of whom 20.3% reported unhealthy alcohol use. Taking opioid medications too often was reported among a larger proportion of the sample with unhealthy alcohol use (34.3%) compared to those without (22.1%, $p=0.04$). Further, among respondents with unhealthy alcohol use, illicit drug use (Adjusted odds ratio [AOR]=12.14, 95% Confidence Interval [CI]= 1.64–89.72) and PTSD (AOR=9.77, 95% CI= 1.70–56.11) were associated with increased odds for opioid medication misuse.

Conclusion—Results suggest respondents with unhealthy alcohol use had distinct health profiles, which may place them at risk for opioid misuse and adverse events, such as overdose. Continued research must work to further understand these relationships and identify intervention and treatment strategies.

Correspondence to: Gerald Cochran.

^dUniversity of Utah, School of Medicine, 30 N. 1900 E Salt Lake City, Utah 84132.

^eVA Salt Lake City Healthcare System, 500 Foothill Dr, Salt Lake City, UT 84148.

Keywords

Opioid medication misuse; unhealthy drinking; community pharmacy

1.1 Introduction

Approximately 4 million people in the US in 2015 were found to have current opioid medication misuse (SAMHSA, 2016). Individuals who misuse opioid medications have a number of attendant behavioral health conditions, such as depression, anxiety, posttraumatic stress disorder (PTSD), illicit drug use (Amari et al., 2011; Becker et al., 2008; Smith et al., 2016; Sullivan et al., 2010), severe pain (Amari et al., 2011; Hudson et al., 2008; Novak et al., 2009; Sullivan et al., 2010); and general poor health (Becker et al., 2008; Hudson et al., 2008).

Unhealthy alcohol consumption is among these attendant health issues (Wang et al., 2013). One-third of persons with opioid use disorder drink alcohol in excess of NIH guidelines for unhealthy alcohol consumption (Nolan et al., 2016). In 2010, 18.5% (n=81,365) of emergency department visits related to opioid pain medications also involved patient consumption of alcohol (Jones et al., 2014). Unhealthy alcohol use is a risk factor for opioid overdose (CDC, 2016; Cochran et al., 2016): overdose deaths rates are 0.63 per 100,000 lives for those who used both opioid medications and alcohol (Calcaterra et al., 2013).

Given that both opioids and alcohol act upon μ -opioid receptors (Amato et al., 2011), simultaneous use of these substances can potentiate the analgesic and side-effects of opioids (Brands et al., 2008). These effects include increased drowsiness, increased dizziness, impaired motor control, unusual behaviors, memory problems, slowed or difficulty breathing, and increased risk of overdose (Kuerbis et al., 2014). Furthermore, some prescription opioids contain acetaminophen, which can worsen liver damage associated with alcohol intake (Brands et al., 2008).

Information available in the field regarding those who use/misuse opioid medications and engage in unhealthy alcohol use is limited primarily to pharmacological interactions and epidemiological descriptions. Among those with unhealthy alcohol use and use/misuse of opioid medications, little information is available describing their behavioral, mental, and physical health and demographic characteristics as well as increased misuse risks associated with these characteristics. Therefore, among a sample of those filling opioid prescriptions, we sought to (1) describe patient demographics, health characteristics, and opioid medication misuse patterns and (2) estimate opioid medication misuse risks for patients with unhealthy alcohol use and for those without. These data have the potential to assist clinical researchers and practitioners to more effectively understand this population and work toward reducing unnecessary risks.

2.1 Materials and Methods

We surveyed a convenience sample of patients filling opioid pain medications in 5 community pharmacy settings (3 urban and 2 rural) in southwestern Pennsylvania from

September 2014 to June 2015. This study was designated as exempt by the University of Pittsburgh Institutional Review Board.

2.2 Sample

Pharmacists and pharmacy staff members were trained by the principal investigator in the study procedure. When patients dropped off prescriptions identified by pharmacy staff as opioid pain medications, they were asked if they were interested in participating in a brief health survey while waiting. Interested patients were handed an iPad tablet that contained 3 initial screening questions to ensure patients were 18 years of age, were not receiving treatment for a cancer diagnosis, and had not previously completed the survey instrument. Participants were provided with a \$20 gift card for their time.

2.3 Instruments

The survey captured demographics (age, sex, education, work status, and pharmacy rural/urban location), behavioral and mental health, physical functioning, and opioid medication type. The Prescription Opioid Misuse Index assessed current opioid medication misuse, with a score of 2 affirmative responses indicating misuse (Knisely et al., 2008). Unhealthy alcohol use was assessed using the Alcohol Use Disorders Identification Test-C. A score of 3 for women and 4 for men are cut-offs for unhealthy alcohol consumption (Bradley et al., 2007; Bush et al., 1998; Gordon et al., 2001; Williams et al., 2012). Illicit drug use severity in the last year was assessed using the Drug Abuse Screening Test-10 (DAST-10). A score of 1 indicates the need for intervention (Yudko et al., 2007).

Depression screening was conducted using the 2-item Patient Health Questionnaire-2 (PHQ-2). A score of 3 indicates a positive screen (Corson et al., 2004; Kroenke et al., 2003). PTSD was assessed using the 4-item Primary Care Post-Traumatic Stress Disorder (PC-PTSD) screen, with a score of 3 indicating PTSD (Ouimette et al., 2008; Prins et al., 2003; van Dam et al., 2010).

Physical health functioning was assessed using the Short-Form Survey-12 (SF-12). Two single-item subscales within this measure ask patients to indicate on 5-point Likert scales their level of general health (5=poor, 4=fair, 3=good, 2=very good, 1=excellent) and pain that interferes with work (5=extremely, 4=quite a bit, 3=moderately, 2=a little bit, 1=not at all).

Opioid medication type was captured by asking patients, "What is the name of your pain medication?" Patients were given a textbox to enter in the medication name. Medications were coded into generic names for comparability and dichotomized into binary indicators for patients filling more than one opioid.

2.4 Statistical Analyses

To characterize differences among patients who were positive for unhealthy alcohol use and those who were not among patients filling opioid medications, we conducted univariate descriptive analyses for proportional and mean differences using chi-square and t-test statistical analyses. We also report predicted probabilities of engaging in unhealthy alcohol

use for those engaged in opioid misuse. Finally, we developed two logistic regression models to assess relationships between behavioral, mental, and physical health status and a positive screen for opioid medication misuse among (1) those positive for unhealthy alcohol use and (2) those who were not involved in unhealthy alcohol use. Models were adjusted for respondent demographic characteristics. Analyses were conducted using Stata 14.2 (StataCorp, 2016).

3.1 Results

A total of 344 patients completed the survey. The average response rate across the 5 pharmacies was 75.8% (rates: rural pharmacy A= 94.2% [98 completed/104 approached]; rural pharmacy B= 13.3% [75 completed/565 approached], urban pharmacy A=87.7% [100 completed/114 approached]; urban pharmacy B=92.3% [60 completed/65 approached]), and urban pharmacy C=91.7 [11 completed/12 approached]). Analyses showed no statistically significant differences for rural pharmacy B compared to the other pharmacies on any demographic or health indicator, with the exception that rural pharmacy B respondents were underrepresented among those with more than a high school education (29.3%, standardized residual= -1.99) compared to the other pharmacy locations (49.1%, standardized residual=1.06, $p=0.002$, results not shown).

3.2 Demographics

A total of 22.3% of the sample ($n=70$) screened positive for unhealthy alcohol use. No differences for unhealthy alcohol use were detected for age (Mean [M]=49.1, standard deviation[SD]=12.4), gender (female=56.1%), or education (high school or less education=57.6%). A significantly larger portion of the sample with a positive screen for unhealthy alcohol use also reported living in urban settings (61.4%) compared to negative respondents (45.9%, $p=0.02$).

3.3 Opioid Misuse, Behaviors, and Medications

The predicted probability of engaging in unhealthy drinking was 0.21 (95% CI=0.09–0.39) for those who misused their opioid medication and was 0.20 (95% CI=0.14–0.25) for those who did not misuse their opioid medication. Table 1 displays misuse and health characteristics of the sample overall and for those engaged in unhealthy and non-unhealthy alcohol use. Overall, 15.9% of the sample screened positive for opioid medication misuse. Of these, 20.3% were engaged in unhealthy alcohol use. The most common misuse behavior, taking medications too often (24.8%), was reported among a larger proportion of the sample with unhealthy alcohol use (34.3%) compared to those without unhealthy alcohol use (22.1%, $p=0.04$). All other misuse behaviors, although not statistically significant, were reported by a larger portion of the sample with unhealthy alcohol use compared to those without.

Data also suggested hydrocodone was the most common opioid medication filled among those positive for unhealthy alcohol use (47.1%) compared to non-unhealthy consumers of alcohol (34% $p=0.05$). The second largest proportion of the sample with unhealthy alcohol use reported using oxycodone (32.9%).

3.4 Health Characteristics

Significantly larger portions of the sample with unhealthy alcohol use reported illicit drug use (17.1% vs. 6.6%, $p=0.01$), and significantly lower levels of pain ($M=3.4$, $SD=1.2$ vs. $M=3.7$, $SD=1.1$; $p=0.02$) and poor health ($M=3.2$, $SD=1.1$ vs. $M=3.7$, $SD=0.9$; $p<0.001$) compared to those with non-unhealthy use.

3.5 Opioid Misuse Risk

Table 2 displays the two logistic regression models that report the relationship between opioid misuse and patient demographics and behavioral, mental, and physical health indicators for those with unhealthy alcohol use and those without. The multivariate association with the largest observed magnitude was for those with illicit drug use in the last year and opioid misuse among those who were positive for unhealthy alcohol use (Adjusted odds ratio [AOR]=12.14, 95% Confidence Interval [CI]= 1.64–89.72). Furthermore, those within the sample with unhealthy alcohol use were more likely to screen positive for PTSD (AOR=9.77, 95%CI= 1.70–56.11). Among those without unhealthy alcohol use, illicit drug use (AOR=5.20, 95%CI= 1.47–18.43), depression (AOR=3.10, 95%CI=1.23–7.78), and urban pharmacy location (rural living status AOR=0.39, 95%CI= 0.16–0.93) were associated with a positive screen for opioid misuse.

4.1 Discussion

Previous research that describes the individual clinical characteristics of those with unhealthy drinking patterns engaged in opioid use/misuse is limited. Findings from the current study showed patients filling opioid prescriptions with unhealthy alcohol use were more likely to take opioid medications more often than directed by their healthcare provider. Such behavior is problematic as consuming high opioid doses has the potential to elevate overdose risk (Dasgupta et al., 2015). Risk of overdose is increased among patients engaged in using opioids, alcohol, and illicit substances (Calcaterra et al., 2013; Cochran et al., 2016). Additionally, given that hydrocodone and oxycodone were the two most common opioids among those who engage in unhealthy alcohol use, increased risk of liver damage is of real concern considering these are commonly prescribed as combination products with acetaminophen. Clinicians must be vigilant therefore to carefully assess alcohol use when prescribing and filling opioid medications to ensure patients follow safety guidelines for their use.

Our results also suggest a possible important distinction for opioid misuse risk among those with positive screens for depression and PTSD. Specifically, while depression (SAMHSA, 2014) and PTSD (Meier et al., 2014; Smith et al., 2016) have been found to have clear associations with opioid medication misuse in previous investigations, it appears for those engaged in unhealthy alcohol use there is a particular link between a positive screen for PTSD and opioid medication misuse. Previous research supports these results, which has shown heightened concurrence of PTSD and alcohol dependence among opioid dependent patients receiving substitution therapy (Peirce et al., 2008). It should be noted, however, that regardless of unhealthy alcohol use, depression and PTSD were somewhat common among

the sample, which conditions of themselves increase vulnerability for misuse and must be attended to by providers.

4.2 Limitations

This survey was conducted in community pharmacy locations in southwestern Pennsylvania; study results may not generalize to other settings or geographic locations. However, the concordance of our results with previously published works regarding opioid misuse supports the external validity of these findings. Our results extend the current knowledgebase in the field regarding the clinical characteristics of patients who use opioid medications and are involved in unhealthy alcohol use as well as the unique risks for opioid misuse faced by this population. Furthermore, because of the smaller number of those positive for unhealthy alcohol use who completed the survey, we observed somewhat wide confidence intervals for some estimates, which estimates would likely decrease were a larger sample of those with unhealthy alcohol use recruited. Finally, given that all study health indicators were captured using brief screening surveys to minimize patient/pharmacy site burden, lengthier diagnostics may provide a richer description of patient health.

5.1 Conclusion

The results of this study suggest that those with unhealthy alcohol use had distinct health profiles from those without unhealthy use, which may place them at higher risk for opioid misuse and adverse events, such as overdose. Continued research must work to further understand these relationships as well as identify needed intervention and treatment strategies at the provider- and health system-level. Efforts aimed at development and implementation of clinical tools for assessment, intervention, and system-level patient management guidelines have the potential to protect patient health and wellbeing.

References

- Amari E, Rehm J, Goldner E, Fischer B. Nonmedical prescription opioid use and mental health and pain comorbidities: a narrative review. *Can J Psychiatry*. 2011; 56:495–502. [PubMed: 21878161]
- Amato L, Minozzi S, Davoli M, Vecchi S. Psychosocial and pharmacological treatments versus pharmacological treatments for opioid detoxification. *The Cochrane database of systematic reviews*. 2011: Cd005031. [PubMed: 21901695]
- Becker WC, Sullivan LE, Tetrault JM, Desai RA, Fiellin DA. Non-medical use, abuse and dependence on prescription opioids among U.S. adults: Psychiatric, medical and substance use correlates. *Drug Alcohol Depend*. 2008; 94:38–47. [PubMed: 18063321]
- Bradley KA, DeBenedetti AF, Volk RJ, Williams EC, Frank D, Kivlahan DR. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol Clin Exp Res*. 2007; 31:1208–1217. [PubMed: 17451397]
- Brands B, Blake J, Marsh DC, Sproule B, Jeyapalan R, Li S. The impact of benzodiazepine use on methadone maintenance treatment outcomes. *J Addict Dis*. 2008; 27:37–48.
- Bush K, Kivlahan DR, McDonell MB, Fihn SD, Bradley KA. The AUDIT alcohol consumption questions (AUDIT-C): an effective brief screening test for problem drinking. *Ambulatory Care Quality Improvement Project (ACQUIP) Alcohol Use Disorders Identification Test Arch Intern Med*. 1998; 158:1789–1795. [PubMed: 9738608]
- Calcaterra S, Glanz J, Binswanger IA. National trends in pharmaceutical opioid related overdose deaths compared to other substance related overdose deaths: 1999–2009. *Drug Alcohol Depend*. 2013; 131:263–270. [PubMed: 23294765]

- CDC. [Accessed on September 7 2016] Drug overdose deaths in the United States hit record numbers in 2014. 2016. <http://www.cdc.gov/drugoverdose/epidemic/index.html>
- Cochran G, Gordon AJ, Lo-Ciganic WH, Gellad WF, Frazier W, Lobo C, Chang CH, Zheng P, Donohue JM. An Examination of Claims-based Predictors of Overdose from a Large Medicaid Program. *Med Care*. 2016; 55:291–298.
- Corson K, Gerrity MS, Dobscha SK. Screening for depression and suicidality in a VA primary care setting: 2 items are better than 1 item. *Am J Manag Care*. 2004; 10:839–845. [PubMed: 15609737]
- Dasgupta N, Funk MJ, Proescholdbell S, Hirsch A, Ribisl KM, Marshall S. Cohort Study of the Impact of High-dose Opioid Analgesics on Overdose Mortality. *Pain Med*. 2015
- Gordon AJ, Maisto SA, McNeil M, Kraemer KL, Conigliaro RL, Kelley ME, Conigliaro J. Three questions can detect hazardous drinkers. *J Fam Pract*. 2001; 50:313–320. [PubMed: 11300982]
- Hudson TJ, Edlund MJ, Steffick DE, Tripathi SP, Sullivan MD. Epidemiology of regular prescribed opioid use: results from a national, population-based survey. *J Pain Symptom Manage*. 2008; 36:280–288. [PubMed: 18619768]
- Jones CM, Paulozzi LJ, Mack KA. Alcohol involvement in opioid pain reliever and benzodiazepine drug abuse-related emergency department visits and drug-related deaths - United States, 2010. *MMWR*. 2014; 63:881–885. [PubMed: 25299603]
- Knisely JS, Wunsch MJ, Cropsey KL, Campbell ED. Prescription Opioid Misuse Index: a brief questionnaire to assess misuse. *J Subst Abuse Treat*. 2008; 35:380–386. [PubMed: 18657935]
- Kroenke K, Spitzer RL, Williams JBW. The Patient Health Questionnaire-2: Validity of a Two-Item Depression Screener. *Med Care*. 2003; 41:1284–1292. [PubMed: 14583691]
- Kuerbis A, Sacco P, Blazer DG, Moore AA. Substance abuse among older adults. *Clin Geriatr Med*. 2014; 30:629–654. [PubMed: 25037298]
- Meier A, Lambert-Harris C, McGovern MP, Xie H, An M, McLeman B. Co-occurring prescription opioid use problems and posttraumatic stress disorder symptom severity. *Am J Drug Alcohol Abuse*. 2014; 40:304–311. [PubMed: 24809229]
- Nolan S, Klimas J, Wood E. Alcohol use in opioid agonist treatment. *Addict Sci Clin Pract*. 2016; 11:17. [PubMed: 27931253]
- Novak SP, Herman-Stahl M, Flannery B, Zimmerman M. Physical pain, common psychiatric and substance use disorders, and the non-medical use of prescription analgesics in the United States. *Drug Alcohol Depend*. 2009; 100:63–70. [PubMed: 19010611]
- Ouimette P, Wade M, Prins A, Schohn M. Identifying PTSD in primary care: Comparison of the Primary Care-PTSD Screen (PC-PTSD) and the General Health Questionnaire-12 (GHQ). *J Anxiety Disord*. 2008; 22:337–343. [PubMed: 17383853]
- Peirce JM, Kindbom KA, Waesche MC, Yuscavage AS, Brooner RK. Posttraumatic stress disorder, gender, and problem profiles in substance dependent patients. *Subst Use Misuse*. 2008; 43:596–611. [PubMed: 18393079]
- Prins A, Ouimette P, Kimerling R, Cameron RP, Hugelshofer DS, Shaw-Hegwer J, Thrailkill A, Gusman FD, Sheikh J. The primary care PTSD screen (PC-PTSD): Development and operating characteristics. *Primary Care Psychiatry*. 2003; 9:9–14.
- SAMHSA. Results from the 2013 National Survey on Drug Use and Health: Mental Health Detailed Tables. Substance Abuse and Mental Health Services Administration; Rockville, MD: 2014.
- SAMHSA. Key Substance Use and Mental Health Indicators in the United States: Results from the 2015 National Survey on Drug Use and Health. Substance Abuse and Mental Health Services Administration; Rockville, MD: 2016.
- Smith KZ, Smith PH, Cercone SA, McKee SA, Homish GG. Past year non-medical opioid use and abuse and PTSD diagnosis: interactions with sex and associations with symptom clusters. *Addict Behav*. 2016; 58:167–174. [PubMed: 26946448]
- StataCorp. Stata Statistical Software: Release 14.2. StataCorp LP; College Station, TX: 2016.
- Sullivan MD, Edlund MJ, Fan MY, Devries A, Brennan Braden J, Martin BC. Risks for possible and probable opioid misuse among recipients of chronic opioid therapy in commercial and medicaid insurance plans: The TROUP Study. *Pain*. 2010; 150:332–339. [PubMed: 20554392]

- Van Dam D, Ehrling T, Vedel E, Emmelkamp PMG. Validation of the Primary Care Posttraumatic Stress Disorder screening questionnaire (PC-PTSD) in civilian substance use disorder patients. *J Subst Abuse Treat.* 2010; 39:105–113. [PubMed: 20598826]
- Wang KH, Becker WC, Fiellin DA. Prevalence and correlates for nonmedical use of prescription opioids among urban and rural residents. *Drug Alcohol Depend.* 2013; 127:156–162. [PubMed: 22819293]
- Williams EC, Lapham GT, Hawkins EJ, Rubinsky AD, Morales LS, Young BA, Bradley KA. Variation in documented care for unhealthy alcohol consumption across race/ethnicity in the Department of Veterans Affairs Healthcare System. *Alcohol Clin Exp Res.* 2012; 36:1614–1622. [PubMed: 22404130]
- Yudko E, Lozhkina O, Fouts A. A comprehensive review of the psychometric properties of the Drug Abuse Screening Test. *J Subst Abuse Treat.* 2007; 32:189–198. [PubMed: 17306727]

Table 1

Opioid misuse, misuse behaviors, medications, and health characteristics (N=344)

	Unhealthy alcohol use					
	Total %	(n)	Yes %	(n)	No %	(n)
Opioid Misuse						
						χ^2
						p
Positive misuse status	15.9	(49)	20.3	(14)	14.6	(35)
						1.3
						0.25
Misuse Behaviors						
Taking more medication than prescribed	12.8	(40)	15.7	(11)	11.9	(29)
						0.70
						0.40
Taking medication too often	24.8	(78)	34.3	(24)	22.1	(54)
						4.30
						0.04
Early Refills	12.5	(39)	15.7	(11)	11.5	(28)
						0.88
						0.35
Medication Buzz	9.3	(29)	12.9	(9)	8.2	(20)
						1.38
						0.24
Medication to deal with problems	3.9	(12)	4.3	(3)	3.7	(9)
						0.05
						0.74 ^a
Doctor shopping	2.2	(7)	2.9	(2)	2.1	(5)
						0.18
						0.65 ^a
Type of medication						
Hydrocodone	36.9	(116)	47.1	(33)	34.0	(83)
						4.02
						0.05
Hydromorphone	3.5	(11)	1.4	(1)	4.1	(10)
						1.14
						0.47 ^a
Oxycodone	40.5	(127)	32.9	(23)	42.6	(104)
						2.15
						0.14
Oxymorphone	5.7	(18)	0.0	(0)	7.4	(18)
						5.47
						0.02 ^a
Morphine	7.6	(24)	1.4	(1)	9.4	(23)
						4.92
						0.02 ^a
Fentanyl	3.8	(12)	1.4	(1)	4.5	(11)
						1.40
						0.48 ^a
Methadone	3.2	(10)	1.4	(1)	3.7	(9)
						0.90
						0.47 ^a
Health characteristics						
Illicit drug use in the last year	8.9	(28)	17.1	(12)	6.6	(16)
						7.50
						0.01
Pain interferes with daily activities ^b	3.6	(1.1)	3.4	(1.2)	3.7	(1.1)
						2.00
						0.02
Poor general health ^b	3.6	(0.9)	3.2	(1.1)	3.7	(0.9)
						4.09
						<.001
Depression	27.4	(86)	27.1	(19)	27.5	(67)
						0.00
						0.96

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Unhealthy alcohol use						
	Yes %	(n)	No %	(n)	χ^2	<i>p</i>
Post-traumatic stress disorder	22.1	(15)	16.8	(40)	0.99	0.32
Total	18.0	(55)				

q Fisher exact test.

b Mean (Standard deviation) t-value

Table 2

Comparison of opioid misuse risk factors for non-hazardous and hazardous drinkers

Health characteristics	Non-unhealthy drinkers				Unhealthy drinkers			
	AOR ^a	SE ^b	p ^c	95% CI ^d	AOR	SE	p	95% CI
Illicit drug use in the last year	5.20	3.4	0.01	(1.47–18.43)	12.14	12.4	0.01	(1.64–89.72)
Depression	3.10	1.5	0.02	(1.23–7.78)	0.97	0.9	0.98	(0.15–6.47)
Post-traumatic stress disorder	2.32	1.2	0.09	(0.88–6.15)	9.77	8.7	0.01	(1.70–56.11)
Poor general health	0.94	0.3	0.81	(0.55–1.60)	1.39	0.7	0.48	(0.56–3.50)
Pain interferes with daily activities	1.37	0.3	0.18	(0.86–2.18)	1.31	0.6	0.56	(0.53–3.21)
Demographic characteristics								
Age	0.99	0.0	0.60	(0.95–1.03)	0.98	0.0	0.48	(0.91–1.04)
Gender	0.77	0.3	0.55	(0.33–1.82)	1.46	1.2	0.65	(0.29–7.40)
Rural living	0.39	0.2	0.03	(0.16–0.93)	0.76	0.7	0.76	(0.12–4.66)
>High school	0.53	0.3	0.18	(0.21–1.34)	1.39	1.2	0.70	(0.25–7.67)
Employed	1.50	0.8	0.44	(0.53–4.21)	1.84	1.7	0.50	(0.31–10.90)

^a AOR=adjusted odds ratio.^b SE= standard error.^c p=probability value.^d 95% CI=95% confidence interval