Published in final edited form as:

Chem Phys Lett. 2017 September 1; 683: 276–280. doi:10.1016/j.cplett.2017.02.049.

Improved Treatment of Nucleosides and Nucleotides in the OPLS-AA Force Field

Michael J. Robertson, Julian Tirado-Rives, and William L. Jorgensen*

Department of Chemistry, Yale University, New Haven, Connecticut 06520-8107, United States

Abstract

DFT calculations have been used to develop improved descriptions of the torsional energetics for nucleosides and nucleotides in the OPLS-AA force field. Scans of nucleotide dihedral angles (γ , χ , and β) and methyl phosphates provided the bases for the new torsional parameters. In addition, the angle-bending parameters of phosphodiester and ribose were updated, and adjustments were made to existing carbohydrate torsions to better capture the sugar puckering landscape of ribose. MD simulations of nucleosides with the new parameters demonstrate a significant improvement in the ribose sugar puckering and χ angle distributions. Additionally, energy-minimization of protein-nucleotide crystal structures with the new parameters produced accurate poses.

Graphical Abstract

Keywords

Molecular Dynamics; Nucleoside; Nucleotide; Force Field

Introduction

Molecular dynamics (MD) and Monte Carlo statistical mechanics simulations are being increasingly used for characterization of biomolecular systems and for structure-based drug design. To enhance the accuracy of the modeling, the description of torsional energetics for proteins was recently improved in the OPLS-AA force field.¹ The revised force field is

*Corresponding Author: william.jorgensen@yale.edu. Phone: 203-432-6278. Fax: 203-432-6299.

Notes

The authors declare no competing financial interest.

Supporting Information

A word document containing a full description of the simulation details along with CHARMM-formatted topology and parameter files. This material is available free of charge via the Internet at <http://pubs.acs.org>.

referred to as OPLS-AA/M, and it was shown to provide significant gains in performance in MD simulations of proteins and protein-ligand complexes in comparison to NMR data.^{1,2} However, systems of interest often include additional biomolecular components, including membranes, polynucleotides, and carbohydrates. Simple nucleotides such as ATP also interact with numerous important drug targets including kinases³ and Ras proteins.⁴ OPLS-AA force field parameters for nucleoside bases and polynucleotides were developed in the 1990s. However, parameters for the torsional energetics were simply adopted from small organic molecules, and the only published testing on polynucleotides was in simulations of a duplex DNA dodecamer.^{5,6} In the spirit of OPLS-AA/M, results of density functional theory (DFT) calculations have also been used as a basis for improved torsional energetics for nucleotides. These efforts are described here along with results of testing for conformer populations in MD simulations.

Methods

Molecular dynamics simulations were run in NAMD.⁷ All simulations were held at a temperature of 300 K and a pressure of 1 atm using a Langevin thermostat with a damping coefficient of 1 ps⁻¹ and a Nose-Hoover Langevin piston barostat with a piston dampening timescale of 50 fs and period of 100 fs. Non-bonded interactions were smoothed from 9 Å to a cutoff at 11 Å, with particle mesh Ewald for the long-range electrostatics. SHAKE and SETTLE were employed so that a 2-fs time step could be used. Each nucleoside was solvated in a 33-Å periodic cube of TIP3P water.⁸ All energy-minimizations of the systems were performed with NAMD and used 500 steps of the conjugate gradient procedure. In the case of the protein-nucleotide system, crystallographic magnesium and waters were retained, and the water molecules were given random initial orientations.

Quantum mechanical dihedral scans were performed using Gaussian 09.⁹ Relaxed gas-phase scans were carried out for dihedral angles β , γ , and χ in ribose and derivatives, as illustrated in Figure 1, as well as for all dihedral angles in methyl di- and triphosphate (**1C**, **1D**). The scans used an increment of 15°; full details are in the Supplementary Information. As in the development of OPLS-AA/M for proteins,¹ each scan point was calculated at the B2PLYP-D3BJ/aug-cc-pVTZ// ω B97X-D/6-311++G(d,p) level of theory.¹⁰⁻¹³ Corresponding molecular mechanics scans were performed with the BOSS program¹⁴ and OPLS-AA force fields.¹⁵

Equations for the angle bending and torsional portions of OPLS force fields are provided in Equations 1 and 2:

$$E_{angles} = \sum_i k_{\theta} (\theta - \theta_0)^2 \quad (1)$$

$$E_{torsion} = \sum_i \frac{V_1^i}{2} [1 + \cos(\phi_i)] + \frac{V_2^i}{2} [1 - \cos(2\phi_i)] + \frac{V_3^i}{2} [1 + \cos(3\phi_i)] + \frac{V_4^i}{2} [1 - \cos(4\phi_i)] \quad (2)$$

Here, θ is a bond angle, θ_0 is the normal reference value, k_θ is the bending force constant, ϕ is a dihedral angle, and V_1, V_2, V_3, V_4 are Fourier coefficients. The equilibrium bond angles and force constants and the first three Fourier coefficients were considered in the parameterization; V_4 terms are not needed for the present cases. The five angle bending parameters for ribose in the CHARMM36¹⁶ force field were adopted, and a new OPLS atom type 'C5' was added to allow implementation in the BOSS software. Previously, angle bending for the atoms in five-membered rings was treated the same as for normal alkanes and ethers. The original torsional parameters developed for glucose¹⁷ were largely retained for ribose, with a few exceptions. The γ dihedral parameters for both a 5' hydroxyl and a 5' phosphodiester were fit to the results of the DFT scans along with χ parameters. The fitting was done to minimize a Boltzmann-weighted error function with a weighting temperature of 2000 K, which was previously found to work well for peptides (Equation 3),¹

$$Error = \sqrt{\frac{\sum (E_{MM} - E_{QM})^2 e^{-\frac{E_{QM}}{k_B T}}}{n}} \quad (3)$$

where E_{MM} and E_{QM} are the molecular and quantum mechanics energy values for a given geometry, T is the temperature used in the fitting to bias low-energy regions of the surface, and k_B is the Boltzmann constant.

The full two-dimensional sugar-puckering surface was calculated in BOSS, and the effect of each Fourier component (V_1, V_2, V_3) for each sugar dihedral parameter (CT-CT-CT-O?, OH-CT-CT-OS, OH-CT-CT-OH, and CT-CT-CT-CT, where O? = OH or OS) on the potential energy surface was determined. Terms that were found to affect only the relative populations of C3' and C2' endo geometries or affect only the barrier between the two states were identified and empirically modified to produce the correct distribution in solution.

Results

Parameterization efforts began with adenosine, particularly focusing on the ribose sugar puckering. Aqueous-phase MD simulations were performed with OPLS-AA parameters for nucleosides,⁵ and the populations of the sugar puckering pseudorotation angles were calculated. Analogous simulations were performed with the CHARMM36 force field for adenosine. As presented in Figure 2, the results with OPLS-AA are qualitatively incorrect, with a single broad peak centered at a pseudorotation angle of 60°, corresponding to neither the C2' or C3' endo state (165° and 15°, respectively). An attempt was made to fit new dihedral parameters to match the quantum mechanical sugar puckering surface of Huang *et al.*¹⁸ This approach, while improving the geometries, resulted in almost entirely the C3' endo conformation. Normally, such gas-phase data has transferred well over to the condensed phase. The failure in this case, even after the angle bending terms were modified, was unusual. This could be a result of the atoms being in a ring, which may exacerbate issues with the standard description of nonbonded energetics. This includes ignoring 1,2- and 1,3-interactions and scaling 1,4-interactions by 0.5. Further complicating matters is that

the gas-phase QM scan¹⁸ had all alcohol hydrogens pointing away from hydrogen bond acceptors, which is not the case in the solution structures.

As noted for OPLS-AA, the angle bending terms in the ribose ring were utilizing the acyclic alkane and ether reference angles (e.g., 112.7° for CT-CT-CT). In contrast, both the CHARMM parameters and the DFT structures have significantly different equilibrium angles; e.g., ca. 100° for CT-CT-CT. When the five ribose angle bending parameters from CHARMM were adopted with OPLS-AA, two distinct sugar conformations do arise with the correct pseudorotation angles (Figure 2). However, the C3' endo conformer is still overly favored. New parameters for the 5'-hydroxyl γ angle (depicted in Figure 1A where R = OH) were fit based upon DFT scans with both sugar puckerings, which resulted in small improvements in the ring puckering populations, but further refinement was desired. The effect of each component of each dihedral parameter on the molecular mechanics sugar puckering potential energy surface was quantified, and parameters that affected the relative energies of the C2' and C3' endo conformations were identified. A minimal number of terms was then adjusted to improve the populations, primarily the V_1 and V_2 of the C5-C5'-O? parameter. This parameter applies to all torsions involving alcohol (OH) and ether (OS) oxygen atoms in ribose, with the exception of the 5'-exocyclic OH torsion.

These improved ribose parameters were applied to all other nucleosides, and new glycosidic χ dihedral parameters were fit from DFT scans with one set of parameters for purines and one set for pyrimidines. The results of MD simulations for all nucleosides are presented in Table 1 along with experimental NMR data for sugar puckering and glycosidic dihedral populations.¹⁹ The improvement was significant in all cases. The one failing of the new parameters is that purines and pyrimidines have roughly the same sugar puckering preferences, whereas experimentally the pyrimidines have a higher population of the C3' endo state than the purines. This most likely is also the reason why the new glycosidic dihedral parameters for cytosine and uracil somewhat underestimate the population of conformers with the χ dihedral *anti*, as there is correlation between the two degrees of freedom.

Next, the dihedral angle β was addressed. There is precedent in the literature for parameterizing β based upon scans with ethyl methyl phosphate (Figure 1B), holding all other dihedral angles fixed.²⁰ Alternatively, β and γ are often scanned as a two-dimensional surface with the construct in Figure 1A possessing methyl phosphate at the 5' end (R = PO₄CH₃) and a ribose sugar.^{20,21} Initially, we chose to explore how well parameterizing β based on scans without the ribose would perform for nucleotides. Thus, the β parameters were fit to reproduce torsion scans of ethyl methyl phosphate (**1B**). DFT scans were also performed for **1A** (R = ⁻OPO₃CH₃, X = H) with several different ribose puckering and phosphate dihedral states. The results of these DFT scans of **1A** were compared to molecular mechanics scans of **1A** with the β parameters generated from **1B**. The new parameters failed to produce well the conformational energetics for **1A**. However, scans with the existing OPLS-AA parameters were found to perform well, and so these parameters were kept. For γ , one dimensional scans were made with construct **1A** (R = ⁻OPO₃CH₃, X = H), holding the sugar puckering and α angle fixed at four different sets of values. Parameterization with this approach was straightforward.

The final step was to consider methyl di- and tri-phosphate (**1C** and **1D**). A viable approach for torsional parameterization of these molecules was developed previously,²² and that procedure was largely adopted. Again, B2PLYP-D3BJ/aug-cc-pVTZ// ω B97X-D/6-311++G(d,p) was utilized in the present scans. In addition, modifications were made to several partial charge terms to incorporate di- and triphosphates into the OPLS-AA force field, while maintaining integer charges and optimizing transferability between the different molecules considered here (Supplementary Information).

Once new parameters were fit, the conformers of **1C** (MDP) and **1D** (MTP) reported by Meagher *et al.*²² were examined. Energy-minimizations were performed at the B2PLYP-D3BJ/aug-cc-pVTZ// ω B97X-D/6-311++G(d,p) level, although MTP conformer 1 (Figure 3) proved to be too shallow a minimum to reproduce. The overall results for relative energies, bond angles, and dihedral angles from the ω B97X-D/6-311++G(d,p) optimizations for MTP conformations 2, 3, and 4 are compared with those from OPLS-AA and the present OPLS-AA/M in Table 2. The improvements in both energies and geometries are substantial using OPLS-AA/M. The strong preference for conformer 2 is now reproduced, where the *gauche* orientation of the methoxy group can be attributed to an anomeric interaction with a σ^*_{PO} orbital.

The finalized OPLS-AA/M parameters were then placed in CHARMM-formatted topology and parameter files, which are provided as Supplementary Information. For further validation, the crystal structures of four protein-nucleotide complexes were energy-minimized using NAMD (Figure 4). Though further testing in MD simulations is desired, the new parameters performed very well in reproducing the bound poses for the range of nucleotides. This includes both the conformation of the sugar and the phosphate tail. The cases where the force field produced less optimal results either had the nucleotide highly exposed to solvent or there was a lack of a coordinating metal ion in the structure.

Conclusions

Significant improvements have been made for the representation of ribose and its nucleosides and nucleotides in the OPLS-AA force field. The essential parameterization featured fitting to results of numerous torsional energy scans using DFT calculations for the key dihedral angles in the sugar, nucleosides, and polyphosphates (Figure 1). MD simulations for the four RNA nucleosides in water with the new force field showed significant improvement for the populations of the ribose ring conformers and the orientation of the nucleoside bases. Additional scanning for methyl diphosphate and methyl triphosphate provided the necessary torsional parameters for representation of important nucleotides including GDP, ATP, CTP, and UTP. Finally, energy minimizations for these four nucleotides bound to different proteins were shown to converge to structures close to those observed by X-ray crystallography. The foundation is now much firmer for additional testing and application in simulations of nucleotides, polynucleotides and their complexes with other biomolecules using the OPLS-AA/M force field.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Gratitude is expressed to the National Institutes of Health (GM32136) for primary support, to the National Science Foundation for a Graduate Research Fellowship under Grant No. DGE-1122492 (M.J.R.), and to the Yale Center for Research Computing for computer time. Prof. Alex MacKerell is sincerely thanked for generously providing the CHARMM-formatted files that were modified for OPLS-AA/M and for checking their implementation.

References

1. Robertson MJ, Tirado-Rives J, Jorgensen WL. Improved Peptide and Protein Torsional Energetics with the OPLS-AA Force Field. *J Chem Theory Comput.* 2015; 11:3499–3509. [PubMed: 26190950]
2. Robertson MJ, Tirado-Rives J, Jorgensen WL. Performance of Protein-Ligand Force Fields for the Flavodoxin-Flavin Mononucleotide System. *J Phys Chem Lett.* 2016; 15:3032–3036.
3. Cohen P, Alessi DR. Kinase Drug Discovery – What’s Next in the Field? 2013. *ACS Chem Biol.* 2013; 8:96–104. [PubMed: 23276252]
4. Cox AD, Fesik SW, Kimmelman AC, Luo J, Der CJ. Drugging the undruggable RAS: Mission Possible? *Nature Reviews Drug Discovery.* 2014; 13:828–851. [PubMed: 25323927]
5. Pranata J, Wierschke SG, Jorgensen WL. OPLS Potential Functions for Nucleotide Bases. Relative Association Constants of Hydrogen-Bonded Base Pairs in Chloroform. *J Am Chem Soc.* 1991; 113:2810–2819.
6. Ulmschneider JP, Jorgensen WL. Monte Carlo Backbone Sampling for Nucleic Acids Using Concerted Rotations Including Variable Bond Angles. *J Phys Chem B.* 2004; 108:16883–16892.
7. Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kale L, Schulten K. Scalable molecular dynamics with NAMD. *J Comput Chem.* 2005; 26:1781–1802. [PubMed: 16222654]
8. Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. Comparison of Simple Potential Functions for Simulating Liquid Water. *J Chem Phys.* 1983; 79:926–935.
9. Frisch, MJ., Trucks, GW., Schlegel, HB., Scuseria, GE., Robb, MA., Cheeseman, JR., Scalmani, G., Barone, V., Mennucci, B., Petersson, GA., Nakatsuji, H., Caricato, M., Li, X., Hratchian, HP., Izmaylov, AF., Bloino, J., Zheng, G., Sonnenberg, JL., Hada, M., Ehara, M., Toyota, K., Fukuda, R., Hasegawa, J., Ishida, M., Nakajima, T., Honda, Y., Kitao, O., Nakai, H., Vreven, T., Montgomery, JA., Jr, Peralta, JE., Ogliaro, F., Bearpark, M., Heyd, JJ., Brothers, E., Kudin, KN., Staroverov, VN., Kobayashi, R., Normand, J., Raghavachari, K., Rendell, A., Burant, JC., Iyengar, SS., Tomasi, J., Cossi, M., Rega, N., Millam, NJ., Klene, M., Knox, JE., Cross, JB., Bakken, V., Adamo, C., Jaramillo, J., Gomperts, R., Stratmann, RE., Yazyev, O., Austin, AJ., Cammi, R., Pomelli, C., Ochterski, JW., Martin, RL., Morokuma, K., Zakrzewski, VG., Voth, GA., Salvador, P., Dannenberg, JJ., Dapprich, S., Daniels, AD., Farkas, Ö., Foresman, JB., Ortiz, JV., Cioslowski, J., Fox, DJ. Gaussian 09: revision D.01. Gaussian, Inc; Wallingford, CT: 2009.
10. Grimme S. Semiempirical hybrid density functional with perturbative second-order correlation. *J Chem Phys.* 2006:124.
11. Grimme S, Ehrlich S, Goerigk L. Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *J Comput Chem.* 2011; 32:1456–1465. [PubMed: 21370243]
12. Kendall RA, Dunning TH Jr, Harrison RJ. Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions. *J Chem Phys.* 1992; 96:6796–6806.
13. Chai J, Head-Gordon M. Long-range corrected hybrid density functionals with damped atom-atom dispersion corrections. *Phys Chem Chem Phys.* 2008; 10:6615–6620. [PubMed: 18989472]
14. Jorgensen WL, Tirado-Rives J. Molecular modeling of organic and biomolecular systems using BOSS and MCPRO. *J Comput Chem.* 2005; 26:1689–1700. [PubMed: 16200637]

15. Jorgensen WL, Maxwell DS, Tirado-Rives J. Development and testing of the OPLS all-atom force field on conformational energetics and properties of organic liquids. *J Am Chem Soc.* 1996; 118:11225–11236.
16. Denning EJ, Priyakumar UD, Nilsson L, MacKerell AD. Impact of 2'-hydroxyl sampling on the conformational properties of RNA: Update of the CHARMM all-atom additive force field for RNA. *J Comput Chem.* 2011; 32:1929–1943. [PubMed: 21469161]
17. Damm W, Frontera A, Tirado-Rives J, Jorgensen WL. OPLS All-Atom Force Field for Carbohydrates. *J Comput Chem.* 1997; 16:1955–1970.
18. Huang M, Giese TJ, Lee TS, York DM. Improvement of DNA and RNA Sugar Pucker Profiles from Semiempirical Quantum Methods. *J Chem Theory Comput.* 2014; 10:1538–1545. [PubMed: 24803866]
19. Yildirim I, Stern HA, Kennedy SD, Tubbs JD, Turner DH. Reparameterization for RNA χ Torsion Parameters for the AMBER Force Field and Comparison to NMR Spectra for Cytidine and Uridine. *J Chem Theory Comput.* 2010; 6:1520–1531. [PubMed: 20463845]
20. Yildirim I, Kennedy SD, Stern HA, Hart JM, Kierzek R, Turner DH. Revision of AMBER Torsional Parameters for RNA Improved Free Energy Predictions for Tetramer Duplexes with GC and iG/C Base Pairs. *J Chem Theory Comput.* 2011; 8:172–181. [PubMed: 22249447]
21. Pérez A, Marchán I, Svozil D, Sponer J, Cheatham TE III, Laughton CA, Orozco M. Refinement of the AMBER Force Field for Nucleic Acids: Improving the Description of α/γ Conformers. *Biophys J.* 2007; 92:3817–3829. [PubMed: 17351000]
22. Meagher KL, Redman LT, Carlson HA. Development of Polyphosphate Parameters for Use with the AMBER Force Field. *J Comput Chem.* 2003; 24:1016–1025. [PubMed: 12759902]
23. Hunter JC, Gurbani D, Ficarro SB, Carrasco MA, Lim SM, Choi HG, Xie T, Marto JA, Chen Z, Gray NS, Westover KD. In Situ Selectivity Profiling and Crystal Structure of SML-8-73-1: an Active Site Inhibitor of Oncogenic K-Ras G12C. *Proc Natl Acad Sci.* 2014; 111:8895–8900. [PubMed: 24889603]
24. Hopfner KP, Karcher A, Shin DS, Craig L, Arthur LM, Carney JP, Tainer JA. Structural Biology of Rad50 ATPase: ATP-driven Conformational Control in DNA Double-Strand Break Repair and the ABC-ATPase Superfamily. *Cell.* 2000; 101:789–800. [PubMed: 10892749]
25. Björkelid C, Bergfors T, Henriksson LM, Stern AL, Unge T, Mowbray SL, Jones TA. Structural and Functional Studies of Mycobacterial IspD Enzymes. *Acta Crystallogr Sect D-Biol Crystallogr.* 2011; 67:403–414. [PubMed: 21543842]
26. Mori G, Chiarelli LR, Esposito M, Makarov V, Bellinzoni M, Hartkoorn RC, Degiamcomi G, Boldrin F, Ekins S, de Jesus Lopes Riberio AL, Marino LB, Centárová I, Svetlíková Z, Blaško J, Kazakova E, Lepioshkin A, Barilone N, Zanoni G, Porta A, Fondi M, Fani R, Baulard AR, Mikušová K, Alzari PM, Manganelli R, de Carvalho LP, Riccardi G, Cole ST, Pasca MR. *Thiophenecarboxamide* Derivatives Activated by EthA Kill Mycobacterium Tuberculosis by Inhibiting the CTP Synthetase PyrG. *Chem Biol.* 2015; 22:917–927. [PubMed: 26097035]

Figure 1.
Schematic of the treated molecules and scanned dihedral angles.

Figure 2.
MD results for the ribose pseudorotation angle populations of adenosine.

Figure 3.
Structures of MTP conformers 1 (A), 2 (B), 3 (C), and 4 (D).

Figure 4.

Comparisons of crystal structures and minimized molecular mechanics structures of nucleotides bound to proteins. A) GDP bound to KRas, PDB: 4OBE.²³ B) ATP bound to RAD50, PDB: 1F2U.²⁴ C) CTP bound to IspD, PDB: 2XWL.²⁵ D) UTP bound to PyrG, 4ZDJ.²⁶ In all images the crystal structure is in cyan and the minimized structure is in green. Water molecules have been removed for clarity.

Table 1
Conformer Populations from MD Simulations of Nucleosides and Experimental Data.

		OPLS-AA	OPLS-AA New Angle	OPLS-AA/M	Expt ^a
Population C3' endo (%)	Adenosine	0 ^b	82	44	37
	Guanosine	0 ^b	83	45	41
	Cytosine	0 ^b	77	46	60
	Uracil	0 ^b	65	40	56
Population <i>anti</i> (%)	Adenosine	71	71	10	
	Guanosine	49	67	8	
	Cytosine	98	100	62	87
	Uracil	95	97	63	93

^aSee Ref. 19.

^bPopulation corresponds to neither C3' endo or C2' endo, see Figure 2.

Table 2

Comparison of Energies and Geometries for Conformations of MDP and MTP.

	MDP		MTP					
	Energy (kcal/mol)		Energy (kcal/mol)			Geometry		
	C1	C2	C1	C2	C3	C4	Dihedral RMSD (°) ^a	Angle RMSD (°) ^a
OPLS-AA	0	0.67	0	-0.4	1.4	1.55	27.5	10.5
OPLS-AA/M	0	4.6	0	0.5	4.9	5.1	8.5	9.5
QM ^b	0	6.8	0 ^c	0	6.9	7.6	0	0

^aRelative to ω B97X-D/6-311++G(d,p) results.^bB2PLYP-D3BJ/aug-cc-pVTZ// ω B97X-D/6-311++G(d,p).^cSeparate C1 and C2 minima were not located.