opinion, none of the previous conditions invalidate qSOFA as a prompt tool for risk assessment, mainly when it is embedded in a clinical decision-making framework at the emergency department, such as illustrated in our original Figure 1 (1). It is important to note that our study was neither designed for nor aimed to assess whether qSOFA performs differently in respiratory infections compared with other sources of infection.

When designing our study, we found it remarkable that qSOFA, derived by a data-driven approach using large databases, was constituted of the same three domains described in 2003 for both the Confusion, Respiratory Rate, and Blood Pressure (CRB) and Confusion, Urea, Respiratory Rate, Blood Pressure, and Age (CURB-65) scores (1–3). However, touching on what the authors have raised in their letter, CRB is more restrictive, using a respiratory rate $\geq 30/\text{min}$ compared with qSOFA ($\geq 22/\text{min}$), with direct implications on prevalence and sensitivity/specificity measures, as we discussed in our article (1). We observed that in our study population, there were 4,175 (61%) patients with respiratory rate $\geq 22/\text{min}$, and 1,766 (26%) using the CRB criteria ($\geq 30/\text{min}$). Therefore, two in five patients with CAP did not present a respiratory rate $\geq 22/\text{min}$ in our data from two tertiary hospitals. Moreover, it has been shown that the association between respiratory rate and outcomes in CAP is not linear, likely being U-shaped, as analyzed in detail in 705,928 patients in Germany (4).

We did not evaluate whether the respiratory rate was responsible for the overestimation observed for qSOFA when evaluating in-hospital mortality, although it might be a plausible explanation. It is important to note that even with certain miscalibration (5), qSOFA outperformed the systemic inflammatory response syndrome criteria in all scenarios for risk-stratification in our study, including the decision-curve analysis. As we discussed, CRB is an alternative prompt tool for the initial assessment of CAP at the emergency department (1).

We do not agree with the authors regarding the concern about selection bias. Our research question was to evaluate the performance of qSOFA and other scores in patients with CAP. Therefore, we enrolled patients who were representative of the population to which we are making inferences: patients with CAP, from two cohorts that included prospectively consecutive patients (1). Selection bias for our research question could have occurred if we had decided to include or exclude patients with CAP with a certain characteristic, such as some symptom (e.g., dyspnea) or previous condition (e.g., chronic respiratory disorder). In addition, we observed very few missing values and losses to follow-up, thereby decreasing the remaining potential for selection bias. As we acknowledged, regarding the limitations and conclusions of our study, the fact of evaluating only CAP might decrease the generalizability of our findings to other infection sources, but not the internal consistency, thereby guaranteeing the reliability of our findings. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Otávio T. Ranzani, M.D., M.Sc.*
University of Barcelona
Barcelona, Spain
and
Universidade de São Paulo
São Paulo, Brazil

Antoni Torres, M.D., Ph.D., F.E.R.S.*
University of Barcelona
Barcelona, Spain

On behalf of all authors

*These authors contributed equally to this work.

ORCID ID: 0000-0002-4677-6862 (O.T.R.).

References

1. Ranzani OT, Prina E, Menéndez R, Ceccato A, Cilloniz C, Méndez R, et al. New sepsis definition (Sepsis-3) and community-acquired pneumonia mortality: a validation and clinical decision-making study. *Am J Respir Crit Care Med* 2017;196:1287–1297.
2. Kolditz M, Scherag A, Rohde G, Ewig S, Welte T, Pletz M; CAPNETZ Study Group. Comparison of the qSOFA and CRB-65 for risk prediction in patients with community-acquired pneumonia. *Intensive Care Med* 2016;42:2108–2110.
3. Lim WS, van der Eerden MM, Laing R, Boersma WG, Karalus N, Town GI, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 2003;58:377–382.
4. Strauß R, Ewig S, Richter K, König T, Heller G, Bauer TT. The prognostic significance of respiratory rate in patients with pneumonia: a retrospective analysis of data from 705,928 hospitalized patients in Germany from 2010–2012. *Dtsch Arztebl Int* 2014;111:503–508.
5. Van Calster B, Vickers AJ. Calibration of risk prediction models: impact on decision-analytic performance. *Med Decis Making* 2015;35:162–169.

Copyright © 2018 by the American Thoracic Society

Low Lung Function in Young Adult Life Is Associated with Early Mortality

To the Editor:

The Tucson data showing early sustained and increased mortality from the second and third decade of life in those with the lowest tercile of FEV₁ are truly challenging (1). There is no biological plausibility for such a minor impairment of lung development of itself to translate into early death. The only possible explanation is that FEV₁ is the canary in the mine (2), warning of the early onset of the premature multimorbidity that is characteristic of later chronic obstructive lung disease (3). Supporting this concept is a recent study of birth weight, itself an important determinant of FEV₁ (4), which demonstrated that 15% of birth weight variance was genetically determined. Pathway analyses demonstrated that genes within birth weight-associated regions were enriched for processes likely important in long-term mortality risk, including type 2 diabetes, fasting glucose, triglycerides, total cholesterol, coronary artery

A.B. is a National Institute for Health Research (NIHR) Senior Investigator and, in addition, was supported by the NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton and Harefield National Health Service Foundation Trust and Imperial College London.

Originally Published in Press as DOI: 10.1164/rccm.201707-1416LE on September 19, 2017

disease, and systolic, mean, and diastolic blood pressure (5). The next step is to perform a similar genetic analysis for FEV₁. The implications of the Tucson findings are profound. First, it is not merely the airway disease of chronic obstructive lung disease that has its origins in early life (6) but also the comorbidities, and it is in early life that the remedies must be found. Second, there is a strong case for performing spirometry at early school age, as well as the routine of measuring weight and height, to identify and target a high-risk population for not just airway but also systemic disease. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Andrew Bush, M.B. B.S., M.A., M.D.

Imperial College London

London, United Kingdom

National Heart and Lung Institute

London, United Kingdom

and

Royal Brompton Harefield NHS Foundation Trust

London, United Kingdom

ORCID ID: 0000-0001-6756-9822 (A.B.).

References

1. Vazquez MM, Zhou M, Hu C, Martinez FD, Guerra S. Low lung function in young adult life is associated with early mortality. *Am J Respir Crit Care Med* 2017;195:1399–1401.
2. Povey H, Cameron D. The bird it was that died. *Thorax* 2013;68:1175, 1179.
3. Vanfleteren LE, Spruit MA, Groenen M, Gaffron S, van Empel VP, Bruijnzeel PL, et al. Clusters of comorbidities based on validated objective measurements and systemic inflammation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2013;187:728–735.
4. Cai Y, Shaheen SO, Hardy R, Kuh D, Hansell AL. Birth weight, early childhood growth and lung function in middle to early old age: 1946 British birth cohort. *Thorax* 2016;71:916–922.
5. Horikoshi M, Beaumont RN, Day FR, Warrington NM, Kooijman MN, Fernandez-Tajes J, et al.; CHARGE Consortium Hematology Working Group. Genome-wide associations for birth weight and correlations with adult disease. *Nature* 2016;538:248–252.
6. Bush A. Lung development and aging. *Ann Am Thorac Soc* 2016;13:S438–S446.

Copyright © 2018 by the American Thoracic Society

Reply to Bush

From the Authors:

We appreciate the cogent comments by Dr. Bush and agree that a possible explanation for the association we reported between early FEV₁ impairment and subsequent mortality risk may be related to a common comorbidity predisposition (1). We propose that a scenario in which FEV₁ deficits *per se* may have a direct effect on mortality risk is also possible. In our study, participants in the low lung function

This study was supported by awards HL107188 and HL095021 from the NHLBI, NIH.

Originally Published in Press as DOI: 10.1164/rccm.201708-1647LE on September 19, 2017

group had FEV₁ levels that were on average 18% lower than those of participants in the highest tertile. Although subclinical, the magnitude of these FEV₁ deficits may have been sufficient to make participants with low lung function more susceptible (or susceptible at a younger age) to cardiopulmonary risk factors, leading to an increased risk for early mortality. Indeed, we argue that these two scenarios are to some extent complementary because it is likely that adverse social and environmental factors in early life, on a background of genetic susceptibility, may alter normal developmental trajectories and predispose to a variety of adverse outcomes in adult life, which may ultimately affect and worsen each other. These outcomes are almost invariably associated with chronic systemic inflammation and accelerated organ function decline during senescence. Therefore, we definitely agree with Bush on the importance of targeting early life and childhood as critical developmental windows for an effective prevention not only of chronic obstructive pulmonary disease but also its comorbidities. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Monica M. Vasquez, Ph.D., M.P.H.

Muhan Zhou, M.P.H.

Chengcheng Hu, Ph.D.

Fernando D. Martinez, M.D.

University of Arizona

Tucson, Arizona

Stefano Guerra, M.D., Ph.D.

University of Arizona

Tucson, Arizona

and

ISGlobal

Barcelona, Spain

Reference

1. Vazquez MM, Zhou M, Hu C, Martinez FD, Guerra S. Low lung function in young adult life is associated with early mortality. *Am J Respir Crit Care Med* 2017;195:1399–1401.

Copyright © 2018 by the American Thoracic Society

Recurrent Respiratory Papillomatosis and Bevacizumab Treatment

To the Editor:

We read with great interest the article by Dr. Kalanjeri and colleagues (1) describing a case of diffuse tracheal papillomatosis. In the case they presented, therapeutic bronchoscopy resulted in significant symptomatic improvement and airway patency, but

Author Contributions: Conception and design, data analysis, and redaction and final revision of the manuscript: S.F.-B., G.L., and M.B. Data analysis and interpretation, and redaction and final revision of the manuscript: M.R.V. Data analysis, editing of the manuscript for important intellectual content, and redaction and final revision of the manuscript: A.M., R.S., H.J.M., and M.J.

Originally Published in Press as DOI: 10.1164/rccm.201702-0279LE on October 2, 2017