Published in final edited form as:

Hematol Oncol Clin North Am. 2018 April ; 32(2): 343–352. doi:10.1016/j.hoc.2017.11.002.

Emerging Therapies

Amaliris Gonzalez^a, Khaled M. Musallam^b, Ali T. Taher^c, and Stefano Rivella^{a,d}

^aDepartment of Pediatrics, Division of Hematology, Children's Hospital of Philadelphia (CHOP), Philadelphia, PA

^bInternational Network of Hematology, London, UK

^cDepartment of Internal Medicine, American University of Beirut Medical Center, Beirut, Lebanon

^dCell and Molecular Biology Graduate Group (CAMB), University of Pennsylvania, Philadelphia, PA

Abstract

At present, the only definitive cure for β -thalassemia is a bone marrow transplant (BMT), however HLA-blood matched donors are scarcely available. Current therapies undergoing clinical investigation with most potential for therapeutic benefit are the β -globin gene transfer (GT) of patient specific hematopoietic stem cells (HSCs) followed by autologous BMT. Other emerging therapies deliver exogenous regulators of several key modulators of erythropoiesis or iron homeostasis. This review focuses on current approaches for the treatment of hemoglobinopathies caused by disruptions of β -globin.

Keywords

β -thalassemia; β -Globin; Gene transfer; Hemichromes; New Therapies; Trap Ligands

Introduction

Hemoglobinopathies arise from genetic mutations in HBA1, HBA2 and HBB that compromise the structure-function of the α - and β -globin chains of hemoglobin (Hb). Collectively, mutations in β -globin gene are referred to as β -thalassemia (BT) ¹. Mutations in the HBB are remarkably heterogeneous at the molecular level with over 300 variations identified (<http://globin.cse.psu.edu/>). Despite the diversity of mutations most cause disruption in the α -, β -globin balance resulting in long-term ineffective erythropoiesis (IE) and extramedullary hematopoiesis (EMH) of the spleen and liver. Ineffective erythropoiesis

Corresponding Author: Stefano Rivella, 3615 Civic Center Blvd, Philadelphia, PA 19104, rivellas@email.chop.edu.

Conflict of Interest statement

K.M. received honoraria from Novartis, Celgene, and CRISPR Therapeutics. A.T. received honoraria and research support from Novartis and Celgene. S.R. is a member of scientific advisory board of Ionis Pharmaceuticals. S.R. has grants from Ionis and NIH. A.G. declares no conflict of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

leads to various degrees of chronic hemolytic anemia and transfusion dependence, as well as upregulation of iron absorption. Strategies to treat BT are based on gene therapy strategies as well as pharmacological treatments to target pathways that modulate erythropoiesis and iron homeostasis. Here, we review emerging therapies in clinical trials and pre-clinical stages of development.

Emerging Therapies in Clinical Trials

Sotatercept (ACE-011) and Luspatercept (ACE-536) are two compounds that showed success in clinical trials (Table 1). Sotatercept was originally developed to treat bone-loss disorders, but clinical studies unexpectedly revealed increased hematocrit and Hb levels in treated patients². Structurally designed to compete with the extracellular domains (ECDs) of Activin Receptor A Type 2A (ACVR2A) or 2B, these peptides act as ligand traps for TGFβ-like molecules^{3,4}. Growth differentiation factor 11 (GDF11), a member of TGFβ superfamily, is the identified ligand target for luspatercept. Treatment with luspatercept is thought to remove GDF11 from circulation and cause subsequent increases of Hb levels⁵. GDF11 is well established as a ligand capable of activating the SMAD2/3 pathway through ACVR2A or 2B⁶. Administration of the luspatercept mouse analog, RAP-536, to BT mice ameliorates intracellular accumulation of hemichromes (HCM), oxidative stress and splenomegaly, while also inducing late-stage erythroid progenitor differentiation⁷. In phase 1 studies of healthy volunteers, luspatercept and sotatercept showed a dose-dependent and sustainable increase in hemoglobin level and were well tolerated.^{8,9} Luspatercept subsequently entered a phase 2, open-label, dose-ranging study in adults with BT (NCT01749540, completed) including an ongoing 2-year extension (NCT02268409). Patients received luspatercept at doses of 0.2–1.25 mg/kg administered subcutaneously every 3 weeks. Available data indicate that luspatercept was generally well tolerated and had a favorable safety profile. Luspatercept reduced transfusion requirements and liver iron concentration among patients with transfusion-dependent (TD) BT, and increased hemoglobin levels, reduced liver iron concentration, and improved patient reported outcomes among those with non-transfusion-dependent (NTD) BT^{10,11}. A double-blind, randomized, placebo-controlled phase 3 study (BELIEVE) has begun to evaluate the efficacy and safety of luspatercept among adults with TD BT. Demonstration of efficacy will require at least a 33% improvement in the number of transfused red blood cell units from baseline (NCT02604433). In the phase 2a study of sotatercept among adults with TD or NTD BT (NCT01571635, completed), data indicated increase in hemoglobin levels, a reduced transfusion burden, and a favorable safety profile¹², but no phase 3 studies are publicly announced.

The JAK2 pathway has been long thought of as a potential link between erythropoiesis and iron metabolism¹³. Negating effects of EPO overstimulation in BT by inhibiting the JAK2/STAT3 pathway has been an attractive area of therapy exploration. One of the first agents to be approved by the FDA as a JAK2 inhibitor was ruxolitinib¹⁴. Studies on mice models of BT, show that the JAK2 inhibitor reduces ineffective erythropoiesis and splenomegaly^{13,15}. A single-arm, phase 2 study (NCT02049450, completed) to evaluate the efficacy and safety of ruxolitinib administered orally at a starting dose of 10 mg twice daily among adults (n=30) with TD BT and splenomegaly has been completed¹⁶. Ruxolitinib was associated

with a slight increase in pre-transfusional hemoglobin levels (by 0.5 g/L increase) and a trend towards reduced transfusion requirements (by 45 mL of hematocrit-adjusted red blood cell volume per 4 weeks) following 30 weeks of treatment. Mean spleen volume also decreased during ruxolitinib treatment. No major adverse effects were reported.

The only cure available for patients suffering from BT is a BMT with cells harboring a functional HBB gene, however finding a donor source is challenging. GT technology and autologous BMT provides an alternative with potential to reach more patients. The development of safe lentiviral vectors and discovery of the β -globin's LCR, has made GT a possibility for BT. The first clinical study (LG001) in a BT patient tested the HPV569 vector^{17–20} harboring a critical amino acid modification (T87Q) that allowed researchers to successfully identify modified cells and accurately measure viral integration and chimerism²¹. The patient became transfusion independent one year after GT, exhibiting increased levels of several Hb variants including HbF with equal proportions of HbA containing the T87Q mutation. New vectors are currently in various stages of clinical trials (NCT02633943, NCT02453477, NCT03275051, NCT02453477, NCT01745120, NCT02151526, NCT03207009, NCT02906202, NCT01639690) (Table 1). Interim data from a phase 1/2 study of autologous hematopoietic stem cells transduced ex vivo with a lentiviral vector among patients with TD BT (NCT01745120) suggest increased levels of HbA and reduced transfusion requirements²². New strategies for allogeneic BMT and reduced intensity conditioning (RIC) are also in clinical trials (NCT00408447, NCT02038478, NCT02165007), which could eventually also aid the success of GT (Table 1).

Innovative Therapies in Pre-Clinical Stages

Patients with high levels of endogenous HbF do not exhibit the symptomology associated with BT. Chemical agents such as hydroxyurea are used to induce HbF in BT patients²³, however these agents are not effective in all patients^{24,25}. Genome-wide associated studies²⁶ identified *BCL11a* as a regulator of HbF. Knock-down studies of *BCL11a* using RNAi^{24,27,28} proved it to be rheostat for silencing of the γ -globin gene. One novel approach to induce HbF by silencing *BCL11a* uses an erythroid specific promoter to drive a microRNA-adapted shRNAs (shRNAmiR)^{29,30}. By using an erythroid specific HS2 and HS3 regions of the β -globin LCR alongside its promoter, the toxicity and engraftment impairments characteristic of Pol III/II driven shRNA and microRNAs are circumvented while simultaneously inducing HbF in erythroid cells³⁰.

Other strategies for *BCL11a* inhibition in preclinical stages employ gene editing (GE) strategies: zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs) and clustered regularly interspaced palindromic repeats (CRISP) in association with Cas 9 (CRISP/Cas9). All GE strategies have successfully targeted *BCL11a*^{31–34} decreasing protein expression of *BCL11a* and causing expression of γ -globin. Several studies using ZFNs, TALENs, and CRISP/Cas9 systems have also shown the potential of editing HBB in both patient derived CD34+ cells and induces pluripotent stem cells (iPSC)^{35–38}. Although questions concerning off-target events remain, latest studies show GE as a strategy with clinical potential^{34,39}.

Targeting iron regulation by direct action on the main iron transporter, transferrin has had promising results. Studies on BT mice treated with exogenous apo-transferrin resulted in a net decrease of iron uptake by RBC precursors through TfR1 and subsequent reduction of HMCs⁴⁰. As HMCs were reduced, the lifespan of RBCs and total amount of Hb in circulation increased. Additionally, haplo-insufficient TfR1 mice also resulted in improved erythropoiesis with results closely mirroring findings of exogenous apo-transferrin treatment⁴¹. Increasing circulating hepcidin and manipulating the hepcidin pathway by targeting its regulators may provide patients with more effective options for iron overload management. Preclinical studies of administration of exogenous sources of hepcidin, such as minihepcidins and other hepcidin agonists, and well as inhibiting negative regulators of *Hamp* (TMPRSS6) expression show to successfully prevent iron overload, improve anemia and ameliorate ineffective erythropoiesis in mice models of BT^{42–46}

Novel tools for drug development and therapy discover

Novel therapeutics are not possible without innovation at the bench side. New animal models, *in vitro* systems with potential to more closely mimic *in vivo* conditions and tools for gathering *in vivo* data in mice models are necessary for the advancement of BT therapies. A Thal-Biobank has been established with goals to supply researchers with the samples of the same cells for studies aimed at solving the plethora of biological and biomedical issues involved in using patient cell cultures⁴⁷. Erythroid pre-cursors collected from 72 BT patients, were expanded, cryopreserved, characterized with respects their individual Hb profile and tested for abilities to produce HbF by hydroxyurea. Furthermore, several laboratories characterized samples from the same donors, showing that data were reproducible using protocols developed by the Thal-bank.

In the last decade immortalizing human progenitor cells has become possible. Two cell lines in particular stand out, Human Immortalized Erythroid Progenitor Cell Lines (HUDEPs)⁴⁸ and the newly established Mui009⁴⁹. HUDEPs were generated from human umbilical cord blood and immortalized with conditional (doxycycline-inducible) expression of Human Papilloma Virus (HPV)-derived E6/E7 proteins. The cells can be expanded infinitely and characterization shows them to be arrested at the pro-erythroblast stage. The line has several clones available, each with its own unique gene expression fingerprint of erythroid specific markers. These cells can be differentiated into enucleated RBCs, providing a useful tool to study erythropoiesis. The Mui0009 cell lines were generated from 32-year-old male with homozygous β -globin deletion and heterozygous α -thalassemia. Peripheral blood mononuclear cells were enriched for CD34+ cells and reprogrammed into iPSC cells. Resultant cells exhibited embryonic stem cell characteristics, including the capability of differentiating into the endodermal, ectodermal and mesodermal germinal layers. However, work is still needed to determine if the cells are suitable for studies requiring RBC differentiation.

Measuring potential therapeutics for inhibition of iron overload has been limited by the inability to measure iron loading on live animal models. The use of magnetic resonance imaging (MRI) offers a clinically relevant assessment tool to measure the presence of iron loading in humans, and recently has been optimized for the use on small animal models⁵⁰.

Tissue resonance relaxation rates, spleen volumetric and cardiac function are now quantifiable in live BT mouse models during treatment course.

Discussion

Despite the success of GT in the last decade, improvements of both the transduction efficiency of lentiviruses and HSC expression of the β -globin transgene are needed. Patients treated with GT display a spectrum of phenotypic heterogeneity in endogenous and GT-delivered β -globin variants. Although some patients treated reach therapeutic HbA thresholds, a majority are left TD^{17,21,51}. The potential of GE and reactivation of HbF production awaits future data from human studies.

Therapeutic strategies aiming to modulate iron metabolism are also showing promise in reducing IE, ROS and ameliorating anemia. Exogenous administration of apo-transferrin ascertains to be a therapeutic with much potential. The safety profile and Hb increasing abilities in clinical trials has been previously reported in treating hypotransferrinemia⁵². RNAi technology has been used in innovative ways to target modulators of both erythropoiesis. Activating HbF by using RNAi is pushing the bounds of discovery by creating new molecules capable of intervening at the level of transcriptional regulation.

Sotatercept and luspatercept are front runners as emerging therapies for BT management. Although they were designed to target different TGF β ligands, results of clinical studies show similar outcomes. The unknown role of the TGF β ligands family on erythropoiesis and their potential contributions of BT treatment make the initial serendipitous sotatercept and luspatercept findings especially exciting. As very little is known about the role the activin pathways play on normal hematopoietic function and current mechanisms suggested do not completely explain the effects of sotatercept and luspatercept^{5,7,8,53}. Further characterization of the TGF β ligands will undoubtedly lead to new insights into both fundamental hematological mechanisms and therapies.

Acknowledgments

Work related to this article was funded by grants from the NIH-NIDDK- R01 DK095112 and R01 DK090554 (SR). We extend special thanks to Ping La (Children's Hospital of Philadelphia) for helpful discussion and support.

References

1. Weatherall DJ, Clegg JB. Inherited haemoglobin disorders: an increasing global health problem. *Bull World Health Organ.* 2001; 79(8):704–712. [PubMed: 11545326]
2. Carrancio S, Markovics J, Wong P, et al. An activin receptor IIA ligand trap promotes erythropoiesis resulting in a rapid induction of red blood cells and haemoglobin. *Br J Haematol.* 2014; 165(6):870–882. [PubMed: 24635723]
3. Pearsall RS, Canalis E, Cornwall-Brady M, et al. A soluble activin type IIA receptor induces bone formation and improves skeletal integrity. *Proc Natl Acad Sci U S A.* 2008; 105(19):7082–7087. [PubMed: 18460605]
4. Sako D, Grinberg AV, Liu J, et al. Characterization of the ligand binding functionality of the extracellular domain of activin receptor type IIb. *J Biol Chem.* 2010; 285(27):21037–21048. [PubMed: 20385559]

5. Suragani RN, Cadena SM, Cawley SM, et al. Transforming growth factor-beta superfamily ligand trap ACE-536 corrects anemia by promoting late-stage erythropoiesis. *Nat Med.* 2014; 20(4):408–414. [PubMed: 24658078]
6. Oh SP, Yeo CY, Lee Y, Schrewe H, Whitman M, Li E. Activin type IIA and IIB receptors mediate Gdf11 signaling in axial vertebral patterning. *Genes Dev.* 2002; 16(21):2749–2754. [PubMed: 12414726]
7. Suragani RN, Cawley SM, Li R, et al. Modified activin receptor IIB ligand trap mitigates ineffective erythropoiesis and disease complications in murine beta-thalassemia. *Blood.* 2014; 123(25):3864–3872. [PubMed: 24795345]
8. Attie KM, Allison MJ, McClure T, et al. A phase 1 study of ACE-536, a regulator of erythroid differentiation, in healthy volunteers. *Am J Hematol.* 2014; 89(7):766–770. [PubMed: 24715706]
9. Sherman ML, Borgstein NG, Mook L, et al. Multiple-dose, safety, pharmacokinetic, and pharmacodynamic study of sotatercept (ActRIIA-IgG1), a novel erythropoietic agent, in healthy postmenopausal women. *J Clin Pharmacol.* 2013; 53(11):1121–1130. [PubMed: 23939631]
10. Antonio Piga SP, Melpignano Angela, Borgna-Pignatti Caterina, Rita Gamberini M, Voskaridou Ersi, Caruso Vincenzo, Ricchi Paolo, Pietrangelo Antonello, Zhang Xiaosha, Wilson Dawn, Bellevue Ashley, Laadem Abderrahmane, Sherman Matthew, Attie Kenneth. Luspatercept Decreases Transfusion Burden and Liver Iron Concentration in Regularly Transfused Adults with Beta-Thalassemia (S836). *Haematologica* (101). 2016; (s1):338–339.
11. Antonio G, Piga IT, Gamberini Rita, Voskaridou Ersi, Melpignano Angela, Ricchi Paolo, Caruso Vincenzo, Pietrangelo Antonello, Zhang Xiaosha, Wilson Dawn M, Leneus Ashley, Laadem Abderrahmane, Sherman Matthew L, Attie Kenneth M. Luspatercept Increases Hemoglobin, Decreases Transfusion Burden and Improves Iron Overload in Adults with Beta-Thalassemia. *Blood.* 2016; 128(851)
12. Cappellini MDPI, Origa R, Forni GL, Voskaridou E, Taher AT, et al. Interim results from a Phase 2a, open-label, dose-finding study of sotatercept (ACE-011) in adult patients with beta-thalassemia. *Haematologica* (100). 2015; (s1):17–18. [PubMed: 25552679]
13. Rivella S. Ineffective erythropoiesis and thalassemias. *Curr Opin Hematol.* 2009; 16(3):187–194. [PubMed: 19318943]
14. Ostojic A, Vrhovac R, Verstovsek S. Ruxolitinib: a new JAK1/2 inhibitor that offers promising options for treatment of myelofibrosis. *Future Oncol.* 2011; 7(9):1035–1043. [PubMed: 21919691]
15. Casu C, Rivella S. Iron age: novel targets for iron overload. *Hematology Am Soc Hematol Educ Program.* 2014; 2014(1):216–221. [PubMed: 25696858]
16. Aydinok YKZ, Cassinerio E, Siritanaratkul N, Kattamis A, Maggio A, et al. Efficacy and Safety of Ruxolitinib in Regularly Transfused Patients with Thalassemia: Results from Single-Arm, Multicenter, Phase 2a Truth Study. *Blood.* 2016; (128):852. [PubMed: 27281793]
17. Negre O, Eggimann AV, Beuzard Y, et al. Gene Therapy of the beta-Hemoglobinopathies by Lentiviral Transfer of the beta(A(T87Q))-Globin Gene. *Hum Gene Ther.* 2016; 27(2):148–165. [PubMed: 26886832]
18. Negre O, Bartholomae C, Beuzard Y, et al. Preclinical evaluation of efficacy and safety of an improved lentiviral vector for the treatment of beta-thalassemia and sickle cell disease. *Curr Gene Ther.* 2015; 15(1):64–81. [PubMed: 25429463]
19. Cavazzana-Calvo M, Payen E, Negre O, et al. Transfusion independence and HMGA2 activation after gene therapy of human beta-thalassaemia. *Nature.* 2010; 467(7313):318–322. [PubMed: 20844535]
20. Ribeil JA, Hacein-Bey-Abina S, Payen E, et al. Gene Therapy in a Patient with Sickle Cell Disease. *N Engl J Med.* 2017; 376(9):848–855. [PubMed: 28249145]
21. Bank A, Dorazio R, Leboulch P. A phase I/II clinical trial of beta-globin gene therapy for beta-thalassemia. *Ann N Y Acad Sci.* 2005; 1054:308–316. [PubMed: 16339679]
22. Thompson AA, Kwiatkowski J, Rasko J, Hongeng S, Schiller GJ, Anurathapan U, Cavazzana M, Ho PJ, von Kalle C, Kletzel M, Leboulch P, Vichinsky E, Petrusich A, Asmal M, Walters MC. Lentiglobin Gene Therapy for Transfusion-Dependent β -Thalassemia: Update from the Northstar Hgb-204 Phase 1/2 Clinical Study. *Blood.* 2016; 128(22):1175.

23. Arruda VR, Lima CS, Saad ST, Costa FF. Successful use of hydroxyurea in beta-thalassemia major. *N Engl J Med*. 1997; 336(13):964.
24. Basak A, Hancarova M, Ulirsch JC, et al. BCL11A deletions result in fetal hemoglobin persistence and neurodevelopmental alterations. *J Clin Invest*. 2015; 125(6):2363–2368. [PubMed: 25938782]
25. Musallam KM, Taher AT, Cappellini MD, Sankaran VG. Clinical experience with fetal hemoglobin induction therapy in patients with beta-thalassemia. *Blood*. 2013; 121(12):2199–2212. quiz 2372. [PubMed: 23315167]
26. Uda M, Galanello R, Sanna S, et al. Genome-wide association study shows BCL11A associated with persistent fetal hemoglobin and amelioration of the phenotype of beta-thalassemia. *Proc Natl Acad Sci U S A*. 2008; 105(5):1620–1625. [PubMed: 18245381]
27. Giani FC, Fiorini C, Wakabayashi A, et al. Targeted Application of Human Genetic Variation Can Improve Red Blood Cell Production from Stem Cells. *Cell Stem Cell*. 2016; 18(1):73–78. [PubMed: 26607381]
28. Sankaran VG, Menne TF, Xu J, et al. Human fetal hemoglobin expression is regulated by the developmental stage-specific repressor BCL11A. *Science*. 2008; 322(5909):1839–1842. [PubMed: 19056937]
29. Guda S, Brendel C, Renella R, et al. miRNA-embedded shRNAs for Lineage-specific BCL11A Knockdown and Hemoglobin F Induction. *Mol Ther*. 2015; 23(9):1465–1474. [PubMed: 26080908]
30. Brendel C, Guda S, Renella R, et al. Lineage-specific BCL11A knockdown circumvents toxicities and reverses sickle phenotype. *J Clin Invest*. 2016; 126(10):3868–3878. [PubMed: 27599293]
31. Bjurstrom CF, Mojadidi M, Phillips J, et al. Reactivating Fetal Hemoglobin Expression in Human Adult Erythroblasts Through BCL11A Knockdown Using Targeted Endonucleases. *Mol Ther Nucleic Acids*. 2016; 5:e351. [PubMed: 28131278]
32. Cavazzana M, Antoniani C, Miccio A. Gene Therapy for beta-Hemoglobinopathies. *Mol Ther*. 2017; 25(5):1142–1154. [PubMed: 28377044]
33. Makis A, Hatzimichael E, Papassotiriou I, Voskaridou E. 2017 Clinical trials update in new treatments of beta-thalassemia. *Am J Hematol*. 2016; 91(11):1135–1145. [PubMed: 27502996]
34. Genovese P, Schirotti G, Escobar G, et al. Targeted genome editing in human repopulating haematopoietic stem cells. *Nature*. 2014; 510(7504):235–240. [PubMed: 24870228]
35. Xu P, Tong Y, Liu XZ, et al. Both TALENs and CRISPR/Cas9 directly target the HBB IVS2-654 (C > T) mutation in beta-thalassemia-derived iPSCs. *Sci Rep*. 2015; 5:12065. [PubMed: 26156589]
36. Finotti A, Borgatti M, Gambari R. Ground state naive pluripotent stem cells and CRISPR/Cas9 gene correction for beta-thalassemia. *Stem Cell Investig*. 2016; 3:66.
37. Chattong S, Ruangwattanasuk O, Yindeedee W, Setpakdee A, Manotham K. CD34+ cells from dental pulp stem cells with a ZFN-mediated and homology-driven repair-mediated locus-specific knock-in of an artificial beta-globin gene. *Gene Ther*. 2017; 24(7):425–432. [PubMed: 28530652]
38. DeWitt MA, Magis W, Bray NL, et al. Selection-free genome editing of the sickle mutation in human adult hematopoietic stem/progenitor cells. *Sci Transl Med*. 2016; 8(360):360ra134.
39. Dever DP, Bak RO, Reinisch A, et al. CRISPR/Cas9 beta-globin gene targeting in human hematopoietic stem cells. *Nature*. 2016; 539(7629):384–389. [PubMed: 27820943]
40. Li H, Rybicki AC, Suzuka SM, et al. Transferrin therapy ameliorates disease in beta-thalassemic mice. *Nat Med*. 2010; 16(2):177–182. [PubMed: 20098432]
41. Li H, Choesang T, Bao W, et al. Decreasing TfR1 expression reverses anemia and hepcidin suppression in beta-thalassemic mice. *Blood*. 2017; 129(11):1514–1526. [PubMed: 28151426]
42. Casu C, Oikonomidou PR, Chen H, et al. Minihepcidin peptides as disease modifiers in mice affected by beta-thalassemia and polycythemia vera. *Blood*. 2016; 128(2):265–276. [PubMed: 27154187]
43. Liu J, Sun B, Yin H, Liu S. Hepcidin: A Promising Therapeutic Target for Iron Disorders: A Systematic Review. *Medicine (Baltimore)*. 2016; 95(14):e3150. [PubMed: 27057839]
44. Nai A, Rubio A, Campanella A, et al. Limiting hepatic Bmp-Smad signaling by matriptase-2 is required for erythropoietin-mediated hepcidin suppression in mice. *Blood*. 2016; 127(19):2327–2336. [PubMed: 26755707]

45. Schmidt PJ, Toudjarska I, Sendamarai AK, et al. An RNAi therapeutic targeting Tmprss6 decreases iron overload in Hfe(−/−) mice and ameliorates anemia and iron overload in murine beta-thalassemia intermedia. *Blood*. 2013; 121(7):1200–1208. [PubMed: 23223430]
46. Casu C, Aghajan M, Oikonomidou PR, Guo S, Monia BP, Rivella S. Combination of Tmprss6-ASO and the iron chelator deferiprone improves erythropoiesis and reduces iron overload in a mouse model of beta-thalassemia intermedia. *Haematologica*. 2016; 101(1):e8–e11. [PubMed: 26405152]
47. Cosenza LC, Breda L, Breveglieri G, et al. A validated cellular biobank for beta-thalassemia. *J Transl Med*. 2016; 14:255. [PubMed: 27590532]
48. Kurita R, Suda N, Sudo K, et al. Establishment of immortalized human erythroid progenitor cell lines able to produce enucleated red blood cells. *PLoS One*. 2013; 8(3):e59890. [PubMed: 23533656]
49. Wongkumool W, Maneepitasut W, Tong-Ngam P, et al. Establishment of MUi009 - A human induced pluripotent stem cells from a 32year old male with homozygous beta degrees -thalassemia coinherited with heterozygous alpha-thalassemia 2. *Stem Cell Res*. 2017; 20:80–83. [PubMed: 28395745]
50. Jackson LH, Vlachodimitropoulou E, Shangaris P, et al. Non-invasive MRI biomarkers for the early assessment of iron overload in a humanized mouse model of beta-thalassemia. *Sci Rep*. 2017; 7:43439. [PubMed: 28240317]
51. Mansilla-Soto J, Riviere I, Boulad F, Sadelain M. Cell and Gene Therapy for the Beta-Thalassemias: Advances and Prospects. *Hum Gene Ther*. 2016; 27(4):295–304. [PubMed: 27021486]
52. Boshuizen M, van der Ploeg K, von Bonsdorff L, et al. Therapeutic use of transferrin to modulate anemia and conditions of iron toxicity. *Blood Rev*. 2017
53. Zhang YH, Cheng F, Du XT, et al. GDF11/BMP11 activates both smad1/5/8 and smad2/3 signals but shows no significant effect on proliferation and migration of human umbilical vein endothelial cells. *Oncotarget*. 2016; 7(11):12063–12074. [PubMed: 26919250]

Key Points

- At present, the only definitive cure for β -thalassemia is a bone marrow transplant (BMT), however HLA-blood matched donors are scarcely available.
- Current therapies undergoing clinical investigation with most potential for therapeutic benefit are the β -globin gene transfer (GT) of patient specific hematopoietic stem cells (HSCs) followed by autologous BMT.
- Other emerging therapies deliver exogenous regulators of several key modulators of erythropoiesis or iron homeostasis.
- This review focuses on current approaches for the treatment of hemoglobinopathies caused by disruptions of β -globin.

Table 1 Emerging therapies in clinical trials for treatment of β -Thalassemia.

Strategy	Phase	NCT Number	Study Title	Interventions	Status
Trap-ligand targeting TGF β -ligands	Phase 2	NCT01571635	Study to Determine the Safety and Tolerability of Sotatercept (ACE-011) in Adults With Beta(β)-Thalassemia.	Sotatercept	Active, not recruiting
	Phase 2	NCT02268409	ACE-536 Extension Study - Beta Thalassemia	Luspatercept	Active, not recruiting
	Phase 3	NCT02604433	An Efficacy and Safety Study of Luspatercept (ACE-536) Versus Placebo in Adults Who Require Regular Red Blood Cell Transfusions Due to Beta (β) Thalassemia	Luspatercept	Active, not recruiting
HbF induction	Early Phase 1	NCT02981329	Fetal Hemoglobin Induction Treatment Metformin	Metformin	Recruiting
RIC	Phase 2	NCT00408447	Stem Cell Transplant in Sickle Cell Disease and Thalassemia	Busulfan, Fludarabine, Alemtuzumab	Recruiting
	Phase 2	NCT02038478	Allograft for Sickle Cell Disease and Thalassemia	Donor Stem Cell Transplantation	Recruiting
	Phase 1	NCT02165007	Haploidentical Hematopoietic Stem Cell Transplantation for Children With Sickle Cell Disease and Thalassemia Using CD34+ Positive Selected Grafts	Peripheral blood stem cell graft that are CD34+ selected	Recruiting
					Recruiting
JAK2 inhibition	Phase 2	NCT02049450	Study of Efficacy and Safety of INC424 in Regularly Transfused Patients With Thalassemia.	Ruxolitinib	Completed
GT	Early Phase 1	NCT02633943	Longterm Follow-up of Subjects With Hemoglobinopathies Treated With Ex Vivo Gene Therapy	Safety and efficacy assessments	Enrolling by invitation
	Phase 2	NCT03275051	Long-term Safety and Efficacy Follow-up of Subject Treated With GSK2696277 for Transfusion Dependent Beta-thalassemia in San Raffaele Telethon Institute of Gene Therapy-Beta Thalassemia (TIGET-BTHAL) Study	Safety and efficacy assessments	Not yet recruiting
	Phase 1/Phase 2	NCT02453477	Gene Therapy for Transfusion Dependent Beta-thalassemia	GLOBE	Recruiting
	Phase 1	NCT01639690	β -Thalassemia Major With Autologous CD34+ Hematopoietic Progenitor Cells Transduced With TNS9.3.55 a Lentiviral Vector Encoding the Normal Human β -Globin Gene	TNS9.3.55	Active, not recruiting

Strategy	Phase	NCT Number	Study Title	Interventions	Status
	Phase 1/Phase 2	NCT02453477	Gene Therapy for Transfusion Dependent Beta-thalassemia	LentiGlobin BB305	Recruiting
	Phase 1/Phase 2	NCT01745120	A Study Evaluating the Safety and Efficacy of the LentiGlobin® BB305 Drug Product in Beta-Thalassemia Major Subjects		Active, not recruiting
	Phase 1/Phase 2	NCT02151526	A Study Evaluating the Efficacy and Safety of the LentiGlobin BB305 Drug Product in Beta-Thalassemia Major and Sickle Cell Disease		Active, not recruiting
	Phase 3	NCT03207009	A Study Evaluating the Efficacy and Safety of the LentiGlobin® BB305 Drug Product in Subjects With Transfusion-Dependent β -Thalassemia, Who Have a $\beta\text{0}/\beta\text{0}$ Genotype		Recruiting
	Phase 3	NCT02906202	A Study Evaluating the Efficacy and Safety of the LentiGlobin® BB305 Drug Product in Subjects With Transfusion-Dependent β -Thalassemia		Recruiting