Published in final edited form as:

Qual Life Res. 2018 February ; 27(2): 291–319. doi:10.1007/s11136-017-1692-4.

Child and adolescent self-report symptom measurement in pediatric oncology research: a systematic literature review

Laura C. Pinheiro¹, Molly McFatrigh¹¹, Nicole Lucas², Jennifer S. Walker³, Janice S. Withycombe⁴, Pamela S. Hinds⁵, Lillian Sung^{6,7}, Deborah Tomlinson⁶, David R. Freyer⁸, Jennifer W. Mack⁹, Justin N. Baker¹⁰, and Bryce B. Reeve¹¹

¹Division of General Internal Medicine, Department of Medicine, Weill Cornell Medicine, New York, NY 10064, USA

²Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

³Health Sciences Library, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

⁴Nell Hodgson Woodruff School of Nursing, Emory University, Atlanta, GA, USA

⁵Department of Nursing Research and Quality Outcomes, Children's National Health System, Department of Pediatrics, George Washington University, Washington, DC, USA

⁶Child Health Evaluative Sciences, The Hospital for Sick Children, Toronto, Canada

⁷Division of Haematology/Oncology, The Hospital for Sick Children, Toronto, Canada

⁸Children's Center for Cancer and Blood Diseases, Children's Hospital Los Angeles, Los Angeles, CA, USA

⁹Dana-Farber Cancer Institute, Boston, MA, USA

¹⁰Division of Quality of Life and Palliative Care, Department of Oncology, St. Jude Children's Research Hospital, Memphis, TN, USA

¹¹Center for Health Measurement, Department of Population Health Sciences, Duke University School of Medicine, Durham, NC, USA

Abstract

Objective—Previous work in pediatric oncology has found that clinicians and parents tend to under-report the frequency and severity of treatment-related symptoms compared to child self-

Correspondence to: Laura C. Pinheiro.

Compliance with ethical standards

Conflict of interest All authors declare that they have no conflicts of interest to disclose.

Informed consent Informed consent from study subjects was not needed as the University of North Carolina at Chapel Hill IRB granted this research exemption from review.

Electronic supplementary material The online version of this article (doi:10.1007/s11136-017-1692-4) contains supplementary material, which is available to authorized users.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. This article does not contain any studies with animals performed by any of the authors.

report. As such, there is a need to identify high-quality self-report instruments to be used in pediatric oncology research studies. This study's objective was to conduct a systematic literature review of existing English language instruments used to measure self-reported symptoms in children and adolescents undergoing cancer treatment.

Methods—A comprehensive literature search was conducted in MEDLINE/PubMed, EMBASE, CINAHL, and PsycINFO to identify relevant articles published through November 10, 2016. Using pre-specified inclusion/exclusion criteria, six trained reviewers carefully screened abstracts and full-text articles for eligibility.

Results—There were 7738 non-duplicate articles identified in the literature search. Forty articles met our eligibility criteria, and within these articles, there were 38 self-report English symptom instruments. Most studies evaluated only cross-sectional psychometric properties, such as reliability or validity. Ten studies assessed an instrument's responsiveness or ability to detect changes in symptoms over time. Eight instruments met our criteria for use in future longitudinal pediatric oncology studies.

Conclusions—This systematic review aids pediatric oncology researchers in identifying and selecting appropriate symptom measures with strong psychometric evidence for their studies. Enhancing the child's voice in pediatric oncology research studies allows us to better understand the impact of cancer and its treatment on the lives of children.

Keywords

Pediatric oncology; Self-report instruments; Adverse event self-report

Introduction

In 2016, an estimated 10,380 children and adolescents between the ages of 0 and 14 years were diagnosed with cancer in the United States [1]. Each year, approximately 40,000 children undergo cancer treatment [2] and more than 60% participate in a clinical trial [3, 4]. Treatments are associated with high degrees of symptom burden, which have both subtle and visible effects on children. For example, nearly 50% of children experience treatment-related fatigue and 30% experience treatment-related pain, nausea, cough, lack of appetite, and psychological deteriorations [5]. Unpleasant side effects may result in significant decrements in physical, mental, emotional, and social health domains as well as disruptions in primary and secondary education, which are of great concern in this young patient population [6–12].

In the past 20 years in pediatric oncology, there has been an increasing recognition of self-reporting being the gold standard for subjective health status indicators, including symptom burden and health-related quality of life (HRQOL) [13, 14]. Objective indicators of symptom burden may not accurately reflect unobservable symptoms that could be better captured by subjective reports of symptom experiences. That is, an objective symptom experience such as a skin ulceration may be more amenable to parent or clinician report, but an unobservable symptom experience such as nausea may be better represented by a child's own self-report. Several pediatric oncology studies have found that compared to child self-report, clinicians often under-report both the prevalence and severity of subjective treatment-related symptoms experienced by children [15–18]. Studies on the agreement between

children and parent proxy symptom reports have found that parents' scores only agree fairly to moderately with their children's self-reports with correlations ranging from 0.30 to 0.49 [8, 9, 15, 19–23]. These research findings indicate that clinician or parental proxy reports of cancer treatment burden for the ill child may not accurately reflect views from the child's perspective [24]. As such, it is imperative for us to better understand the child's experience in order to provide more patient-centered care in pediatric oncology.

Several self-report symptom instruments have been developed and used in pediatric oncology. However, these instruments differ in important ways including: symptoms assessed, number of questions asked, reference period used, phrasing of questions, and number and type of response options. Differences often reflect the respective study's target population, with consideration of type of cancer, developmental stage, and/or the chronological age of the children being studied. There is also variation in the type and quality of psychometric evidence supporting existing pediatric self-report instruments. For example, some instruments used in pediatric oncology have not actually been validated in children (only in adults) whereas others have been validated in children, but not those with cancer (Fig. 1).

The objective of the current study was to conduct a systematic literature review to identify and evaluate available English language instruments for measuring self-reported symptoms in children and adolescents undergoing cancer treatment. We evaluated the evidence regarding the psychometric properties of each identified instrument, whether or not the evidence derived was from cross-sectional or longitudinal evaluations, and if the instrument was used in a clinical trial. Based on the evidence, we offer recommendations regarding which self-report instruments are ready for use in pediatric oncology research. Findings from our comprehensive systematic review will help inform longitudinal studies, comparative effectiveness research studies, patient-centered outcomes research studies, and clinical trials that are interested in using self-report symptom instruments in pediatric oncology.

Methods

Literature search strategy

A comprehensive literature search was conducted in MEDLINE/PubMed, EMBASE, CINAHL, and PsycINFO, to identify relevant articles from each database's inception through November 10, 2016. The literature search included Medical Subject Headings (MeSH), Emtree headings, and related text and keyword searches when appropriate, focusing on terms used to describe self-reported symptom measurements in children and adolescents with cancer (Appendix A).

The study's search strategy was developed with input from members of the research team, and an experienced librarian who conducted the searches. Details of the search strategy are presented in Appendix A.

Inclusion and exclusion criteria

Criteria for inclusion of studies were: (1) only empirical studies with children and adolescent self-report instruments; (2) in the English language; (3) designed or evaluated in children or adolescents with cancer less than 21 years of age; (4) focused on measuring symptoms and either reported scores for symptoms individually or in aggregate form (e.g., overall toxicity or symptom burden score); (5) reporting results for participants that were on treatment separately from those who were off treatment, as we were specifically interested in children currently on treatment; and (6) reporting psychometric evidence for the instrument in at least one publication in a pediatric cancer population. Psychometric evidence included cross-sectional or longitudinal studies that assessed reliability (internal consistency, test—retest) and validity (content, construct, responsiveness) of the self-report instrument. Studies that only correlated child assessments with parent/proxy assessments, but did not evaluate other psychometric properties did not meet our criteria for psychometric evaluation.

HRQOL instruments were eligible if they also measured symptoms and reported scores for symptoms (individually or aggregated), but not if HRQOL scores were summarized as an overall HRQOL or well-being score. Studies were excluded if instruments only reported parent or clinician proxy measures.

Data abstraction: cross-sectional psychometric properties

Two trained members of the study team evaluated each eligible study's design (e.g., cross-sectional or longitudinal), inclusion/exclusion criteria of the study sample (e.g., English speaking/reading only), participant characteristics (e.g., sample size, mean age, percent female, cancer types included), reliability of each measure (e.g., internal consistency and test—retest), and how validity was assessed (e.g., content, construct). In particular, an instrument's validity had to be assessed using an instrument that has been validated in a pediatric oncology population. Two senior members of the study team assessed all data abstraction for quality control and consistency. The psychometric properties of the eligible instruments are presented in Table 2.

Data abstraction: longitudinal psychometric properties

Two members of the study team further evaluated studies that were deemed eligible for the psychometric property assessment to determine whether they also assessed the pediatric self-report instrument's responsiveness or ability to detect changes over time. Eligible studies included ones that performed a psychometric evaluation of responsiveness or studies that simply collected and reported change scores for the instrument at two time points. Longitudinal study attributes including study design (e.g., observational, randomized control trial), participant characteristics (e.g., cancer types, mean age, sample size, percent female), and conclusions regarding the instrument's responsiveness are presented in Table 3.

Data abstraction: use of instrument in clinical trials

We were interested in which pediatric self-report symptom measurement instruments were currently being used in oncology clinical trials. We conducted a review of the [ClinicalTrials.gov](https://clinicaltrials.gov) database (last checked September 2016) using these search criteria: (1) instrument name (or abbreviated name) identified from the first two searches; (2) cancer;

and (3) child (limited to birth—21 years of age). We collected information on the study status (i.e., completed, recruiting), study type (i.e., interventional or observational), intervention type, phase of trial, estimated sample size, age of children, and if self-report symptoms were measured as a primary, secondary, or exploratory endpoint. Findings are presented in Supplemental Table 4.

Criteria for the symptom instruments that are ready for pediatric oncology longitudinal research

Informed by published criteria on the use of patient-reported outcomes (PROs) for outcomes research, [26–29] we identified the following three criteria to select an instrument to be ready for pediatric oncology research with longitudinal data collection:

1. There must be evidence of the instrument's reliability ($\alpha > .70$) in children receiving treatment for cancer. Reliability can include either internal consistency or test—retest reliability.
2. There must be evidence for the validity of the measure in children undergoing treatment for cancer. Validity includes assessments of content and construct validity.
3. The symptom measure must show evidence of the measure's ability to capture symptom change over time in children undergoing treatment for cancer.

Results

Literature search results

A total of 9482 articles were identified through database searching, of which 7738 were non-duplicates. A total of 7166 articles were excluded during the initial abstract screening phase leaving us with 567 articles eligible for full-text review. An additional 516 articles were excluded during the full-text review phase, leaving 53 articles that met all eligibility criteria for inclusion in this study. For cross-sectional psychometric studies, we included 40 empirical studies. For longitudinal studies, we included 20 empirical studies.

Study selection

A minimum of two trained members of the research team independently screened all titles and abstracts for inclusion using the eligibility criteria described above. Six trained reviewers screened the initial 7738 abstracts. Inter-rater reliability among all six reviewers was kappa of 0.72 (95% CI 0.61–0.84), estimated from a subset of 100 articles that all six team members reviewed. Studies with titles and abstracts that met inclusion criteria or lacked adequate information to determine inclusion or exclusion underwent full-text review. A senior member of the review team resolved conflicts.

During the full-text review, a minimum of two trained members of the research team independently reviewed each full-text article for inclusion or exclusion based on the eligibility criteria described above. If both reviewers agreed that a study did not meet eligibility criteria, the study was excluded. If the two reviewers disagreed, a senior member

of the review team resolved conflicts. Five reviewers participated in the full-text article screening.

Table 1 presents characteristics of the 38 self-report English symptom instruments used in children and adolescents undergoing cancer treatment that met our eligibility criteria. Table 1 includes a summary of the instruments' age range, types of symptoms assessed, attributes of the symptoms measured (e.g., severity, frequency, interference), recall period (e.g., past 7 days), number of questions included, and response options.

Pediatric self-report instrument characteristics

The youngest recommended age was 4 years; instruments intended for younger ages rely on faces scales for response options (Table 1). Most instrument age ranges were from 8 to 18 years (Table 1). Some instruments were dedicated to measuring one symptom while others included a range of symptoms experienced by children and adolescents undergoing cancer treatment. Reference periods for symptom recall ranged between "right now" (or today), 7 days (or the past week) up to one year prior. Response options varied considerably and most instruments used a version of the Likert scale. The most commonly assessed symptoms were nausea, pain, fatigue, depression, and anxiety. The 38 self-report symptom instruments included in this review measured approximately 81 different symptoms. The most comprehensive instrument was the Memorial Symptom Assessment Scale (MSAS) 10–18, assessing 30 symptoms [5]. Whereas the Pediatric Patient-Reported Outcomes version of the Common Terminology Criteria for Adverse Events (PRO-CTCAE) is an item library assessing up to 62 symptomatic toxicities [30].

Psychometric studies

Table 2 summarizes the 40 psychometric studies of the 38 symptom instruments included in Table 1. The majority of the studies assessed cross-sectional psychometric properties with 10 instruments being assessed longitudinally. Sample sizes ranged from 6 to 291 children/adolescents, but nearly 60% of eligible studies included fewer than 100 children. Instruments generally used a diverse sample of children and adolescents with respect to cancer type, age, and gender.

Instruments varied in terms of reporting scores for individual symptoms versus aggregating across symptoms to report an overall toxicity score. As such, some studies reported one internal consistency score, whereas others reported Cronbach's alphas for instrument subscales or specific symptoms. Overall, 20 validated instruments reported reliability measures above 0.70 when assessed at single time point (internal consistency) or multiple time points (test—retest). There were 15 instruments that did not report any measures of reliability and three instruments whose reliability did not meet our 0.70 threshold.

The types of evidence for the validity of the instruments also varied across instruments. Many studies evaluated an instrument's content validity using focus groups, in-depth interviews, or expert opinion (e.g., clinicians). Other studies tested known group's validity by comparing children on and off treatment or children with cancer to other chronic diseases or to healthy children their age. Convergent validity was often evaluated by comparing pediatric self-report scores to parent proxies as well as clinicians and nurses. Other studies

assessed convergent validity by comparing the new instrument with an established PRO instrument that was known to measure the same or similar concept of interest. Finally, some studies compared self-report scores to other more objective measures such as sleep duration or number of emetic episodes.

Longitudinal studies

Table 3 presents longitudinal studies (not test—retest studies) that examined an instrument's responsiveness or sensitivity to changes over time. Ten instruments were used to assess changes over time in children undergoing cancer treatment. Each study include 2–5 assessment points spanning two or more 24-h sequential periods and several months up to one year. We also present data on the timing of assessment points to highlight the variation across studies. Many of the eligible studies were not designed as psychometric studies, but rather were studies that evaluated symptoms over time as part of their larger study objectives (e.g., evaluating the effectiveness of an intervention).

Instruments used in clinical trials

Supplemental Table 4 includes instruments that have been used in studies that are registered in clinicaltrials.gov. Twenty-two instruments were identified in this search. The PedsQL 4.0 Generic Core Scales, PedsQL Multidimensional Fatigue Scale, PedsQL 3.0 Cancer Module, Fatigue Scale (Adolescent and Child version), and the Faces Pain Scale-Revised were the instruments most often cited in clinicaltrials.gov.

Discussion

This systematic review identified 38 self-report symptom instruments used in pediatric oncology research with at least one psychometric validation study in children with cancer undergoing active treatment. These English language instruments varied considerably in terms of symptoms measured, phrasing and format of questions asked, and the amount of psychometric evidence supporting each instrument.

Ten of the instruments were included in a study with the intention of measuring change in a symptom or aggregate of symptoms over time. However, a majority of the longitudinal studies were not designed as a psychometric study to evaluate the responsiveness of the instrument. For example, some studies collected self-report symptom and HRQOL data over time, but did not characterize the trends over time. The following instruments met our three assessment criteria for use in longitudinal pediatric oncology studies (described above): Children's Depression Inventory, State-Trait Anxiety Inventory for Children, Children's International Mucositis Evaluation Scale, Fatigue Scale (Child and Adolescent versions), Pain Squad App, PedsQL 4.0 Generic Core Scales, and PedsQL 3.0 Cancer Module. These eight instruments have been psychometrically evaluated (both reliability and validity) and able to detect significant changes over time. All of the instruments, except the Pain Squad App, are currently being used in a pediatric oncology clinical trial.

The NIH's Patient-Reported Outcomes Measurement Information System (PROMIS) Pediatric instruments include measures of symptoms such as anxiety, depressive symptoms, fatigue, and pain interference. At the time of this article's publication, cross-sectional data

supporting the validity and reliability of the instruments in children with cancer have been published [31, 32]. In addition, these measures have been found to be responsive over time [14]. It is the opinion of the authors that the PROMIS pediatric measures show adequate evidence for their use in longitudinal pediatric oncology studies.

Methods for summarization of patient-reported data as scores also varied considerably, an issue particularly important for symptom research. The use of summary scores (i.e., aggregating across multiple symptoms) to represent overall patient well-being or symptom burden can often mask domain- or symptom-specific decrements. Summarizing data as individual symptom scores allows researchers and clinicians to examine the effectiveness of an intervention on a specific symptom or adverse event.

Some studies did not show statistically significant changes in symptoms over time. We believe that some of the non-significant findings are due to specific features of the study design. For example, issues of small sample sizes and not selecting time points that capture an instrument's sensitivity (i.e., assessments selected at times when symptoms change in intensity) were seen across most studies that did not observe differences over time. Factors such as broad study participant inclusion criteria and ambiguous time points likely contributed to small effect sizes and insignificant results. If researchers intend to measure an instrument's responsiveness, they must be thoughtful and deliberate when selecting time points, being careful to choose time points where it is reasonable to expect a change in the symptom experience. Another possibility is to ask the patient directly if they felt a change in their symptoms between time points, which could be done by administering a global impression of change scale. These considerations are especially important for social or behavioral interventions where the anticipated effect size may already be small.

Additionally, a few studies, including clinical trials, collected longitudinal data using self-report symptom instruments, but investigators did not present findings in their publications. Researchers should consider including results from these types of instruments in their published findings to add further to the evidence base. Further, for studies that include different instrument versions for child, adolescent, and adult reports of symptoms, treatment toxicities, and quality of life, we recommend that the findings be distinctly reported for each instrument version and not as one total group.

Limitations

There were limitations to this systematic review. First, our search was limited to English language instruments because verifying the quality of the translation of the instrument and the further content evaluation was outside the scope of this review. Including only English language instruments limits the geographical scope of this review. Next, inconsistent use of the instrument's name challenged our search within clinicaltrials.gov. As such, if an investigator entered only a portion of the instrument's name or abbreviation, it is possible the instrument does not appear in Supplemental Table 4. In addition, while we did assess the quality of the instrument (i.e., validity and reliability were assessed), we did not examine the quality of each study. We recommend that future studies examine this important issue. Lastly, we focused only on symptom measures for children currently undergoing active

cancer treatment, but further studies should also consider self-report instruments for children and adolescents in palliative care as well as survivors of childhood cancer.

Conclusion

Investigators should consider several issues in the selection of the appropriate self-report instrument for their pediatric oncology study. Before considering the use of an instrument, the researcher should have a comprehensive understanding of the characteristics of the target population in terms of age, cancer type, and disease status. They should know what outcomes or domains they want to measure, which should be driven by the study's research aims. The selected instrument should be designed to be relevant for the age of the children under study, and take into account a child's literacy and cognitive abilities. In addition, the instrument should have undergone psychometric evaluation of reliability, validity, and responsiveness, ideally with the study's target population. If the particular domains are primary study endpoints, choosing an instrument with strong reliability and validity data is optimal.

This systematic review serves as guidance for pediatric oncology researchers to select appropriate symptom measures with strong psychometric evidence. The overarching motivation behind this work is to enhance the child's voice in pediatric oncology research studies in order to obtain a better understanding of the impact of cancer and its treatment on the lives of children.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank Natalie Ernecoff, Kathryn Jackson, Susan Keller, and Sejin Lee for their help with this study.

Funding This research was supported by the National Cancer Institute of the National Institutes of Health under award number R01CA175759.

References

1. American Cancer Society. Cancer facts & figures 2016. Atlanta: American Cancer Society; 2016. p. 11
2. CureSearch Childhood Cancer Statistics. <http://curesearch.org/Childhood-Cancer-Statistics>
3. Sateren WB, Trimble EL, Abrams J, Brawley O, Breen N, Ford L, et al. How sociodemographics, presence of oncology specialists, and hospital cancer programs affect accrual to cancer treatment trials. *Journal of Clinical Oncology*. 2002; 20(8):2109–2117. [PubMed: 11956272]
4. Bleyer A, Budd T, Montello M. Adolescents and young adults with cancer: The scope of the problem and criticality of clinical trials. *Cancer*. 2006; 107(7 Suppl):1645–1655. DOI: 10.1002/cncr.22102 [PubMed: 16906507]
5. Collins JJ, Byrnes ME, Dunkel IJ, Lapin J, Nadel T, Thaler HT, et al. The measurement of symptoms in children with cancer. *Journal of Pain and Symptom Management*. 2000; 19(5):363–377. [PubMed: 10869877]
6. Docherty SL. Symptom experiences of children and adolescents with cancer. *Annual Review of Nursing Research*. 2003; 21:123–149.

7. Eiser C, Hill JJ, Vance YH. Examining the psychological consequences of surviving childhood cancer: Systematic review as a research method in pediatric psychology. *Journal of Pediatric Psychology*. 2000; 25(6):449–460. [PubMed: 10980049]
8. Linder LA. Measuring physical symptoms in children and adolescents with cancer. *Cancer Nursing*. 2005; 28(1):16–26. [PubMed: 15681978]
9. Pickard AS, Topfer LA, Feeny DH. A structured review of studies on health-related quality of life and economic evaluation in pediatric acute lymphoblastic leukemia. *Journal of the National Cancer Institute Monographs*. 2004; 2004(33):102–125. DOI: 10.1093/jncimonographs/lgh002
10. Ruland CM, Hamilton GA, Schjodt-Osmo B. The complexity of symptoms and problems experienced in children with cancer: A review of the literature. *Journal of Pain and Symptom Management*. 2009; 37(3):403–418. DOI: 10.1016/j.jpainsymman.2008.03.009 [PubMed: 18694633]
11. Vance YH, Eiser C. The school experience of the child with cancer. *Child: Care, Health and Development*. 2002; 28(1):5–19.
12. Waters E, Stewart-Brown S, Fitzpatrick R. Agreement between adolescent self-report and parent reports of health and well-being: Results of an epidemiological study. *Child: Care, Health and Development*. 2003; 29(6):501–509.
13. F. A. D. A. U.S. Department of Health and Human Services. , editor. U.S. Department of Health and Human Services, F. A. D. A.. Guidance for industry: Patient-reported outcome measures: Use in medical product development to support labeling claims. 2009.
14. Hinds, P. S. (2016). In B. B. Reeve (Ed.), (Email ed.).
15. Hockenberry MJ, Hinds PS, Barrera P, Bryant R, Adams-McNeill J, Hooke C, et al. Three instruments to assess fatigue in children with cancer: The child, parent and staff perspectives. *Journal of Pain and Symptom Management*. 2003; 25(4):319–328. [PubMed: 12691683]
16. Le Gales C, Costet N, Gentet JC, Kalifa C, Frappaz D, Edan C, et al. Cross-cultural adaptation of a health status classification system in children with cancer. First results of the French adaptation of the Health Utilities Index Marks 2 and 3. *International Journal of Cancer Supplement*. 1999; 12:112–118.
17. Glaser AW, Davies K, Walker D, Brazier D. Influence of proxy respondents and mode of administration on health status assessment following central nervous system tumours in childhood. *Quality of Life Research*. 1997; 6(1):43–53. [PubMed: 9062441]
18. Parsons SK, Barlow SE, Levy SL, Supran SE, Kaplan SH. Health-related quality of life in pediatric bone marrow transplant survivors: According to whom? *International Journal of Cancer Supplement*. 1999; 12:46–51.
19. Collins JJ, Devine TD, Dick GS, Johnson EA, Kilham HA, Pinkerton CR, et al. The measurement of symptoms in young children with cancer: The validation of the Memorial Symptom Assessment Scale in children aged 7–12. *Journal of Pain and Symptom Management*. 2002; 23(1):10–16. [PubMed: 11779663]
20. Yeh CH, Chang CW, Chang PC. Evaluating quality of life in children with cancer using children's self-reports and parent-proxy reports. *Nursing Research*. 2005; 54(5):354–362. [PubMed: 16224322]
21. Upton P, Lawford J, Eiser C. Parent-child agreement across child health-related quality of life instruments: A review of the literature. *Quality of Life Research*. 2008; 17(6):895–913. DOI: 10.1007/s11136-008-9350-5 [PubMed: 18521721]
22. Russell KM, Hudson M, Long A, Phipps S. Assessment of health-related quality of life in children with cancer: Consistency and agreement between parent and child reports. *Cancer*. 2006; 106(10): 2267–2274. DOI: 10.1002/cncr.21871 [PubMed: 16604563]
23. Levi RB, Drotar D. Health-related quality of life in childhood cancer: Discrepancy in parent-child reports. *International Journal of Cancer Supplement*. 1999; 12:58–64.
24. Sawyer M, Antoniou G, Toogood I, Rice M. A comparison of parent and adolescent reports describing the health-related quality of life of adolescents treated for cancer. *International Journal of Cancer Supplement*. 1999; 12:39–45.

25. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009; 6(7):e1000097.doi: 10.1371/journal.pmed1000097 [PubMed: 19621072]
26. Reeve BB, Withycombe JS, Baker JN, Hooke MC, Lyons JC, Mowbray C, et al. The first step to integrating the child's voice in adverse event reporting in oncology trials: A content validation study among pediatric oncology clinicians. *Pediatric Blood & Cancer.* 2013; 60(7):1231–1236. DOI: 10.1002/pbc.24463 [PubMed: 23335328]
27. Mokkink LB, Terwee CB, Knol DL, Stratford PW, Alonso J, Patrick DL, et al. The COSMIN checklist for evaluating the methodological quality of studies on measurement properties: A clarification of its content. *BMC Medical Research Methodology.* 2010; 10:22.doi: 10.1186/1471-2288-10-22 [PubMed: 20298572]
28. Matza LS, Patrick DL, Riley AW, Alexander JJ, Rajmil L, Pleil AM, et al. Pediatric patient-reported outcome instruments for research to support medical product labeling: Report of the ISPOR PRO good research practices for the assessment of children and adolescents task force. *Value Health.* 2013; 16(4):461–479. DOI: 10.1016/j.jval.2013.04.004 [PubMed: 23796280]
29. Aaronson N, Alonso J, Burnam A, Lohr KN, Patrick DL, Perrin E, et al. Assessing health status and quality-of-life instruments: Attributes and review criteria. *Quality of Life Research.* 2002; 11(3):193–205. [PubMed: 12074258]
30. Reeve BB, McFatrigh M, Pinheiro LC, Weaver MS, Sung L, Withycombe JS, et al. Eliciting the child's voice in adverse event reporting in oncology trials: Cognitive interview findings from the pediatric patient-reported outcomes version of the common terminology criteria for adverse events initiative. *Pediatric Blood & Cancer.* 2017; doi: 10.1002/pbc.26261
31. Hinds PS, Nuss SL, Ruccione KS, Withycombe JS, Jacobs S, DeLuca H, et al. PROMIS pediatric measures in pediatric oncology: Valid and clinically feasible indicators of patient-reported outcomes. *Pediatric Blood & Cancer.* 2013; 60(3):402–408. DOI: 10.1002/pbc.24233 [PubMed: 22829446]
32. DeWalt DA, Gross HE, Gipson DS, Selewski DT, DeWitt EM, Dampier CD, et al. PROMIS((R)) pediatric self-report scales distinguish subgroups of children within and across six common pediatric chronic health conditions. *Quality of Life Research.* 2015; 24(9):2195–2208. DOI: 10.1007/s11136-015-0953-3 [PubMed: 25715946]
33. Walker AJ, Johnson KP, Miaskowski C, Lee KA, Gedaly-Duff V. Sleep quality and sleep hygiene behaviors of adolescents during chemotherapy. *Journal of Clinical Sleep Medicine.* 2010; 6(5): 439–444. [PubMed: 20957843]
34. Sawyer M, Antoniou G, Toogood I, Rice M. A comparison of parent and adolescent reports describing the health-related quality of life of adolescents treated for cancer. *International Journal of Cancer, Suppl.* 1999; 12:39–45.
35. Hockenberry-Eaton M, Manteuffel B, Bottomley S. Development of two instruments examining stress and adjustment in children with cancer. *Journal of Pediatric Oncology Nursing.* 1997; 14(3): 178–185. [PubMed: 9260409]
36. Noll RB, Gartstein MA, Vannatta K, Correll J, Bukowski WM, Davies WH. Social, emotional, and behavioral functioning of children with cancer. *Pediatrics.* 1999; 103(1):71–78. [PubMed: 9917442]
37. Jacobs S, Baggott C, Agarwal R, Hesser T, Schechter T, Judd P, et al. Validation of the Children's International Mucositis Evaluation Scale (ChIMES) in paediatric cancer and SCT. *British Journal of Cancer.* 2013; 109(10):2515–2522. DOI: 10.1038/bjc.2013.618 [PubMed: 24129238]
38. Tomlinson D, Gibson F, Treister N, Baggott C, Judd P, Hendershot E, et al. Refinement of the Children's International Mucositis Evaluation Scale (ChIMES): Child and parent perspectives on understandability, content validity and acceptability. *European Journal of Oncology Nursing.* 2010; 14(1):29–41. DOI: 10.1016/j.ejon.2009.10.004 [PubMed: 19955021]
39. Tomlinson D, Hesser T, Maloney AM, Ross S, Naqvi A, Sung L. Development and initial evaluation of electronic Children's International Mucositis Evaluation Scale (eChIMES) for children with cancer. *Supportive Care in Cancer.* 2014; 22(1):115–119. DOI: 10.1007/s00520-013-1953-x [PubMed: 24000043]

40. Woolery M, Carroll E, Fenn E, Wieland H, Jarosinski P, Corey B, et al. A constipation assessment scale for use in pediatric oncology. *Journal of Pediatric Oncology Nursing*. 2006; 23(2):65–74. DOI: 10.1177/1043454205285874 [PubMed: 16476780]
41. Hinds PS, Hockenberry M, Rai SN, Zhang L, Razzouk BI, Cremer L, et al. Clinical field testing of an enhanced-activity intervention in hospitalized children with cancer. *Journal of Pain and Symptom Management*. 2007; 33(6):686–697. [PubMed: 17360151]
42. Hinds PS, Hockenberry M, Tong X, Rai SN, Gattuso JS, McCarthy K, et al. Validity and reliability of a new instrument to measure cancer-related fatigue in adolescents. *Journal of Pain and Symptom Management*. 2007; 34(6):607–618. DOI: 10.1016/j.jpainsymman.2007.01.009 [PubMed: 17629669]
43. Hinds PS, Yang J, Gattuso JS, Hockenberry M, Jones H, Zupanec S, et al. Psychometric and clinical assessment of the 10-item reduced version of the Fatigue Scale-Child instrument. *Journal of Pain and Symptom Management*. 2010; 39(3):572–578. DOI: 10.1016/j.jpainsymman.2009.07.015 [PubMed: 20303031]
44. Mandrell BN, Yang J, Hooke MC, Wang C, Gattuso JS, Hockenberry M, et al. Psychometric and clinical assessment of the 13-item reduced version of the fatigue scale-adolescent instrument. *Journal of Pediatric Oncology Nursing*. 2011; 28(5):287–294. DOI: 10.1177/1043454211418667 [PubMed: 21844243]
45. Bhatia S, Jenney ME, Bogue MK, Rockwood TH, Feusner JH, Friedman DL, et al. The minneapolis-manchester quality of life instrument: Reliability and validity of the adolescent form. *Journal of Clinical Oncology*. 2002; 20(24):4692–4698. DOI: 10.1200/JCO.2002.05.103 [PubMed: 12488415]
46. Wu E, Robison LL, Jenney ME, Rockwood TH, Feusner J, Friedman D, et al. Assessment of health-related quality of life of adolescent cancer patients using the Minneapolis-Manchester Quality of Life Adolescent Questionnaire. *Pediatric Blood & Cancer*. 2007; 48(7):678–686. DOI: 10.1002/pbc.20874 [PubMed: 16628553]
47. Bhatia S, Jenney ME, Wu E, Bogue MK, Rockwood TH, Feusner JH, et al. The minneapolis-manchester quality of life instrument: Reliability and validity of the youth form. *Journal of Pediatrics*. 2004; 145(1):39–46. DOI: 10.1016/j.jpeds.2004.02.034 [PubMed: 15238904]
48. Shankar S, Robison L, Jenney ME, Rockwood TH, Wu E, Feusner J, et al. Health-related quality of life in young survivors of childhood cancer using the minneapolis-manchester quality of life-youth form. *Pediatrics*. 2005; 115(2):435–442. DOI: 10.1542/peds.2004-0649 [PubMed: 15687453]
49. Manji A, Tomlinson D, Ethier MC, Gassas A, Maloney AM, Sung L. Psychometric properties of the oral mucositis daily questionnaire for child self-report and importance of mucositis in children treated with chemotherapy. *Supportive Care in Cancer*. 2012; 20(6):1251–1258. DOI: 10.1007/s00520-011-1211-z [PubMed: 21681385]
50. Tomlinson D, Isitt JJ, Barron RL, Doyle J, Judd P, Gassas A, et al. Determining the understandability and acceptability of an oral mucositis daily questionnaire. *Journal of Pediatric Oncology Nursing*. 2008; 25(2):107–111. DOI: 10.1177/1043454208314458 [PubMed: 18270307]
51. Stinson JN, Jibb LA, Nguyen C, Nathan PC, Maloney AM, Dupuis LL, et al. Construct validity and reliability of a real-time multidimensional smartphone app to assess pain in children and adolescents with cancer. *Pain*. 2015; 156(12):2607–2615. DOI: 10.1097/j.pain.0000000000000385 [PubMed: 26580680]
52. Morley TE, Cataudella D, Fernandez CV, Sung L, Johnston DL, Nesin A, et al. Development of the Pediatric Advanced Care Quality of Life Scale (PAC-QoL): evaluating comprehension of items and response options. *Pediatric Blood & Cancer*. 2014; 61(10):1835–1839. DOI: 10.1002/pbc.25111 [PubMed: 24947009]
53. Seid M, Varni JW, Rode CA, Katz ER. The pediatric cancer quality of life inventory: a modular approach to measuring health-related quality of life in children with cancer. *International Journal of Cancer Supplement*. 1999; 12:71–76.
54. Varni JW, Katz ER, Seid M, Quiggins DJ, Friedman-Bender A. The pediatric cancer quality of life inventory-32 (PCQL-32): I. Reliability and validity. *Cancer*. 1998; 82(6):1184–1196. [PubMed: 9506367]

55. Lai JS, Cella D, Peterman A, Barocas J, Goldman S. Anorexia/cachexia-related quality of life for children with cancer. *Cancer*. 2005; 104(7):1531–1539. DOI: 10.1002/cncr.21315 [PubMed: 16088963]
56. Dupuis LL, Taddio A, Kerr EN, Kelly A, MacKeigan L. Development and validation of the pediatric nausea assessment tool for use in children receiving antineoplastic agents. *Pharmacotherapy*. 2006; 26(9):1221–1231. DOI: 10.1592/phco.26.9.1221 [PubMed: 16945043]
57. Roddenberry A, Renk K. Quality of life in pediatric cancer patients: The relationships among parents' characteristics, children's characteristics, and informant concordance. *Journal of Child and Family Studies*. 2008; 17(3):402–426. DOI: 10.1007/s10826-007-9155-0
58. Reeve BB, McFatrigh M, Pinheiro LC, Weaver MS, Sung L, Withycombe JS, et al. Eliciting the child's voice in adverse event reporting in oncology trials: Cognitive interview findings from the pediatric patient-reported outcomes version of the common terminology criteria for adverse events initiative. *Pediatric Blood & Cancer*. 2017; doi: 10.1002/pbc.26261
59. Fortier MA, Wahi A, Bruce C, Maurer EL, Stevenson R. Pain management at home in children with cancer: A daily diary study. *Pediatric Blood & Cancer*. 2014; 61(6):1029–1033. DOI: 10.1002/pbc.24907 [PubMed: 24376073]
60. Parsons SK, Fairclough DL, Wang J, Hinds PS. Comparing longitudinal assessments of quality of life by patient and parent in newly diagnosed children with cancer: The value of both raters' perspectives. *Quality of Life Research*. 2012; 21(5):915–923. DOI: 10.1007/s11136-011-9986-4 [PubMed: 21822735]
61. Penn A, Lowis SP, Hunt LP, Shortman RI, Stevens MC, McCarter RL, et al. Health related quality of life in the first year after diagnosis in children with brain tumours compared with matched healthy controls; a prospective longitudinal study. *European Journal of Cancer*. 2008; 44(9):1243–1252. DOI: 10.1016/j.ejca.2007.09.015 [PubMed: 17997300]
62. Razzouk BI, Hord JD, Hockenberry M, Hinds PS, Feusner J, Williams D, et al. Double-blind, placebo-controlled study of quality of life, hematologic end points, and safety of weekly epoetin alfa in children with cancer receiving myelosuppressive chemotherapy. *Journal of Clinical Oncology*. 2006; 24(22):3583–3589. DOI: 10.1200/JCO.2005.03.4371 [PubMed: 16877725]
63. Varni JW, Burwinkle TM, Katz ER, Meeske K, Dickinson P. The PedsQL in pediatric cancer: reliability and validity of the Pediatric Quality of Life Inventory Generic Core Scales, Multidimensional Fatigue Scale, and Cancer Module. *Cancer*. 2002; 94(7):2090–2106. [PubMed: 11932914]
64. Wesley KM, Zelikovsky N, Schwartz LA. Physical symptoms, perceived social support, and affect in adolescents with cancer. *Journal of Psychosocial Oncology*. 2013; 31(4):451–467. DOI: 10.1080/07347332.2013.798761 [PubMed: 23844924]
65. Phipps S, Steele R. Repressive adaptive style in children with chronic illness. *Psychosomatic Medicine*. 2002; 64(1):34–42. [PubMed: 11818584]
66. Schneider SM, Workman ML. Effects of virtual reality on symptom distress in children receiving chemotherapy. *Cyberpsychology & Behavior*. 1999; 2(2):125–134. DOI: 10.1089/cpb.1999.2.125 [PubMed: 19178248]
67. O'Sullivan C, Dupuis LL, Gibson P, Johnston DL, Baggott C, Portwine C, et al. Refinement of the symptom screening in pediatrics tool (SSPedi). *British Journal of Cancer*. 2014; 111(7):1262–1268. DOI: 10.1038/bjc.2014.445 [PubMed: 25101569]
68. Redd WH, Jacobsen PB, Die-Trill M, Dermatis H, McEvoy M, Holland JC. Cognitive/attentional distraction in the control of conditioned nausea in pediatric cancer patients receiving chemotherapy. *Journal of Consulting and Clinical Psychology*. 1987; 55(3):391–395. [PubMed: 3597954]
69. Mulhern RK, Fairclough D, Douglas SM, Smith B. Physical distress and depressive symptomatology among children with cancer. *Children's Health Care*. 1994; 23(3):167–179. DOI: 10.1207/s15326888chc2303_2
70. Hinds PS, Hockenberry MJ, Gattuso JS, Srivastava DK, Tong X, Jones H, et al. Dexamethasone alters sleep and fatigue in pediatric patients with acute lymphoblastic leukemia. *Cancer*. 2007; 110(10):2321–2330. DOI: 10.1002/cncr.23039 [PubMed: 17926333]

71. Hinds PS, Hockenberry M, Rai SN, Zhang L, Razzouk BI, McCarthy K, et al. Nocturnal awakenings, sleep environment interruptions, and fatigue in hospitalized children with cancer. *Oncology Nursing Forum*. 2007; 34(2):393–402. DOI: 10.1188/07.ONF.393-402 [PubMed: 17573303]
72. Hooke MC, Garwick AW, Gross CR. Fatigue and physical performance in children and adolescents receiving chemotherapy. *Oncology Nursing Forum*. 2011; 38(6):649–657. DOI: 10.1188/11.ONF.649-657 [PubMed: 22037328]
73. Hooke MC, McCarthy K, Taylor O, Hockenberry MJ. Fatigue and carnitine levels over multiple cycles of chemotherapy in children and adolescents. *European Journal of Oncology Nursing*. 2015; 19(1):7–12. DOI: 10.1016/j.ejon.2014.07.015 [PubMed: 25260954]
74. Hooke MC, Gilchrist L, Tanner L, Hart N, Withycombe JS. Use of a fitness tracker to promote physical activity in children with acute lymphoblastic leukemia. *Pediatric Blood & Cancer*. 2016; 63(4):684–689. DOI: 10.1002/pbc.25860 [PubMed: 26756736]
75. Jacobs S, Mowbray C, Cates LM, Baylor A, Gable C, Skora E, et al. Pilot study of massage to improve sleep and fatigue in hospitalized adolescents with cancer. *Pediatric Blood & Cancer*. 2016; 63(5):880–886. DOI: 10.1002/pbc.25902 [PubMed: 26757318]
76. Penn A, Lowis SP, Stevens MC, Hunt LP, Shortman RI, McCarter RJ, et al. Family, demographic and illness-related determinants of HRQL in children with brain tumours in the first year after diagnosis. *Pediatric Blood & Cancer*. 2009; 53(6):1092–1099. DOI: 10.1002/pbc.22157 [PubMed: 19743518]
77. Penn A, Shortman RI, Lowis SP, Stevens MC, Hunt LP, McCarter RJ, et al. Child-related determinants of health-related quality of life in children with brain tumours 1 year after diagnosis. *Pediatric Blood & Cancer*. 2010; 55(7):1377–1385. DOI: 10.1002/pbc.22743 [PubMed: 20981692]
78. Klaassen RJ, Krahn M, Gaboury I, Hughes J, Anderson R, Grundy P, et al. Evaluating the ability to detect change of health-related quality of life in children with Hodgkin disease. *Cancer*. 2010; 116(6):1608–1614. DOI: 10.1002/cncr.24883 [PubMed: 20143448]
79. Tyc VL, Fairclough D, Fletcher B, Leigh L, Mulhern RK. Children's distress during magnetic resonance imaging procedures. *Child Health Care*. 1995; 24(1):5–19. DOI: 10.1207/s15326888chc2401_2 [PubMed: 10142086]
80. Wall VJ, Womack W. Hypnotic versus active cognitive strategies for alleviation of procedural distress in pediatric oncology patients. *American Journal of Clinical Hypnosis*. 1989; 31(3):181–191. DOI: 10.1080/00029157.1989.10402887 [PubMed: 2919572]

Fig. 1. PRISMA [25] diagram of studies of pediatric self-report symptom measures in children undergoing active cancer treatment

Table 1

Characteristics of pediatric self-report symptom measures in cancer research

Instrument name (alphabetical order)	Age range (years)	Symptoms (and HRQOL) domains assessed	Symptom attributes (severity, frequency, etc.)	Reference period	Number of questions	Number of response options
Adolescent Sleep Wake Scale (ASWS)	12–18	Sleep quality	Frequency	"Last week"	28	6
Child Health Questionnaire (CHQ)	10–18	Physical functioning, pain, sadness, fatigue, mobility, concentration, anger, frightened, loneliness, Nervousness, body image, worry, headache, insomnia, overall health, behavior issues	Interference, Frequency, Severity	Varied by question: "Past 4 weeks"; "In general"; "Compared to one year ago"	87	4–6
Childhood Cancer Stressors Inventory (CCSI)	7–13	Fatigue, vomiting, pain, worry, insomnia, anorexia, hair loss, weight change	Presence, Frequency, Intensity	Since the diagnosis of cancer	18	5
Children's Depression Inventory (CDI)	7–17	Depression	Presence, Severity	"Past 2 weeks"	28	3
Children's International Mucositis Evaluation Scale (ChIMES)	8–18	Mucositis (mouth/throat pain, dysphagia, mouth sores)	Severity, Interference, Presence	"Today"	6–7 (conditional questions)	2–6
Constipation Assessment Scale (CAS)	9–21	Abdominal distension (bloating), constipation, diarrhea, fecal incontinence, rectal pain, change in flatulence	Presence, Severity	"Compared to your usual pattern"	8	3
Faces Pain Scale-Revised (FPS-R)	4–16	Pain	Presence, Severity, Intensity	"Right now"	1	6
Fatigue Scale-Adolescents (FS-A)	13–18	Fatigue	Frequency, Interference	"Past 7 days"	13	5
Fatigue Scale-Children (FS-C)	7–12	Fatigue	Frequency, Interference	"Past 7 days"	10	2–5
Memorial Symptom Assessment Scale (MSAS) 7–12	7–12	Fatigue, sadness, Itch, pain, worry, anorexia, nausea, insomnia	Presence, Frequency, Severity, Distress	"Last 2 days"	8 (Additional conditional questions)	2–4
Memorial Symptom Assessment Scale (MSAS) 10–18	10–18	Fatigue, pain, concentration, cough, dry mouth, nausea, worry, neuropathy, insomnia, diarrhea, sadness, vomiting, problems with urination, nervousness, difficulty swallowing, anorexia, hair loss, weight change, dizziness, changes in skin, sweating, itch, headache, irritability, dyspnea, drowsiness,	Presence, Frequency, Severity, Distress	"Past week"	30 (Additional conditional questions)	2–5

Instrument name (alphabetical order)	Age range (years)	Symptoms (and HRQOL) domains assessed	Symptom attributes (severity, frequency, etc.)	Reference period	Number of questions	Number of response options
Minneapolis-Manchester Quality of Life-Adolescent Form (MMQL-AF)	13–20	dysgeusia, mucositis, constipation, lymphedema Physical functioning- (such as fatigue, mobility), Psychological functioning- (such as sadness, worry, anger, nervousness, loneliness, fear), Cognitive functioning- (such as concentration, memory), Appearance- (such as body image), social functioning	Presence Distress Frequency	Participants are asked to choose a statement that describes themselves or how they feel “in general”	46	2–5
Minneapolis-Manchester Quality of Life-Youth Form (MMQL-YF)	8–12	Physical symptoms- (such as blurry vision, insomnia, headache, pain, muscle aches, difficulty hearing, difficulty speaking), physical functioning- (such as fatigue, mobility), psychological functioning- (such as sadness, worry, anger, fear, loneliness), family dynamics and outlook on life	Presence Distress Frequency	Participants are asked to choose a statement that describes themselves or how they feel “in general”	32	4–5
Oral Mucositis Daily Questionnaire (OMDQ)	12–18	Mouth and throat soreness, diarrhea, overall health	Severity Impact Interference	“Past 24 h”	3 (Additional conditional questions)	4
Pain Squad App	8–18	Pain	Intensity, Interference, Unpleasantness	“Last 12 h”	22 (Additional conditional questions)	Varies from 0–12; Some open-ended questions
Patient-Reported Outcomes Measurement Information System (PROMIS) Pediatric Measures	8–17	Depression, Anxiety, Anger, Pain Interference, Fatigue, Mobility, Upper extremity functioning (see www.HealthMeasures.net for a full list of PROMIS Pediatric domains)	Frequency	“Past 7 days”	PROMIS allows different lengths of measures; Tested are 8 items per symptom except fatigue with 10 items.	5
Pediatric Advanced Care Quality of Life Scale (PAC-QoL)	8–12 child 13–18 adolescent	Physical Comfort- (such as pain, constipation, diarrhea, fatigue, nausea, dyspnea, anorexia, headache, nosebleeds, blurry vision, mobility, upper extremity functioning, difficulty speaking), psychological well-being- (such as emotional distress, worry, loneliness,	Frequency	“Past 7 days”	57–65 depending on version	4

Instrument name (alphabetical order)	Age range (years)	Symptoms (and HRQOL) domains assessed	Symptom attributes (severity, frequency, etc.)	Reference period	Number of questions	Number of response options
Pediatric Cancer Quality of Life Inventory (PCQL-32) (two versions based on age)	8–12 child 13–18 adolescent	nervousness, anger, fear, sadness, insomnia, memory) Physical functioning- (such as mobility), disease and treatment-related symptoms/problems- (such as anxiety, nausea, dysgeusia, muscle aches, pain), psychological functioning- (such as sadness, fear, worry), social functioning- (such as behavior issues), cognitive functioning- (such as concentration, memory)	Interference, Frequency	“Past month”	32	4
Pediatric-Functional Assessment of Cancer Therapy-Pediatric Anorexia/Cachexia (Peds-FAACT)	7–12 version; 12–18 version;	Fall, fatigue, mobility, Muscle weakness, pain, depression, worry, anorexia, dysgeusia, nausea, vomiting, abdominal pain, loneliness, nervousness	Frequency, Interference	“Past 7 days”	37–40 depending on version	5
Pediatric-Functional Assessment of Anorexia and Cachexia Therapy (Peds-FAACT)	12–18	Anorexia, nausea, abdominal pain, vomiting	Frequency	“Past 7 days”	10	5
10 Item version Pediatric Nausea Assessment Tool (PeNAT)	4–18	Nausea	Intensity	“Right now”	1: Plus script to determine term used by family to refer to nausea and vomiting	4
Pediatric Oncology Quality of Life Scale (POQOLS)	8–19	Physical function- (such as fatigue, mobility, pain), Reaction to current treatment- (such as hair loss, weight change, nausea, vomiting), Emotional distress- (such as sadness, anger, insomnia, fear)	Frequency	“Past two weeks”	21	7
Pediatric Patient-Reported Outcomes Version of the Common Terminology Criteria for Adverse Events (Pediatric PRO-CTCAE)	7–15	Library including 62 symptomatic adverse events; core items include: abdominal pain, anorexia, anxiety, constipation, cough, depression, diarrhea, fatigue, headache, insomnia, mucositis oral, nausea, pain, peripheral sensory neuropathy, vomiting, (and 47 additional less prevalent items)	Frequency, severity, interference	“Past 7 days”	Variable depending on the symptoms selected from the library	4

Instrument name (alphabetical order)	Age range (years)	Symptoms (and HRQOL) domains assessed	Symptom attributes (severity, frequency, etc.)	Reference period	Number of questions	Number of response options
Pediatric Quality of Life Inventory-Cancer Module (4 versions based on age)	5–7 version; 8–12 version; 13–18 version; 18–25 version	Pain and hurt, nausea, procedural anxiety, treatment anxiety, worry, cognitive problems (such as concentration), body image	Severity	“Past one month”	27; (5–7 version 26)	5; 5–7 version has 3 response options
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL)	5–7 version; 8–12 version; 13–18 version; 18–25 version	Physical functioning- (such as pain, fatigue, mobility), emotional functioning- (such as depression, insomnia, anger, scared), Social functioning, School functioning- (such as concentration problems)	Severity	Acute version- “Past 7 days” Standard version- “Past 1 month”	23	5; 5–7 version has 3 response options
Pediatric Quality of Life Inventory-Multidimensional Fatigue Scale (3 versions based on age)	5–7 version; 8–12 version; 13–18 version	General fatigue, sleep/rest fatigue (such as insomnia), cognitive fatigue (such as concentration problems)	Severity	“Past one month”	18	5; 5–7 version has 3 response options
Revised Children’s Manifest Anxiety Scale (RCMNS)	6–19	Physiological anxiety- (such as nausea, fatigue, insomnia, dyspnea), worry and oversensitivity- (such as nervousness, fear, irritability, loneliness), concentration anxiety- (such as restlessness)	Presence	“What I think and feel”	37	2
State-Trait Anxiety Inventory for Children (STAIC State)	9–12	Apprehension, tension, worry	Intensity	“Feelings now, at this moment”	20	4
State-Trait Anxiety Inventory for Children (STAIC Trait)	5–13	Anxiety	Frequency	“How you generally/ usually feel”	20	4
Symptom Distress Scale (SDS)	10–17	Nausea, anorexia, insomnia, pain, fatigue, bowel discomfort, concentration, dyspnea, cough	Severity	“Lately”	13	5
Symptom Screening in Pediatrics Tool (SSPedi)	8–18	Sadness, worry, anger, fatigue, mouth sores, pain, headache, neuropathy, anorexia, vomiting, nausea, diarrhea, constipation, concentration, memory, dysgeusia	Interference	“Yesterday or Today”	15	5
10-cm Self-Report Visual Analog Scale (VAS) Nausea	9–20	Nausea	Intensity	“Right now”	1	Left endpoint- No nausea

Instrument name (alphabetical order)	Age range (years)	Symptoms (and HRQOL) domains assessed	Symptom attributes (severity, frequency, etc.)	Reference period	Number of questions	Number of response options
10-cm Self-Report Visual Analog Scale (VAS) Anxiety	9–18	Nausea Anxiety	Intensity	“Right now”	2	Right endpoint- Nausea as bad as it could be Left endpoint- No Nausea/ Anxiety Right endpoint- Nausea/ Anxiety as bad as it could be

The seven validated PROMIS domains are considered as seven separate instruments

Table 2

Studies that evaluated psychometric properties of pediatric self-report symptom measures in cancer research

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{1-r})	Measure Validity: content, construct
Adolescent Sleep Wake Scale (ASWS) [33]	Acute lymphocytic leukemia (41%), acute myelogenous leukemia (4%), Hodgkin lymphoma (8%), non-Hodgkin lymphoma (8%), bone tumor (15%), brain tumor (12%), other (12%)	14.2 (10–19) years	$N = 51$, 43% female	Full ASWS $\alpha = 0.83$. Subscales α ranges (0.66–0.78)	Known groups validity: Adolescents on chemotherapy (CTX) reported significantly lower ASWS scores than healthy adolescents ($p < 0.01$). Adolescents on CTX reported higher ASWS scores than those with chronic pain ($p < 0.01$)
Child Health Questionnaire (CHQ) [34]	Acute lymphocytic leukemia (44.3%), Hodgkin's lymphoma (11.4%), non-Hodgkin's lymphoma (4.3%), Wilms tumor (5.7%), Acute myeloid leukemia (5.7%), medulloblastoma (2.9%), Ewing's tumor (4.3%), rhabdomyosarcoma (2.9%), other (18.5%), 66% off treatment, treatment: chemo, radiation (when necessary), bone marrow transplant (leukemia)	13.6 (10–18) years	$N = 70$, 47% female	Not assessed.	Known groups validity: Comparison between on treatment and off treatment for CHQ, the subscales of General Health Perceptions, Role/Social-Behavior, Self-Esteem, and Mental Health were significant ($p < 0.05$). Concurrent validity: There was significant correlation between parent report and child report with time since diagnosis for Physical Functioning, Role/Social-Physical, and General Health Perceptions
Child Health Questionnaire (CHQ) [22]	On active tx: 31% ALL, 8.5% other leukemia, 14.1% HD/NHL, 29.6% solid tumor, 16.9% brain tumor	11.7 (7–18) years	$N = 71$, % female not reported	Not assessed	Known groups validity: On-treatment group reported worst HRQOL than off-treatment and control group (best HRQOL) on 7 of the 10 subscales. The on- and off-treatment groups did not differ from each other in the role/social limitations-physical scale and the general health perceptions subscales Concurrent validity Moderate significant ($p = 0.05$) correlations between the child and parent reports on 10 HRQL scales for active tx group (range 0.344–0.592)
Childhood Cancer Stressors Inventory (CCSI) [35]	64% leukemia/lymphoma, 36% solid tumor	Mean not reported (7–13 years)	$N = 75$, 45% female	CCSI $\alpha = 0.82$; Intensity subscale $\alpha = 0.83$	Content validity: 5 pediatric oncology nurses reviewed the instrument and rated the items relevant, and the wording understandable. Children age 7–13 evaluated readability of the instrument Concurrent validity: The number of school days missed correlated with the frequency of perceived stressors ($r = 0.34$, $p < 0.01$) and the intensity of stressors ($r = 0.23$, $p = 0.03$)
Children's Depression (CDI) [36]	Children on active treatment who did not have CNS and were not enrolled in full-time	11.5 (8–15) years	$N = 70$, % female not reported	Not assessed	Known groups validity: No significant differences found between cancer and non-cancer group in CDI score (7.42 in both with $p = 0.99$)

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Children's International Mucositis Evaluation Scale (ChIMES) [37]	special education, Leukemia, lymphomas and solid tumors $n = 36$ leukemia/lymphoma, $n = 38$ solid tumor, $n = 7$ brain tumor, $n = 6$ other	12.2 (8–18) years	$N = 87$, 37% female	Test-retest reliability of the total score is $r_{t-r} = 0.854$ ($p < 0.01$) and percentage score $r_{t-r} = 0.852$ ($p < 0.01$) for all children age 12–18 years old. Total score for all children 8–12 $r_{t-r} = 0.902$ ($p < 0.01$) and percentage score $r_{t-r} = 0.902$ ($p < 0.01$)	Convergent validity: ChIMES correlated with WHO Mucositis, VAS Mucositis, CTCAE grading of Mucositis, and Oral Mucositis Daily Questionnaire (OMDQ) items. For ages 12–18 correlation for ChIMES total score ranged from 0.585 to 0.904 ($p < 0.01$ for all), percentage score $r = 0.587$ to 0.905 ($p < 0.01$ for all). For ages 8 to <12 ChIMES total score ranged from 0.551 to 0.928 ($p < 0.01$ for all), percentage score $r = 0.549$ to 0.926 ($p < 0.01$ for all) Responsiveness: see Table 3
Children's International Mucositis Evaluation Scale (ChIMES) [38]	All children were receiving or had received chemotherapy for cancer, including HSCT	12.5 (8–18) years	$N = 82$, % female not reported	Not assessed	Content validity: Cognitive interviews with children and parents. ChIMES mostly rated as “very easy” or “easy” to understand. Most rated the overall instrument as “good” or “okay.”
Children's International Mucositis Evaluation Scale (ChIMES) [39]	First phase: Leukemia/lymphoma (80%), Solid tumor (10%), Brain tumor (10%) Second phase: Leukemia/lymphoma (47.5%), Solid tumor (45%), Brain tumor (7.5%)	Phase 1: median: 15.3 (9.2–17.7) years Phase 2: median: 12.4 (8.0–17.8) years	Phase 1 $N = 10$, 40% female Phase 2 $N = 40$, 70% female	Not assessed	Content validity: Cognitive interviews for first phase (instructions) and second phase (use, understandability, and suitability). Overall found that the instrument was easy to use, easy to understand and suitable to assess mucositis
Constipation Assessment Scale (CAS) [40]—modified version for children	Ewing's sarcoma (38.1%) Rhabdomyosarcoma (23.8%) Osteosarcoma (14.3%) Acute lymphocytic leukemia (4.8%) Ependymoma (4.8%) Lymphoma/leukemia (4.8%) Malignant peripheral nerve sheath tumor (4.8%) Stomach (4.8%)	15.7 (9–21) years	$N = 21$, 52% female	$r_{t-r} = 0.93$, $p < .01$ CAS α range (0.60–0.79) during weeks 1 & week 2. CAS α range (0.28–0.82) in weeks 3–5	Known groups validity: Significant difference in mean CAS scores ($t = 4.4$, $p < .01$) for subjects designated as not constipated (2.83; SD, ± 1.9) versus constipated (7.80; SD, ± 2.6) Concurrent validity: CAS was associated with constipation noted in the medical record and the physical examination
Fatigue Scale-Children (FS-C); Fatigue Scale-Adolescents (FS-A) [41]	Solid tumors (86%) and AML (14%).	12.5 (7–18) years	$N = 29$, 59% female	FS-C α range (0.64–0.72), FS-A α range (0.76–0.96)	Known groups validity: FS-C and FS-A scores did not differ significantly between the experimental and standard care arms Responsiveness: see Table 3
Fatigue Scale-Adolescents (FS-A) [42]	Leukemia, Lymphoma, ALL (60.9%)	15.3 (13–18) years	Focus groups: $N = 15$, 67% female Quantitative study: $N = 64$, 44% female	α range (0.67–0.95) across 4 studies	Content validity: Individual and focus group interviews to learn about the fatigue experience, and expert panel review with health care professionals and adolescents with cancer Known groups validity: Anemic patients had significantly higher fatigue than non-anemic patients, but article only cites parent report here, ($p = 0.04$)

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Fatigue Scale-Children (FS-C) [43]	Leukemia (67.9%), Lymphoma (7.2%), Brain tumor (4.1%) Solid tumor (19.9%)	9.2 (6.6–13) years	Study 1 $N = 53$, 42% female. Study 2 $N = 150$, 44% female. Study 3 $N = 18$, 78% female	10-item FS-C $\alpha = 0.76$, 14-item FS-C $\alpha = 0.81$	Concurrent validity: Adolescent scores were significantly correlated with parent scores on 2 out of 4 studies and then when aggregating study data Responsiveness: see Table 3 Structural validity Factor analysis showed good fit: CFI = 0.94 (>0.90 threshold), RMSEA = 0.028 (<0.05 threshold). Concurrent validity: FS-C correlated with FS-P (parent) responses ($r = 0.44$, $p < .01$)
Revised 13-item Fatigue Scale-Adolescents (FS-A) [44]	37.68% ALL, 2.89% AML, 37.68% HL/lymphoma, 18.1% solid tumor, 3.62% germ cell tumor	15.5 (12.7–18.8) years	$N = 138$, 44% female	13-item $\alpha = 0.87$, 14-item $\alpha = 0.86$	Structural validity: Confirmatory factor analysis yielded reasonable fit of 4 factors; goodness-of-fit index 0.8551; Root Mean Square residual (RMSR) = 0.08 Concurrent validity: Spearman correlation in the subset of 75 patient and parent dyads was 0.347 ($p < .01$)
Memorial Symptom Assessment Scale (MSAS) (10–18) [5]	Leukemia, lymphoma, other solid tumor, CNS tumor, rare malignancies of childhood	14 (10–18.2) years	$N = 159$, 42% female	α ranged from 0.83–0.87 Test-retest: 26 of 30 symptoms had significant correlation	Content validity: Experts in developmental pediatrics and pain management modified the adult version of the instrument. Then instrument was run through the readability tests in MS Word (SMOG, FRY). Finally, the modified instrument was pilot-tested in 10 patients around 10 years old Convergent validity: MSAS correlated with Memorial Pain Assessment Card-pediatric (MPAC-ped), and Investigator-created, face-valid VAS to assess domains of nausea, global physical, global psychological. There was strong agreement between children and parents for 24 of 30 symptoms Known groups validity: Higher symptom number and distress among inpatients than outpatients. Significant differences between recent treatment (2 weeks–1 month) vs. off-treatment (>4 months) groups for MSAS total and all sub scores except the psychological subscale
Memorial Symptom Assessment Scale (MSAS) (7–12) [19]	Acute lymphoblastic leukemia, CNS tumor, Non-Hodgkin's lymphoma, Rhabdomyosarcoma, Wilms' tumor, Hodgkin's lymphoma, Acute myelogenous leukemia, Neuroblastoma, Ewing's sarcoma, Hepatoblastoma, Osteosarcoma	9.6 (7–12) years	$N = 149$, 47% female	$r_{t-r} = 0.67$	Content validity: Creation of MSAS (7–12) based on data from the MSAS (10–18) study. Instrument ran through the readability tests in MS Word (SMOG, FRY) Convergent validity: Investigator-created, face-valid VAS to assess domains of pain, nausea, sadness, and body showed significant correlations ranging from 0.70–0.76 ($p < 0.01$). Higher MSAS symptom number and distress among inpatients than outpatients. Fair to moderate agreement between children and parents: nausea ($k = 0.46$), pain ($k = 0.46$), and lethargy ($k = 0.42$).

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Minneapolis-Manchester Quality of Life-Adolescent Form (MMQL-AF) * [45]	ALL, AML, Hodgkin's, Non-Hodgkin's, Brain tumors, Wilms' tumor, Neuroblastoma, Rhabdomyosarcoma, Osteosarcoma, Ewing's sarcoma, Hepatoblastoma, Other cancer	16.7 (13–20.9) years	$N = 110$, 45% female	Overall $r_{t-r} = 0.71$. Subscale r_{t-r} ranged from 0.60–0.90. Overall $\alpha = 0.92$, subscale α ranges (0.67–0.89)	Fair agreement for anorexia ($k = 0.35$), sadness ($k = 0.33$), and insomnia ($k = 0.20$). Poor agreement for itch ($k = 0.11$) and worry ($k = 0.16$) Content validity Focus group of 20 individuals including, patients, parents, nurses, doctors. Then 20 individual "open interviews" with survivors conducted to identify domains. Two rounds of cognitive interviews each with 10 adolescent survivors of cancer Known groups validity: Significant differences between 3 groups (children with cancer, both on and off treatment, and those without cancer): Physical, Cognitive, Psychological factors. But MMQL-AF could not discriminate between 3 groups for body image, intimate relationships, outlook on life domains
Minneapolis-Manchester Quality of Life-Adolescent Form (MMQL-AF) * [46]	Leukemia (44%), Lymphoma (19%), Brain tumors (4%), other solid tumors (33%).	15.4 (5.3–20.0) years	$N = 136$, 40% female	Not assessed	Known groups validity: Scores for those on treatment (3.77), survivor (3.96) and healthy controls (4.05) were significantly different ($p < .01$). On-therapy patients were at increased risk for poor QOL (defined at a score below the 25th percentile); OR 3.3, $p < 0.01$. On-therapy patients were at increased risk for poor physical functioning OR 11.8, $p < .01$
Minneapolis-Manchester Quality of Life-Youth Form (MMQL-YF) * [47]	Leukemia, Lymphoma, Brain Tumors, Other solid tumors (on-treatment sample must be patients actively undergoing treatment for cancer for at least 2 months)	10.2 (8–12) years	$N = 72$, 42% female	r_{t-r} ranged from 0.56–0.79 ($p < 0.05$). Overall $\alpha = 0.85$, subscales α range (0.72–0.80)	Content validity: Focus group of 10 survivors to identify concerns. Then "open interviews" were conducted with 20 survivors to identify questions to include in the questionnaire. Two rounds of cognitive interviews each with 20 survivors Known groups validity: MMQL-YF was able to distinguish between 3 groups, but On-treatment patients had the best scores of all three groups (when they were expected to have the worst) for Physical Symptoms and Psychological Functioning domains Convergent validity: MMQL-YF had some moderate, significant correlations when compared with Child Health Questionnaire-87 (compared across multiple sub-domains)
Minneapolis-Manchester Quality of Life-Youth Form (MMQL-YF) * [48]	Children 8–12 actively undergoing treatment for cancer for at least 2 months. Leukemia (64%), lymphoma (6%), brain tumors (8%), other solid tumors (22%)	9.1 (1.9–12.5) y	$N = 76$, 42% female	Not assessed	Known groups validity: Children on treatment had lower MMQL-YF scores than healthy controls ($p = 0.01$). Children on therapy also had lower physical functioning ($p = 0.01$) and outlook on life ($p < 0.01$)
Oral Mucositis Daily Questionnaire (OMDQ) [49]	Children undergoing induction or consolidation chemotherapy for AML.	Median: 16.7 (12–18) years	$N = 15$, 47% female	All questions, but diarrhea exhibited very good correlation ($r_{t-r} > .$	Convergent validity: OMDQ had strong to moderate significant correlations with WHO mucositis scale, a mucositis VAS, and the oral component of the

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Oral Mucositis Daily Questionnaire (OMDQ) [50]	relapsed ALL, or advanced lymphoma or receiving myeloablative SCT (87% of children in this study diagnosed with Leukemia/Lymphoma)	15 (13–17) years	$N = 6$, 17% female	80, Diarrhea question $r_{t-r} = 0.73$	Functional Assessment of Cancer Therapy Esophageal Cancer Subscale (FACT-ECS) on questions about pain (0.90, 0.81, -0.62 – 0.77), swallow (0.65, 0.67, -0.60 – 0.71), drink (0.73, 0.57, -0.65 – 0.77) and eat (0.83, 0.470.82–0.85) Sleep and talk questions had fair correlations with WHO, VAS mucositis and FACT-ECS (between 0.35–0.59; -0.35 – 0.59). The diarrhea question had poor correlations ranging from 0.04–0.16
Pain Squad App [51]	Diagnosis not discussed, 100% received chemotherapy		Not assessed		Content validity: Cognitive interviews: 5 out of 6 children rated the first and third questions very easy to understand, 2 out of 6 rated the second question very easy to understand. Children found this instrument easy to use overall
	Study 1: ALL (45%), AML (9%), Non-Hodgkin Lymphoma (9%), Sarcomas (16%), CNS tumors (8%), Other (13%)	Study 1: 13.1 (8–18) years	Study 1: $N = 92$, 49% female Study 2: $N = 14$, 50% female	$\alpha = 0.96$	Discriminant Validity: PedsQL Generic Module had weak to moderate, significant correlations ($p < 0.01$) with Pain Squad app weekly pain indices' scores for intensity ($r = -0.3$), unpleasantness ($r = -0.38$), interference ($r = -0.46$). But no significant correlations with PedsQL Cancer Module
Pediatric Advanced Care Quality of Life Scale (PAC-QoL) [52]	Doesn't specify (only says "malignancy")	12.6 (9–17) years	$N = 28$, 64% female	Not assessed	Responsiveness: see Table 3
Pediatric Cancer Quality of Life Inventory (PCQL-32) [53]	Cancer type not specified. Included on and off treatment. Excluded: Non-English speaking, comorbid disease, and major development disorders	11.8 (8–18) years	$N = 281$, % female not reported	Core form $\alpha = 0.83$, Subscales: Pain $\alpha = 0.83$, Nausea $\alpha = 0.71$.	Content validity: Cognitive interviews demonstrated that the scale was easy to understand and self-report
Pediatric Cancer Quality of Life Inventory (PCQL-32) [54]	Acute lymphocytic leukemia, acute myelogenous leukemia, other leukemias, Hodgkin's lymphoma, non-Hodgkin's lymphoma, brain tumor, Wilms tumor, neuroblastoma, rhabdomyosarcoma, osteosarcoma, Ewing's sarcoma, retinoblastoma, hepatoma	11.8 (8–18) years	$N = 291$, 35% female	Overall $\alpha = 0.91$, Subscales: Symptoms/Problems, $\alpha = 0.83$, Physical Functioning $\alpha = 0.78$, Psychological $\alpha = 0.76$, Social $\alpha = 0.69$, Cognitive $\alpha = 0.81$.	Known groups validity: Mean scores for on vs off treatment. Core: 50.8 vs. 49.4 ($p = 0.03$). Pain: 51.2 vs. 49.1 ($p = 0.02$). Nausea: 52.3 vs. 48.2 ($p < 0.01$) Concurrent validity: Significant correlations for the core and nausea scales with number of days missed from school
Pediatric-Functional Assessment of Anorexia and Cachexia Therapy (Peds-FAACT) [55]	Leukemia (43%), brain/spinal cord tumor (26%).	11.3 (7–17) years	$N = 96$, 50% female	For adolescents (12–17): $\alpha = 0.80$ For children (7–11): $\alpha = 0.47$ 10-item version $\alpha = 0.83$ for children 10 years and older	Known groups validity: On-treatment had significantly worse health/symptom scores than off-treatment groups for instrument as a whole, as well as Disease/Treatment Symptoms and Physical Functioning, and Psychological Functioning subscales Convergent Validity: Moderate correlations ($p < 0.05$) between instrument with Children's Depression Inventory-32 (CDI), State-Trait Anxiety Inventory-32 (STAI-C), Self-Perception Profile for Children (SPPC) and Adolescents (SPPA)
					Convergent validity: 6-item & 10-item peds-FAACT correlated ($r = .51$, $r = .46$) with the PedsQL Cancer module Nausea subscale. Known groups validity: 6-item peds-FAACT statistically differentiated Patients from 2 functional performance groups (Lansky and Kamofsky) ($p < 0.05$). But, no significant difference found between patients with

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Pediatric Nausea Assessment Tool (PeNAT) [56]	Diagnosis not described, but children were divided into two groups-2 groups receiving chemotherapy, and 2 not receiving chemotherapy (Exclusion criteria: children who were developmentally delayed, non-English speaking)	10.3 (4–17.8) years	$N=177$, 39% female	Test-retest (1 h in-between assessment): $r_{t-r}=0.649$, $p<0.01$	acceptable BMI and patients who were either underweight or overweight ($p=0.87$) Structural validity: Scale unidimensionality confirmed by factor analysis Content validity: PeNAT reviewed by clinicians, parents. Feasibility testing in 15 children receiving chemo followed by interviews (for adolescents 16) on understanding instructions and ease of administration (but not understanding of construct) Convergent validity: First PeNAT score and number of emetic episodes (Spearman $\rho=0.322$, $p=0.12$); First PeNAT score and dietary intake (Spearman $\rho=-0.217$, $p=0.096$) Known groups validity: Children undergoing HSCT higher median scores than children with cancer not receiving chemo ($p<.01$). HSCT group significantly higher when compared with general pediatrics ($p=0.04$). Concurrent validity: Moderate correlation observed between children's PeNAT scores and the parents' assessment of their child's nausea (Spearman's $\rho=0.442$, $p<0.0001$)
Pediatric Oncology Quality of Life Scale (POQOLS), Children's Depression Inventory (CDI), Revised Children's Manifest Anxiety Scale (RCMAS) [57]	Leukemia (39%), brain tumor (6%), Hodgkin's disease (28%) and other types of cancer (27%).	13.7 (8–19) years	$N=19$, % female not reported	POQOLS $\alpha=0.83$, CDI $\alpha=0.81$, RCMAS $\alpha=0.30$, symptom checklist $\alpha=0.86$	Concurrent validity: Children's scores on POQOLS were positively and significantly correlated with parents' scores
Pediatric Patient-Reported Outcomes Version of the Common Terminology Criteria for Adverse Events (Pediatric PRO-CTCAE) [58]	Round 1: Leukemia (49%), Lymphoma (18%), Sarcoma (18%), Other solid tumor (13%), Brain (2%) Round 2: Leukemia (50%), Lymphoma (19%), Sarcoma (28%), Other solid tumor (3%)	Round 1: Mean not reported (7–12 years) Round 2: Mean not reported (7–15 years)	Round 1: $N=45$, 58% female Round 2: $N=36$, 42% female	Not assessed	Content validity: Two rounds of cognitive interviews with children/adolescents on active treatment and their caregivers. Assessed symptom AE terms, question structure, response options, and recall period. Overall instrument performed well and found to be easy to complete
Pediatric Quality of Life Inventory-4.0 Generic Score (PedsQL) Pediatric Quality of Life Inventory-Cancer Module State-Trait Anxiety Inventory for Children (STAIC) Faces Pain Scale-Revised (FPS-R) [59]	51% leukemia, 17% CNS tumor, 17% sarcoma, 10% lymphoma, 5% other (germ cell, Wilms)	8.6 (4–17) years	$N=45$; 38% female	Not assessed	Known groups validity: Children who received medication (compared with those not on medication) reported lower QOL as measured by the PedsQL Cancer Module ($p=0.02$) and lower general HRQOL as measured by the PedsQL ($p=0.01$) and greater pain severity as measured by FPS-R ($p<0.01$)
Pediatric Quality of Life Inventory-4.0 Generic Score (PedsQL) [60]	Solid tumors, Hodgkin's disease, ALL, and NHL	11.6 (5–18) years	$N=222$, 46% female	Physical $\alpha=0.90$, Emotional $\alpha=0.84$, Social $\alpha=0.71$, School $\alpha=0.82$ r_{t-r} ranged 0.53–0.67	Concurrent validity: Agreement between child and parents ranged from 0.61 (social function) to 0.86 (physical function)

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range); years	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL) [61]	Intracranial tumors. Children alive at one year after diagnosis were included in analyses	Median 9.4 (1.8–16.6) years	$N = 26$, % female not reported	Not assessed	Known groups validity: at T1 there was significant difference between patients and controls for all summary scores ($p = 0.03$), but not emotional/social domains ($p = 0.12$). At T6, total score and summary scores were significantly different between patients and controls ($p = 0.03$). At T12, only the physical summary score was significantly different between the two groups ($p = 0.02$). Concurrent validity: at T1, child/parent agreement was 0.65 to 0.79. At T6, parent/child agreement ranged from 0.53 to 0.86. At T12, parent/child agreement ranged from 0.66 to 0.87 Responsiveness: see Table 3
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL); Pediatric Quality of Life Inventory-Cancer Module [62]	Solid tumor, Hodgkin's disease, Acute lymphocytic leukemia, Non-Hodgkin's lymphoma	11.6 (5–18) years	$N = 222$, 45% female	Not assessed	Known groups validity: Mean PedsQL scores between EPO and placebo were not significant ($p = 0.76$) Concurrent validity: For the general score, child/parents agreement ranged from 0.59 to 0.77 ($p < 0.01$) and for the cancer module it ranged from 0.34 to 0.59 ($p < 0.01$). Correlation between changes in hemoglobin and PedsQL-GCS ($r = 0.15$, $p = 0.03$) Responsiveness: see Table 3
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL); Acute version, Pediatric Quality of Life Inventory-Multidimensional Fatigue Scale: Acute version, Pediatric Quality of Life Inventory-Cancer Module: Acute version [63]	Children with ALL (50%), Brain tumor (7%), NHL (6%), HL (3%), Wilm's tumor (6%), other cancers (28%)	10.9 (8–18) years	$N = 222$, 44% female	Total PedsQL $\alpha = 0.88$. Subscales: Physical health $\alpha = 0.81$, Psychosocial health $\alpha = 0.83$, Emotional health $\alpha = 0.73$, Social functioning $\alpha = 0.70$ PedsQL Multidimensional Fatigue Total fatigue $\alpha = 0.89$. Subscales: General fatigue $\alpha = 0.77$, Sleep/rest fatigue $\alpha = 0.78$, Cognitive fatigue $\alpha = 0.83$ PedsQL 3.0 Cancer Module overall: $\alpha = 0.72$, Pain and hurt: $\alpha = 0.70$, Nausea $\alpha = 0.79$, Procedural anxiety $\alpha = 0.82$, Treatment anxiety $\alpha = 0.79$, Worry $\alpha = 0.74$, Cognitive problems $\alpha = 0.76$, Perceived physical appearance $\alpha = 0.49$, Communication $\alpha = 0.66$	Known groups validity: For PedsQL Oncology Sample reported worse HROQL than the Healthy Sample ($p < 0.01$). For PedsQL Multidimensional Fatigue Scale, the Oncology Sample reported worse fatigue than the Healthy Sample ($p < 0.05$). For PedsQL 3.0 Cancer Module, children on treatment reported worse Nausea, Treatment Anxiety, Worry compared to children off treatment (>12 months) Convergent validity: Examined through an analysis of the inter-correlations among the PedsQL 4.0 Generic Core Total Scale score with the PedsQL Multidimensional Fatigue Total Score and the PedsQL Cancer Module Scales show medium effect size

Instrument being validated	Cancer Type (treatment type, inclusion/exclusion criteria)	Child Age (mean, range);	Sample Size; % female	Measure Reliability: internal consistency (α), test-retest (r_{t-r})	Measure Validity: content, construct
Pediatric Quality of Life Inventory-Cancer Module; Pediatric Quality of Life Inventory-Multidimensional Fatigue Scale [64]	Not reported	15.6 (13–19) years	$N = 102$, 49% female	Not assessed	Discriminant validity: Correlation between pain and fatigue is 0.62 ($p < 0.01$), pain and nausea is 0.33 ($p < 0.01$), fatigue and nausea is 0.47 ($p < 0.01$)
PROMIS Pediatric Measures: Physical Functioning—Mobility, Physical Functioning—Upper Extremity, Pain Interference, Fatigue, Depressive Symptoms Anxiety, Peer Relationships, and Anger [31]	Currently receiving curative cancer treatment (60% were diagnosed with leukemia or lymphoma)	12.5 (8–17) years	$N = 200$, 45%	Not assessed	Known group validity: Active treatment vs. survivorship compared across 8 domains (Depression, Anxiety, Anger, Peer Relationships, Pain interference, Fatigue, Upper extremity, Mobility); Children in active tx group had statistically significantly lower (worse) mean scores compared to survivors in all categories except anger
State-Trait Anxiety Inventory for Children (STAIC) [65]	Cancer group: ALL (25.4%), other Leukemia (6.9%), HD/NHL (28.5%), Solid Tumor (30.0%), Brain tumor (9.2%)	13.1 (7–18) years	$N = 130$, 41% female	Not assessed	Known groups validity Cancer group had the lowest anxiety score (34.3) compared to chronically ill (35.1) and control (37.0), $F(2,597) = 8.0$, $p < .01$
Symptom Distress Scale (SDS), State-Trait Anxiety Inventory for Children (STAIC) [66]	Leukemia (64%) and Hodgkin's Disease (36%).	Mean not reported (10–17) years	$N = 11$, 45% female	SDS $\alpha = 0.85$, STAIC-1 $\alpha = 0.86$	Responsiveness: see Table 3
Symptom Screening in Pediatrics Tool (SSPedi) [67]	Included children under active treatment. Excluded those who were severely ill, had cognitive disabilities or visual impairment	Mean not reported (8–18) years	$N = 30$, 33% female	Not assessed	Content validity Cognitive interviews found the SSPedi was easy to complete and understandable among children undergoing active treatment for cancer
10-cm Self-Report Visual Analog Scale (VAS) Nausea 10-cm Self-Report Visual Analog Scale (VAS) Anxiety [68]	Leukemia (40%), lymphoma (27%), brain tumor (7%), sarcoma (27%).	12.7 (9–18) years;	$N = 15$, 27% female	Not assessed	Known groups validity: To assess nausea VAS: compared children on antiemetics reported more nausea than children with no antiemetics. Assessing nausea and anxiety VAS: compared to clinical anchors (pulse rate, systolic, and diastolic blood pressure, antiemetic v. no antiemetic) comparison with significance: systolic blood pressure increased significantly ($m = 104.80$) from previous intervention level ($m = 98.27$) Responsiveness: see Table 3

* Intended use with cancer survivors

VAS Visual Analog Scale

Longitudinal studies providing evidence for the responsiveness of the pediatric symptom measures

Table 3

Instrument validated	Cancer type (treatment type, inclusion/exclusion criteria)	Child age (mean, range); years	Sample size; % female	Responsiveness	Number and timing of assessments
Children's Depression Inventory (CDI) [69]	Children with Leukemia (41%) or solid tumors (59%). Exclusion criteria included patients with progressive disease, individuals diagnosed <2 weeks ago, and those in the intensive care unit	Median: 12.9 (8–16.8) years	<i>N</i> = 99, 39% female	At T1: median CDI for children was 6.0 and 3.5 at T2 ($p < 0.01$). The 23-item version of the CDI had a median of 4.0 at time 1 and 3.5 at time 2 ($p < 0.01$). Both the CDI and CDI-23 should statistically significant decline from T1 to T2	Two time points T1: At study enrollment and T2: 4–6 weeks after
Children's International Mucositis Evaluation Scale (ChIMES) [37]	<i>n</i> = 36 Leukemia/Lymphoma, <i>n</i> = 38 Solid tumor, <i>n</i> = 7 Brain tumor, <i>n</i> = 6 Other	Median: 14.7 (8–18) years	<i>N</i> = 87, 37% female	Among children who reported mucositis on the 14th day, Children ages 12–18 years: at baseline reported 0.9 and at 14 days reported 10.6 ($p < 0.01$). Children ages 8–12 years: at baseline reported 0.9 and at 14 days reported 12.6 ($p = 0.06$). Overall, the CHIMES showed significant increases in mucositis between baseline and day 14	14 time points T1: baseline is 2–5 days and then again daily for days 7–17
Fatigue Scale-Adolescent (FS-A) [42]	ALL (60.9%), AML (4.7%), HD/lymphoma (9.3%), Solid tumor (25%)	15.3 (13–18) years	<i>N</i> = 64, 44% female	Combined scores across 4 studies. FS-A scores increased significantly between T1 and T2 ($p = 0.01$)	Two time points. MFCC study, T1: first day of re-induction, T2: end of re-induction. SLEEP study, T1: 2nd day of the first 5-day period, T2: 2nd day of the second 5-day period. SLEEP2 study, T1: date of admission to hospital, T2: 3rd day of hospitalization. CLUSTERS study, T1: 1st day of chemotherapy, T2: week after final dose of chemotherapy.
Fatigue Scale-Adolescents (FS-A), Fatigue Scale-Children (FS-C) [41]	Solid tumors (86%) and AML (14%). Eligibility: 7–18 years, inpatient, able to understand English, and able to give assent. Children with CNS tumors, being treated for recurrent disease, scheduled for rehab due to amputated limb, and those experiencing uncontrolled pain (>3 on a 5-point scale) were ineligible for the study.	12.5 (7–18) years	<i>N</i> = 29, 59% female	FS-C and FS-A scores did not vary significantly over time ($F = 0.50$, $p = 0.61$)	Four time points T1: Date of hospital admission and fatigue was collected the following three days (T2, T3, and T4)
Fatigue Scale-Adolescents (FS-A), Fatigue Scale-Children (FS-C) [70]	ALL (low risk, standard risk (63%); 50 weeks after diagnosis (on continuation therapy)	9.2 (5–18) years	<i>N</i> = 100, 38% female	Children ages 7–12 years completing the FS-C reported significantly more fatigue during the on-dex period than during the off-dex period ($p < 0.01$). Findings were similar for adolescents ages 13–18 completing the FS-A ($p < 0.01$)	Four time points (2 consecutive 5-day periods). T1 (day 2) and T2 (day 5): Patients not taking dexamehasone. T3 (day 7) and T4 (day 10): Patients taking dexamehasone
Fatigue Scale-Adolescents (FS-A),	Solid tumor (86%), acute myeloid leukemia (14%)	12.5 (7.4–18.2) years	<i>N</i> = 29, 59% female	Mean FS-C was 16.59 at T1, 22.69 at T2, 23.22 at T3, and 23.33 at T4. Mean FS-A was	Four time points

Instrument validated	Cancer type (treatment type, inclusion/exclusion criteria)	Child age (mean, range);	Sample size; % female	Responsiveness	Number and timing of assessments
Fatigue Scale-Children (FS-C) [71]				25 at T1, 31.67 at T2, 33.78 at T3, and 37 at T4. Authors do not present whether this was a statistically significant change over time	T1: Date of admission, fatigue was collected for the following 3 days (T2, T3, and T4)
Fatigue Scale-Adolescents (FS-A), Fatigue Scale-Children (FS-C) [72]	ALL (6 children, 3 adolescents); AML (1 adolescent); lymphoma (5 children, 7 adolescents); solid tumor (5 children, 3 adolescents). Newly diagnosed within 3 cycles of chemo	No mean (6–18 years)	N = 30, 30% female	FS-C at T1: 18 and at T2: 10.5 ($p = 0.05$) and FS-A at T1: 23.5 and at T3: 20.5 ($p = 0.15$). Using the FS-C, fatigue significantly decreased from cycle 1 to 3, but there was no significant change using the FS-A	Two time points T1: first cycle of chemotherapy between days 15–29 of the cycle, T2: third cycle of chemotherapy between days 15–29 of the cycle
Fatigue Scale-Adolescents (FS-A), Fatigue Scale-Children (FS-C) [73]	Solid tumor (59%), Lymphoma (41%)	No mean (7–18 years)	N = 45, 35% female	CFS showed significant decreases in overall fatigue among children 7–12 years ($p = 0.04$), but not among adolescents 13–18 years ($p = 0.94$). The CFS also showed improvements over time in the energy ($p = 0.01$) and function domains ($p = 0.04$), indicating that fatigue declines	Four time points, T1: During the 2nd cycle of chemotherapy. Measured again every 2 chemotherapy cycles for up to 4 cycles (8 cycles total)
Fatigue Scale-Adolescents (CFS-A), Fatigue Scale-Children (CFS-C) [74]	Acute Lymphoblastic Leukemia	8.7 (6–18) years	N = 16, 69% female	Fatigue levels did not change significantly over the 2 week study period ($p = 0.42$)	Three assessment points over 2 weeks
Fatigue Scale-Adolescents (CFS-A), State-Trait Anxiety Inventory for Children (STAIC) [75]	Leukemia, Bone tumor, Brain tumor, Lymphoma	15.7 (12–21) years	N = 34, 29% female	Anxiety levels did not change significantly over time in the control group ($p = 0.4701$) or the intervention group ($p = 0.8291$) Fatigue levels did not change significantly over time in the control group ($p = 0.5337$) or the intervention group ($p = 0.9949$)	Fatigue and anxiety collected daily during hospitalization. Pre-intervention (days 1–2) and post-intervention (days 3–4+)
Pain Squad App [51]	Osteosarcoma (57%), Ewing sarcoma (29%), Other (14%)	Study 2: 14.8 (9–18) years	Study 2: N = 14, 50% female	Detect change in pain scores after surgery using F scores: The increase in pain intensity from week 1 to week 2 was not significant. However, decrease in pain intensity from week 2 to week 3 was significant ($p = 0.03$). Pain interference significantly increased from week 1 to week 2 ($p = 0.005$), but pain interference did not significantly decrease from week 2 to week 3. Median effect sizes were large for pain unpleasantness (Cohen's effect size = .96, 95% CI 0.14 to 1.79) and pain interference from week 1 to 2 (Cohen's effect size = 1.08, 95% CI 0.16 to 2.00)	Daily assessments for 3 weeks: Participants started completing daily assessments 1 week before scheduled surgery. Then continued to complete daily assessments the next two weeks after surgery
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL) [61]	Primary intracranial tumors. Children alive at one year after diagnosis were included in analyses	Median: 9.4 (1.8–16.6) years	N = 26, does not report % female	There were statistically significant improvements in HRQOL over time for the total and physical summary scores ($p = 0.01$), but not for the psychosocial summary, emotional, social or school domains ($p = 0.28$). HRQOL improved over time	Three time points. T1: 1-month post-diagnosis, T2: 6-month post-diagnosis, T3: 12-month post-diagnosis

Instrument validated	Cancer type (treatment type, inclusion/exclusion criteria)	Child age (mean, range); years	Sample size; % female	Responsiveness	Number and timing of assessments
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL) [76]	Primary intracranial tumors	Median: 9.1 (1.5–16.4) years	<i>N</i> = 35, 49% female	Mean score at T1: 73.4, mean score at T2: 78.4, and mean score at T3: 81.3. Authors do not report on the statistical significance of these increases over time	Three time points. T1: 1-month post-diagnosis, T2: 6-month post-diagnosis, T3: 12-month post-diagnosis
Pediatric Quality of Life Inventory-4.0 Generic Core (PedsQL) (self- and parent-report versions) [77]	<i>n</i> = 5 Ependymoma, <i>n</i> = 19 Astrocytoma, <i>n</i> = 3 Germ cell, <i>n</i> = 3 Medulloblastoma, <i>n</i> = 5 Other (children must completed the 12 month assessment to be included in the study)	Median: 9.1 (1.5–16.4) years	<i>N</i> = 35, 50% female	PedsQL at T1: 73.4 and at T2 81.3 (no statistically significant difference)	Two time points. T1: 1-month post-diagnosis, T2: 12-month post-diagnosis
Pediatric Quality of Life Inventory-4.0 Generic Score (PedsQL) Pediatric Quality of Life Inventory-Cancer Module [78]	Hodgkin disease (Inclusion: Patients newly diagnosed, had to be enrolled by the second week after starting chemotherapy-all stages could participate)	14.7 (8–18) years	<i>N</i> = 49, 55% female	Responsiveness between T1 and T4 (in subset of participants who reported improvement in health based on global impression of change): Effect sizes were large for all instruments. However, when analyzing the changes between individual time points 2–3 and 3–4, The PedsQL Cancer Module had small to moderate effect sizes. The PedsQL had moderate to large effect sizes. Between T1 and T4: PedsQL Cancer Module: Significant change over time ($r = 5.83$, $p < 0.05$). Cohen's effect size = 1.22; PedsQL 4.0 ($r = 7.38$, $p < 0.05$). Cohen's effect size = 1.69	Four time points. T1: 2 weeks after beginning 1st course of chemo, T2: Day 3 of 2nd course of chemo, T3: 3rd week of radiation, T4: 1 year post-diagnosis. Children who did not receive radiation therapy did not complete T3 questionnaires
Pediatric Quality of Life Inventory-4.0 Generic Score (PedsQL) Pediatric Quality of Life Inventory-Cancer Module [62]	Confirmed diagnosis of solid tumor, Hodgkin's disease, Acute Lymphocytic Leukemia, Non-Hodgkin's lymphoma	11.6 (5–18) years	<i>N</i> = 222, 45% female	Two groups (EPO and placebo) of 111 children in each. Mean PedsQL-GCS at the final visit was significantly greater in the EPO group (compared to mean baseline score) among kids 5–7 years (mean 88.0 vs. 78.1, $p = 0.043$), but no statistically significant differences over time in the 8–12 or 13–18 year old groups or in the placebo group	Two time points. T1: baseline, T2 = final visit
State-Trait Anxiety Inventory for Children (STAIC) [79]	56% Central Nervous System neoplasm, 36% Non-CNS solid tumor, 7.2% Leukemia.	Median: 11.9 (8.1–21.9) years	<i>N</i> = 55, 42% female	Children's ratings of distress decreased significantly pre/post magnetic resonance imaging procedure ($p < 0.01$)	Two time points (pre/post). T1: Immediately pre magnetic resonance imaging procedure T2: immediately after
State-Trait Anxiety Inventory for Children (STAIC) [80]	Not reported	No mean (5–18 years)	<i>N</i> = 20, does not report % female	No significant differences in STAIC anxiety	Two time points (pre/post). T1: pre-intervention and T2: post-intervention
Symptom Distress Scale (SDS), State-Trait Anxiety Inventory for Children (STAIC) [66]	Leukemia (64%) and Hodgkin's Disease (36%). Inclusion: aged 10–17 years, first diagnosis of cancer, within first 4 months of initial chemo regimen, need at least three more chemo cycles to complete their treatment, able to read/write in English, no metastatic disease to brain and no history of motion sickness	No mean (10–17 years)	<i>N</i> = 11, 45% female	Immediately after the chemotherapy treatments, there were significant differences in the SDS ($p = 0.06$), but no differences in the STAIC-1 ($p = 0.11$). When the instruments were used 48 h after the chemotherapy was given, they found that STAIC-1 scores differed over time ($p = 0.07$), but there were no statistically significant differences in the SDS ($p = 0.19$)	Two time points. T1: Immediately after chemotherapy treatments and T2: Within 48 h of chemotherapy being given

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Instrument validated	Cancer type (treatment type, inclusion/exclusion criteria)	Child age (mean, range);	Sample size; % female	Responsiveness	Number and timing of assessments
10-cm Self-Report Visual Analog Scale (VAS) Nausea [68]	Experiment one: Leukemia (27%), Lymphoma (22%), Sarcoma (50%), Teratoma (4%), Brain tumor (4%)	15.4 (9–20) years	$N = 26$, 27% female	hypothesis that the visual distraction would have a lasting effect on helping manage chemotherapy-induced symptoms was not supported In the experiment (stratified by antiemetic use), there was a significant decline in nausea observed ($p = 0.02$)	Two time points T1: Before video game intervention, T2: after video game intervention